1   Characterizing spatiotemporal variation in transmission heterogeneity
2   during the 2022 mpox outbreak in the USA
3

4   Jay Love*[1], Cormac R. LaPrete*[2], Theresa R. Sheets[2], George G. Vega Yon[1], Alun Thomas[1],
5   Matthew H. Samore[1], Lindsay T. Keegan[1], Frederick R. Adler[2,3], Rachel B. Slayton[4], Ian H.
6   Spicknall[5], Damon J.A. Toth[1]
7
8   *Contributed equally
9
[1] Division of Epidemiology, University of Utah, Salt Lake City, Utah
[2] Department of Mathematics, University of Utah, Salt Lake City, Utah
[3] School of Biological Sciences, University of Utah, Salt Lake City, Utah
[4] Division of Healthcare Quality and Promotion, National Center for Emerging and Zoonotic
Infectious Diseases, Centers for Disease Control and Prevention, Atlanta, Georgia
[5] Division of STD Prevention, Centers for Disease Control and Prevention, National Center for
HIV/AIDS, Viral Hepatitis, STD and TB Prevention, Atlanta, Georgia

17
18
19
20 Abstract
21

Transmission heterogeneity plays a critical role in the dynamics of an epidemic. During an
outbreak of an emerging infectious disease, efforts to characterize transmission heterogeneity are
generally limited to quantifications during a small outbreak or a limited number of generations of
a larger outbreak. Understanding how transmission heterogeneity itself varies over the course of
a large enduring outbreak not only improves understanding of observed disease dynamics but
also informs public health strategy and response. In this study, we employ a method, adaptable to
other emerging infectious disease outbreaks, to quantify spatiotemporal variation in transmission
heterogeneity for the 2022 mpox outbreak in the United States. Based on past research on mpox
and following reports of potential superspreading events early in this outbreak, we expected to
find high transmission heterogeneity as quantified by the dispersion parameter of the offspring
distribution, k. Our methods use maximum likelihood estimation to fit a negative binomial
distribution to transmission chain offspring distributions informed by a large mpox contact
tracing dataset. We find that, while estimates of transmission heterogeneity varied across the
outbreak with spatiotemporal pockets of higher heterogeneity, overall transmission heterogeneity
was low. When testing our methods on simulated data from an outbreak with high transmission
heterogeneity, k estimate accuracy depended on the contact tracing data completeness. Because
the actual contact tracing data had high incompleteness, our values of k estimated from the
empirical data may be artificially high. However, it is also possible that our estimates accurately
reflect low transmission heterogeneity for the United States mpox outbreak.

41
42
43 Introduction
44
In an infectious disease outbreak, transmission heterogeneity occurs when there is variation in
the number of secondary infections per infected host, which can be caused by underlying

mechanistic heterogeneity. This variation can manifest at spatial (May and Anderson 1984),
temporal (Bjørnstad, Finkenstädt, and Grenfell 2002), and/or individual (Lloyd-Smith et al.
2005; Ke et al. 2022) scales. These different components of transmission heterogeneity interact
with one another, creating the potential for complex patterns of transmission heterogeneity over
the course of an outbreak. Because transmission heterogeneity plays a critical role in epidemic
spread, characterizing how it changes over an outbreak can help to illuminate the causes behind
observed epidemic dynamics, informing measures taken by public health decision makers.
54
Prior methods have simultaneously estimated the mean transmission (R) and the dispersion (k)
parameters from data on the number of secondary infections per primary infection (*i.e.*, the
offspring distribution) in an outbreak (Lloyd-Smith et al. 2005; Lloyd-Smith 2007; Blumberg
and Lloyd-Smith 2013; Toth et al. 2015). These methods have been applied both retrospectively
and during ongoing outbreaks (Toth et al. 2016; Kucharski and Althaus 2015; Nishiura et al.
2015) for several diseases. Past research on transmission heterogeneity has largely focused on
small outbreaks or has used data from early transmission generations in larger outbreaks (Lloyd-
Smith et al. 2005; Lloyd-Smith 2007; Blumberg and Lloyd-Smith 2013; Toth et al. 2015; 2016;
Kucharski and Althaus 2015; Nishiura et al. 2015). Some research has explored how the first
transmission generation may differ from subsequent generations in R and k (Toth et al. 2016),
but work to understand how individual transmission heterogeneity may vary over the component
outbreaks of a large epidemic is more rare (Lau et al. 2020; Tkachenko et al. 2021; Miyama,
Jung, and Nishiura 2022).
68
Among those diseases for which transmission heterogeneity has been characterized in the past is
mpox (formerly 'Monkeypox' (World Health Organization 2022b)). Monkeypox virus (MPXV)
is a member of the *Orthopoxvirus* genus of pathogens and is endemic in West and Central
Africa. Mpox was first detected in humans in 1970, shortly after the eradication of smallpox
(Breman et al. 1980), and has subsequently caused occasional outbreaks thought to result from
repeated spillover events (Nalca et al. 2005; Yinka-Ogunleye et al. 2018; Bunge et al. 2022;
Durski et al. 2018). While the exact reservoir host is still unknown, the virus has been found in
several rodent species native to central and west African countries (Durski et al. 2018). Mpox is
known to spread through direct contact with mpox rash, scabs, or bodily fluids; touching objects
that have been used by someone with mpox; contact with respiratory secretions; and sexual or
intimate contact with an infected person (Centers for Disease Control and Prevention 2022).
80
Past outbreaks have been largely contained to a modest number of cases. However, regional
outbreaks have been growing in recent years (Sklenovská and Van Ranst 2018; Beer and
Bhargavi Rao 2019; Rimoin et al. 2010). Past work to define the basic reproductive number (R_0,
the expected mean transmissions per infection in a large, fully susceptible population), for mpox
outbreaks have produced estimates below 1, the critical threshold above which an infectious
disease will continue to spread and below which an outbreak will eventually die out. However,
more recent work has accounted for declining cross-protective immunity from smallpox to reach
an estimate of $R_0 = 2.13$ (uncertainty bounds: 1.46-2.67) for mpox in the Democratic Republic of
Congo (Grant, Nguyen, and Breban 2020). Past estimates of transmission heterogeneity,
quantified by the inversely varying dispersion parameter (low values of k indicate high
transmission heterogeneity) of the offspring distribution (k), for mpox outbreaks have been 0.58
(90% confidence interval [CI]: 0.32-3.57) (Lloyd-Smith et al. 2005), 0.36 (0.16-1.47), and 0.33

(0.19-0.64) (Blumberg and Lloyd-Smith 2013), indicating high transmission heterogeneity.
94
In April 2022, MPXV was identified outside of the countries where the virus is endemic and
quickly spread to over 100 countries (World Health Organization 2022a). The World Health
Organization (WHO) declared this mpox epidemic a Public Health Emergency of International
Concern on July 23, 2022. This rapid global spread was not characteristic of an infectious
disease with a basic reproductive number below 1. Contact tracing conducted during the 2022
outbreak indicated that the disease was initially spreading by sexual contact between gay and
bisexual men, and nonbinary and transgender people, often at large gatherings (Philpott et al.
2022). In light of this, public health messaging focused on reducing spread among this group of
people, especially at large gatherings (Delaney et al. 2022). Potentially due to a novel, sexually
mediated, dominant transmission route, the discrepancy between estimates of the basic
reproductive number for mpox outbreaks became globally relevant. Estimates of the 2022
outbreak vary by region and range from 1.56 to 3.8 (Branda, Pierini, and Mazzoli 2022) or
potentially higher (Endo et al. 2022).
108
Because the degree distribution of sexual contact networks can be overdispersed, having greater
variability than expected, transmission through sexual contact networks can result in
heterogeneous patterns of secondary transmission across the infected population (Handcock and
Jones 2006). Spreading of the disease through large gatherings, termed 'superspreader events',
can also cause high levels of transmission heterogeneity (Althouse et al. 2020). Such
heterogeneous transmission has been shown to influence outbreak dynamics (Toth et al. 2015;
Elie, Selinger, and Alizon 2022) and therefore may have contributed substantially to the rapid
global spread of mpox. Modeling studies have provided support for the idea that heterogeneous
transmission through overdispersed contact networks could partially explain the dynamics of the
2022 mpox outbreak (Endo et al. 2022; Spicknall et al. 2022). To date, empirical estimates of
transmission heterogeneity for the present outbreak have not been published.
120
Here, we estimate the mean transmission and transmission heterogeneity for the 2022 mpox
epidemic in the United States using contact tracing data fortified with algorithmically determined
probable case pairs. We estimate these parameters directly from the empirical data and, through
simulation, assess the sensitivity of our approach. We stratify our analysis by geographic
location and time. In this way, we characterize transmission heterogeneity on spatial, temporal,
and individual scales. Our aims are to (1) better understand the dynamics of the 2022 mpox
outbreak and (2) advance the theory of transmission heterogeneity.
128
129   Methods
130
Overview
We quantified the mean transmission using R, defined as the mean number of secondary
transmissions per infected individual at a given time period and location and quantified as the
mean of the offspring distribution generated from a series of transmissions within that period and
location. We quantified transmission heterogeneity using k, the dispersion parameter of the
negative binomial distribution, with lower values indicating greater heterogeneity (Fig. 1a). We
generated estimates of R and k for the United States and its public health jurisdictions at different
periods of time based on offspring distributions of algorithmically generated numbers of

secondary infections per case. We then used a maximum likelihood approach to fit negative
binomial distributions to the offspring distributions of given regions and time periods and
estimated the 95% confidence intervals (CI) for both parameters (*detailed below*). Transmission
trees were constructed using methods similar to those in previous works (e.g., Haydon et al 2003;
Wallinga and Teunis 2004; Heijne et al 2012), though our method differs in notable ways, as
detailed below.
145

a.

146
TEXT BOX: Figure 1. (a) Frequencies of negative binomially distributed discrete data, smoothed
with a bandwidth of .75 and normalized by total samples for visual clarity. These negative
binomial pseudo-distributions have R (mean) = 2 and varying k (dispersion). When k is infinity
(red), the negative binomial distribution is equivalent to the Poisson. (b-d) Visualizations of the
contact network used in simulations, with simulated jurisdictions shown in colored polygons. See
text for description of methods used to generate networks (b) shows two jurisdictions, (c) shows
three jurisdictions, inclusive of the two in (b), and (d) shows the full contact network.
Overlapping polygons in (d) are visual artifacts of the large network – each node is associated
with only one jurisdiction.

Data

We used data collected through U.S. Centers for Disease Control and Prevention (CDC) case detection and contact tracing in the United States from May 1st to November 7th, 2022. This dataset includes, for each probable and confirmed case, information on date of symptom onset, date of reporting, date of positive test, reporting jurisdiction, any recent travel history, and any known contacts with other probable or confirmed cases.

Processing

To compile the offspring distribution, we started by including contact-tracing identified probable transmission events and their associated cases. For each case where a probable infector has not been identified by contact tracing, we used dates of symptom onset included in the dataset and published serial interval estimates to compile a pool of possible infectors (Fig. 2). We defined a possible infector of a case as one with overlapping location (US county) and whose date of symptom onset is within one maximum serial interval, 21 days, of the case's date of symptom onset (UK Health Security Agency 2022). Because the pool of possible infectors generally includes more than one possible infector, we then sampled one possible infector from each pool to serve as the chosen infector for each case. In this sampling process, the serial interval was calculated for each possible transmission, and the probability that a serial interval was sampled is equal to the gamma distribution density, parameterized according to published estimates of the serial interval distribution for mpox (shape = 2.9, scale = 2.9; (Madewell et al. 2022)). The case producing the sampled serial interval was assigned to be the most probable infector for a case. From this dataset, containing both contact-tracing identified and algorithmically determined transmission pairs, we calculated the number of secondary infections produced for each case and fit a negative binomial distribution to the resulting offspring distribution, finding the best-fit parameter values and their 95% CIs. This entire process was repeated 100 times with replacement, producing a sample of the possible offspring distributions of an outbreak.

187
188
TEXT BOX: Figure 2. A representation of data processing to build a complete transmission tree.
(a) Pre-processed case data points from two public health jurisdictions with four contact tracing
identified links. (b) Arrows added to show pools of possible infectors for each case, drawn based
on jurisdiction and time (each column of points represents one time step where, for example,
possible infectors come from only the most recent two time steps). (c) Chosen infectors sampled
and other possible infectors disregarded. This is the post-processing data set used for analysis.

195
196   We compiled a separate set of offspring distributions using only the contact tracing-identified
197   transmission pairs. Because this set of data is much smaller than the hybrid dataset described
198   above (see results), we employ it only at the national level. Parameter estimates from the two
199   datasets are compared as a validity check.
200
Parameter estimation
To fit a negative binomial distribution to the offspring distribution, we used a maximum
likelihood approach, following established methods (Lloyd-Smith et al. 2005; Toth et al. 2016).
This approach is sensitive to the number of individuals with mpox assumed to transmit to no
other individuals, which we identify as terminal cases. Because a portion of the individuals in
our dataset with no reported offspring may have actually transmitted the disease to others
without being detected, including these false terminal cases would artificially inflate the number
of zeros in the offspring distribution. We used the zero-inflated negative binomial to test the
sensitivity of our methods to this type of potential bias, complementing past work that accounts
for underreporting of terminal cases (Lloyd-Smith 2007).
211
The zero-inflated negative binomial differs from the negative binomial distribution in its
inclusion of a parameter p_0, the probability that an individual's onward transmission data are
absent, independent of the number of individuals they infected, leaving $1 - p_0$ as the probability
that the individual's transmission data are complete. A lower p_0 indicates greater confidence in
the accuracy of the data, and the zero-inflated negative binomial reduces to the negative binomial
when $p_0 = 0$. Similar to past work that fits modified negative binomial distributions to discrete
data that undercounts zeros (Ridout, Demetrio, and Hinde 1998; Lloyd-Smith 2007), accounting
for p_0 helps to correct for missingness in the data and make more reliable estimates of R and k.
The probability mass function (pmf) for the zero-inflated negative binomial can be defined as
follows, with the reported number of offspring for each infected individual distributed based on
whether or not the data point, y, is a zero, where we assume the true number of offspring is
negative binomially distributed:
224

$$
f(y) = \begin{cases} p_0 + (1 - p_0) \left(\frac{k}{k+R} \right)^k & \text{if } y = 0 \\ (1 - p_0) \begin{pmatrix} k + y - 1 \\ y \end{pmatrix} \left(\frac{R}{k+R} \right)^y \left(\frac{k}{k+R} \right)^k & \text{if } y \neq 0 \end{cases}
$$

225
226
In the negative binomial estimates, the logarithms of negative binomially distributed
probabilities of each data point are a function of R and k, while in the zero-inflated estimates the
logarithms of zero-inflated negative binomially distributed probabilities of each data point are a
function of R, k, and p_0 . We used maximum likelihood estimation by taking the log-likelihood
function of the entire data set. We then used multivariate optimization to find R and k for the
negative binomial estimates and R, k, and p_0 for the zero-inflated estimates. We used the
likelihood ratio test to find 95% CIs for these parameters.
234
Sliding and telescoping window approaches
To evaluate change in parameter estimates over time, we conducted our fitting methods on
offspring distributions generated across both sliding and telescoping windows of time. In the

sliding window approach, offspring distributions were generated consecutively over 14-day
periods, with each consecutive window starting 7 days after the start of the previous window.
The sliding window approach generates estimates that can be likened to parameters R_t and k_t
described elsewhere (Schneckenreither et al. 2022; Zhang, Britton, and Zhou 2022; Wallinga and
Teunis 2004), though to avoid confusion regarding definitions and assumptions that differ from
those of our estimated parameters, we simply use R and k. In the telescoping window approach,
consecutive windows are inclusive of the previous window; the end date for each consecutive
window is advanced 7 days while the start date remains the same, May 1st, 2022. Estimates of R
and k generated from the telescoping window approach describe the mean and dispersion of
secondary transmission from the beginning of the outbreak up to the end date of a given time
window. In this way, they can be thought of as the effective R and k parameters, but again we
simply use our previously defined R and k terms for the mean and dispersion of the offspring
distribution.
251
Simulations
To characterize the reliability of our methods to estimate transmission heterogeneity, we
repeated them on simulated data generated with a model of infectious disease spread over a static
contact network. We conducted simulations on undirected scale-free networks, shown to produce
relatively heterogeneous transmission (Bansal, Grenfell, and Meyers 2007). We initialized the
scale-free network with a Barabási-Albert algorithm of preferential attachment (Barabási and
Albert 1999), allowing for the network to serve as a proxy for a population with highly
heterogeneous contact rates where simulated outbreaks can be expected to result in an
overdispersed offspring distribution. The network was generated using the *barabasi.game*
function from the *igraph* package in the *R* programming environment (Csardi and Nepusz 2006),
using a population of 50,000 and power of 1.
263
Each outbreak was seeded with a randomly chosen single infection. Disease dynamics were
modeled following a *Susceptible – Exposed – Infectious – Recovered* paradigm, parameterized
for mpox. The latent period was a random variable drawn from a lognormal distribution with log
mean = 1.5 and log standard deviation = 0.7 (Madewell et al. 2022). For lack of detailed
information on the duration of infectiousness for mpox, the infectious period in our simulations
was a random variable drawn from a gamma distribution with shape = 21 days and scale = 1.
This distribution is consistent with the reported serial interval distribution (Madewell et al.
2022), clinical guidance indicating symptoms last 2-4 weeks (Word Health Organization 2022c),
and investigations reporting that some hospitalized cases show symptoms for much longer
periods (Adler et al. 2022). However, it should be noted that more detailed study may reveal that
the distribution we used does not describe the infectious period for mpox well. Contacts of
infectious nodes were infected with probability of 0.15 per contact day.
276
In order to replicate our methods on the simulated data, we assigned each node a "jurisdiction"
using a modularity optimization algorithm that identifies community structure in networks
(Blondel et al. 2008), as depicted in Figure 1b. We then identified the most probable infector for
each case, as detailed above, and estimated the parameters R and k by fitting a negative binomial
distribution, as detailed above. In order to test the sensitivity of our methods to incompleteness in
detailed contact information, we also conducted a batch of simulations with the same level of
incompleteness as the empirical dataset.

284
285   All computational methods were conducted in R statistical software (R Core Team 2017).
286
287   <u>Results</u>

288
TEXT BOX: Figure 3. Temporal variation in transmission parameters. Left column shows

estimates made with a sliding window, right column shows estimates made with a telescoping
window, starting on May 1st, 2022. Time points for the sliding window approach (left column)
are the start of the 14-day sliding windows, time points for the telescoping window approach
(right column) are the end of the telescoping windows. Top row shows empirical parameter
estimates, middle and bottom rows show estimates made from simulated outbreaks. In the
middle and bottom rows, the x axis is units of days since the start of the outbreak simulation. In
all plots, black points represent values of $1/k$, plotted on the right vertical axes, whereas blue
points represent values of R, plotted on the left vertical axes. (a,b) Estimated median and 95%
CIs for $1/k$ (black, right-hand vertical axis) and R (blue, left-hand vertical axis), displayed over
daily new mpox cases(pink) for the 2022 mpox epidemic in the United States. (c-f) Parameter
values estimated on simulated data from a simulated outbreak which exhibited strongly
heterogeneous transmission. (c-d) Estimates from simulated data where probable transmission
pairs were determined with complete information on true individual contacts for 3.6% of the
population, with the remainder being determined only with simulated dates and geographic
locations. This reflects the condition of the empirical data in a,b. (e-f) Estimates from simulated
data where all pairs were determined with complete information on true individual contacts. In
all plots, values of $1/k$, instead of k, are displayed to ease visualization.
307
308   *National estimates*
309   There were 22,391 probable and confirmed mpox cases in the empirical dataset, after excluding
310   5,074 (18.47%) cases with missing data on symptom onset date. In the exclusive dataset, 820
311   (3.66%) of these cases had contact tracing-identified connections to other probable and
312   confirmed cases, compared to 12 (0.24%) of those excluded cases.
313
R and k estimated at the national level changed over time. Using a sliding 2-week window (Fig.
3a), we observe that, after early increase to a peak, estimated R declines over the course of the
epidemic, dropping below 1 around the peak of new infections on approximately August 1st, as
is expected to occur when an outbreak changes from growth to decline. Median estimated k
remains above 1 for all time windows, though with high variability – a pattern which does not
indicate high transmission heterogeneity.
320
Applying a telescoping window approach allows us to evaluate R and k taken cumulatively at
different points during the epidemic (Fig 3b). Using the time window beginning at the start of the
outbreak, May 1st, and ending at the peak of new cases, August 1st, we find that $R = 1.25$ (95%
CI: 1.21-1.28) and $k = 12.18$ (95% CI: 8.18-21.27). Using a zero-inflated negative binomial
distribution returned the same values of R and k for this period, with a p_0 value of 0.001 (95% CI:
0.00-0.02) (see supplemental results).
327
328
329   TEXT BOX: Figure 4: Spatiotemporal variation in estimated values of R across US states during
330   the middle of the 2022 mpox epidemic. Unshaded regions indicate insufficient data to conduct

331   parameter estimations for that region and time period.

332
Jurisdiction estimates
Estimates of R varied across U.S. State and over time (Fig. 4). Estimates of k also vary across
public health jurisdictions and over time (Table S1). While median estimates of k are over 1 at all
time points for every jurisdiction, the variation in k estimates across iterations is substantial for
some points. Estimates of p_0 made using the zero-inflated negative binomial were highly variable
across jurisdictions and timepoints (see supplemental results).
339
Simulations
Conducting the methods on data resulting from an outbreak simulation over a scale-free contact
network showed reliability in detecting overdispersed offspring distributions when detailed
contact information was used, reflected by estimates of k below 1 (Fig. 3e,f).When using only
network structure-based jurisdiction clusters for 96.4% of the population, and using detailed
contact information for the remaining 3.6%, estimates of k were much higher, staying above 1
(Fig. 3c,d).
347
348   <u>Discussion</u>
349
Our results suggest that transmission of mpox during the 2022 outbreak in the United States was
not highly heterogeneous. Instead, estimates of $k \gg 1$ across many jurisdictions and time periods
indicate that transmission was approximately what would be expected in an evenly mixing
population of people with equal inherent transmissibility. We do find some indication of
overdispersed transmission in certain locations during certain time periods (Table S1). Such a
pattern could be consistent with explanations that invoke individual level transmission
heterogeneity, resulting from individuals who spread to far more than the average infected
individual and/or event-based superspreading. Whatever the case, our methods do not return
evidence that such heterogeneity has been widespread. However, conducting our methods on
simulated data indicates that we only expect to recover a strong pattern of heterogeneity if we
have very high-quality data. Because the data quality in our empirical dataset likely varied by
case, geographic location, and through time, we cannot rule out the possibility that our estimates
of k for the 2022 US mpox epidemic are artificially high, and care must be taken with any
interpretation.
364

There are points during the data collection process when errors may be introduced and
underreporting may occur, resulting in incomplete and/or inaccurate contact tracing data. Our
methods operated under a number of assumptions related to missing data. First, it was assumed
that the data are a representative sample of all mpox cases. If there is an association between case
detection and secondary infections produced, then this assumption could impact our estimates.
For example, diagnostic testing can be biased towards cases of more severe disease (Tsang et al.
2021), which could produce more secondary infections through an increased duration of viral
shedding (Fielding et al. 2014). Second, we excluded the cases lacking symptom onset dates, and
we assumed that the cases for which date of symptom onset exist constitute a random sample
from the population. If there is a relationship between having the symptom onset date for a case
and some other factor (e.g., information on contacts), then our results could be impacted.
376
Incomplete contact tracing data can be the result of various errors introduced during the data
collection process. For the 2022 mpox outbreak we focused on the undercounting of secondary
transmissions manifesting as extra zeros in the offspring distribution, which are mainly
introduced in the following three ways. First, inconsistencies in the dates associated with cases
could impact our analyses. Date of reporting could occur at any time during the course of an
infection, even after the patient is no longer infectious, and often includes "data dumps" where
particular jurisdictions may have posted large portions of their data on a given day. While we use
date of symptom onset, it is possible that these dates were influenced by behavioral or systematic
processes in healthcare and disease progression . Even for cases with a date of symptom onset
we cannot be sure those dates are consistent with the start of the infectious period. For the most
recent cases in the dataset, it may be too close to the end of the dataset to know whether any of
their contacts had become infected. Second, there are barriers to a suspected case becoming a
probable or confirmed case – tests may only be provided to verified contacts of another probable
or confirmed case or to those in high-risk subpopulations. In the extreme, this feature of the
contact tracing process can result in missing entire transmission chains. Third, identifying case-
by-case contacts and their exact dates can be difficult in sexual/social networks such as those
where mpox has been shown to spread. Infected individuals may not know or choose to disclose
the identities of their close contacts and additionally may delay seeking care due to historical
discrimination and stigmatization of the LGBTQ+ community in the healthcare setting (Ennab et
al. 2022; Rodríguez et al. 2022; The Lancet Regional Health – Europe 2022). The spatiotemporal
variability in estimates of p_0 may reflect variation in data incompleteness related to this potential
zero inflation. However, overall at the national level, estimates of p_0 were consistently low, and
our sensitivity analysis using the zero-inflated negative binomial distribution returned very
similar parameter estimates to those returned using the negative binomial distribution. In a large
population, the influence of missing case data on estimates of R and k may be minimized or
obscured.
One reason that transmission heterogeneity is of potential interest to public health decision
making is that it can indicate that within-host processes during infection result in inherent
differences in the probability of infection given contact across infected people (Ke et al. 2022).
In such a situation, equal rates of contact between infected and susceptible people will result in
different numbers of secondary infections, and interventions, when implemented, may differ in
their efficacy relative to expectations. Additionally, we should expect to find consistently low
values of k across spatiotemporal dimensions. In the case of the 2022 mpox outbreak in the
United States, we did not find evidence that this sort of individual-level transmission

heterogeneity was broadly present. Instead, the rare occurrences of highly heterogeneous
transmission that we observe likely resulted from differences in contact rates, either due to
standing individual variation in contact rates or to the occurrence of superspreading events. In
this case, targeted interventions could be employed to reduce the spread of disease. Without high
heterogeneity in transmission, the rate of disease spread is well-characterized by an R_0 value
greater than the threshold 1, as has been suggested by other recent studies of R_0 for the 2022
mpox epidemic (Branda, Pierini, and Mazzoli 2022; Endo et al. 2022). These results are
consistent with the theory that cross-protective immunity from smallpox may be waning,
resulting in higher rates of overall transmission (Grant, Nguyen, and Breban 2020).
420
In light of past work which estimated low values of k for mpox outbreaks, it is surprising that we
did not recover strong evidence of high transmission heterogeneity for this mpox epidemic. The
population structure, timing, environmental characteristics, and primary mechanisms of
transmission differed between the 2022 mpox outbreak in the United States and the outbreaks
described in other studies. It is possible that these differences explain the differences in estimates
of k between our study and the others. The timing of the US outbreak relative to the beginning of
the global epidemic may also have afforded individuals with a higher potential risk of infection
the time necessary to take precautions against infection. This may be especially true as it relates
to large events, which were implicated in the initial spread outside the United States and were the
focus of targeted public health messaging in the United States (Delaney et al. 2022). Indeed,
survey results suggest that many US gay and bisexual men, and nonbinary and transgender
people took active measures to reduce the risk of exposure to mpox (Delaney et al. 2022), and so
this could explain the values of k we observed. It is important to recognize that, while it is
plausible that the k estimates recovered by our methodology accurately reflect real patterns of
transmission heterogeneity, the limited ability to detect cases and track their contacts during the
2022 mpox outbreak in the United States could impact our estimates of both R and k.
437
Molecular, phylogenetic, and in-vivo studies have indicated that monkeypox virus samples from
the outbreak in this study are significantly divergent from those of past outbreaks (Isidro et al.
2022, Americo et al. 2023). The outbreak investigated in the current study is caused by a virus
that might have quite different epidemiological characteristics than the virus associated with
outbreaks upon which past estimates of epidemiological parameters have been based. Our
methods can be extended to clarify the differences between our estimates of k and those of past
studies of the same disease.
445
Our study contributes to a small but growing literature on spatiotemporal variation in
transmission heterogeneity of human infectious diseases. Past research on transmission
heterogeneity has largely focused on small outbreaks or has used data from the first generations
of infections in larger outbreaks. Related research on the influence of contact networks on the
spread of infectious diseases has illuminated important concepts regarding transmission
heterogeneity. For example, as the variation in the degree distribution of the infected population
changes over the course of an epidemic simulated on a network with a particular underlying
structure (Bansal, Grenfell, and Meyers 2007), transmission heterogeneity should be expected to
change over time.
455
456   Our methods used a straightforward algorithm to determine likely infectors for cases without a

contact tracing-identified infector, and we subjected these methods to a limited sensitivity
analysis using simulated data. In the process, we determined that our methods can be expected to
accurately estimate k with high-quality data on the direct contacts of each case. In some
instances, our empirical data may meet these qualifications, but in other instances they likely do
not. So, it is possible for our methods to mask high transmission heterogeneity under some
conditions. One study has found that different models used to measure transmission
heterogeneity may result in different estimates with starkly different interpretations, with their
instant-individual heterogeneity model recovering high heterogeneity when alternate models do
not (Zhang, Britton, and Zhou 2022). The methods we present here are adaptable to incidence
time series data, holding promise that transmission heterogeneity may be reliably and easily
estimated, but care must be taken to contextualize findings within the caveats of the data and the
potential limitations of data gathered during emerging outbreaks.

Acknowledgements

The findings and conclusions in this report are those of the authors and do not necessarily
represent the views of the Centers for Disease Control and Prevention.

Literature Cited

Adler H, Gould S, Hine P, Snell LB, Wong W, Houlihan CF, et al. 2022 "Clinical features and
　　management of human monkeypox: a retrospective observational study in the UK." *The*
　　Lancet Infectious Diseases. 22(8):1153–62.

Americo JL, Earl PL, Moss B. 2023 "Virulence differences of mpox (monkeypox) virus clades I,
　　IIa, and IIb.1 in a small animal model." *Proc Natl Acad Sci USA*.
　　21;120(8):e2220415120.

Althouse, Benjamin M., Edward A. Wenger, Joel C. Miller, Samuel V. Scarpino, Antoine Allard,
　　Laurent Hebert-Dufresne, and Hao Hu. 2020. "Superspreading Events in the
　　Transmission Dynamics of SARS-CoV-2: Opportunities for Interventions and Control."
　　PLoS Biology 18 (11): 1–13. https://doi.org/10.1371/journal.pbio.3000897.

Bansal, Shweta, Bryan T. Grenfell, and Lauren Ancel Meyers. 2007. "When Individual
　　Behaviour Matters: Homogeneous and Network Models in Epidemiology." *Journal of the*
　　Royal Society Interface 4 (16): 879–91. https://doi.org/10.1098/rsif.2007.1100.

Barabási, Albert-László, and Réka Albert. 1999. "Emergence of Scaling in Random Networks."
　　Science 286 (5439): 509–12. https://doi.org/10.1126/science.286.5439.509.

Beer, Ellen M., and V. Bhargavi Rao. 2019. "A Systematic Review of the Epidemiology of
　　Human Monkeypox Outbreaks and Implications for Outbreak Strategy." *PLoS Neglected*
　　Tropical Diseases 13 (10): 1–20. https://doi.org/10.1371/journal.pntd.0007791.

Bjørnstad, Ottar N., Bärbel F. Finkenstädt, and Bryan T. Grenfell. 2002. "Dynamics of Measles
　　Epidemics: Estimating Scaling of Transmission Rates Using a Time Series SIR Model."
　　Ecological Monographs 72 (2): 169–84. https://doi.org/10.1890/0012-
　　9615(2002)072[0169:DOMEES]2.0.CO;2.

Blondel, Vincent D, Jean-Loup Guillaume, Renaud Lambiotte, and Etienne Lefebvre. 2008.
　　"Fast Unfolding of Communities in Large Networks." *Journal of Statistical Mechanics:*
　　Theory and Experiment 2008 (10): P10008. https://doi.org/10.1088/1742-

5468/2008/10/P10008.

Blumberg, Seth, and J. O. Lloyd-Smith. 2013. "Inference of R0 and Transmission Heterogeneity
from the Size Distribution of Stuttering Chains." *PLoS Computational Biology* 9 (5): 1–
17. https://doi.org/10.1371/journal.pcbi.1002993.

Branda, Francesco, Massimo Pierini, and Sandra Mazzoli. 2022. "Monkeypox: Early Estimation
of Basic Reproduction Number R 0 in Europe." *Journal of Medical Virology*.
https://doi.org/10.1002/jmv.28270.

Breman, J. G., K. Ruti, M. V. Steniowski, E. Zanotto, A.I. Gromyko, and I. Arita. 1980. "Human
Monkeypox, 1970-79." *Bulletin of the World Health Organization* 58 (2): 165–82.

Bunge, Eveline M., Bernard Hoet, Liddy Chen, Florian Lienert, Heinz Weidenthaler, Lorraine R.
Baer, and Robert Steffen. 2022. "The Changing Epidemiology of Human Monkeypox—
A Potential Threat? A Systematic Review." *PLoS Neglected Tropical Diseases* 16 (2): 1–
20. https://doi.org/10.1371/journal.pntd.0010141.

Centers for Disease Control and Prevention. 2022. "CDC Monkeypox Response: Transmission -
Media Statement," 1–2.

______. 2022. "Case Definitions for Use in the 2022 Mpox Response"
https://www.cdc.gov/poxvirus/mpox/clinicians/case-definition.html

Csardi G, Nepusz T. 2006. "The igraph software package for complex network research."
InterJournal, Complex Systems, 1695. https://igraph.org.

Delaney, Kevin P., Travis Sanchez, Marissa Hannah, O. Winslow Edwards, Thomas Carpino,
Christine Agnew-Brune, Kaytlin Renfro, et al. 2022. "Strategies Adopted by Gay,
Bisexual, and Other Men Who Have Sex with Men to Prevent Monkeypox Virus
Transmission — United States, August 2022." *MMWR. Morbidity and Mortality Weekly*
Report 71 (35): 1126–30. https://doi.org/10.15585/mmwr.mm7135e1.

Durski, Kara N, Andrea M Mccollum, Yoshinori Nakazawa, Brett W Petersen, and Mary G
Reynolds. 2018. "Emergence of Monkeypox in West Africa and Central Africa, 1970–
2017." *Releve Epidemiologique Hebdomadaire* 93 (11): 125–32.

Elie, Baptiste, Christian Selinger, and Samuel Alizon. 2022. "The Source of Individual
Heterogeneity Shapes Infectious Disease Outbreaks." *Proceedings of the Royal Society*
B: Biological Sciences 289 (1974). https://doi.org/10.1098/rspb.2022.0232.

Endo, Akira, Hiroaki Murayama, Sam Abbott, Ruwan Ratnayake, Carl A. B. Pearson, W. John
Edmunds, Elizabeth Fearon, and Sebastian Funk. 2022. "Heavy-Tailed Sexual Contact
Networks and the Epidemiology of Monkeypox Outbreak in Non-Endemic Regions, May
2022." *MedRxiv*, no. May: 2022.06.13.22276353.

Ennab, Farah, Faisal A. Nawaz, Kapil Narain, Goodluck Nchasi, and Mohammad Yasir Essar.
2022. "Rise of Monkeypox: Lessons from COVID-19 Pandemic to Mitigate Global
Health Crises." *Annals of Medicine and Surgery* 79 (June): 14–17.
https://doi.org/10.1016/j.amsu.2022.104049.

Grant, Rebecca, Liem Binh Luong Nguyen, and Romulus Breban. 2020. "Modelling Human-to-
Human Transmission of Monkeypox." *Bulletin of the World Health Organization* 98 (9):
638–40. https://doi.org/10.2471/BLT.19.242347.

Handcock, Mark S., and James Holland Jones. 2006. "Interval Estimates for Epidemic
Thresholds in Two-Sex Network Models." *Theoretical Population Biology* 70 (2): 125–
34. https://doi.org/10.1016/j.tpb.2006.02.004.

Haydon, D.T., M. Chase-Topping, D. J. Shaw, L. Matthews, J. K. Friar, J. Wilesmith and M. E.
J. Woolhouse. 2003. "The Construction and Analysis of Epidemic Trees with Reference

to the 2001 UK Footand-Mouth Outbreak." *Proceedings of the Royal Society: B*
270(1511): 121-127. https://doi.org/10.1098/rspb.2002.2191

Heijne, Jannek C.M., Marc Rondy, Linda Verhoef, Jacco Wallinga, Mirjam Kretzschmar, Nicola
Low, Marion Koopmans and Peter F. M. Teunis. 2012. "Quantifying Transmission of
Norovirus During an Outbreak." *Epidemiology* 23:277-284.
https://doi.org/10.1097/EDE.0b013e3182456ee6

Isidro J, Borges V, Pinto M, Sobral D, Santos JD, Nunes A, et al. 2022. "Phylogenomic
characterization and signs of microevolution in the 2022 multi-country outbreak of
monkeypox virus." *Nat Med.* 28(8):1569–72.

Ke, Ruian, Pamela P. Martinez, Rebecca L. Smith, Laura L. Gibson, Agha Mirza, Madison
Conte, Nicholas Gallagher, et al. 2022. "Daily Longitudinal Sampling of SARS-CoV-2
Infection Reveals Substantial Heterogeneity in Infectiousness." *Nature Microbiology* 7
(5): 640–52. https://doi.org/10.1038/s41564-022-01105-z.

Kucharski, A. J., and C. L. Althaus. 2015. "The Role of Superspreading in Middle East
Respiratory Syndrome Coronavirus (MERS-CoV) Transmission." *Eurosurveillance* 20
(25): 14–18. https://doi.org/10.2807/1560-7917.ES2015.20.25.21167.

Lau, Max S.Y., Bryan Grenfell, Michael Thomas, Michael Bryan, Kristin Nelson, and Ben
Lopman. 2020. "Characterizing Superspreading Events and Age-Specific Infectiousness
of SARS-CoV-2 Transmission in Georgia, USA." *Proceedings of the National Academy*
of Sciences of the United States of America 117 (36): 22430–35.
https://doi.org/10.1073/pnas.2011802117.

Lloyd-Smith, J. O. 2007. "Maximum Likelihood Estimation of the Negative Binomial
Dispersion Parameter for Highly Overdispersed Data, with Applications to Infectious
Diseases." *PLoS ONE* 2 (2): 1–8. https://doi.org/10.1371/journal.pone.0000180.

Lloyd-Smith, J. O., S. J. Schreiber, P. E. Kopp, and W. M. Getz. 2005. "Superspreading and the
Effect of Individual Variation on Disease Emergence." *Nature* 438 (7066): 355–59.
https://doi.org/10.1038/nature04153.

Madewell, Zachary J, Kelly Charniga, Nina Masters, and Et. Al. 2022. "Serial Interval and
Incubation Period Estimates of Monkeypox Virus Infection in 12 U.S. Jurisdictions, May
– August 2022." *MedRxiv*. https://doi.org/10.1101/2022.10.26.22281516.

May, Robert M., and Roy M. Anderson. 1984. "Spatial Heterogeneity and the Design of
Immunization Programs." *Mathematical Biosciences* 72 (1): 83–111.
https://doi.org/10.1016/0025-5564(84)90063-4.

Miyama, Takeshi, Sung-mok Jung, and Hiroshi Nishiura. 2022. "Decrease in Overdispersed
Secondary Transmission of COVID-19 over Time in Japan." *Epidemiology and Infection*,
November, 1–29. https://doi.org/10.1017/S0950268822001789.

Nalca, Aysegul, Anne W. Rimoin, Sina Bavari, and Chris A. Whitehouse. 2005. "Reemergence
of Monkeypox: Prevalence, Diagnostics, and Countermeasures." *Clinical Infectious*
Diseases 41 (12): 1765–71. https://doi.org/10.1086/498155.

Nishiura, H., Y. Miyamatsu, G. Chowell, and M. Saitoh. 2015. "Assessing the Risk of Observing
Multiple Generations of Middle East Respiratory Syndrome (MERS) Cases given an
Imported Case." *Eurosurveillance* 20 (27): 6–11. https://doi.org/10.2807/1560-
7917.ES2015.20.27.21181.

Philpott, David, Christine M Hughes, Karen A Alroy, Janna L Kerins, Jessica Pavlick, and
Lenore Asbel. 2022. "Epidemiologic and Clinical Characteristics of Monkeypox Cases
— United States, May 17-July 22, 2022." *Morbidity and Mortality Weekly Report* 71

(32): 1018–22.

R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R
Foundation for Statistical Computing; 2017.

Ridout, Martin, Clarice G B Demetrio, and John Hinde. 1998. "Models for Count Data with
Many Zeros." In *Proceedings of the XIXth International Biometric Conference*, 19:179–
92. International Biometric Society Invited Papers. Cape Town, South Africa.

Rimoin, Anne W., Prime M. Mulembakani, Sara C. Johnston, James O. Lloyd Smith, Neville K.
Kisalu, Timothee L. Kinkela, Seth Blumberg, et al. 2010. "Major Increase in Human
Monkeypox Incidence 30 Years after Smallpox Vaccination Campaigns Cease in the
Democratic Republic of Congo." *Proceedings of the National Academy of Sciences of the
United States of America* 107 (37): 16262–67. https://doi.org/10.1073/pnas.1005769107.

Rodríguez, Berta Suárez, Bernardo R Guzmán Herrador, Asunción Díaz Franco, María Paz
Sánchez-seco Fariñas, Amo Valero, Adrián Hugo, Aginagalde Llorente, et al. 2022.
"Epidemiologic Features and Control Measures during Monkeypox" 28 (9): 1847–51.

Schneckenreither, Günter, Lukas Herrmann, Rafael Reisenhofer, Nikolas Popper, and Philipp
Grohs. 2022. "Assessing the Heterogeneity in the Transmission of Infectious Diseases
from Time Series of Epidemiological Data." *Na* NA (NA): NA-NA.

Sklenovská, Nikola, and Marc Van Ranst. 2018. "Emergence of Monkeypox as the Most
Important Orthopoxvirus Infection in Humans." *Frontiers in Public Health* 6
(September): 1–12. https://doi.org/10.3389/fpubh.2018.00241.

Spicknall, Ian H., Emily D. Pollock, Patrick A. Clay, Alexandra M. Oster, Kelly Charniga, Nina
Masters, Yoshinori J. Nakazawa, Gabriel Rainisch, Adi V. Gundlapalli, and Thomas L.
Gift. 2022. "Modeling the Impact of Sexual Networks in the Transmission of Monkeypox
Virus Among Gay, Bisexual, and Other Men Who Have Sex with Men — United States,
2022." *MMWR. Morbidity and Mortality Weekly Report* 71 (35): 1131–35.
https://doi.org/10.15585/mmwr.mm7135e2.

The Lancet Regional Health – Europe. 2022. "Lessons from COVID-19 Are Shaping the
Response to Monkeypox Outbreak." *The Lancet Regional Health - Europe* 18: 100463.
https://doi.org/10.1016/j.lanepe.2022.100463.

Tkachenko, Alexei V., Sergei Maslov, Ahmed Elbanna, George N. Wong, Zachary J. Weiner,
and Nigel Goldenfeld. 2021. "Time-Dependent Heterogeneity Leads to Transient
Suppression of the COVID-19 Epidemic, Not Herd Immunity." *Proceedings of the
National Academy of Sciences* 118 (17): e2015972118.
https://doi.org/10.1073/pnas.2015972118.

Toth, Damon J.A., Molly Leecaster, Warren B.P. Pettey, Adi V. Gundlapalli, Hongjiang Gao,
Jeanette J. Rainey, Amra Uzicanin, and Matthew H. Samore. 2015. "The Role of
Heterogeneity in Contact Timing and Duration in Network Models of Influenza Spread in
Schools." *Journal of the Royal Society Interface* 12 (108).
https://doi.org/10.1098/rsif.2015.0279.

Toth, Damon J.A., Windy D. Tanner, Karim Khader, and Adi V. Gundlapalli. 2016. "Estimates
of the Risk of Large or Long-Lasting Outbreaks of Middle East Respiratory Syndrome
after Importations Outside the Arabian Peninsula." *Epidemics* 16: 27–32.
https://doi.org/10.1016/j.epidem.2016.04.002.

Tsang, Tim K., Can Wang, Bingyi Yang, Simon Cauchemez, and Benjamin J. Cowling. 2021.
"Using Secondary Cases to Characterize the Severity of an Emerging or Re-Emerging
Infection." *Nature Communications* 12 (1): 6372. https://doi.org/10.1038/s41467-021-

26709-7.

UK Health Security Agency. 2022. "Investigation into Monkeypox Outbreak in England:
Technical Briefing 1." https://www.gov.uk/government/publications/monkeypox-
outbreak-technical-briefings/investigation-into-monkeypox-outbreak-in-england-
technical-briefing-1.

Wallinga, J., and P. Teunis. 2004. "Different Epidemic Curves for Severe Acute Respiratory
Syndrome Reveal Similar Impacts of Control Measures." *American Journal of*
Epidemiology 160 (6): 509–16. https://doi.org/10.1093/aje/kwh255.

World Health Organization. 2022a. "No Title." 2022 Mpox Outbreak: Global Trends. 2022.
https://worldhealthorg.shinyapps.io/mpx_global/.

———. 2022b. "WHO Recommends New Name for Monkeypox Disease." 2022.
https://www.who.int/news/item/28-11-2022-who-recommends-new-name-for-
monkeypox-disease.

______. 2022c. "Monkeypox." 2022 Fact Sheets. https://www.who.int/news-room/fact-
sheets/detail/monkeypox

Yinka-Ogunleye, Adesola, Olusola Aruna, Dimie Ogoina, Neni Aworabhi, Womi Eteng, Sikiru
Badaru, Amina Mohammed, et al. 2018. "Reemergence of Human Monkeypox in
Nigeria, 2017." *Emerging Infectious Diseases* 24 (6): 1149–51.
https://doi.org/10.3201/eid2406.180017.

Zhang, Y, T Britton, and X Zhou. 2022. "Monitoring Real-Time Transmission Heterogeneity
from Incidence Data," *PLoS Comput Biol 18(12): e1010078.*
https://doi.org/10.1371/journal.pcbi.1010078

664