Characterizing spatiotemporal variation in transmission heterogeneity

during the 2022 mpox outbreak in the USA

Jay Love[+][*][1], Cormac R. LaPrete[*][1], Theresa R. Sheets[1], George G. Vega Yon[1][#], Alun Thomas[1],

Matthew H. Samore[1], Lindsay T. Keegan[1], Frederick R. Adler[1], Rachel B. Slayton[2], Ian H.

Spicknall[2], Damon J.A. Toth[1]

[+]Corresponding Author

[*]Contributed equally

[1]University of Utah, Salt Lake City, Utah

[2]Centers for Disease Control and Prevention, Atlanta, Georgia

[#]Current affiliation: Booz Allen Hamilton, Inc.

<u>Keywords</u>

transmission heterogeneity, superspreading, mpox, monkeypox virus, MPXV

<u>Running Title</u>

Transmission heterogeneity of mpox, 2022, US

<u>Biographical Sketch</u>

Jay Love, Ph.D., conducts academic research in Infectious Disease Epidemiology and Biology.

He is currently a Research Associate in the Division of Epidemiology at the University of Utah

and a Guest Researcher at the Centers for Disease Control and Prevention.

Abstract

Understanding how transmission heterogeneity varies over the course of an enduring

infectious disease outbreak improves understanding of observed disease dynamics and informs

public health strategy. We quantified the spatiotemporal variation in transmission heterogeneity

for the 2022 mpox outbreak in the US using the dispersion parameter of the offspring

distribution, k. Our methods fit negative binomial distributions to transmission chain offspring

distributions informed by a large mpox contact tracing dataset. We found that estimates of

transmission heterogeneity varied across the outbreak, but overall estimated transmission

heterogeneity was low. When testing our methods on simulated data, estimate accuracy

depended on contact tracing data accuracy and completeness. Because the actual contact tracing

data had high incompleteness, the values of k estimated from the empirical data may therefore be

artificially high. Through simulation, we explore a method to correct estimated k for data

incompleteness and, further, explore baseline expectations for temporal dynamics of k.

Introduction

In an infectious disease outbreak, transmission heterogeneity occurs when there is

variation in the number of secondary infections per infected host, which can be caused by

underlying mechanistic heterogeneity. This variation can manifest at spatial (May and Anderson

1984), temporal (Bjørnstad, Finkenstädt, and Grenfell 2002), and/or individual (Lloyd-Smith et

al. 2005; Ke et al. 2022) scales. These different components of transmission heterogeneity

interact with one another, creating the potential for complex patterns of transmission

heterogeneity over the course of an outbreak. Because transmission heterogeneity plays a critical

role in epidemic spread, characterizing how it changes over an outbreak can help to illuminate

the causes behind observed epidemic dynamics and inform measures taken by public health

decision makers, such as the utility of targeting potentially high-risk individuals or locations for

transmission mitigation (Shen et al. 2023; Nielsen et al. 2023).

Prior methods have simultaneously estimated the mean transmission (R) and the

dispersion (k) parameters from data on the number of secondary infections per primary infection

(*i.e.*, the offspring distribution) in an outbreak (Lloyd-Smith et al. 2005; Lloyd-Smith 2007;

Blumberg and Lloyd-Smith 2013; Toth et al. 2015). These methods have been applied both

retrospectively and during ongoing outbreaks (Toth et al. 2016; Kucharski and Althaus 2015;

Nishiura et al. 2015) for several diseases. Past research on transmission heterogeneity has largely

focused on small outbreaks or has used data from early transmission generations in larger

outbreaks (Lloyd-Smith et al. 2005; Lloyd-Smith 2007; Blumberg and Lloyd-Smith 2013; Toth

et al. 2015; 2016; Kucharski and Althaus 2015; Nishiura et al. 2015). Some research has

explored how the first transmission generation may differ from subsequent generations in R and

k (Toth et al. 2016), but work to understand how individual transmission heterogeneity may vary

over the component outbreaks of a large epidemic is more rare (Lau et al. 2020; Tkachenko et al.

2021; Miyama, Jung, and Nishiura 2022).

Among those diseases for which transmission heterogeneity has been characterized in the

past is mpox (formerly 'Monkeypox' (World Health Organization 2022b)). Monkeypox virus

(MPXV), the causative pathogen of mpox disease, is a member of the *Orthopoxvirus* genus of

pathogens and is endemic in West and Central Africa. Mpox was first detected in humans in

1970, shortly after the eradication of smallpox (Breman et al. 1980), and has subsequently

caused occasional outbreaks thought to result from repeated spillover events (Nalca et al. 2005;

Yinka-Ogunleye et al. 2018; Bunge et al. 2022; Durski et al. 2018). While the exact reservoir

host is still unknown, the virus has been found in several rodent species native to central and

west African countries (Durski et al. 2018). Mpox is known to spread through direct contact with

mpox rash, scabs, or bodily fluids; touching objects that have been used by someone with mpox;

contact with respiratory secretions; and sexual or intimate contact with an infected person

(Centers for Disease Control and Prevention 2022).

Regional outbreaks have been growing in recent years (Sklenovská and Van Ranst 2018;

Beer and Bhargavi Rao 2019; Rimoin et al. 2010). Past work to define the basic reproductive

number (R_0, the expected number of transmissions from an average infected person in a large,

fully susceptible population), for mpox outbreaks have produced estimates below 1, the critical

threshold above which an infectious disease will continue to spread and below which an outbreak

will eventually die out. However, more recent work has accounted for declining cross-protective

immunity from smallpox to reach an estimate of $R_0 = 2.13$ (uncertainty bounds: 1.46-2.67) for

mpox in the Democratic Republic of Congo (Grant, Nguyen, and Breban 2020). Past estimates of

transmission heterogeneity, quantified by the inversely varying dispersion parameter (low values

indicate high transmission heterogeneity) of the offspring distribution (k), for mpox outbreaks

have been 0.58 (90% confidence interval [CI]: 0.32-3.57) (Lloyd-Smith et al. 2005), 0.36 (0.16-

1.47), and 0.33 (0.19-0.64) (Blumberg and Lloyd-Smith 2013), indicating high transmission

heterogeneity.

In April 2022, MPXV was identified outside of the countries where the virus is endemic

and quickly spread to over 100 countries (World Health Organization 2022a). The World Health

Organization (WHO) declared this mpox epidemic a Public Health Emergency of International

Concern on July 23, 2022. This rapid global spread was not characteristic of an infectious

disease with a basic reproductive number below 1. Contact tracing conducted during the 2022

outbreak indicated that the disease was primarily spreading by sexual contact between gay,

bisexual, and other men who have sex with men, often at large gatherings (Philpott et al. 2022).

In light of this, public health messaging focused on reducing spread among these groups,

especially at large gatherings (Delaney et al. 2022). Potentially due to a novel sexually mediated

dominant transmission route, the discrepancy between estimates of the basic reproductive

number for mpox outbreaks became globally relevant. Estimates of the 2022 outbreak vary by

region and range from 1.56 to 3.8 (Branda, Pierini, and Mazzoli 2022) or potentially higher

(Endo et al. 2022).

Because the degree distribution of sexual contact networks can be overdispersed,

indicating high variability in the number of contacts per individual, transmission through sexual

contact networks can result in heterogeneous patterns of secondary transmission across an

infected population (Handcock and Jones 2006). Spreading of the disease through large

gatherings, termed 'superspreader events', can also cause high levels of transmission

heterogeneity (Althouse et al. 2020). Such heterogeneous transmission has been shown to

influence outbreak dynamics (Toth et al. 2015; Elie, Selinger, and Alizon 2022) and therefore

may have contributed substantially to the rapid global spread of mpox. Modeling studies have

provided support for the idea that heterogeneous transmission through contact networks could

partially explain the dynamics of the 2022 mpox outbreak (Endo et al. 2022; Spicknall et al.

2022).

Here, we estimate the mean transmission and transmission heterogeneity for the 2022

mpox epidemic in the United States using contact tracing data fortified with algorithmically

determined probable case pairs. We estimate these parameters directly from the empirical data

and, through simulation, assess the sensitivity of our approach relative to data incompleteness

and explore opportunities to correct estimates influenced by data incompleteness. We stratify our

analysis by geographic location and time. In this way, we characterize transmission

heterogeneity on spatial, temporal, and individual scales. Our aims are to (1) better understand

the dynamics of the 2022 mpox outbreak and (2) advance the theory of transmission

heterogeneity.

<u>Methods</u>

Overview

We used mpox case detection and contact tracing data to algorithmically generate

offspring distributions, the number of secondary infections from each case, for given time

periods and locations in the United States using methods described in more detail below. Then

we used a maximum likelihood approach to fit negative binomial models to those offspring

distributions. The negative binomial model fittings provide estimates for the mean R and

dispersion parameter k, with lower values of k indicating greater transmission heterogeneity. We

also tested our method using synthetic transmission data generated from contact networks known

to produce high and low levels of transmission heterogeneity.

Data

We used data collected through U.S. Centers for Disease Control and Prevention (CDC)

mpox case detection and contact tracing in the United States from May 1st to November 7th,

2022. This dataset includes, for each probable and confirmed case, information on date of

symptom onset, date of reporting, date of positive test, reporting jurisdiction, any recent travel

history, and any known contacts with other probable or confirmed cases.

Algorithmic generation of offspring distributions

To compile an offspring distribution associated with a particular subset of the data, we

applied an algorithm to incorporate probable transmission events and their associated infected

individuals with unobserved probable transmission events, relying on published mpox serial

interval estimates and the dates of symptom onset for infected individuals in the dataset (Figure

1). These transmission trees were constructed using methods similar to those in previous works

(e.g., Haydon et al. 2003; Wallinga and Teunis 2004; Heijne et al. 2012), though our method

differs in notable ways (see supplemental methods).

Parameter estimation

To fit a negative binomial distribution to the offspring distribution, we used a maximum

likelihood approach, following established methods (Lloyd-Smith et al. 2005; Toth et al. 2016).

This approach is sensitive to the number of individuals with mpox assumed to transmit to no

other individuals, which we identify as terminal cases. Because a portion of the individuals in

our dataset with no reported offspring may have actually transmitted the disease to others

without being detected, including these false terminal cases would artificially inflate the number

of zeros in the offspring distribution. To test the sensitivity of our methods to this type of

potential bias, we repeated our methods using the zero-inflated negative binomial (see

supplemental methods), complementing past work that accounts for underreporting of terminal

cases (Lloyd-Smith 2007).

In the negative binomial estimates, the logarithms of negative binomially distributed

probabilities of each data point are a function of R and k, while in the zero-inflated estimates the

logarithms of zero-inflated negative binomially distributed probabilities of each data point are a

function of R, k, and p_0. We used maximum likelihood estimation by taking the log-likelihood

function of the entire data set. We then used multivariate optimization to find R and k for the

negative binomial estimates and R, k, and p_0 for the zero-inflated estimates. We used the

likelihood ratio test to find 95% confidence intervals for these parameters.

Sliding and telescoping approaches

To evaluate change in parameter estimates over time, we conducted our fitting methods

on offspring distributions generated across both sliding (fixed width) and telescoping (varying

width) windows of time (see supplemental methods).

Simulations

To characterize the reliability of our methods to estimate transmission heterogeneity, we

repeated them on simulated data generated with a model of infectious disease spread over a static

contact network. We conducted simulations on undirected scale-free networks, shown to produce

heterogeneous transmission (Bansal, Grenfell, and Meyers 2007). We initialized the scale-free

network with a Barabási-Albert algorithm of preferential attachment (Barabási and Albert 1999),

allowing for the network to serve as a proxy for a population with highly heterogeneous contact

rates where simulated outbreaks can be expected to result in an overdispersed offspring

distribution. The network was generated using the *barabasi.game* function from the *igraph* R

package (Csardi and Nepusz 2006), using a population of 50,000 and power of 1.

Each outbreak was seeded with a randomly chosen single infection. Disease dynamics

were modeled following a *Susceptible – Exposed – Infectious – Recovered* paradigm,

parameterized for mpox. The latent period was a random variable drawn from a lognormal

distribution with log mean = 1.5 and log standard deviation = 0.7 (Madewell et al. 2022). For

lack of detailed information on the duration of infectiousness for mpox, the infectious period in

our simulations was a random variable drawn from a gamma distribution with shape = 21 days

and scale = 1. This distribution is consistent with the reported serial interval distribution

(Madewell et al. 2022), clinical guidance indicating symptoms last 2-4 weeks (Word Health

Organization 2022c), and investigations reporting that some hospitalized cases show symptoms

for much longer periods (Adler et al. 2022). However, it should be noted that more detailed study

may reveal that the distribution we used may not describe the infectious period for mpox.

Contacts of infectious nodes were infected with probability of 0.15 per contact day.

In order to replicate our methods on the simulated data, we assigned each node a

"jurisdiction" using a modularity optimization algorithm that identifies community structure in

networks (Blondel et al. 2008), as depicted in Figure 2. We then identified the most probable

infector for each case, as detailed above, and estimated the parameters R and k by fitting a

negative binomial distribution, as detailed above.

In order to test the sensitivity of our methods to incompleteness in detailed contact

information, we repeated the methods with different levels of incompleteness. Proportions of

data completeness (f) ranged from 0 to 1, and we explored two types of incompleteness. For the

first type of incompleteness ("missing nodes"), individuals for which detailed contact

information was completely excluded were randomly sampled from the simulated population

with equal probability. For the second type of incompleteness ("missing edges"), connections

between individuals to be excluded from the dataset were randomly sampled with equal

probability. Methods were repeated on each level of completeness, and the resulting k estimates

at the end of the outbreak using a telescoping window were compared. The relationship between

f and k was described by fitting linear and exponential model to the data (log transforming the

response variable for the exponential model) using the 'lm' function and goodness-of-fit was

assessed with the Akaike Information Criterion using the 'AIC' function in R statistical software

(R Core Team 2017).

To evaluate the effectiveness of our method in estimating k from data on outbreaks where

high transmission heterogeneity is not expected, we simulated an outbreak over a version of the

contact network used in the first simulation modified by connecting all nodes within jurisdiction

clusters with edges and maintaining the existing cross-jurisdiction edges. The modified network

therefore has very high connectivity within jurisdiction clusters (i.e., jurisdiction subgraphs are

complete, representing very well mixed local populations), but connectivity across jurisdiction

clusters is equal to that of the unmodified network. We repeated our methods as detailed above,

estimating parameters R and k across different levels of incompleteness in detailed contact

information.

All computational methods were conducted in R statistical software (R Core Team 2017).

<u>Results</u>

National estimates

There were 22,391 probable and confirmed mpox cases in the empirical dataset, after

excluding 5,074 (18.47%) cases with missing data on symptom onset date. In the exclusive

dataset, 820 (3.66%) of these cases had contact tracing-identified connections to other probable

and confirmed cases, compared to 12 (0.24%) of those excluded cases.

R and k estimated at the national level changed over time. Using a sliding 2-week

window (Fig. 3a), we observe that, after early increase to a peak, estimated R declines over the

course of the epidemic, dropping below 1 around the peak of new infections on approximately

August 1st, as is expected to occur when an outbreak changes from growth to decline. Median

estimated k remains above 1 for all time windows, though with high variability – a pattern which

does not indicate high transmission heterogeneity.

Applying a telescoping window approach allows us to evaluate R and k taken

cumulatively at different points during the epidemic (Fig 3b). Using the time window beginning

at the start of the outbreak, May 1[st], and ending at the peak of new cases, August 1[st], we find that

$R = 1.25$ (95% CI: 1.21-1.28) and $k = 12.18$ (95% CI: 8.18-21.27). Using a zero-inflated negative

binomial distribution returned the same values of R and k for this period, with a p_0 value of 0.001

(95% CI: 0.00-0.02).

Jurisdiction estimates

Estimates of R varied across U.S. State and over time (Fig. 4). Estimates of k also vary

across public health jurisdictions and over time (Table S1). While median estimates of k are over

1 at all time points for every jurisdiction, the variation in k estimates across iterations is

substantial for some points, indicating a greater potential for higher transmission heterogeneity at

that time and place. Estimates of p_0 made using the zero-inflated negative binomial were highly

variable across jurisdictions and timepoints (see supplemental results).

Simulations

Conducting the estimation methods on data resulting from an outbreak simulation over a

scale-free contact network and with a gamma-distributed infectious period indicated

heterogeneous transmission, as anticipated, with k estimates well below 1 with complete data

(Fig 3e,f; Fig 5a-d). Time-varying k estimates acquired through the use of a sliding window were

low at the start and increased over the course of the outbreak. Conducting the methods on data

from an outbreak over the modified network showed that transmission was less heterogeneous

overall, but time-varying k estimates showed pulses of values less than 1, indicating temporary

periods of high transmission heterogeneity after which heterogeneity declines (Fig 5e,f).

The method showed reliability in detecting overdispersed offspring distributions when

detailed contact information was abundant, reflected by estimates of k below 1 when f was high

(Fig. 3e,f). However, under the missing node paradigm with $f = 0.036$ (i.e., when data

missingness was equivalent to the estimated missingness in the empirical dataset), estimates of k

were much higher, staying above 1 (Fig. 3c,d). By varying the proportion of the simulated

population for which detailed contact information was used, we explored the relationship

between k estimate accuracy and data completeness (f). This relationship is approximately linear

(AIC: –19.3 versus –4.59 for best-fit exponential model) and can be described by the equation

$$\left(\frac{1}{k}\right)(f) = 2.33\,f + 0.150$$

271   where $1/k$ is the value of $1/k$ at the end of the simulation using a telescoping window, and f is the

272   fraction of data that is complete (Fig. 6a). Under the missing edge paradigm with $f = 0.036$,

273   estimates of k were also above 1 (Fig. 6c). Varying f under this paradigm, we recover an

274   exponential relationship (AIC = –39.2 versus 2.23 for best-fit linear model) between k and f as:

$$\left(\frac{1}{k}\right)(f) = 0.248\ (11.2^f)$$

When the methods to estimate k were conducted on data from an outbreak simulated over

a modified contact network with high within-jurisdiction connectivity (see methods), estimates

of k were around 1 (Fig. 5e). $k = 1$ is a special condition of the negative binomial: the geometric

distribution. A geometric offspring distribution is expected under the assumption of constant

infectiousness and a constant recovery rate across infected individuals in a well-mixed

population (Grassly and Fraser 2008). At times, estimates of k drop well below 1 (observed as

spikes in $1/k$ in Fig. 5e). These instances may occur simultaneously with pathogen invasion of a

previously uninfected jurisdiction cluster, and the transient jump in heterogeneity may reflect a

sudden increase in the effective susceptible population followed quickly by a sharp decline in the

same. Estimates of k acquired with a telescoping window approach tended towards 0.83 by the

end of the outbreak (Fig. 5f), reflecting the cumulative influence of these temporary spikes in

transmission heterogeneity over the course of the outbreak. In contrast to the results from the

scale-free network-based simulation (Fig. 5a-d), k estimates are very near their true values

regardless of the degree of data incompleteness (Fig. 5e,f). Fitting a linear model produces

equation

$$\left(\frac{1}{k}\right)(f) = 0.0369\ f + 1.17$$

with the small slope indicating constancy (AIC = –82.2 versus –77.9 for exponential).

Correcting empirical estimates to account for data incompleteness

Ideally, the empirical dataset in this study would be sufficient to derive a correction

function, as with the simulated data, above. However, with only 3.6% completeness and

substantial noise (discussed below), a model fit to a set of k values estimated from datasets with

artificially increasing incompleteness may be unreliable. To explore possible values for corrected

k estimates, we proceeded to use a correction factor derived from simulated data on our estimates

from empirical data.

299   Under the assumption that the 2022 mpox outbreak in the United States occurred over a

contact network that was roughly scale-free (i.e., where the degree distribution follows a power

law) (Murayama et al. 2023), the function derived from our simulation results can be used to

correct our estimates from empirical data to account for data incompleteness. Doing so yields

much lower estimates of k. Using the slope of the linear model from the high heterogeneity

simulations under the missing node paradigm, we can use our data point ($f = 0.036$, $k = 7.06$) to

find equation

$$\left(\frac{1}{k}\right)(f) = 2.33\,f + 0.056$$

Therefore, when $f = 1$ (i.e., under complete detailed contact information), $k = 0.418$. This value

of k indicates high heterogeneity and is in line with past estimates of k for mpox (Lloyd-Smith et

al. 2005; Blumberg and Lloyd-Smith 2013). The validity of this corrected estimate is highly

contingent upon the assumption that the contact network properties of the real mpox outbreak are

similar to the simulated network. While there is some evidence that this could be the case

(Murayama et al. 2023), care must be taken with any applied interpretation. If the contact

network of the real outbreak was, instead, evenly mixing within jurisdictions (as in simulations

over a modified contact network, Fig. 5e,f), correcting for incompleteness would impact k

estimates far less, and the corrected k estimate may not indicate high heterogeneity.

Discussion

Our estimates of transmission heterogeneity over the course of the 2022 mpox outbreak

in the United States were low, but these estimates may be biased due to data incompleteness. If

taken at face value, our estimates of k garnered from direct analysis of the real data suggest that

transmission of mpox during the 2022 outbreak in the United States was not highly

heterogeneous; estimates of $k > 1$ across many jurisdictions and time periods would seem to

indicate that transmission was approximately what would be expected in an evenly mixing

population of people with equal inherent transmissibility, if not lower. We do, however, find

some indication of overdispersed transmission in certain locations during certain time periods

(Table S1), which could be consistent with explanations that invoke individual level transmission

heterogeneity.

However, if the context of data incompleteness is taken into account and it is assumed

that contact network structure was roughly scale-free, corrected estimates of k could, under

certain conditions of data missingness, indicate that transmission of mpox was highly

heterogeneous. This second possible conclusion hinges heavily upon the assumption that the

contact network of the real mpox outbreak was roughly scale-free. If it is not, then the function

to correct k based on data missingness would have a low slope, the effect of data incompleteness

on k estimates may be much smaller (as in fig 5e,f), and therefore the empirical estimate could be

seen as accurate.

The contact network structure and the data quality in our empirical dataset likely varied

by geographic location and through time. Therefore, we cannot rule out the possibility that our

estimates of k, corrected or otherwise, for the 2022 US mpox epidemic are artificially high in

some cases while artificially low in others, and care must be taken with any interpretation. Still,

our procedure to correct for data incompleteness under the assumption of a proposed contact

network structure leads our k estimates to be more in line with past estimates of transmission

heterogeneity for mpox (e.g., Lloyd-Smith et al. 2005; Blumberg and Lloyd-Smith 2013).

Our study contributes to a small but growing literature on spatiotemporal variation in

transmission heterogeneity of human infectious diseases. Past research on transmission

heterogeneity has largely focused on small outbreaks or has used data from the first generations

of infections in larger outbreaks. Related research on the influence of contact networks on the

spread of infectious diseases has illuminated important concepts regarding transmission

heterogeneity. For example, as the variation in the degree distribution of the infected population

changes over the course of an epidemic simulated on a network with a particular underlying

structure (Bansal, Grenfell, and Meyers 2007), transmission heterogeneity should be expected to

change over time. Our study provides a description of the baseline expected temporal change in

transmission heterogeneity for outbreaks on scale-free contact networks with non-constant

infectious period: generally, transmission heterogeneity reaches an early peak and then declines

over the course of an outbreak.

These methods are adaptable to other infectious disease outbreaks. They could provide

utility in informing public health decision makers about time-varying transmission heterogeneity,

with a lag determined largely by the time from infection to reporting. The method we present to

correct for incompleteness is robust to variation in transmission mode and contact network

structure across outbreaks, given contact information for a sufficiently large proportion of the

population. The degree of incompleteness in the empirical dataset in the present study did not

reach a threshold that allowed us to derive a reliable correction function from that data, so we

used our simulated data, parameterized for mpox, as a proxy to derive a correction function to

apply to the empirical data. Ultimately, this data limitation reduces our confidence in our

corrected estimate, and care must be taken with interpretation. Future work to apply the method

we present here to ongoing infectious disease outbreaks will require solutions to these data

limitations.

<u>Acknowledgements</u>

The findings and conclusions in this report are those of the authors and do not necessarily

represent the views of the Centers for Disease Control and Prevention. JL, TRS, GGVY, AT,

MHS, LTK, and DJT acknowledge research support by the Centers for Disease Control and

Prevention (1U01CK000585; 75D30121F00003)

<u>Literature Cited</u>

Adler H, Gould S, Hine P, Snell LB, Wong W, Houlihan CF, et al. 2022 "Clinical features and

management of human monkeypox: a retrospective observational study in the UK." *The*

Lancet Infectious Diseases. 22(8):1153–62.

Americo JL, Earl PL, Moss B. 2023 "Virulence differences of mpox (monkeypox) virus clades I,

IIa, and IIb.1 in a small animal model." *Proc Natl Acad Sci USA*.

21;120(8):e2220415120.

Althouse, Benjamin M., Edward A. Wenger, Joel C. Miller, Samuel V. Scarpino, Antoine Allard,

Laurent Hebert-Dufresne, and Hao Hu. 2020. "Superspreading Events in the

Transmission Dynamics of SARS-CoV-2: Opportunities for Interventions and Control."

PLoS Biology 18 (11): 1–13. https://doi.org/10.1371/journal.pbio.3000897.

Bansal, Shweta, Bryan T. Grenfell, and Lauren Ancel Meyers. 2007. "When Individual

Behaviour Matters: Homogeneous and Network Models in Epidemiology." *Journal of the*

Royal Society Interface 4 (16): 879–91. https://doi.org/10.1098/rsif.2007.1100.

Barabási, Albert-László, and Réka Albert. 1999. "Emergence of Scaling in Random Networks."

Science 286 (5439): 509–12. https://doi.org/10.1126/science.286.5439.509.

Beer, Ellen M., and V. Bhargavi Rao. 2019. "A Systematic Review of the Epidemiology of

Human Monkeypox Outbreaks and Implications for Outbreak Strategy." *PLoS Neglected*

Tropical Diseases 13 (10): 1–20. https://doi.org/10.1371/journal.pntd.0007791.

Bjørnstad, Ottar N., Bärbel F. Finkenstädt, and Bryan T. Grenfell. 2002. "Dynamics of Measles

Epidemics: Estimating Scaling of Transmission Rates Using a Time Series SIR Model."

Ecological Monographs 72 (2): 169–84. https://doi.org/10.1890/0012-

9615(2002)072[0169:DOMEES]2.0.CO;2.

Blondel, Vincent D, Jean-Loup Guillaume, Renaud Lambiotte, and Etienne Lefebvre. 2008.

"Fast Unfolding of Communities in Large Networks." *Journal of Statistical Mechanics:*

Theory and Experiment 2008 (10): P10008. https://doi.org/10.1088/1742-

5468/2008/10/P10008.

Blumberg, Seth, and J. O. Lloyd-Smith. 2013. "Inference of R0 and Transmission Heterogeneity

from the Size Distribution of Stuttering Chains." *PLoS Computational Biology* 9 (5): 1–

17. https://doi.org/10.1371/journal.pcbi.1002993.

Branda, Francesco, Massimo Pierini, and Sandra Mazzoli. 2022. "Monkeypox: Early Estimation

of Basic Reproduction Number R 0 in Europe." *Journal of Medical Virology*.

https://doi.org/10.1002/jmv.28270.

Breman, J. G., K. Ruti, M. V. Steniowski, E. Zanotto, A.I. Gromyko, and I. Arita. 1980. "Human

Monkeypox, 1970-79." *Bulletin of the World Health Organization* 58 (2): 165–82.

Bunge, Eveline M., Bernard Hoet, Liddy Chen, Florian Lienert, Heinz Weidenthaler, Lorraine R.

Baer, and Robert Steffen. 2022. "The Changing Epidemiology of Human Monkeypox—

A Potential Threat? A Systematic Review." *PLoS Neglected Tropical Diseases* 16 (2): 1–

20. https://doi.org/10.1371/journal.pntd.0010141.

Centers for Disease Control and Prevention. 2022. "CDC Monkeypox Response: Transmission -

Media Statement," 1–2.

______. 2022. "Case Definitions for Use in the 2022 Mpox Response"

https://www.cdc.gov/poxvirus/mpox/clinicians/case-definition.html

Csardi G, Nepusz T. 2006. "The igraph software package for complex network research."

InterJournal, Complex Systems, 1695. https://igraph.org.

Delaney, Kevin P., Travis Sanchez, Marissa Hannah, O. Winslow Edwards, Thomas Carpino,

Christine Agnew-Brune, Kaytlin Renfro, et al. 2022. "Strategies Adopted by Gay,

Bisexual, and Other Men Who Have Sex with Men to Prevent Monkeypox Virus

Transmission — United States, August 2022." *MMWR. Morbidity and Mortality Weekly*

Report 71 (35): 1126–30. https://doi.org/10.15585/mmwr.mm7135e1.

Durski, Kara N, Andrea M Mccollum, Yoshinori Nakazawa, Brett W Petersen, and Mary G

Reynolds. 2018. "Emergence of Monkeypox in West Africa and Central Africa, 1970–

2017." *Releve Epidemiologique Hebdomadaire* 93 (11): 125–32.

Elie, Baptiste, Christian Selinger, and Samuel Alizon. 2022. "The Source of Individual

Heterogeneity Shapes Infectious Disease Outbreaks." *Proceedings of the Royal Society*

B: Biological Sciences 289 (1974). https://doi.org/10.1098/rspb.2022.0232.

Endo, Akira, Hiroaki Murayama, Sam Abbott, Ruwan Ratnayake, Carl A. B. Pearson, W. John

Edmunds, Elizabeth Fearon, and Sebastian Funk. 2022. "Heavy-Tailed Sexual Contact

Networks and the Epidemiology of Monkeypox Outbreak in Non-Endemic Regions, May

2022." *MedRxiv*, no. May: 2022.06.13.22276353.

Ennab, Farah, Faisal A. Nawaz, Kapil Narain, Goodluck Nchasi, and Mohammad Yasir Essar.

2022. "Rise of Monkeypox: Lessons from COVID-19 Pandemic to Mitigate Global

Health Crises." *Annals of Medicine and Surgery* 79 (June): 14–17.

https://doi.org/10.1016/j.amsu.2022.104049.

Grant, Rebecca, Liem Binh Luong Nguyen, and Romulus Breban. 2020. "Modelling Human-to-

Human Transmission of Monkeypox." *Bulletin of the World Health Organization* 98 (9):

638–40. https://doi.org/10.2471/BLT.19.242347.

Grassly NC, Fraser C. 2008. "Mathematical models of infectious disease transmission." *Nat Rev*

Microbiol 6(6):477–87.

Handcock, Mark S., and James Holland Jones. 2006. "Interval Estimates for Epidemic

Thresholds in Two-Sex Network Models." *Theoretical Population Biology* 70 (2): 125–

34. https://doi.org/10.1016/j.tpb.2006.02.004.

Haydon, D.T., M. Chase-Topping, D. J. Shaw, L. Matthews, J. K. Friar, J. Wilesmith and M. E.

J. Woolhouse. 2003. "The Construction and Analysis of Epidemic Trees with Reference

to the 2001 UK Footand-Mouth Outbreak." *Proceedings of the Royal Society: B*

270(1511): 121-127. https://doi.org/10.1098/rspb.2002.2191

Heijne, Jannek C.M., Marc Rondy, Linda Verhoef, Jacco Wallinga, Mirjam Kretzschmar, Nicola

Low, Marion Koopmans and Peter F. M. Teunis. 2012. "Quantifying Transmission of

Norovirus During an Outbreak." *Epidemiology* 23:277-284.

https://doi.org/10.1097/EDE.0b013e3182456ee6

Isidro J, Borges V, Pinto M, Sobral D, Santos JD, Nunes A, et al. 2022. "Phylogenomic

characterization and signs of microevolution in the 2022 multi-country outbreak of

monkeypox virus." *Nat Med*. 28(8):1569–72.

Ke, Ruian, Pamela P. Martinez, Rebecca L. Smith, Laura L. Gibson, Agha Mirza, Madison

Conte, Nicholas Gallagher, et al. 2022. "Daily Longitudinal Sampling of SARS-CoV-2

Infection Reveals Substantial Heterogeneity in Infectiousness." *Nature Microbiology* 7

(5): 640–52. https://doi.org/10.1038/s41564-022-01105-z.

Kucharski, A. J., and C. L. Althaus. 2015. "The Role of Superspreading in Middle East

Respiratory Syndrome Coronavirus (MERS-CoV) Transmission." *Eurosurveillance* 20

(25): 14–18. https://doi.org/10.2807/1560-7917.ES2015.20.25.21167.

Lau, Max S.Y., Bryan Grenfell, Michael Thomas, Michael Bryan, Kristin Nelson, and Ben

Lopman. 2020. "Characterizing Superspreading Events and Age-Specific Infectiousness

of SARS-CoV-2 Transmission in Georgia, USA." *Proceedings of the National Academy*

of Sciences of the United States of America 117 (36): 22430–35.

https://doi.org/10.1073/pnas.2011802117.

Lloyd-Smith, J. O. 2007. "Maximum Likelihood Estimation of the Negative Binomial

Dispersion Parameter for Highly Overdispersed Data, with Applications to Infectious

Diseases." *PLoS ONE* 2 (2): 1–8. https://doi.org/10.1371/journal.pone.0000180.

Lloyd-Smith, J. O., S. J. Schreiber, P. E. Kopp, and W. M. Getz. 2005. "Superspreading and the

Effect of Individual Variation on Disease Emergence." *Nature* 438 (7066): 355–59.

https://doi.org/10.1038/nature04153.

Madewell, Zachary J, Kelly Charniga, Nina Masters, and Et. Al. 2022. "Serial Interval and

Incubation Period Estimates of Monkeypox Virus Infection in 12 U.S. Jurisdictions, May

– August 2022." *MedRxiv*. https://doi.org/10.1101/2022.10.26.22281516.

May, Robert M., and Roy M. Anderson. 1984. "Spatial Heterogeneity and the Design of

Immunization Programs." *Mathematical Biosciences* 72 (1): 83–111.

https://doi.org/10.1016/0025-5564(84)90063-4.

Miyama T, Jung S, and Nishiura H. 2022. "Decrease in Overdispersed Secondary Transmission

of COVID-19 over Time in Japan." *Epidemiology and Infection*, November, 1–29.

https://doi.org/10.1017/S0950268822001789.

Murayama H, Pearson CAB, Abbott S, Miura F, Jung S, Fearon E, et al. 2023. "Accumulation of

Immunity in Heavy-Tailed Sexual Contact Networks Shapes Mpox Outbreak Sizes." *The*

Journal of Infectious Diseases 254. https://doi.org/10.1093/infdis/jiad254

Nalca, Aysegul, Anne W. Rimoin, Sina Bavari, and Chris A. Whitehouse. 2005. "Reemergence

of Monkeypox: Prevalence, Diagnostics, and Countermeasures." *Clinical Infectious*

Diseases 41 (12): 1765–71. https://doi.org/10.1086/498155.

Neilsen BF, Sneppen K, Simonsen L. 2023. "The counterintuitive implications of superspreading

diseases" *Nature Communications* (14): 6954. https://doi.org/10.1038/s41467-023-

42612-9

Nishiura, H., Y. Miyamatsu, G. Chowell, and M. Saitoh. 2015. "Assessing the Risk of Observing

Multiple Generations of Middle East Respiratory Syndrome (MERS) Cases given an

Imported Case." *Eurosurveillance* 20 (27): 6–11. https://doi.org/10.2807/1560-

7917.ES2015.20.27.21181.

Philpott, David, Christine M Hughes, Karen A Alroy, Janna L Kerins, Jessica Pavlick, and
Lenore Asbel. 2022. "Epidemiologic and Clinical Characteristics of Monkeypox Cases
— United States, May 17-July 22, 2022." *Morbidity and Mortality Weekly Report* 71
(32): 1018–22.

R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R
Foundation for Statistical Computing; 2017.

Ridout, Martin, Clarice G B Demetrio, and John Hinde. 1998. "Models for Count Data with
Many Zeros." In *Proceedings of the XIXth International Biometric Conference*, 19:179–
92. International Biometric Society Invited Papers. Cape Town, South Africa.

Rimoin, Anne W., Prime M. Mulembakani, Sara C. Johnston, James O. Lloyd Smith, Neville K.
Kisalu, Timothee L. Kinkela, Seth Blumberg, et al. 2010. "Major Increase in Human
Monkeypox Incidence 30 Years after Smallpox Vaccination Campaigns Cease in the
Democratic Republic of Congo." *Proceedings of the National Academy of Sciences of the
United States of America* 107 (37): 16262–67. https://doi.org/10.1073/pnas.1005769107.

Rodríguez, Berta Suárez, Bernardo R Guzmán Herrador, Asunción Díaz Franco, María Paz
Sánchez-seco Fariñas, Amo Valero, Adrián Hugo, Aginagalde Llorente, et al. 2022.
"Epidemiologic Features and Control Measures during Monkeypox" 28 (9): 1847–51.

Schneckenreither, Günter, Lukas Herrmann, Rafael Reisenhofer, Nikolas Popper, and Philipp
Grohs. 2022. "Assessing the Heterogeneity in the Transmission of Infectious Diseases
from Time Series of Epidemiological Data." *PLOS ONE* 18(5): e0286012.
https://doi.org/10.1371/journal.pone.0286012

Shen, Mingwang, Hao Lai, and Lei Zhang. 2023. "Targeted interventions for individuals at high

523   risk is essential for mpox control." *The Lancet Infectious Diseases* 24(1):8-9

524 Sklenovská, Nikola, and Marc Van Ranst. 2018. "Emergence of Monkeypox as the Most

525   Important Orthopoxvirus Infection in Humans." *Frontiers in Public Health* 6

526   (September): 1–12. https://doi.org/10.3389/fpubh.2018.00241.

527 Spicknall, Ian H., Emily D. Pollock, Patrick A. Clay, Alexandra M. Oster, Kelly Charniga, Nina

528   Masters, Yoshinori J. Nakazawa, Gabriel Rainisch, Adi V. Gundlapalli, and Thomas L.

529   Gift. 2022. "Modeling the Impact of Sexual Networks in the Transmission of Monkeypox

530   Virus Among Gay, Bisexual, and Other Men Who Have Sex with Men — United States,

531   2022." *MMWR. Morbidity and Mortality Weekly Report* 71(35): 1131–35.

532   https://doi.org/10.15585/mmwr.mm7135e2.

533 The Lancet Regional Health – Europe. 2022. "Lessons from COVID-19 Are Shaping the

534   Response to Monkeypox Outbreak." *The Lancet Regional Health - Europe* 18: 100463.

535   https://doi.org/10.1016/j.lanepe.2022.100463.

536 Tkachenko, Alexei V., Sergei Maslov, Ahmed Elbanna, George N. Wong, Zachary J. Weiner,

537   and Nigel Goldenfeld. 2021. "Time-Dependent Heterogeneity Leads to Transient

538   Suppression of the COVID-19 Epidemic, Not Herd Immunity." *Proceedings of the*

539   *National Academy of Sciences* 118 (17): e2015972118.

540   https://doi.org/10.1073/pnas.2015972118.

541 Toth, Damon J.A., Molly Leecaster, Warren B.P. Pettey, Adi V. Gundlapalli, Hongjiang Gao,

542   Jeanette J. Rainey, Amra Uzicanin, and Matthew H. Samore. 2015. "The Role of

543   Heterogeneity in Contact Timing and Duration in Network Models of Influenza Spread in

544   Schools." *Journal of the Royal Society Interface* 12 (108).

545   https://doi.org/10.1098/rsif.2015.0279.

Toth, Damon J.A., Windy D. Tanner, Karim Khader, and Adi V. Gundlapalli. 2016. "Estimates

of the Risk of Large or Long-Lasting Outbreaks of Middle East Respiratory Syndrome

after Importations Outside the Arabian Peninsula." *Epidemics* 16: 27–32.

https://doi.org/10.1016/j.epidem.2016.04.002.

Tsang, Tim K., Can Wang, Bingyi Yang, Simon Cauchemez, and Benjamin J. Cowling. 2021.

"Using Secondary Cases to Characterize the Severity of an Emerging or Re-Emerging

Infection." *Nature Communications* 12 (1): 6372. https://doi.org/10.1038/s41467-021-

26709-7.

UK Health Security Agency. 2022. "Investigation into Monkeypox Outbreak in England:

Technical Briefing 1." https://www.gov.uk/government/publications/monkeypox-

outbreak-technical-briefings/investigation-into-monkeypox-outbreak-in-england-

technical-briefing-1.

Wallinga, J., and P. Teunis. 2004. "Different Epidemic Curves for Severe Acute Respiratory

Syndrome Reveal Similar Impacts of Control Measures." *American Journal of*

Epidemiology 160 (6): 509–16. https://doi.org/10.1093/aje/kwh255.

World Health Organization. 2022a. "No Title." 2022 Mpox Outbreak: Global Trends. 2022.

https://worldhealthorg.shinyapps.io/mpx_global/.

———. 2022b. "WHO Recommends New Name for Monkeypox Disease." 2022.

https://www.who.int/news/item/28-11-2022-who-recommends-new-name-for-

monkeypox-disease.

______. 2022c. "Monkeypox." 2022 Fact Sheets. https://www.who.int/news-room/fact-

sheets/detail/monkeypox

Yinka-Ogunleye, Adesola, Olusola Aruna, Dimie Ogoina, Neni Aworabhi, Womi Eteng, Sikiru

Badaru, Amina Mohammed, et al. 2018. "Reemergence of Human Monkeypox in

Nigeria, 2017." *Emerging Infectious Diseases* 24 (6): 1149–51.

https://doi.org/10.3201/eid2406.180017.

Zhang, Y, T Britton, and X Zhou. 2022. "Monitoring Real-Time Transmission Heterogeneity

from Incidence Data," *PLoS Comput Biol 18(12): e1010078.*

https://doi.org/10.1371/journal.pcbi.1010078

FIGURE LEGENDS

Figure 1. A representation of data processing to build a complete transmission tree. (a) Pre-

processed case data points from two public health jurisdictions with four contact tracing

identified links. (b) Arrows added to show pools of possible infectors for each case, drawn based

on jurisdiction and time (each column of points represents one time step where, for example,

possible infectors come from only the most recent two time steps). (c) Chosen infectors sampled

and other possible infectors disregarded. This is the post-processing data set used for analysis.

Figure 2. Visualizations of the contact network used in simulations, with simulated jurisdictions

shown in colored polygons. See text for description of methods used to generate networks; (a)

shows two jurisdictions, (b) shows three jurisdictions, inclusive of the two in (a), and (c) shows

the full contact network. Overlapping polygons in (c) are visual artifacts of the large network –

each node is associated with only one jurisdiction.

Figure 3. Temporal variation in transmission parameters. Left column shows estimates made

with a sliding window, right column shows estimates made with a telescoping window, starting

on May 1st, 2022. Time points for the sliding window approach (left column) are the start of the

14-day sliding windows, time points for the telescoping window approach (right column) are the

end of the telescoping windows. Top row shows empirical parameter estimates, middle and

bottom rows show estimates made from outbreaks simulated on a scale-free contact network. In

the middle and bottom rows, the horizontal axis is units of days since the start of the outbreak

simulation. In all plots, black points represent values of $1/k$, plotted on the right vertical axes,

and blue points represent values of R, plotted on the left vertical axes. (a-b) Estimated median

and 95% CIs for $1/k$ (black, right-hand vertical axis) and R (blue, left-hand vertical axis),

displayed over daily new mpox cases (pink, peak of cases in empirical dataset in panels a,b is

506 on August 1st, 2022) for the 2022 mpox epidemic in the United States. (c-f) Parameter values

estimated on simulated data from a simulated outbreak which exhibited strongly heterogeneous

transmission. Peak of cases (pink) in simulated dataset is 410 on day 151 of the simulation. (c-d)

Estimates from simulated data where probable transmission pairs were determined with complete

information on true individual contacts for 3.6% of the population, with the remainder being

determined only with simulated dates and geographic locations. This approximates the condition

of the empirical data in a-b. (e-f) Estimates from simulated data where all pairs were determined

with complete information on true individual contacts. In all plots, values of $1/k$, instead of k, are

displayed to ease visualization.

Figure 4: Spatiotemporal variation in estimated values of R across US states during the middle of

the 2022 mpox epidemic. Unshaded regions indicate insufficient data to conduct parameter

estimations for that region and time period.

Figure 5: Estimates of k over time from a simulated outbreak over a scale-free (a-d) and a

modified contact network with high within-jurisdiction connectivity (e,f), using sliding (left

column) and telescoping (right column) analysis windows. Panels a-b show "missing node"

mode of missingness, panels c-d show "missing edge" mode.

Figure 6. Estimates of 1/k on highly heterogeneous outbreaks over a scale-free contact network

(a,c) and low heterogeneous outbreaks over a modified contact network (b,d), at the end of a

simulation using methods applied to various fractions of complete detailed contact information

plotted as black points, with red dotted line describing best-fit models (1/k)(f). Panels a-b show

"missing node" mode of missingness, panels c-d show "missing edge" mode.