Risk factors affecting polygenic score performance across diverse cohorts

Daniel Hui[1], Scott Dudek[1], Krzysztof Kiryluk[2], Theresa L. Walunas[3], Iftikhar J. Kullo[4], Wei-Qi Wei[5], Hemant K. Tiwari[6], Josh F. Peterson[5], Wendy K. Chung[7], Brittney Davis[8], Atlas Khan[2], Leah Kottyan[9], Nita A. Limdi[8], Qiping Feng[10], Megan J. Puckelwartz[11], Chunhua Weng[12], Johanna L. Smith[4], Elizabeth W. Karlson[13], Regeneron Genetics Center[14], Gail P. Jarvik[15], Marylyn D. Ritchie[1]

[1]Department of Genetics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA.
[2]Division of Nephrology, Department of Medicine, Columbia University, NY, New York.
[3]Department of Preventive Medicine, Northwestern University Feinberg School of Medicine, Chicago, Illinois, USA.
[4]Department of Cardiovascular Medicine, Mayo Clinic, Rochester, MN.
[5]Department of Biomedical Informatics, Vanderbilt University Medical Center, Nashville, TN.
[6]Department of Pediatrics, University of Alabama at Birmingham, Birmingham, AL.
[7]Departments of Pediatrics and Medicine, Columbia University Irving Medical Center, Columbia University, New York, NY.
[8]Department of Neurology, School of Medicine, University of Alabama at Birmingham, Birmingham, AL.
[9]The Center for Autoimmune Genomics and Etiology, Division of Human Genetics, Cincinnati Children's Hospital Medical Center, Cincinnati, OH.
[10]Division of Clinical Pharmacology, Department of Medicine, Vanderbilt University Medical Center, Nashville, TN, USA.
[11]Center for Genetic Medicine, Northwestern University Feinberg School of Medicine, Chicago, IL.
[12]Department of Biomedical Informatics, Vagelos College of Physicians & Surgeons, Columbia University, New York, NY.
[13]Division of Rheumatology, Inflammation, and Immunity, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA.
[14]Regeneron Pharmaceuticals Inc., Tarrytown, NY.
[15]Departments of Medicine (Medical Genetics) and Genome Sciences, University of Washington Medical Center, Seattle, WA.

Correspondence to:
Marylyn D. Ritchie, PhD
marylyn@pennmedicine.upenn.edu

Abstract

Apart from ancestry, personal or environmental covariates may contribute to differences in polygenic score (PGS) performance. We analyzed effects of covariate stratification and interaction on body mass index (BMI) PGS (PGS_{BMI}) across four cohorts of European (N=491,111) and African (N=21,612) ancestry. Stratifying on binary covariates and quintiles for continuous covariates, 18/62 covariates had significant and replicable R^2 differences among strata. Covariates with the largest differences included age, sex, blood lipids, physical activity, and alcohol consumption, with R^2 being nearly double between best and worst performing quintiles for certain covariates. 28 covariates had significant PGS_{BMI}-covariate interaction effects, modifying PGS_{BMI} effects by nearly 20% per standard deviation change. We observed overlap with covariates that had significant R^2 differences between strata and interaction effects – across all covariates, their main effects on BMI were correlated with maximum R^2 differences and interaction effects (0.56 and 0.58, respectively), suggesting high-PGS-score individuals have highest R^2 and PGS effect increases. Given significant and replicable evidence for context-specific PGS_{BMI} performance and effects, we investigated ways to increase model performance taking into account non-linear effects. Machine learning models (neural networks) increased relative model R^2 (mean 23%) across datasets. Finally, creating PGS_{BMI} directly from GxAge GWAS effects increased relative R^2 by 7.8%. These results demonstrate that certain covariates, especially those most associated with BMI, significantly affect both PGS_{BMI} performance and effects across diverse cohorts and ancestries, and we provide avenues to improve model performance that consider these effects.

Introduction

Polygenic scores (PGS) provide individualized genetic predictors of a phenotype by aggregating genetic effects across hundreds or thousands of loci, typically from genome-wide association studies (GWAS). In recent years it has become increasingly apparent that the transferability of PGS performance across different cohorts is poor (1). Most analyses to date have focused on ancestry differences as the main driver of this lack of transferability (2–4). However, a growing body of evidence has demonstrated that PGS performance and effect estimates are influenced by differences in certain environmental (classically termed "gene-environment" effects or interactions) or personal-level covariates – different phenotypes seem to be differently affected by these covariates, with most evidence being for adiposity traits such as body mass index (BMI) (5–14).

There are several gaps in current knowledge about these covariate-specific effects. Many analyses have assessed only a handful of these covariates due to the myriad of choices possible in typical large-scale biobanks. Little investigation has been done to systematically understand why certain covariates affect PGS performance, with such knowledge being useful to reduce the potential search for variables that impart context-specific effects. Furthermore, most studies investigating PGS-covariate interactions have been in European ancestry individuals; unfortunately, comparing differences in PGS performance and prediction while controlling for differences in ancestry versus differences in context has not been assessed in previous studies. Moreover, covariate-specific effects are notorious for having poor replicability in human genetics studies, and previous PGS-covariate interaction studies have been predominantly performed in the UK Biobank (UKBB) (15), where the majority of individuals are aged 40-69 (i.e., excluding young adults), are overall healthier than those from other e.g., hospital-based cohorts, and are predominantly European ancestry. Additionally, PGS performance is often assessed using linear models and in isolation of clinical covariates, which in practice would often be available. Machine learning models can have increased performance over linear models and are capable of modeling complex relationships and interactions between variables, which may serve to increase predictive performance, especially given evidence for PGS-covariate specific effects. Finally, given evidence for context-specific effects, it should be possible to directly incorporate SNP-covariate interaction effects from a GWAS directly to improve prediction performance, instead of relying on post-hoc interactions from a typical PGS calculated from main GWAS effects.

In this study we aimed to address the previously mentioned concerns. Using genetic data with linked-phenotypic information from electronic health records, we estimated the effects of covariate stratification and interaction on performance and effect estimates of PGS for BMI (PGS_{BMI}) – Figure 1 is a flowchart of the project. These analyses were done across four datasets: UK Biobank (UKBB), Penn Medicine BioBank (PMBB) (15), Electronic Medical Records and Genomics (eMERGE) network dataset (16), and Genetic Epidemiology Research on Adult Health and Aging (GERA). These datasets include participants from two ancestry groups (N=491,111 European ancestry (EUR), N=21,612 African ancestry (AFR)), and 62 covariates (25 present in multiple datasets) representing laboratory, survey, and biometric data types typically associated with cardiometabolic health and adiposity (including blood lipids, socioeconomic status, blood pressure, diet, physical activity, alcohol intake, smoking, and lung function, among others). We calculated the difference in PGS_{BMI} R^2 among individuals stratified by different quintiles of these variables (different groups for binary variables) and assessed whether a variable's association with BMI could explain its effect on PGS_{BMI} performance across

groups. We also assessed the significance of PGS_{BMI}-covariate interaction terms, and their
increases to model R^2 over models only using main effects, as well as correlation of main effect,
interaction effect, and R^2 differences. Replication and comparisons across all datasets and
ancestries were conducted. We then used machine learning models and evaluated their increase
in performance over linear models, even those that included regularization and interaction terms.
Finally, we conducted BMI GWAS using GxAge interaction terms, created PGS_{BMI} directly
using these effects, and assessed performance gains over using typical main effect GWAS. This
study addresses a plethora of open issues considering performance and effects of PGS on
individuals from diverse backgrounds.
148
Results
<u>Effect of covariate stratification on PGS_{BMI} performance</u>
We assessed 62 covariates for R^2 differences (25 present, or suitable proxies, in multiple datasets)
after stratifying on binary covariates and quintiles for continuous covariates. With UKBB EUR
as discovery (N=376,729), 18 covariates had significant differences (Bonferroni p<.05/62) in R^2
among groups (Figure 2a), including age, sex, alcohol consumption, different physical activity
measurements, Townsend deprivation index, different dietary measurements, lipids, blood
pressure, and A1c, with 40 covariates having suggestive (p<.05) evidence of R^2 differences.
From an original PGS_{BMI} R^2 of .076, R^2 increased to .094, .093, and .088 for those in the bottom
physical activity, alcohol intake, and high-density lipoprotein (HDL) cholesterol quintiles, and
decreased to .067, .57, and .049 for those in the top quintile (71%, 61%, and 56% relative R^2),
respectively, comparable to differences due to ancestry (1). We note that the differences in R^2
due to alcohol intake and HDL were larger than those of any physical activity phenotype, despite
physical activity having one of the oldest and most replicable evidence of interaction with
genetic effects of BMI (33,34). Despite considerable published evidence suggesting covariate-
specific genetic effects between BMI and smoking behaviors (6,8), we were only able to find
suggestive evidence for R^2 differences when stratifying individuals across several smoking
phenotypes (minimum p=0.016, for smoking pack years). R^2 differences due to educational
attainment were also only suggestive (p=0.015), of which published evidence is conflicting (35–
37).
We replicated these analyses in three additional large-scale cohorts of European and
African ancestry individuals (Figure 2b, Supplemental Table 3), as well as in African ancestry
UKBB individuals. For covariates with significant performance differences in the discovery
analysis, we were able to replicate significant (p<.05) R^2 differences for age, HDL cholesterol,
alcohol intake frequency, physical activity, and HbA1c, despite much smaller sample sizes. We
again observed largely insignificant differences across cohorts and ancestries when stratifying
due to different smoking phenotypes and educational attainment. Combining across cohorts, for
each covariate and ancestry we regressed R^2 values across groupings on covariate values in a
weighted linear regression, weighting by sample size. Slope of a weighted linear regression
across cohorts was directionally inconsistent between ancestries for the same covariate
(triglyceride levels, HbA1c, diastolic blood pressure, and sex), although larger sample sizes may
be needed to confirm these differences.
We noted several observations related to age-specific effects on PGS_{BMI}. First, in the
weighted linear regression of all R^2 across ancestries, expected R^2 for African ancestry
individuals overcomes that of European ancestry individuals among individuals within bottom
and top age quintiles observed in these data. For instance, the predicted R^2 of .048 for 80 year-

old European ancestry individuals would be lower than that of African ancestry individuals aged
24.7 and lower, indicating that certain differences in covariates can overcome differences in
PGS_{BMI} performance due to ancestry (with age being one of the most common variables used for
assessing and determining health). Second, we obtained these results despite the average age of
GWAS individuals being 57.8, which should increase PGS_{BMI} R^2 of individuals closest to this
age (13). This result suggests that in the case of these analyses, PGS performance due to
decreased heritability with age cannot be reconciled by having GWAS individuals of similar age
being used to create the PGS_{BMI}. Finally, we observed that PGS_{BMI} R^2 increases as age decreases,
consistent with published evidence suggesting that the heritability of BMI decreases with age
(38,39).
195
PGS-covariate interaction effects
Next, we estimated difference in PGS effects due to interactions with covariates, by modeling
interaction terms between PGS_{BMI} for each covariate (Methods). We implemented a correction
for shared heritability between covariates of interest and outcome (which can inflate test
statistics (28)) to better measure the environmental component of each covariate, and show that
this correction successfully reduces significance of interaction estimates (Supplemental Figure
1). Again, using UKBB EUR as discovery, we observed 28 covariates with significant
(Bonferroni p<.05/62) PGS-covariate interactions (Table 1), with 38 having suggestive (p<.05)
evidence (Supplemental Table 4). We observed the largest effect of PGS-covariate interaction
with alcohol drinking frequency (20.0% decrease in PGS effect per 1 standard deviation (SD)
increase, p=2.62×10^{-55}), with large effects for different physical activity measures (9.4%-12.5%
decrease/SD, minimum p=3.11×10^{-66}), HDL cholesterol (15.3% decrease/SD, p=1.71×10^{-96}) and
total cholesterol (12.7% decrease/SD, p=1.64×10^{-71}). We observed significant interactions with
diastolic blood pressure (10.8% increase/SD, p=6.06×10^{-60}), but interactions with systolic blood
pressure were much smaller (1.17% increase/SD, p=4.41×10^{-3}). Significant interactions with
HbA1c (4.63% increase/SD, p=5.37×10^{-14}) and type 2 diabetes (27.2% PGS effect increase in
cases, p=1.83×10^{-7}) were also observed. Other significant PGS-covariate interactions included
lung function, age, sex, and LDL cholesterol – various dietary measurements also had significant
interactions, albeit with smaller effects than other significant covariates. We were able to find
significant interaction effects for smoking pack years (4.78% increase/SD, p=3.68×10^{-7}), but
other smoking phenotypes had insignificant interactions after multiple testing correction
(minimum p=2.7×10^{-3}) – interactions with educational attainment were also insignificant (p=
4.54×10^{-2}).
219   We replicated these analyses across ancestries and the other cohorts in this study (Figure
3, Supplemental Table 4). For age and sex, which were available for all cohorts, interactions
were significant (p<.05) and directionally consistent across all cohorts and ancestries (except for
GERA AFR both effects were reversed, albeit with small sample size (N=1,789)). We were able
to test interactions with alcohol intake frequency and physical activity in GERA, and we were
able to replicate significant and directionally consistent association. We observed poor
replication for LDL cholesterol, HbA1c, and smoking pack years, with insignificant and
directionally inconsistent interaction effects across cohorts. Educational attainment was available
in GERA, and interactions were once again insignificant. We observed significant and
directionally consistent interaction effects for TG in eMERGE EUR and PMBB EUR, while the
effect was inconsistent in UKBB EUR despite much larger sample size.

However, despite significance of interaction terms, increases in model R^2 when including
PGS-covariate interaction terms were small. For instance, the maximum increase among all
covariates in UKBB EUR was only .0024 from a base R^2 of .1049 (2.1% relative increase), for
alcohol drink frequency per week. Across all cohorts and ancestries, the maximum increase in R^2
was only .0058 from a base R^2 of 0.09454 (6.1% relative increase), when adding a PGS-age
interaction term for eMERGE EUR ($p=5.40 \times 10^{-46}$) – this was also the largest relative increase
among models with significant interaction terms. This result suggests that, while interaction
effects can significantly modify the effect of PGS_{BMI}, their overall impact on model performance
is relatively small, despite the observation that stratifying by covariates produces large
differences in R^2.
240
<u>Correlations between R^2 differences, interaction effects, and main effects</u>
We next investigated the relationship between interaction effects, maximum R^2 differences
across quintiles, and main effects of covariates on BMI. We first estimated main effects of each
covariate on BMI (Methods, Supplemental Table 5), and then calculated the correlation weighted
by sample size between main effects, maximum PGS_{BMI} R^2 across quintiles, and PGS-covariate
interaction effects (Figure 4) across all cohorts and ancestries – GERA data were excluded from
these analyses due to slightly different phenotype definition (Supplemental Table 6), as were
binary variables. Interaction effects and maximum R^2 differences had a .80 correlation
($p=2.1 \times 10^{-27}$), indicating that variables with the largest interaction effects also had the largest
effect on PGS_{BMI} performance across quintiles, and that covariates that increase PGS_{BMI} effect
also have the largest effect on PGS_{BMI} performance i.e., individuals most at risk for obesity will
have both disproportionately larger PGS_{BMI} effect and R^2. Main effects and maximum R^2
differences had a .56 correlation ($p=1.3 \times 10^{-11}$), while main effects and interaction effects had a
.58 correlation (7.6×10^{-12}) again suggesting that PGS_{BMI} are more predictive in individuals with
higher values of BMI-associated covariates, although predictive to a lower degree than
estimating the interaction effects themselves directly. However, this result demonstrates that
covariates that influence both PGS_{BMI} effect and performance can be predicted just using main
effects of covariates, which are often known for certain phenotypes and easier to calculate, as
genetic data and PGS construction would not be required.
260
<u>Effects of machine learning approaches on predictive performance</u>
Given evidence of PGS-covariate dependence, we aimed to assess increases in R^2 when using
machine learning models (neural networks), which can inherently model interactions and other
nonlinearities, over linear models even with interaction terms. We first included age and sex as
the only covariates (along with genotype PCs and PGS_{BMI}), as age and sex were present in all
datasets and had significant and replicable evidence for PGS-dependence across our analyses.
Three models were assessed – L1-regularized (i.e., LASSO) linear regression without any
interaction terms, LASSO including a PGS-age and PGS-sex interaction term, and neural
networks (without interaction terms). When comparing neural networks to LASSO with
interaction terms, the relative 10-fold cross-validated R^2 increased up to 67% (mean 23%) across
cohorts and ancestries (Figure 5, Supplemental Table 7). The inclusion of interaction terms
increased cross-validated R^2 up to 12% (mean 3.9%) when comparing LASSO including
interaction terms to LASSO with main effects only.
We then modeled all available covariates for each cohort and did similar comparisons.
Cross-validated R^2 increased up to 17.6% (mean 9.5%) when using neural networks versus

LASSO with interaction terms, and up to 7.0% (mean 2.0%) when comparing LASSO with
interaction terms to LASSO with main effects only. Increases in model performance using neural
networks were smaller in UKBB, perhaps due to the age range being smaller than other cohorts
(all participants aged 39-73). This result suggests that additional variance explained through non-
linear effects with age and sex are explained by other variables present in the remainder of the
datasets. Our findings suggest that machine learning methods can improve model R^2 that include
PGS_{BMI} as variables beyond including interaction terms in linear models, even when variable
selection is performed using LASSO.
284
<u>Calculating PGS directly from GxAge GWAS effects</u>
Previous studies (13) have created PGS using GWAS stratified by different personal-level
covariates, but for practical purposes this leads to a large loss of power as the full size of the
GWAS is not utilized for each strata and continuous variables are forced into bins. We developed
a novel strategy where PGS are instead created from a full-size GWAS that includes SNP-
covariate interaction terms (Methods). We focused on age interactions, given their large and
replicable effects based on our results – similar to a previous study, we conducted these analyses
in the European UKBB. We used a 60% random split of study individuals to conduct three sets
of PGS using GWAS of the following designs: main effects only, main effects also with a SNP-
age interaction term, and main effects but stratified into quartiles by age. 20% of the remaining
data were used for training and the final 20% were held-out as a test set. The best performing
PGS created from SNP-age interaction terms (PGS_{GxAge}) increased test R^2 to 0.0771 from 0.0715
(7.8% relative increase) compared to the best performing main effect PGS (Figure 6,
Supplemental Table 9) – age-stratified PGS had much lower performance than both other
strategies (unsurprising given reasons previously mentioned). Including a PGS_{GxAge}-age
interaction term only marginally increased R^2 (.0001 increase), with similar increases for the
other two sets of PGS, further demonstrating that post-hoc modeling of interactions cannot
reconcile performance gained through directly incorporating interaction effects from the original
GWAS. The strategy of creating PGS directly from full-sized SNP-covariate interactions is
potentially quite useful as it increases PGS performance without the need for additional data –
there are almost certainly a variety of points of improvement (Discussion), but we consider their
investigation outside the scope of this study.
307
Discussion
Across four large-scale cohorts of diverse ancestry, we uncover replicable effects of covariates
on both performance and effects of PGS_{BMI}. When stratifying by quintiles of different covariates,
we find certain covariates that have significant and replicable evidence for differences in PGS_{BMI}
R^2, with R^2 being nearly double between top and bottom performing quintiles for covariates with
the largest differences. When testing covariates for PGS-covariate interaction effects, we also
find covariates with significant interaction effects, where, for the largest effect covariates, each
standard deviation change affects PGS_{BMI} effect by nearly 20%. Across analyses, we find age and
sex had the most replicable interaction effects, with levels of serum cholesterol, physical activity,
and alcohol consumption having the largest effects across cohorts. We also observed that
interaction effects and R^2 differences were strongly correlated, with main effects also being
correlated with interaction effects and R^2 differences, suggesting that covariates with the largest
interaction effects also contribute to the largest R^2 differences, with simple main effects also
being predictive of expected differences in R^2 and interaction effects. While in this study we

stratified by quintile, similar trends of performance differences would be apparent in smaller
percentiles. This observation is relevant to current research and clinical use of PGS, as
individuals above a percentile cutoff are designated high risk (40). These analyses imply that
individuals most at-risk for obesity should have both disproportionately higher predicted BMI
and increased BMI prediction performance compared to the general population – more work is
required to understand why, and if similar trends can be observed in other traits relevant to
human health. We also employ machine learning methods for prediction of BMI with models
that include PGS_{BMI} and demonstrate that these methods outperform regularized linear regression
models that include interaction effects. Finally, we employ a novel strategy that directly
incorporates SNP interaction effects into PGS construction, and demonstrate that this strategy
improves PGS performance when modeling SNP-age interactions compared to PGS created only
from main effects.
Future work may include replicating these analyses across additional traits, and further
exploring machine learning and deep learning methods on other phenotypes to determine if this
trend of inclusion of PGS along with covariate interaction effects outperforms linear models for
risk prediction. Additionally, the method of including a PGS for the covariate to better measure
its environmental effect is potentially worth exploring further, which should improve in the
future as PGS performance continues to increase. A slight limitation of this method in our study
is that for the UKBB analyses the GWAS used for PGS construction were also from UKBB thus
not out-of-sample, although many of the covariates only have GWAS available through UKBB
individuals. Furthermore, a variety of improvements are likely possible when creating PGS
directly from SNP-covariate interaction terms. First, we used the same SNPs that were selected
by pruning and thresholding based on their main effect p-values, but selection of SNPs based on
their interaction p-values should also be possible and would likely improve performance.
Additionally, performance of pruning and thresholding-based strategies have largely been
overtaken by methods that first adjust all SNP effects for LD and don't require exclusion of
SNPs, and a method that could do a similar adjustment for interaction effects would likely
outperform most current methods for traits with significant context-specific effects. Finally,
incorporation of additional SNP-covariate interactions (e.g., SNP-gender) would also likely
further improve prediction performance, although any SNP selection/adjustment procedures may
further complicated by additional interaction terms.
We propose several ideas on why BMI-associated covariates have larger interaction
effects and impact on R^2 for PGS_{BMI}. Age may be a proxy for accumulated gene-environment
interactions as younger individuals have less exposure to environmental influences on weight
compared to older individuals; therefore, one may expect that in younger individuals their
phenotype could be better explained by genetics compared to environment. Secondly, PGS may
more readily explain more extreme phenotypes because they incorporate large effect variants
associated with very high weight or height (41). For BMI, the distribution is skewed toward the
right i.e., effects in trait-increasing variants may have a larger potential to affect trait variation
compared to trait-reducing variants. This explanation would likely be better suited to traits with
left-tailed distributions and is not fully satisfactory as first log-transforming or rank-normal
transforming the phenotype, as was done in this study, may invalidate this explanation.
PGS is a promising technique to stratify individuals for their risk of common, complex
disease. In order to achieve accurate predictions as well as promote equity, further research is
required regarding PGS methods and assessments. This research provides firm evidence on the
context-specific nature of PGS and the incorporation of covariate interaction effects for

predicting BMI and, apart from promoting equitable use of PGS across ancestries and cohorts,
demonstrates ways to improve PGS performance incorporating context-specific effects.
370
371 **Methods**
372 <u>Study datasets</u>
373 Individual inclusion criteria and sample sizes per cohort are described below – additional
374 information is available in Supplemental Table 1.
375
UKBB
Individual-level quality-control and filtering have been described elsewhere (17) for European
ancestry individuals. Briefly, individuals were split by ancestry according to both genetically
inferred ancestry and self-reported ethnicity (15). Individuals with low genotyping quality and
sex mismatch were removed, only unrelated individuals (pi-hat < .250) were retained, and
variants were filtered at INFO > .30 and minor allele frequency (MAF) > .01. For African
ancestry, individuals were first selected based on self-reported ethnicity "Black or Black
British", "Caribbean", "African", or "Any other Black background". Individuals that were low
quality i.e., "Outliers for heterozygosity or missing rate", and that were Caucasian from "Genetic
ethnic grouping" were removed. Of these individuals, those that were +/- 6 standard deviations
from the mean of the first 5 genetic principal components provided by UKBB were excluded.
Finally, only unrelated individuals were retained up to the second degree using plink2 (18) "-
king-cutoff .125". After QC and consideration of phenotype, a total of 7,046 individuals in the
UKBB AFR data who also had BMI available were used for downstream analyses. In total,
383,775 individuals were used for analysis (N_{EUR}=376,729, N_{AFR}=7,046).
391
392 *eMERGE*
393 Ancestry and relatedness inference have been described elsewhere (15). Individuals were split
394 into European and African ancestry cohorts, and related individuals were removed (pi-hat >
395 .250) such that all were unrelated. 35,064 individuals (N_{EUR}=31,961, N_{AFR}=3,103) were used for
396 analysis.
397
398 *GERA*
399 Ancestry inference has been described elsewhere (19), and study individuals were divided into
400 European and African ancestry cohorts. Related individuals were removed using plink2 "-king-
401 cutoff .125". 57,838 individuals (N_{EUR}=56,049, N_{AFR}=1,789) were used for analysis.
402
403 *PMBB*
404 Ancestry inference and relatedness inference have been described elsewhere (20). Individuals
405 were split into European and African ancestry cohorts, and related individuals were removed at
406 pi-hat > .250. 36,046 individuals (N_{EUR}=26,372 , N_{AFR}=9,674) were used for analysis.
407
<u>Choice of covariates</u>
A total of 62 covariates were included in the analyses, 25 of which were present (or similar
proxies) in multiple datasets. These covariates were selected based on relevance to
cardiometabolic health and obesity, and previous evidence of context-specific effects with BMI
(5,6,8,21–23). For UKBB, phenotype values were used from the collection that was closest to
recruitment, and for PMBB the median values were used – for GERA and eMERGE only one

value was available. Additional details on covariate constructions, transformations, filtering, and
cohort availability are in Supplemental Table 2.
416
<u>PGS construction</u>
PGS for BMI (PGS_{BMI}) were constructed using PRS-CSx (24), using GWAS summary statistics
for individuals of European (25), African (26), and East Asian (27) ancestry that were out-of-
sample of study participants. A set of 1.29 million HapMap3 SNPs provided by PRS-CSx was
used for PGS calculation, which are generally well-imputed and variable frequency across global
populations. Default settings for PRS-CSx (downloaded November 22, 2021) were used, which
have been demonstrated to perform well for highly polygenic traits such as BMI (list of
parameters in Supplemental Table 8). The final PGS_{BMI} per ancestry and cohort was calculated
by regressing log(BMI) on the PGS_{BMI} per ancestry without covariates – the combined, predicted
value was used as a single PGS_{BMI} in downstream analyses.
427
428 For GERA, BMI was not transformed, as it was already binned into a categorical variable with 5
429 levels (18≤, 19-25, 26-29, 30-39, >40). Additionally, for GERA the uncombined ancestry-
430 specific PGS_{BMI} were used in the final models, as it had higher R^2 than using the combined
431 PGS_{BMI} (data not shown).
432
433 <u>PGS_{BMI} performance after covariate stratification</u>
434 Analyses were performed separately for each cohort and ancestry. For each covariate, individuals
435 were binned by binary covariates or quintiles for continuous covariates. Incremental PGS_{BMI} R^2
436 was calculated by taking the difference in R^2 between:
437
438 $\log(BMI) \sim PGS_{BMI} + Age + Sex + PCs_{1\text{-}5}$
439 $\log(BMI) \sim Age + Sex + PCs_{1\text{-}5}$
440
441 We performed 5,000 bootstrap replications to obtain a bootstrapped distribution of R^2. P-values
442 for differences in R^2 were calculated between groups by calculating the proportion of overlap
443 between two normal distributions of the R^2 value using the standard deviations of the bootstrap
444 distributions. Again for GERA, BMI was not transformed.
445
<u>PGS_{BMI} interaction modelling</u>
Evidence for interaction with each covariate with the PGS_{BMI} was evaluated using linear
regression. It has been reported that the inclusion of covariates that share heritability with the
outcome can inflate test statistic estimates (28–30). To assuage these concerns, we introduced a
novel correction, where we first calculated a PGS for the covariate ($PGS_{Covariate}$) and included it
in the model, as well as an interaction term between PGS_{BMI} and $PGS_{Covariate}$. The $PGS_{Covariate}$
terms were calculated using the European ancestry Neale Lab summary statistics (URLs) and
PRS-CS (31). To standardize effect sizes across analyses, PGS_{BMI} and Covariate were first
converted to mean zero and standard deviation of 1 (binary covariates were not standardized).
We demonstrate inclusion of $PGS_{Covariate}$ terms successfully reduced significance of the
PGS_{BMI}*Covariate term (Supplemental Figure 1). The final model used to evaluate PGS_{BMI} and
Covariate interactions was:
458
459 $\log(BMI) \sim PGS_{BMI}*Covariate + PGS_{BMI} + Covariate + PGS_{Covariate} + PGS_{BMI}*PGS_{Covariate} + Age + Sex + PCs_{1\text{-}5}$
460

461 We report the effect estimates of the PGS_{BMI}*Covariate term, and differences in model R^2 with
462 and without the PGS_{BMI}*Covariate term. Again for GERA, BMI was not transformed.
463
464 <u>Correlation between R^2 differences, interaction effects, and main effects</u>
465 We estimated main effects of each covariate on BMI with the following model:
466
467 $\log(BMI) \sim Covariate + Age + Sex + PCs_{1-5}$
468
Note that we ran new models with main effects only, instead of using the main effect from the
interaction models (as the main effects in the interaction models depend on the interaction terms,
and main effects used to create interaction terms are sensitive to centering of variables despite
the scale invariance of linear regression itself (32)). We then estimated the correlation between
main effects, interaction effects, and maximum R^2 differences across all cohorts and ancestries
weighting by sample size, analyzing quantitative and binary variables separately.
475
<u>Machine learning models</u>
UKBB EUR and GERA EUR models were restricted to 30,000 random individuals, for
computational reasons – BMI distributions did not differ from the full-sized datasets
(Kolmogorov-Smirnov p-value of 0.29 and 0.57, respectively). PGS_{BMI} and top five genetic
principal components were included as features in all models. Two sets of models were
evaluated for each cohort and ancestry: including age and sex as features, and including all
available covariates in each cohort as features. 'Ever Smoker' status was used in favor of 'Never'
vs 'Current smoking' status (if present), as individuals with 'Never' vs 'Current' status are a
subset of those with 'Ever Smoker' status. UKBB AFR with all covariates was excluded due to
small sample size (N=53).
Neural networks were used as the model of choice, given their inherent ability model
interactions and other nonlinear dependencies. Prior to modeling, all features were scaled to be
between 0 and 1. We used average 10-fold cross validation R^2 to evaluate model performance.
Separate models were trained using untransformed and log(BMI). L1-regularized linear
regression was used with 18 values of lambda (1.0, 5.0×10^{-1}, 2.0×10^{-1}, 1.0×10^{-1}, 5.0×10^{-2}, 2.0×10^{-2}
^{2}, …, 1.0×10^{-5}, 5.0×10^{-6}, 2.0×10^{-6}). Models were trained without inclusion of interaction terms
(which neural networks can implicitly model), using 1,000 iterations of random search with the
following hyperparameter ranges: size of hidden layers [10, 200], learning rate [.01, .0001], type
of learning rate [constant, inverse scaling], power t [.4, .6], momentum [.80, 1.0], batch size [32,
256], and number of hidden layers [1, 2].
496
497 <u>GxAge PGS_{BMI} creation and assessment</u>
498 Analyses were conducted in the European UKBB (N=376,629), as was done in a study on a
499 similar topic (13). Three sets of analyses were performed, using GWAS conducted in a 60%
500 random split of individuals using the following models (BMI was rank-normal transformed
501 before GWAS):
502
503 1) $BMI \sim SNP + Age + Sex + PCs_{1-5}$
504
505 2) $BMI \sim SNP + Age*SNP + Age + Sex + PCs_{1-5}$
506

3) Using the model in 1) but stratified into quartiles by age. BMI was rank-normal transformed
within each quartile.

Using each set of GWAS, PGS was first calculated in a 20% randomly selected training set of
the dataset using pruning and thresholding using 10 p-value thresholds (.50, .10, …, 5.0×10^{-5},
1.0×10^{-5}) and remaining settings as default in Plink 1.9. For 2), GxAge PGS_{BMI} were calculated
using SNPs clumped by their main effect p-values from 1), and additionally incorporating the
GxAge interaction effects per SNP. In other words, instead of typical PGS construction as:

$PGS_i = \beta_1 k_1 + \beta_2 k_2 + \ldots + \beta_n k_n$

For an individual i's PGS calculation, with main SNP effect β, and n SNPs, PGS incorporating
GxAge effects (PGS_{GxAge}) was calculated as:

$PGS_{GxAge,i} = \beta_1 k_1 + \beta_{GxAge,1} k_1 Age_i + \beta_2 k_2 + \beta_{GxAge,2} k_2 Age_i \ldots + \beta_n k_n + \beta_{GxAge,n} k_n Age_i$

Where β_{GxAge} is the GxAge effect for each SNP n and Age_i is the age for individual i.

For each of the three analyses, the parameters and models resulting in the best performing PGS
(highest incremental R^2, using same main effect covariates as in the three GWAS) from the
training set were evaluated in the remaining 20% of the study individuals. For 3), models were
first trained within each quartile separately. To maintain sense of scale across quartiles (after
rank-normal transformation), R^2 between all predicted values and true values was calculated
together.

URLs
Neale Lab UKBB summary statistics: http://www.nealelab.is/uk-biobank

Data Availability statement
UK Biobank data was accessed under project #32133. eMERGE data is available at dbGaP in
phs001584.v2.p2. GERA data is available at dbGaP in phs000674.v3.p3. Summary statistics for
PMBB are available from authors upon request.

Regeneron Genetics Center Banner Author List and Contribution Statements
<u>RGC Management and Leadership Team</u>
Goncalo Abecasis, PhD , Aris Baras, M.D. , Michael Cantor, M.D. , Giovanni Coppola, M.D. ,
Andrew Deubler , Aris Economides, Ph.D. , Luca A. Lotta, M.D., Ph.D. , John D. Overton, Ph.D.
, Jeffrey G. Reid, Ph.D. , Katherine Siminovitch, M.D. , Alan Shuldiner, M.D.

<u>Sequencing and Lab Operations</u>
Christina Beechert , Caitlin Forsythe, M.S. , Erin D. Fuller , Zhenhua Gu, M.S. , Michael Lattari ,
Alexander Lopez, M.S. , John D. Overton, Ph.D. , Maria Sotiropoulos Padilla, M.S. , Manasi
Pradhan, M.S. , Kia Manoochehri, B.S. , Thomas D. Schleicher, M.S. , Louis Widom , Sarah E.
Wolf, M.S. , Ricardo H. Ulloa, B.S.

<u>Clinical Informatics</u>

Amelia Averitt, Ph.D. , Nilanjana Banerjee, Ph.D. , Michael Cantor, M.D. , Dadong Li, Ph.D. , Sameer Malhotra, M.D. , Deepika Sharma, MHI , Jeffrey Staples , Ph.D.

Genome Informatics
Xiaodong Bai, Ph.D. , Suganthi Balasubramanian, Ph.D. , Suying Bao, Ph.D. , Boris Boutkov, Ph.D. , Siying Chen, Ph.D. , Gisu Eom, B.S. , Lukas Habegger, Ph.D. , Alicia Hawes, B.S. , Shareef Khalid , Olga Krasheninina, M.S. , Rouel Lanche, B.S. , Adam J. Mansfield, B.A. , Evan K. Maxwell, Ph.D. , George Mitra, B.A. , Mona Nafde, M.S. , Sean O'Keeffe, Ph.D. , Max Orelus, B.B.A. , Razvan Panea, Ph.D. , Tommy Polanco, B.A. , Ayesha Rasool, M.S. , Jeffrey G. Reid, Ph.D. , William Salerno, Ph.D. , Jeffrey C. Staples, Ph.D. , Kathie Sun, Ph.D.

Analytical Genomics and Data Science
Goncalo Abecasis, D.Phil. , Joshua Backman, Ph.D. , Amy Damask, Ph.D. , Lee Dobbyn, Ph.D. , Manuel Allen Revez Ferreira, Ph.D. , Arkopravo Ghosh, M.S. , Christopher Gillies, Ph.D. , Lauren Gurski, B.S. , Eric Jorgenson, Ph.D. , Hyun Min Kang, Ph.D. , Michael Kessler, Ph.D. , Jack Kosmicki, Ph.D. , Alexander Li , Ph.D. , Nan Lin, Ph.D. , Daren Liu, M.S. , Adam Locke, Ph.D. , Jonathan Marchini, Ph.D. , Anthony Marcketta, M.S. , Joelle Mbatchou, Ph.D. , Arden Moscati, Ph.D. , Charles Paulding, Ph.D. , Carlo Sidore, Ph.D. , Eli Stahl, Ph.D. , Kyoko Watanabe, Ph.D. , Bin Ye, Ph.D. , Blair Zhang, Ph.D. , Andrey Ziyatdinov, Ph.D.

Therapeutic Area Genetics
Ariane Ayer, B.S. , Aysegul Guvenek, Ph.D. , George Hindy, Ph.D. , Giovanni Coppola, M.D. , Jan Freudenberg, M.D. , Jonas Bovijn M.D. , Katherine Siminovitch, M.D. , Kavita Praveen, Ph.D. , Luca A. Lotta, M.D. , Manav Kapoor, Ph.D. , Mary Haas, Ph.D. , Moeen Riaz , Ph.D. , Niek Verweij, Ph.D. , Olukayode Sosina, Ph.D. , Parsa Akbari, Ph.D. , Priyanka Nakka, Ph.D. , Sahar Gelfman, Ph.D. , Sujit Gokhale, B.E. , Tanima De, Ph.D. , Veera Rajagopal, Ph.D. , Alan Shuldiner, M.D. , Bin Ye, Ph.D. , Gannie Tzoneva, Ph.D. , Juan Rodriguez-Flores, Ph.D.

Research Program Management & Strategic Initiatives
Esteban Chen, M.S. , Marcus B. Jones, Ph.D. , Michelle G. LeBlanc, Ph.D. , Jason Mighty, Ph.D., Lyndon J. Mitnaul, Ph.D. , Nirupama Nishtala, Ph.D. , Nadia Rana, Ph.D. , Jaimee Hernandez

Acknowledgements
We acknowledge David Crosslin for helping clean the eMERGE data.

Funding
M.D.R., S.D., and D.H. are funded by AI077505. W.K.C is funded by grant NIDDK DK52431. J.F.P. is funded by grant U01 HG011166. C.W. is funded by U01 HG008680. This phase of the eMERGE Network was initiated and funded by the NHGRI through the following grants: U01HG008657 (Group Health Cooperative/University of Washington); U01HG008685 (Brigham and Women's Hospital); U01HG008672 (Vanderbilt University Medical Center); U01HG008666 (Cincinnati Children's Hospital Medical Center); U01HG006379 (Mayo Clinic); U01HG008679 (Geisinger Clinic); U01HG008680 (Columbia University Health Sciences); U01HG008684 (Children's Hospital of Philadelphia); U01HG008673 (Northwestern University); U01HG008701 (Vanderbilt University Medical Center serving as the Coordinating

599   Center); U01HG008676 (Partners Healthcare/Broad Institute); U01HG008664 (Baylor College
600   of Medicine); and U54MD007593 (Meharry Medical College).
601

References

1. Martin AR, Kanai M, Kamatani Y, Okada Y, Neale BM, Daly MJ. Clinical use of current
polygenic risk scores may exacerbate health disparities. Nat Genet. 2019 Apr;51(4):584–91.

2. Wang Y, Guo J, Ni G, Yang J, Visscher PM, Yengo L. Theoretical and empirical
quantification of the accuracy of polygenic scores in ancestry divergent populations. Nat
Commun. 2020 Jul 31;11(1):3865.

3. Galinsky KJ, Reshef YA, Finucane HK, Loh PR, Zaitlen N, Patterson NJ, et al. Estimating
cross-population genetic correlations of causal effect sizes. Genet Epidemiol. 2019
Mar;43(2):180–8.

4. Shi H, Gazal S, Kanai M, Koch EM, Schoech AP, Siewert KM, et al. Population-specific
causal disease effect sizes in functionally important regions impacted by selection. Nat
Commun. 2021 Feb 17;12(1):1098.

5. Rask-Andersen M, Karlsson T, Ek WE, Johansson Å. Gene-environment interaction study
for BMI reveals interactions between genetic factors and physical activity, alcohol
consumption and socioeconomic status. PLoS Genet. 2017 Sep;13(9):e1006977.

6. Robinson MR, English G, Moser G, Lloyd-Jones LR, Triplett MA, Zhu Z, et al. Genotype-
covariate interaction effects and the heritability of adult body mass index. Nat Genet. 2017
Aug;49(8):1174–81.

7. Sulc J, Mounier N, Günther F, Winkler T, Wood AR, Frayling TM, et al. Quantification of
the overall contribution of gene-environment interaction for obesity-related traits. Nat
Commun. 2020 Mar 13;11(1):1385.

8. Justice AE, Winkler TW, Feitosa MF, Graff M, Fisher VA, Young K, et al. Genome-wide
meta-analysis of 241,258 adults accounting for smoking behaviour identifies novel loci for
obesity traits. Nat Commun. 2017 Apr 26;8:14977.

9. Helgeland Ø, Vaudel M, Juliusson PB, Lingaas Holmen O, Juodakis J, Bacelis J, et al.
Genome-wide association study reveals dynamic role of genetic variation in infant and early
childhood growth. Nat Commun. 2019 Oct 1;10(1):4448.

10. Vogelezang S, Bradfield JP, Ahluwalia TS, Curtin JA, Lakka TA, Grarup N, et al. Novel loci
for childhood body mass index and shared heritability with adult cardiometabolic traits.
PLoS Genet. 2020 Oct;16(10):e1008718.

11. Couto Alves A, De Silva NMG, Karhunen V, Sovio U, Das S, Taal HR, et al. GWAS on
longitudinal growth traits reveals different genetic factors influencing infant, child, and adult
BMI. Sci Adv. 2019 Sep;5(9):eaaw3095.

12. Choh AC, Lee M, Kent JW, Diego VP, Johnson W, Curran JE, et al. Gene-by-age effects on
BMI from birth to adulthood: the Fels Longitudinal Study. Obes Silver Spring Md. 2014
Mar;22(3):875–81.

13. Mostafavi H, Harpak A, Agarwal I, Conley D, Pritchard JK, Przeworski M. Variable
prediction accuracy of polygenic scores within an ancestry group. eLife. 2020 Jan
30;9:e48376.

14. Elks CE, den Hoed M, Zhao JH, Sharp SJ, Wareham NJ, Loos RJF, et al. Variability in the
heritability of body mass index: a systematic review and meta-regression. Front Endocrinol.
2012;3:29.

15. Bycroft C, Freeman C, Petkova D, Band G, Elliott LT, Sharp K, et al. The UK Biobank
resource with deep phenotyping and genomic data. Nature. 2018 Oct;562(7726):203–9.

16. Stanaway IB, Hall TO, Rosenthal EA, Palmer M, Naranbhai V, Knevel R, et al. The
eMERGE genotype set of 83,717 subjects imputed to ~40 million variants genome wide and
association with the herpes zoster medical record phenotype. Genet Epidemiol. 2019
Feb;43(1):63–81.

17. Zhang X, Lucas AM, Veturi Y, Drivas TG, Bone WP, Verma A, et al. Large-scale genomic
analyses reveal insights into pleiotropy across circulatory system diseases and nervous
system disorders. Nat Commun. 2022 Jun 14;13(1):3428.

18. Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. Second-generation
PLINK: rising to the challenge of larger and richer datasets. GigaScience. 2015;4:7.

19. Banda Y, Kvale MN, Hoffmann TJ, Hesselson SE, Ranatunga D, Tang H, et al.
Characterizing Race/Ethnicity and Genetic Ancestry for 100,000 Subjects in the Genetic
Epidemiology Research on Adult Health and Aging (GERA) Cohort. Genetics. 2015
Aug;200(4):1285–95.

20. Penn Medicine BioBank [Internet]. Available from: https://pmbb.med.upenn.edu/

21. Tyrrell J, Wood AR, Ames RM, Yaghootkar H, Beaumont RN, Jones SE, et al. Gene-
obesogenic environment interactions in the UK Biobank study. Int J Epidemiol. 2017 Apr
1;46(2):559–75.

22. Young AI, Wauthier F, Donnelly P. Multiple novel gene-by-environment interactions modify
the effect of FTO variants on body mass index. Nat Commun. 2016 Sep 6;7:12724.

23. Winkler TW, Justice AE, Graff M, Barata L, Feitosa MF, Chu S, et al. The Influence of Age
and Sex on Genetic Associations with Adult Body Size and Shape: A Large-Scale Genome-
Wide Interaction Study. PLoS Genet. 2015 Oct;11(10):e1005378.

24. Ruan Y, Lin YF, Feng YCA, Chen CY, Lam M, Guo Z, et al. Improving polygenic
prediction in ancestrally diverse populations. Nat Genet. 2022 May;54(5):573–80.

25. Locke AE, Kahali B, Berndt SI, Justice AE, Pers TH, Day FR, et al. Genetic studies of body
mass index yield new insights for obesity biology. Nature. 2015 Feb 12;518(7538):197–206.

26. Ng MCY, Graff M, Lu Y, Justice AE, Mudgal P, Liu CT, et al. Discovery and fine-mapping
of adiposity loci using high density imputation of genome-wide association studies in
individuals of African ancestry: African Ancestry Anthropometry Genetics Consortium.
PLoS Genet. 2017 Apr;13(4):e1006719.

27. Sakaue S, Kanai M, Tanigawa Y, Karjalainen J, Kurki M, Koshiba S, et al. A cross-
population atlas of genetic associations for 220 human phenotypes. Nat Genet. 2021
Oct;53(10):1415–24.

28. Aschard H, Vilhjálmsson BJ, Joshi AD, Price AL, Kraft P. Adjusting for heritable covariates
can bias effect estimates in genome-wide association studies. Am J Hum Genet. 2015 Feb
5;96(2):329–39.

29. Kerin M, Marchini J. Inferring Gene-by-Environment Interactions with a Bayesian Whole-
Genome Regression Model. Am J Hum Genet. 2020 Oct 1;107(4):698–713.

30. Vanderweele TJ, Ko YA, Mukherjee B. Environmental confounding in gene-environment
interaction studies. Am J Epidemiol. 2013 Jul 1;178(1):144–52.

31. Ge T, Chen CY, Ni Y, Feng YCA, Smoller JW. Polygenic prediction via Bayesian regression
and continuous shrinkage priors. Nat Commun. 2019 Apr 16;10(1):1776.

32. Afshartous D, Preston RA. Key Results of Interaction Models with Centering. J Stat Educ.
2011 Nov;19(3):1.

33. Kilpeläinen TO, Qi L, Brage S, Sharp SJ, Sonestedt E, Demerath E, et al. Physical activity
attenuates the influence of FTO variants on obesity risk: a meta-analysis of 218,166 adults
and 19,268 children. PLoS Med. 2011 Nov;8(11):e1001116.

34. Rampersaud E, Mitchell BD, Pollin TI, Fu M, Shen H, O'Connell JR, et al. Physical activity
and the association of common FTO gene variants with body mass index and obesity. Arch
Intern Med. 2008 Sep 8;168(16):1791–7.

35. Amin V, Dunn P, Spector T. Does education attenuate the genetic risk of obesity? Evidence
from U.K. Twins. Econ Hum Biol. 2018 Sep;31:200–8.

36. Li Y, Cai T, Wang H, Guo G. Achieved educational attainment, inherited genetic
endowment for education, and obesity. Biodemography Soc Biol. 2021 Jun;66(2):132–44.

37. Frank M, Dragano N, Arendt M, Forstner AJ, Nöthen MM, Moebus S, et al. A genetic sum
score of risk alleles associated with body mass index interacts with socioeconomic position
in the Heinz Nixdorf Recall Study. PloS One. 2019;14(8):e0221252.

38. Ge T, Chen CY, Neale BM, Sabuncu MR, Smoller JW. Phenome-wide heritability analysis
of the UK Biobank. PLoS Genet. 2017 Apr;13(4):e1006711.

39. Min J, Chiu DT, Wang Y. Variation in the heritability of body mass index based on diverse
twin studies: a systematic review. Obes Rev Off J Int Assoc Study Obes. 2013
Nov;14(11):871–82.

40. Linder JE, Allworth A, Bland ST, Caraballo PJ, Chisholm RL, Clayton EW, et al. Returning
integrated genomic risk and clinical recommendations: The eMERGE study. Genet Med Off
J Am Coll Med Genet. 2023 Jan 6;25(4):100006.

41. Robinson PN, Arteaga-Solis E, Baldock C, Collod-Béroud G, Booms P, De Paepe A, et al.
The molecular genetics of Marfan syndrome and related disorders. J Med Genet. 2006
Oct;43(10):769–87.

749 **Figures**

750
751 Figure 1. A flowchart of the project.

a)

763
764
765 b)

766
767
Figure 2. PGS R^2 stratified by quintiles for quantitative variables and by binary variables. a)
Continuous covariates with significant (p < 7.7x10^{-4}) R^2 differences across quintiles in UKBB

EUR. Pork and processed meat consumption per week were excluded from this plot in favor of
pork and processed meat intake. b) Covariates with significant differences that were available in
multiple cohorts, note actual values instead of percentiles plotted on x-axis (except percentiles
were used for physical activity, alcohol intake frequency, and socioeconomic status, which had
slightly differing phenotype definitions across cohorts). Townsend index and income were used
as variables for socioeconomic status UKBB and GERA, respectively. Note that the sign for
Townsend index was reversed, since increasing Townsend index is lower socioeconomic status,
while increasing income is higher socioeconomic status. Abbreviations: physical activity (PA),
International Physical Activity Questionnaire IPAQ).

Figure 3. Relative percentage changes in PGS effect per unit change in covariate, for covariates that significantly changed PGS effect (i.e., significant interaction beta at Bonferroni $p < 7.7 \times 10^{-4}$ – denoted by asterisks) and were present in multiple cohorts and ancestries. Same covariate groupings and transformations were performed as with Figure 1.

801
802
803 Figure 4. Pearson correlations weighted by sample size between maximum R^2 differences across
804 strata, main effects of covariate on log(BMI), and PGS-covariate interaction effects on log(BMI).
805 Main effect units are in standard deviations, interaction effect units are in PGS standard
806 deviations multiplied by covariate standard deviations. Only continuous variables are plotted and
807 modeled. GERA was excluded due to slightly different phenotype definitions.

Figure 5. Model R^2 across cohorts and ancestries using age and gender as covariates (along with PGS_{BMI} and PCs 1-5). Across all cohorts and ancestries, LASSO with PGS-age and PGS-gender interaction terms had better average 10-fold cross-validation R^2 than LASSO without interaction terms, while neural networks outperformed LASSO models.

838
839 Figure 6. PGS R^2 based on three sets of GWAS. "Main effects" were from a typical main effect
840 GWAS, "GxAge" effects were from a GWAS with a SNP-age interaction term, and "Age
841 stratified" GWAS had main effects only but were conducted in four age quartiles. PGS R^2 was
842 evaluated using two models: one with main effects only, and one with an additional PGS*Age
843 interaction term.

858   <u>Supplemental Figure:</u>

859
860   S Figure 1. PGS-covariate interaction term p-values in UKBB EUR, with and without including
861   the covariate PGS in the model – the mean -$\log_{10}$(p) is reduced from 18.0899 to 14.97072 with
862   their inclusions. Note age and sex PGS were not calculated, and their interaction p-values are
863   excluded from this figure.

872   **Table**

Variable type	Covariate	% change in β_{PGS} per covariate unit change	Interaction P	R^2 increase with interaction term	N
	HDL cholesterol	-15.29	1.71×10^{-96}	0.0012	328719
	Total cholesterol	-12.70	1.64×10^{-71}	0.00082	359221
	IPAQ	-12.50	3.11×10^{-66}	0.001	304951
	Moderate-vigorous PA	-11.41	8.92×10^{-65}	0.001	304951
	Diastolic BP	10.84	6.06×10^{-60}	0.0007	352804
	Townsend Index	6.78	2.86×10^{-58}	0.00089	376283
	Age	-9.02	3.60×10^{-57}	0.00061	376729
	FVC	-9.66	4.69×10^{-56}	0.0008	343467
	Drink frequency/wk	-19.96	2.62×10^{-55}	0.0024	122281
	LDL cholesterol	-9.86	2.63×10^{-51}	0.00058	358556
	N days vigorous PA/wk	-9.37	2.42×10^{-35}	0.0007	299963
Continuous	FEV1	-7.38	7.15×10^{-35}	0.0005	343544
	Mean alcohol consumption	-7.38	7.65×10^{-22}	0.00113	126756
	HbA1c	4.63	5.37×10^{-14}	0.0002	358798
	Mean drinks/wk	-7.66	1.01×10^{-13}	0.0008	112204
	Water intake	4.60	2.97×10^{-13}	0.00014	347472
	Processed meat intake	3.70	2.38×10^{-7}	0.0002	376205
	Starch mean	5.51	3.15×10^{-7}	0.00018	128346
	Smoking pack years	4.78	3.68×10^{-7}	0.0002	114135
	Protein mean	4.82	6.52×10^{-7}	0.00018	128181
	Saturated fat mean	4.92	1.23×10^{-6}	0.00017	127899
	Fat mean	4.40	1.64×10^{-5}	0.00013	128092
	Saturated fat grams/wk	2.46	1.79×10^{-5}	4.00E-05	364629
	Retinol mean	3.77	3.54×10^{-4}	9.00E-05	126029
	IPAQ	-12.68	5.30×10^{-62}	0.0009	304951
Binary	Vigorous PA/wk	-20.55	9.07×10^{-54}	0.0009	304951
	Sex	-11.02	1.41×10^{-24}	0.00025	376729
	Diabetes	27.19	1.83×10^{-7}	0.0004	375903

873
Table 1. Model descriptive statistics on 28 of 62 covariates, which have significant (p<.05/62)
PGS-covariate interaction terms, in UKBB EUR. The third column is the percentage change in
PGS effect per unit change (standard deviations for continuous variables, binary variables
encoded as 0 or 1) in covariate. The fifth column is the increase in model R^2 with a PGS-
covariate interaction term versus a main effects only model. Abbreviations: blood pressure (BP),
physical activity (PA), forced vital capacity (FVC), forced expiratory volume in 1-second
(FEV1), International Physical Activity Questionnaire (IPAQ).