Vitamin D deficiency increases vulnerability to canagliflozin-induced adverse

effects on 1,25-dihydroxyvitamin D and PTH

Zhinous Shahidzadeh Yazdi, Elizabeth A. Streeten, Hilary B. Whitlatch, May E. Montasser,

Amber L. Beitelshees, Simeon I. Taylor

Department of Medicine

Division of Endocrinology, Diabetes, and Nutrition

University of Maryland School of Medicine

Baltimore, MD, USA

Keywords: SGLT2 inhibitor, vitamin D, bone, biomarker, and side effect.

Corresponding author: Simeon I. Taylor, MD, PhD, Department of Medicine, HSRFIII, Room
4182, 670 W Baltimore St, University of Maryland School of Medicine, Baltimore, MD 21201,
USA
Telephone number: +1 301 980 5272
Email: staylor2@som.umaryland.edu

Funding: R01DK118942, R01DK118942-02S1, and T32DK098107

Disclosure summary: SIT serves as a consultant for Ionis Pharmaceuticals and receives an
inventor's share of royalties from NIDDK for metreleptin as a treatment for generalized
lipodystrophy. EAS, MEM, and HBW receive research support (unrelated to this project) from
the Regeneron Genetics Center. ZSY and ALB have no conflicts to disclose.

Abstract

Context. Canagliflozin has been reported to increase the risk of bone fracture – possibly

mediated by decreasing 1,25-dihydroxyvitamin D [$1,25(OH)_2D$] and increasing PTH.

Objective. To investigate whether baseline vitamin D (VitD) deficiency renders individuals

vulnerable to this adverse effect and whether VitD3 supplementation is protective.

Design. This study had a paired design comparing individual participants before and after VitD3

supplementation.

Setting. Community-based outpatient.

Patients. 11 VitD deficient (25-hydroxyvitamin D [25(OH)D] ≤ 20 ng/mL) individuals recruited

from the Amish population in Lancaster PA.

Interventions. Participants underwent two canagliflozin challenge protocols (300 mg daily for five

days): the first before and the second after VitD3 supplementation. In the VitD3 supplementation

protocol, participants received VitD3 supplementation (50,000 IU once or twice a week

depending on BMI for 4-6 weeks) to achieve 25(OH)D ≥ 30 ng/mL.

Main Outcome Measures. Two co-primary endpoints were identified: effects of VitD3

supplementation on canagliflozin-induced changes in $1,25(OH)_2D$ and PTH. Secondary

endpoints included effects of VitD3 supplementation on baseline levels of VitD metabolites and

PTH.

Results. VitD3 supplementation increased mean 25(OH)D from 16.5±1.6 to 44.3±5.5 ng/mL

(p=0.0006) and 24,25-dihydroxyvitamin D [$24,25(OH)_2D$] from 1.0±0.1 to 4.3±0.6 ng/mL

(p=0.0002). Mean $1,25(OH)_2D$ and PTH were unchanged. VitD3 supplementation decreased the

magnitude of canagliflozin-induced changes in $1,25(OH)_2D$ (from -31.3%±4.7% to -9.3%±8.3%;

p=0.04) and PTH (from +36.2%±6.2% to +9.7%±3.7%; p=0.005).

Conclusions. VitD deficiency rendered individuals more vulnerable to adverse effects of

canagliflozin on biomarkers associated with bone health. VitD3 supplementation was protective

against canagliflozin's short-term adverse effects on $1,25(OH)_2D$ and PTH.

Introduction

SGLT2 inhibitors display a very attractive clinical profile including glucose-lowering efficacy, decreased risk of major adverse cardiovascular events, decreased risk of hospitalization for heart failure, and slowed progression of kidney disease (1-7). However, there is controversy about the bone related side effects of these drugs. Canagliflozin and dapagliflozin have been reported to increase the risk of bone fracture in some (4, 8) but not all (1, 2, 9) clinical trials. Canagliflozin has also been reported to accelerate loss of bone mineral density (10), which may contribute to the increased risk of bone fracture – possibly by decreasing 1,25-dihydroxyvitamin D [1,25(OH)$_2$D] and increasing PTH (11, 12). Canagliflozin, dapagliflozin, and empagliflozin have all been reported to increase serum phosphorus and thereby trigger the FGF23/1,25(OH)$_2$D/PTH axis (11, 13, 14), which represent potential endocrine mechanisms that may mediate adverse effects of SGLT2 inhibitors on bone health (12, 15). Although data from most studies assessing the risk of bone fracture in SGLT2 inhibitor users are reassuring, there are important limitations in some of these studies.

Vitamin D (VitD) status is a contributory determinant of bone health (16-18). Low 1,25(OH)$_2$D and increased levels of PTH can work in tandem to negatively impact bone health (19). Moreover, canagliflozin, dapagliflozin, and empagliflozin have all been reported to decrease 1,25(OH)$_2$D and increase PTH (11, 13, 14). We hypothesized that low availability of the substrate 25-hydroxyvitamin D [25(OH)D] might render individuals more vulnerable to the canagliflozin-induced decrease in 1α-hydroxylase activity. In other words, VitD deficiency might amplify the canagliflozin-induced changes in 1,25(OH)$_2$D and PTH. Accordingly, we investigated potential nutrient-drug interactions between VitD and canagliflozin by conducting a clinical trial in otherwise healthy VitD deficient individuals with baseline total serum level of 25(OH)D ≤ 20 ng/mL in whom

89   we assessed the effects of canagliflozin on $1,25(OH)_2D$, PTH, and FGF23 before and after VitD3

90   supplementation.

Materials and Methods

This study ("VitD Sub-study") was part of our ongoing parent "Genetics of Response to Canagliflozin" clinical trial (clinicaltrials.gov Identifier: NCT02891954) in healthy volunteers recruited from the Old Order Amish population in Lancaster PA, who are non-Hispanic White people. Although participants were excluded from the parent study if their 25(OH)D level was ≤ 20 ng/mL, these VitD deficient individuals were eligible to participate in the VitD Sub-study, which investigated the effects of VitD3 supplements on canagliflozin-triggered changes in the phosphate/FGF23/1,25(OH)$_2$D/PTH axis. Participants were required to meet all other exclusion/inclusion criteria for the parent study as listed below.

Inclusion/exclusion criteria

Inclusion criteria:

- Amish descent
- ≥18 years old
- VitD deficiency at baseline with 25(OH)D ≤ 20 ng/mL
- BMI of 18-40 kg/m^2
- Participants were required to be willing not to alter their regimen of vitamins, nutritional supplements, and over-the-counter medications (with the exception of acetaminophen) during the course of the study. If changes needed to be made, the PI or designated research physician would review and approve changes in order to remain in the study.

Exclusion criteria:

- Known allergy to canagliflozin or VitD3
- History of diabetes and/or HbA1c > 6.5%

115	•	History of medical condition which, in the judgment of the PI or designated research physician, would alter the response to canagliflozin and/or increase the risk to the patient
116		
117	•	History of low trauma bone fractures associated with a history of osteoporosis
118	•	History of a urinary tract infection or genital yeast infection within the past year
119	•	Current treatment with diuretics, antihypertensive medication, uric acid lowering medications, or other medication that the investigator judges will make interpretation of the results difficult
120		
121		
122	•	Significant debiliatating chronic cardiac, hepatic, pulmonary, or renal disease or other diseases that the investigator judges will make interpretation of the results difficult or increase the risk of participation
123		
124		
125	•	Seizure disorder
126	•	Unhealed foot ulcer
127	•	Pregnant by self-report, positive urine pregnancy test done by nursing staff prior to administration of study drug, or known pregnancy within three months of the start of the study
128		
129		
130	•	Currently breast feeding or breast feeding within three months of the start of the study
131	•	Estimated glomerular filtration rate <60 mL/min/1.73 m^2
132	•	Liver function tests >2 times the upper limit of normal
133	•	Hematocrit <35%
134	•	Abnormal TSH
135	•	Urinalysis suggesting urinary tract infection.

Study population

Otherwise, healthy participants were recruited from the Old Order Amish population in Lancaster

PA, as described in the Results ("Disposition and baseline characteristics of participants" section).

Rationale for enrolling individuals with 25(OH)D ≤ 20 ng/mL

The Endocrine Society's guidelines proposed thresholds of 25(OH)D ≤20 ng/mL (≤ 50 nmol/L) for

VitD deficiency, 21-29 ng/mL (51-74 nmol/L) for VitD insufficiency, and ≥ 30 ng/mL (≥ 75 nmol/L)

for VitD sufficiency. We applied The Endocrine Society's definition of VitD deficiency (≤ 20 ng/mL)

as a key inclusion criterion and applied the definition of VitD sufficiency (≥ 30 ng/mL) as the

criterion for adequacy of VitD3 supplementation.

Canagliflozin challenge protocol

Research participants underwent two canagliflozin challenges. Canagliflozin challenge #1 was

conducted when participants were judged to be VitD deficient [25(OH)D ≤ 20 ng/mL]; canagliflozin

challenge #2 was conducted after participants had received VitD3 supplementation and 25(OH)D

levels were ≥ 30 ng/mL. Research nurses conducted home visits at approximately 7am to obtain

blood samples on Day 1 (pre-dose), Day 3 (after 2 doses of canagliflozin), and Day 6 (after 5

doses of canagliflozin). To minimize the impact of circadian rhythms, blood samples were

obtained at the time of these home visits. Canagliflozin (300 mg/d, p.o.) was administered every

morning shortly after obtaining 7am blood samples on Days 1-5. Blau et al. (11) previously

reported that maximum responses to canagliflozin in a 5-day study were observed after 48 hours

(Day 3) for serum phosphorus, FGF23, and 1,25(OH)$_2$D and after 120 hours (Day 6) for PTH.

These previously published observations (11) provided the rationale for measuring these

parameters on Days 1, 3, and 6 and focusing on canagliflozin-triggered changes on Day 6 for

PTH, and on Day 3 for the other biomarkers.

VitD3 supplementation protocol

VitD deficient participants received VitD3 capsules (Bio-Tech Pharmacal; Fayetteville, AR) in

doses of either 50,000 IU per week (for participants with BMI <30 kg/m^2) or 50,000 IU twice a

week (for participants with BMI ≥30 kg/m^2) for four weeks, followed by a second 25(OH)D

measurement a week after completion of the four-week VitD3 supplementation protocol. If

screening 25(OH)D levels (Quest Diagnostics) were <30 ng/mL, high-dose VitD3

supplementation was continued. Once screening 25(OH)D (Quest Diagnostics) levels were

confirmed to be ≥30 ng/mL, maintenance-dose VitD3 supplementation (1000 IU/d or 2000 IU/d

for BMI <30 kg/m^2 or BMI ≥30 kg/m^2, respectively) was continued at least until completion of

canagliflozin challenge #2.

Clinical chemistry

Blood samples obtained at home visits were collected in test tubes as appropriate for each assay:

EDTA anticoagulant (purple top tube) for measurement of plasma samples and red top tube for

collecting serum samples. After placing tubes on ice, blood samples were transported to the

clinical laboratory at the Amish Research Clinic (maximum transport time, 2 hours). After

centrifugation (3300 rpm for 10 min), plasma/serum was sent on the same day to a laboratory for

assay.

Most laboratory assays were conducted at Quest Diagnostics except when stated otherwise.

Some specialized assays were conducted at the Cytokine Core Laboratory of the University of

Maryland School of Medicine using the following kits: intact human FGF23 (iFGF23) and C-

terminal FGF23 (cFGF23) (Quidel Corporation, San Diego, CA).

For consistency with the methods of the parent study, screening serum levels of 25(OH)D and all

serum levels of 1,25(OH)$_2$D were assayed at Quest Diagnostics. As requested by the funding

agencies (NIH Office of Dietary Supplements and National Institute of Diabetes and Digestive and

Kidney Disease; National Institutes of Health), plasma levels of 25(OH)D and 24,25-

dihydroxyvitamin D [24,25(OH)$_2$D] were measured by Heartland Assays (Ames, IA). Quest's

25(OH)D levels were used to determine eligibility for the VitD Sub-study. Heartland's 25(OH)D

levels were used to assess the response to VitD3 supplements. All assays of hydroxylated VitD

metabolites provided separate measurements for both VitD2 and VitD3 derivatives. We report

25(OH)D and 24,25(OH)$_2$D in ng/mL and 1,25(OH)$_2$D in pg/mL. One can use the conversion factor

of 2.49 from ng/mL to nmol/L for 25(OH)D.

Co-primary endpoints

We established two co-primary outcomes to investigate whether VitD3 supplements mitigate the

canagliflozin-induced changes in biomarkers related to bone health: effect of VitD3 supplements

on canagliflozin-induced decrease in 1,25(OH)$_2$D and effect of VitD3 supplements on

canagliflozin-induced increase in PTH.

Secondary endpoints

Our secondary endpoints included 2 categories: (a) canagliflozin-dependent endpoints and (b)

canagliflozin-independent endpoints.

Canagliflozin-dependent endpoints

(a) Effect of canagliflozin on serum phosphorus (Day 3 versus Day 1) before and after VitD3

supplementation; we compared the canagliflozin-induced changes in serum phosphorus before

versus after VitD3 supplementation.

(b) Effect of canagliflozin on serum calcium (Day 3 versus Day 1) before and after VitD3

supplementation; we compared the canagliflozin-induced changes in serum calcium before

versus after VitD3 supplementation.

(c) Effect of canagliflozin on intact FGF23 [iFGF23] (Day 3 versus Day 1) before and after VitD3

supplementation; we compared the canagliflozin-induced changes in iFGF23 before versus after

VitD3 supplementation.

(d) Effect of canagliflozin on C-terminal FGF23 [cFGF23] (Day 3 versus Day1) before and after VitD3 supplementation; we compared the canagliflozin-induced changes in cFGF23 before versus after VitD3 supplementation.

We used paired t-tests for all analyses.

Canagliflozin-independent endpoints

(a) We compared baseline values of plasma $1,25(OH)_2D$ and $24,25(OH)_2D$ before versus after VitD3 supplementation, independent of canagliflozin.

(b) We compared baseline values of serum phosphorus, calcium, plasma PTH, iFGF23, and cFGF23, before versus after VitD3 supplementation, independent of canagliflozin.

Statistical analysis

We applied a stepwise approach to test statistical significance for our primary endpoints:

1. Paired t-test to confirm that VitD3 supplements increased levels of 25(OH)D.

2. Paired t-test to confirm that canagliflozin decreased levels of $1,25(OH)_2D$ in VitD deficient patients. This comparison was based on ratios of $1,25(OH)_2D$ levels on Day 3:Day 1.

3. Paired t-test to investigate whether VitD3 supplements affected the magnitude of canagliflozin-induced decrease in levels of $1,25(OH)_2D$. This assessment was based on a comparison of ratios of $1,25(OH)_2D$ levels on Day 3:Day 1 in VitD deficient patients versus the same ratio in patients who had received VitD3 supplements.

4. Paired t-test to confirm that canagliflozin increased levels of PTH in VitD deficient patients. This comparison was based on ratios of PTH levels on Day 6:Day 1.

5. Paired t-test to investigate whether VitD3 supplements affected the magnitude of canagliflozin-induced increase in levels of PTH. This assessment was based on a

comparison of ratios of PTH levels on Day 6:Day 1 in VitD deficient patients versus the

same ratio in patients who had received VitD3 supplements.

These statistical tests of significance were conducted in sequential fashion. If a statistical analysis

yielded a statistically significant result (p-value ≤ 0.05), then we were permitted to proceed to the

next comparison. If a statistical analysis did not yield a statistically significant result (p-value >

0.05), then the subsequent analyses would be viewed as exploratory. A two-sided p-value ≤ 0.05

(paired t-tests) was taken as the threshold for statistical significance without any correction for

multiple comparisons. For two analyses (Figs. 2A and 4A), We pooled all the data before and

after VitD3 supplementation; in those two analyses, we calculated values for Pearson or

Spearman correlation coefficients (Excel).

Study approval

The Clinical Trial Protocol was approved by the University of Maryland Baltimore Institutional

Review Board (FWA00007145). Informed consent was obtained from all participants.

Results

Disposition and baseline characteristics of participants

Research participants (N=11) were recruited from the Old Order Amish population in Lancaster PA. The disposition of participants is summarized in a CONSORT diagram (Fig. 1). The parent study ("Genetics of the Response to Canagliflozin") identified 24 individuals with 25(OH)D levels ≤ 20 ng/mL between September, 2019 and July, 2021 who were assessed for eligibility for this VitD sub-study. Nineteen participants gave informed consent and were enrolled in the study. Three participants withdrew consent and 16 completed the first canagliflozin challenge test. Two participants were subsequently withdrawn because they did not achieve 25(OH)D levels ≥ 30 ng/mL in response to VitD3 supplements. Three participants withdrew because of interruptions associated with the COVID-19 pandemic. Eleven participants (7 males and 4 females) completed the entire study. Baseline characteristics and demographics are shown in Table 1. Our study population had a mean age of 48.4 ± 3.2 years and a mean BMI of 29.0 ± 1.2 kg/m^2 at baseline.

Effect of VitD3 supplementation on VitD metabolites and mineral metabolism biomarkers

Clinical trial participants were required to have baseline 25(OH)D levels ≤ 20 ng/mL at the time of screening. For consistency with the methods in the parent study ("Genetics of Response to Canagliflozin"), we used a 25(OH)D assay provided by Quest Diagnostics to screen people for eligibility for both the parent study and the VitD sub-study. To maximize the ability to compare with data on 24,25(OH)$_2$D levels, Heartland Assays provided assays for both 25(OH)D and 24,25(OH)$_2$D to analyze blood samples collected during the canagliflozin challenge tests conducted as part of the VitD sub-study. Figure 2 demonstrates that the two assays (i.e., Quest Diagnostics and Heartland Assays) yielded essentially identical results for mean levels of 25(OH)D and exhibited good overall correlation for individual participants (r=0.9; p=10^{-8}). Blood samples for screening were collected in advance of (typically one or two weeks before) the

canagliflozin challenge tests. The comparability between the two sets of data is reassuring given

that the samples were not collected on the same day and the assays were conducted in different

laboratories.

Supplementation with VitD3 significantly increased mean (± SEM) 25(OH)D by 2.7-fold from 16.5

± 1.6 ng/mL in the VitD deficient state to 44.3 ± 5.5 ng/mL after VitD3 supplementation (p=0.0006)

(Fig. 3A). Mean $24,25(OH)_2D$ levels increased 4.3-fold from 1.0 ± 0.1 ng/mL in the VitD deficient

state to 4.3 ± 0.6 ng/mL after VitD3 supplementation (p=0.0002) (Fig. 3B). In contrast, mean

plasma levels of $1,25(OH)_2D$ in the VitD deficient state did not change significantly after VitD3

supplementation (43.8 ± 3.6 versus 44.9 ± 4.1 pg/mL; p=0.7) (Fig. 3C).

Mean levels of plasma PTH and FGF23 did not change in response to VitD3 supplementation.

Mean levels of serum phosphorus, serum calcium, plasma PTH, and plasma FGF23 (intact and

C-terminal) were within the normal range when participants were VitD deficient and did not

change with VitD3 supplementation (Table 2). 25(OH)D levels spanned an 8-fold range (11 - 88

ng/mL); PTH levels ranged between 22-85 pg/mL (Fig. 4A) and were not significantly correlated

with 25(OH)D levels in an analysis of our pooled data (Fig. 4A). Furthermore, we did not observe

a consistent effect of VitD3 supplementation on PTH levels of individual participants (Fig. 4B).

Baseline levels of PTH were closely correlated with PTH levels after administration of VitD3

supplements (r=0.83; p=0.001) (Fig. 4C). Although the published literature provides extensive

evidence that VitD deficiency can cause secondary hyperparathyroidism, It is suggested that

secondary hyperparathyroidism is most commonly triggered in the context of more severe VitD

deficiency [e.g., total 25(OH)D levels <10-12 ng/mL] (20-23).

Interaction between baseline VitD status and pharmacodynamic responses to canagliflozin.

When participants were VitD deficient, canagliflozin significantly decreased mean (± SEM)

$1,25(OH)_2D$ levels by 31.3% ± 4.7%: from 43.8 ± 3.6 pg/mL on Day 1 to 29.1 ± 2.3 pg/mL on Day

3 (p=0.0003). In contrast, after participants had received VitD3 supplements, canagliflozin

induced a smaller, non-significant 9.3% ± 8.3% decrease in mean levels of $1,25(OH)_2D$: from 45

± 4.1 pg/mL on Day 1 to 41 ± 5.4 pg/mL on Day 3 (p=0.3). Thus, VitD3 supplementation provided

statistically significant protection from the adverse effect of canagliflozin to decrease mean

plasma levels of $1,25(OH)_2D$ (p=0.04) (Fig. 5A). Similarly, when participants were VitD deficient,

canagliflozin triggered a significant 36.2% ± 6.2% increase in mean levels of PTH: from 47.5 ±

4.8 pg/mL on Day 1 to 58.5 ± 5.4 pg/mL on Day 6 (0.0009). After participants had received VitD3

supplements, canagliflozin induced a smaller mean increase (9.7% ± 3.7%) in mean plasma

levels of PTH: from 48.4 ± 5.1 pg/mL on Day 1 to 53.3 ± 6.1 pg/mL on Day 6 (p=0.02). Therefore,

VitD3 supplementation significantly decreased the magnitude of the canagliflozin-induced

increase in mean levels of PTH (p=0.005) (Fig. 5B).

Effect of VitD on canagliflozin-induced changes in secondary endpoints

In the VitD deficient state, canagliflozin significantly increased mean (± SEM) serum phosphorus

on Day 3 versus Day 1 (3.6 ± 0.1 mg/dL to 3.8 ± 0.1 mg/dL; p=0.007) with the mean percentage

change of +7.0% ± 2.1% (Table 3). In contrast, the percentage change was -0.2% ± 3.8% after

VitD3 supplementation (3.72 ± 0.1 mg/dL to 3.69 ± 0.2 mg/dL; p=0.8). While the canagliflozin-

induced change in serum phosphorus was numerically larger in the VitD deficient state, the impact

of VitD3 supplementation did not achieve statistical significance (p=0.1).

Canagliflozin increased iFGF23 by 21.3% when participants were VitD deficient (from 43.1 ± 5.6

pg/mL on Day 1 to 50.9 ± 6.2 pg/mL on Day 3; p=0.009) and by 17.4% after VitD3 supplementation

(from 46.5 ± 5.0 pg/mL on Day 1 to 54.2 ± 8.9 pg/mL on Day 3; p=0.2) (Table 3). Canagliflozin

also increased cFGF23 by 19.2% (from 76.5 ± 13.2 RU/mL on Day 1 to 86.4 ± 13.2 RU/mL on

Day 3; p=0.08) before VitD3 supplementation and by 15.6% (from 83.0 ± 14.0 RU/mL on Day 1

to 94.3 ± 20.8 RU/mL on Day 3; p=0.3) after VitD3 supplementation (Table 3). $1,25(OH)_2D$ has

been reported to increase the expression of the *FGF23* gene (24). It is possible, therefore, to

envision interactions between VitD and the magnitude of canagliflozin's effect on iFGF23 even

though we did not observe a statistically significant effect of VitD3 supplementation on the

magnitude of the canagliflozin-induced increase in either iFGF23 or cFGF23 – possibly because

the sample size was not sufficiently large.

Discussion

Canagliflozin inhibits SGLT2-mediated sodium and glucose co-transport in the proximal tubule. This increases the electrochemical gradient for Na^+, which in turn increases the drive for sodium-dependent reabsorption of phosphate. The resulting increase in serum phosphorus is associated with an increase in plasma FGF23, which in turn decreases expression of 1α-hydroxylase in the kidney and decreases plasma levels of $1,25(OH)_2D$. This decreases intestinal absorption of dietary calcium, which ultimately increases PTH secretion (11, 12, 15). Low levels of $1,25(OH)_2D$ and secondary hyperparathyroidism work in tandem to adversely impact bone health (19). We hypothesized that low levels of 25(OH)D [the biosynthetic precursor of $1,25(OH)_2D$] would increase vulnerability to the pathophysiological impact of the canagliflozin-induced decrease in 1α-hydroxylase. This clinical trial provides evidence confirming that hypothesis. Furthermore, VitD3 supplementation decreased the magnitude of drug-induced adverse changes in $1,25(OH)_2D$ and PTH.

Several lines of evidence suggest that SGLT2 inhibitors exert adverse effects on bone health. Canagliflozin and dapagliflozin have been reported to increase the risk of treatment-emergent bone fractures (4, 8). Canagliflozin has also been reported to accelerate the loss of bone mineral density (10). Furthermore, canagliflozin, dapagliflozin, and empagliflozin have been reported to trigger the phosphate/FGF23/$1,25(OH)_2D$/PTH axis, which represents a biologically plausible mechanism whereby SGLT2 inhibitors might exert adverse effect on bone health (12-14). On the other hand, several studies did not observe an increased risk of bone fractures (1-3, 25-28). It is not clear how to explain this discrepancy, but it is possible that differences in patient populations might contribute. Our current clinical trial demonstrated that VitD deficiency increased vulnerability of participants to the adverse effects of canagliflozin on pharmacodynamic biomarkers – specifically $1,25(OH)_2D$ and PTH. It is possible that differences in VitD status might have

contributed to discrepant outcomes among clinical trials with respect to the risk of bone fracture

and may help to explain why treatment with SGLT2 inhibitors was associated with increased risk

of bone fractures in some studies but not in others. Interestingly, mean baseline plasma 25(OH)D

levels were relatively low in the studies demonstrating that dapagliflozin and empagliflozin trigger

decreased levels of $1,25(OH)_2D$ and increased levels of PTH. Specifically, 25(OH)D levels were

18 ng/mL in the study with dapagliflozin (13) and 15.6 ng/mL in the study with empagliflozin (14).

It is also possible that patients' baseline bone mineral densities differed among the clinical trials.

For example, patients with higher baseline bone mineral density would be predicted to have

longer lag times before they lose sufficient bone mineral density to be at risk for fragility fractures.

Finally, follow-up times varied among the studies (4, 29-32). As shown in studies with other drugs

(e.g., thiazolidinediones), long lag times may be required before an increase in the risk of fracture

becomes apparent (33). Some studies (especially studies of "real world" experience) were

relatively short (e.g., one year of follow-up) (32); it is likely that longer follow-up might be required

to detect an increased risk of bone fracture.

Study design: strengths and limitations

To maximize statistical power, the study was designed as a crossover study so that we could

conduct paired statistical analyses comparing data for individual participants when they were VitD

deficient to data obtained after participants received VitD3 supplements. Because our principal

endpoints [levels of serum phosphorus, plasma FGF23, plasma $1,25(OH)_2D$, and PTH] all

demonstrate circadian rhythms (11, 34, 35), our study design assured that all blood samples

would be obtained at the same time of day (~7:00 am) to minimize the impact of variation due to

circadian rhythms. Our assays of VitD metabolites (provided by Quest and Heartland)

distinguished between VitD2 and VitD3. We did not detect metabolites of VitD2 in any of our

participants. The absence of VitD2 metabolites avoided potential issues related to differences

between VitD2 and VitD3 (36), which simplified data interpretation. Our preliminary data guided

us to obtain blood samples at the times when canagliflozin exerted maximal effects on the study's

endpoints (11): Day 6 for PTH and Day 3 for serum phosphorus, FGF23, and $1,25(OH)_2D$.

Furthermore, we obtained data on levels of 25(OH)D and $24,25(OH)_2D$ from a company

(Heartland Assays) that specializes in applying liquid chromatography and mass spectroscopy

(LC/MS/LC) to conduct scientifically rigorous and reproducible assays of VitD metabolites.

In addition to these strengths of our study design, the study had important limitations: (a) In our

original study design, we proposed to study 25 participants. The impact of the COVID-19

pandemic necessitated a decrease in the sample size to 11 participants. (b) The study's outcomes

were all based on biomarkers rather than hard clinical outcomes (e.g., fractures). (c) Since none

of our participants exhibited 25(OH)D levels below 10 ng/mL, our analyses were limited to

individuals with mild or moderate degrees of VitD deficiency; we relied on the published literature

for information on individuals with more severe VitD deficiency.

Limitations associated with the relatively small sample size are partially mitigated by our efforts

to assure scientific rigor and reproducibility of the primary data. The crossover study design

increased statistical power by enabling paired statistical comparisons. As a result, many of our

analyses [e.g., VitD-induced changes in 25(OH)D and $24,25(OH)_2D$] demonstrated high levels of

statistical significance. Other analyses [e.g., comparisons for $1,25(OH)_2D$ and PTH] demonstrated

essentially identical mean values before and after VitD3 supplementation.

Conclusions

Taken together, our observations suggest that physicians should consider screening for VitD

deficiency in patients receiving SGLT2 inhibitors. We acknowledge that our study does not

provide definitive proof that VitD supplementation would decrease the risk of bone fracture in

patients receiving SGLT2 inhibitors. Indeed, there is not yet a consensus that treatment with

SGLT2 inhibitors increases the risk of fracture. Unfortunately, it seems quite unlikely that funding

will be available to conduct a large and expensive clinical trial for a rigorous test of the hypothesis

that VitD supplements would protect against adverse effects of SGLT2 inhibitors on bone health.

Thus, especially in light of the relative safety and low cost of over-the-counter VitD supplements,

biomarker data may be sufficient to justify prescription of VitD supplements in this context.

Acknowledgements

The authors gratefully acknowledge research funding provided by NIDDK and the NIH Office of Dietary Supplements: R01DK118942, R01DK118942-02S1, and T32DK098107. We are also grateful to the staff at University of Maryland's Amish Research Clinic and to the members of the Old Order Amish Community in Lancaster, PA who participated in this clinical trial.

Data Availability

Primary data will be made available upon request to qualified academic investigators for research purposes under a Data Transfer Agreement to protect research participants' confidential information.

412 **Table 1. Participants' demographics and baseline characteristics (mean ± SEM).**

	Total	Females	Males
N	11	4	7
Age (years)	48.0 ± 3.2	49 ± 5.3	48 ± 4.4
Height (cm)	170 ± 2.4	163.3 ± 2.0	173.9 ± 2.6
Weight (kg)	81.7 ± 3.1	77.5 ± 5.4	84.5 ± 3.6
BMI (kg/m^2)	29.0 ± 1.2	28.8 ± 1.5	28.9 ± 1.7
HbA1c (%)	5.4 ± 0.08	5.5 ± 0.1	5.3 ± 3.08
eGFR (mL/min/1.73 m^2)	98.5 ± 4.0	100 ± 8.6	98.6 ± 3.5

415 **Table 2. Effect of VitD3 supplementation on levels of serum phosphorus, serum calcium, plasma**

416 **PTH, plasma iFGF23, and cFGF23.** The Table summarizes mean levels (±SEM) when participants

417 (N=11) were VitD deficient and after they received VitD3 supplements. p-values were calculated using

418 Student's t-test for paired comparisons.

	VitD deficient state	VitD sufficient state	p-values
Serum Phosphorus (mg/mL)	3.6 ± 0.1	3.7 ± 0.1	0.3
Serum Calcium (mg/mL)	9.3 ± 0.1	9.2 ± 0.1	0.4
Plasma PTH (pg/mL)	47.5 ± 4.8	48.4 ± 5.1	0.8
Plasma iFGF23 (pg/mL)	43.1 ± 5.6	46.5 ± 5.0	0.2
Plasma cFGF23 (RU/mL)	76.5 ± 13.2	83.0 ± 14.0	0.2

421 **Table 3. Effects of VitD3 supplementation on canagliflozin-induced changes in secondary**

422 **endpoints.** The Table summarizes canagliflozin-induced changes (mean ± SEM) in serum phosphorus,

423 serum calcium, plasma iFGF23, and cFGF23. p-values were calculated using Student's t-test for paired

424 comparisons for the effects of canagliflozin.

		VitD deficient state	VitD sufficient state	p-values
Canagliflozin-induced percent change in secondary endpoints	Phosphorus	+7.0 ± 2.1	-0.2 ± 3.8	0.1
	Calcium	+0.9 ± 1.3	+0.5 ± 1.1	0.9
	iFGF23	+21.3 ± 8.5	+17.5 ± 13.3	0.7
	cFGF23	+19.2 ± 7.3	+15.6 ± 13.0	0.7

Legends to Figures

Figure 1. CONSORT diagram. The figure summarizes recruitment and disposition of research participants in this clinical trial.

Figure 2. Comparison of two commercial assays for 25(OH)D in participants in our clinical trial: Quest Diagnostics and Heartland Assays. Panel A plots individual levels of total 25(OH)D measured by Heartland Assays versus individual levels of total 25(OH)D measured by Quest Diagnostics. Panel B compares measurements of 25(OH)D levels in the two assays at the level of individual participants when they were VitD deficient (gray circles) and after they received VitD3 supplements (black squares). Panel C shows means ± SEM for total 25(OH)D levels in VitD-deficient participants before (gray bars) and after receiving VitD3 supplements (black bars) as measured by Quest Diagnostics and Heartland Assays.

Figure 3. Effect of VitD3 supplementation on mean baseline plasma levels of hydroxylated VitD metabolites: 25(OH)D (panel A), 24,25 (OH)$_2$D (panel B), and 1,25(OH)$_2$D (panel C). Mean (±SEM) baseline plasma levels of VitD metabolites were measured when research participants (N=11) were VitD deficient (black bars) and again after participants received VitD3 supplements (gray bars). Mean levels of VitD metabolites are indicated in white font. Error bars represent standard errors of the mean. Levels of statistical significance are indicated by p-values using Student's t-test for paired comparisons.

Figure 4. Effect of VitD status on PTH levels. Panel A presents pooled data for individual participants when they were VitD deficient (gray circles) and after they received VitD3 supplements (black circles). PTH levels are plotted as a function of total plasma levels of 25(OH)D. Panel B plots levels of PTH when individual participants were VitD deficient versus plasma levels of PTH after they received VitD3 supplements. Panel C plots PTH levels after participants received VitD3 supplements as a function of PTH levels when participants were

VitD deficient. The text boxes in panels A and C present the equations for the linear regression

lines and the Pearson correlation coefficients.

Figure 5. Canagliflozin-induced change in primary endpoints: plasma 1,25(OH)$_2$D and

PTH. Research participants (N=11) underwent two canagliflozin challenge protocols: the first

when they were VitD deficient (black bars) and the second after receiving VitD supplements

(gray bars). Mean canagliflozin-induced percentage changes (depicted in white font) are shown

for Day 6 for PTH and for Day 3 for 1,25(OH)$_2$D. Error bars represent standard errors of the

mean. p-values were calculated using Student's t-test for paired comparisons for the effects of

VitD.

References

1. Perkovic V, Jardine MJ, Neal B, Bompoint S, Heerspink HJL, Charytan DM, et al. Canagliflozin and Renal Outcomes in Type 2 Diabetes and Nephropathy. *N Engl J Med.* 2019;380(24):2295-306.

2. Wiviott SD, Raz I, Bonaca MP, Mosenzon O, Kato ET, Cahn A, et al. Dapagliflozin and Cardiovascular Outcomes in Type 2 Diabetes. *N Engl J Med.* 2019;380(4):347-57.

3. Zinman B, Wanner C, Lachin JM, Fitchett D, Bluhmki E, Hantel S, et al. Empagliflozin, Cardiovascular Outcomes, and Mortality in Type 2 Diabetes. *N Engl J Med.* 2015;373(22):2117-28.

4. Watts NB, Bilezikian JP, Usiskin K, Edwards R, Desai M, Law G, et al. Effects of Canagliflozin on Fracture Risk in Patients With Type 2 Diabetes Mellitus. *J Clin Endocrinol Metab.* 2016;101(1):157-66.

5. Petrie MC, Verma S, Docherty KF, Inzucchi SE, Anand I, Belohlávek J, et al. Effect of Dapagliflozin on Worsening Heart Failure and Cardiovascular Death in Patients With Heart Failure With and Without Diabetes. *Jama.* 2020;323(14):1353-68.

6. Packer M, Anker SD, Butler J, Filippatos G, Pocock SJ, Carson P, et al. Cardiovascular and Renal Outcomes with Empagliflozin in Heart Failure. *N Engl J Med.* 2020;383(15):1413-24.

7. Filippatos G, Butler J, Farmakis D, Zannad F, Ofstad AP, Ferreira JP, et al. Empagliflozin for Heart Failure With Preserved Left Ventricular Ejection Fraction With and Without Diabetes. *Circulation.* 2022;146(9):676-86.

8. Kohan DE, Fioretto P, Tang W, and List JF. Long-term study of patients with type 2 diabetes and moderate renal impairment shows that dapagliflozin reduces weight and blood pressure but does not improve glycemic control. *Kidney Int.* 2014;85(4):962-71.

484   9.   Sarafidis PA, and Tsapas A. Empagliflozin, Cardiovascular Outcomes, and Mortality in

485   Type 2 Diabetes. *N Engl J Med.* 2016;374(11):1092.

486   10.   Bilezikian JP, Watts NB, Usiskin K, Polidori D, Fung A, Sullivan D, et al. Evaluation of

487   Bone Mineral Density and Bone Biomarkers in Patients With Type 2 Diabetes Treated

488   With Canagliflozin. *J Clin Endocrinol Metab.* 2016;101(1):44-51.

489   11.   Blau JE, Bauman V, Conway EM, Piaggi P, Walter MF, Wright EC, et al. Canagliflozin

490   triggers the FGF23/1,25-dihydroxyvitamin D/PTH axis in healthy volunteers in a

491   randomized crossover study. *JCI Insight.* 2018;3(8).

492   12.   Blau JE, and Taylor SI. Adverse effects of SGLT2 inhibitors on bone health. *Nat Rev*

493   *Nephrol.* 2018;14(8):473-4.

494   13.   de Jong MA, Petrykiv SI, Laverman GD, van Herwaarden AE, de Zeeuw D, Bakker SJL,

495   et al. Effects of Dapagliflozin on Circulating Markers of Phosphate Homeostasis. *Clin J*

496   *Am Soc Nephrol.* 2019;14(1):66-73.

497   14.   Rau M, Thiele K, Hartmann NK, Möllmann J, Wied S, Hohl M, et al. Effects of empagliflozin

498   on markers of calcium and phosphate homeostasis in patients with type 2 diabetes - Data

499   from a randomized, placebo-controlled study. *Bone Rep.* 2022;16:101175.

500   15.   Taylor SI, Blau JE, and Rother KI. Possible adverse effects of SGLT2 inhibitors on bone.

501   *Lancet Diabetes Endocrinol.* 2015;3(1):8-10.

502   16.   Carpenter KJ. Harriette Chick and the problem of rickets. *J Nutr.* 2008;138(5):827-32.

503   17.   Chick HD, E.G.; Hume, M.; Mackay, H.M.M.; Smith, H.H.; Wimberger, H. The aetiology of

504   rickets in infants: Prophylactic and curative observations at the Vienna University

505   Kinderklinik. *The Lancet.* 1922:7-11.

506   18.   Rajakumar K. Vitamin D, cod-liver oil, sunlight, and rickets: a historical perspective.

507   *Pediatrics.* 2003;112(2):e132-5.

508   19.   Rodríguez-Ortiz ME, and Rodríguez M. Recent advances in understanding and managing

509   secondary hyperparathyroidism in chronic kidney disease. *F1000Res.* 2020;9.

510   20.   Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, et al.

511   Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society

512   clinical practice guideline. *J Clin Endocrinol Metab.* 2011;96(7):1911-30.

513   21.   Jesudason D, Need AG, Horowitz M, O'Loughlin PD, Morris HA, and Nordin BE.

514   Relationship between serum 25-hydroxyvitamin D and bone resorption markers in vitamin

515   D insufficiency. *Bone.* 2002;31(5):626-30.

516   22.   Rosen CJ, Abrams SA, Aloia JF, Brannon PM, Clinton SK, Durazo-Arvizu RA, et al. IOM

517   committee members respond to Endocrine Society vitamin D guideline. *J Clin Endocrinol*

518   *Metab.* 2012;97(4):1146-52.

519   23.   Souberbielle JC, Cormier C, Kindermans C, Gao P, Cantor T, Forette F, et al. Vitamin D

520   status and redefining serum parathyroid hormone reference range in the elderly. *J Clin*

521   *Endocrinol Metab.* 2001;86(7):3086-90.

522   24.   Kaneko I, Saini RK, Griffin KP, Whitfield GK, Haussler MR, and Jurutka PW. FGF23 gene

523   regulation by 1,25-dihydroxyvitamin D: opposing effects in adipocytes and osteocytes. *J*

524   *Endocrinol.* 2015;226(3):155-66.

525   25.   Fralick M, Kim SC, Schneeweiss S, Kim D, Redelmeier DA, and Patorno E. Fracture Risk

526   After Initiation of Use of Canagliflozin. *Ann Intern Med.* 2019;171(1):80.

527   26.   Ljunggren Ö, Bolinder J, Johansson L, Wilding J, Langkilde AM, Sjöström CD, et al.

528   Dapagliflozin has no effect on markers of bone formation and resorption or bone mineral

529   density in patients with inadequately controlled type 2 diabetes mellitus on metformin.

530   *Diabetes Obes Metab.* 2012;14(11):990-9.

27.   Monteiro P, Bergenstal RM, Toural E, Inzucchi SE, Zinman B, Hantel S, et al. Efficacy and

safety of empagliflozin in older patients in the EMPA-REG OUTCOME® trial. *Age Ageing.*

2019;48(6):859-66.

28.   Zhuo M, Hawley CE, Paik JM, Bessette LG, Wexler DJ, Kim DH, et al. Association of

Sodium-Glucose Cotransporter-2 Inhibitors With Fracture Risk in Older Adults With Type

2 Diabetes. *JAMA Netw Open.* 2021;4(10):e2130762.

29.   Adimadhyam S, Lee TA, Calip GS, Smith Marsh DE, Layden BT, and Schumock GT.

Sodium-glucose co-transporter 2 inhibitors and the risk of fractures: A propensity score-

matched cohort study. *Pharmacoepidemiol Drug Saf.* 2019;28(12):1629-39.

30.   Han SJ, Ha KH, Lee N, and Kim DJ. Effectiveness and safety of sodium-glucose co-

transporter-2 inhibitors compared with dipeptidyl peptidase-4 inhibitors in older adults with

type 2 diabetes: A nationwide population-based study. *Diabetes Obes Metab.*

2021;23(3):682-91.

31.   Sasaki T, Sugawara M, and Fukuda M. Sodium-glucose cotransporter 2 inhibitor-induced

changes in body composition and simultaneous changes in metabolic profile: 52-week

prospective LIGHT (Luseogliflozin: the Components of Weight Loss in Japanese Patients

with Type 2 Diabetes Mellitus) Study. *J Diabetes Investig.* 2019;10(1):108-17.

32.   Ueda P, Svanström H, Melbye M, Eliasson B, Svensson AM, Franzén S, et al. Sodium

glucose cotransporter 2 inhibitors and risk of serious adverse events: nationwide register

based cohort study. *Bmj.* 2018;363:k4365.

33.   Kahn SE, Zinman B, Lachin JM, Haffner SM, Herman WH, Holman RR, et al.

Rosiglitazone-associated fractures in type 2 diabetes: an Analysis from A Diabetes

Outcome Progression Trial (ADOPT). *Diabetes Care.* 2008;31(5):845-51.

34.   el-Hajj Fuleihan G, Klerman EB, Brown EN, Choe Y, Brown EM, and Czeisler CA. The

parathyroid hormone circadian rhythm is truly endogenous--a general clinical research

center study. *J Clin Endocrinol Metab.* 1997;82(1):281-6.

35.   Kawai M, Kinoshita S, Shimba S, Ozono K, and Michigami T. Sympathetic activation

induces skeletal Fgf23 expression in a circadian rhythm-dependent manner. *J Biol Chem.*

2014;289(3):1457-66.

36.   Tripathi A, Ansari M, Dandekar P, and Jain R. Analytical methods for 25-hydroxyvitamin

D: advantages and limitations of the existing assays. *J Nutr Biochem.* 2022;109:109123.

Figure 1. CONSORT diagram

Assessed for Eligibility
9/2019-7/2021
(n = 24)

Declined Participation
(n = 5)

Consented and Enrolled in Clinical Trial
(n =19)

Withdrew Consent
n=3

Completed Canagliflozin Challenge #1
(n =16)

Completed VitD Supplementation Protocol
(n =16)

Withdrew due to Covid-19 Pandemic Interruptions
(n=3)
Did not Attain Required 25(OH)D Sufficient Level
(n = 2)

Completed Canagliflozin Challenge #2
(n = 11)

Analyzed and Reported
(n =11)

Figure 2

Figure 3

A

PTH vs 25(OH)D

y = -0.08x + 50.5
r = 0.1

25(OH)D (ng/mL)

PTH (pg/mL)

B

PTH

PTH (pg/mL)

Vit deficient   VitD sufficient

C

PTH levels (pg/mL) in VitD-sufficient vs VitD-deficient state

y = 0.89x + 6.2
r = 0.83

PTH in VitD-deficient state

PTH in VitD-sufficient state

Figure 5