Correlation between symptoms of attention deficit hyperactivity disorder and life satisfaction in dental students: The mediating role of resilience

Yuwei Sun[1], Lei Miao[2], Siying Chen[3], Zhenya Piao[4] and Chi Tong[5]*

[1]Department of Psychiatry, The First Affiliated Hospital of China Medical University, Shenyang, China.

[2]Department of Periodontics, School and Hospital of Stomatology, China Medical University, Liaoning Provincial Key Laboratory of Oral Diseases, Shenyang, China

[3]Grade 2021, Five-year Program, Clinical Medicine, China Medical University, Shenyang, China

[4]Grade 2021, Five-year Program, Stomatology, China Medical University, Shenyang, China

[5]College of Distance and Continuing Education, China Medical University, Shenyang, China

* Corresponding author

E-mail: ydtongcjy@163.com (CT)

Abstract

Background

Few studies focus on investigating Attention Deficit Hyperactivity Disorder(ADHD) symptoms in dental students. Statistical analysis of demographic factors in this population can assist in supplementing clinical practice. This study was designed to examine the prevalence of ADHD symptoms among dental students, explore the correlation between life satisfaction and ADHD symptoms, and discuss whether resilience mediates the association between life satisfaction and ADHD symptoms.

Methods

This cross-sectional study was conducted in May 2023 at a medical university in China. Self-report questionnaires consisting of the Adult ADHD Self-Report Scale, the Wender Utah Rating Scale, the Life Satisfaction Scale, the Conner-Davidson Resilience Scale, and sociodemographic characteristics. Hierarchical linear regression analyses were used to investigate the effect of ADHD symptoms on life satisfaction, and resampling and asymptotic strategies were used to explore the mediation of resilience.

Results

Totally,291 dental students became final objects. On account of the cutoffs of the scales, 6.87% of the students were likely to have ADHD. There were differences in ADHD prevalence among objects of diverse ages and with varying levels of paternal education. The ADHD asymptomatic group had distinctly higher resilience and life satisfaction degrees than the symptomatic group. Both inattention and hyperactivity

were negatively correlated with resilience and life satisfaction. In the students, resilience was observably positively related to life satisfaction. Resilience plays a mediating role between the two symptoms of ADHD and life satisfaction.

Conclusions

The prevalence of ADHD in Chinese dental students is slightly higher than in general adults. It was revealed that paternal education level was relevant to the morbidity. Hyperactivity was significantly relevant to life satisfaction. Detecting ADHD symptoms earlier is helping improve future dentists' mental health. Resilience intervention programs can enhance dental students' life satisfaction, especially for students with ADHD symptoms.

Introduction

A global study pointed out that mental disorders' burden increased from 1990 to 2019. [1]As one of the most common mental disorders affecting children and adults, attention deficit hyperactivity disorder (ADHD) causes an enormous economic burden worldwide. [2] The main features of ADHD include inattention, hyperactivity, and impulsivity. [3]With the progress of many cohort studies, more and more scholars have recognized that ADHD symptoms usually appear in childhood, and a significant proportion persists into adolescence or adulthood. The prevalence in children is about 5% and is twice as high in boys as in girls. [4]The World Health Organization's World Mental Health Survey points to a prevalence of 2.8% in adults. [5]Chronic symptoms of ADHD have invariably been shown to be associated with

impairment in multiple domains of life, such as occupational achievement, social relationships, substance use, academic performance, and other psychiatric comorbidities. [6–10] Nevertheless, few studies have statistically analyzed the prevalence of ADHD in dental students by searching mainstream databases such as Pubmed and Web of science.

With the continuous development of Psychiatry, Psychology, and Education, more and more people began concentrating on ADHD in college students. However, studies on medical students have mainly focused on the adverse effects, including suicidal behaviour and Internet addiction, accompanying ADHD[11–13]. At the same time, the research on the relationship between ADHD and positive outcomes, including life satisfaction, is minimal. In scientific efforts, clinical practice, and teaching practice, the dimensional approach of ADHD can allow people to obtain more information than the traditional classification method. [14] A study of middle-aged and older Australians showed that inattention was more relevant than hyperactivity in ADHD symptoms in social, academic, occupational functioning, and psychological outcomes, which include life satisfaction. [15] Until now, however, no study has examined this issue among dental students.

Although ADHD symptoms may generate impairment in some related functions, they do not significantly impact some people's lives. According to our conjecture, this group may have positive psychological resources to cope with adversity better. Since positive psychology was proposed to prevent and treat mental health problems and improve people's overall subjective well-being, positive psychology

has been increasingly applied to many people, including children, doctors, and soldiers. [16–18]. As one of the positive psychological resources, resilience can reduce the negative impact of psychiatric symptoms in ADHD patients and improve their life satisfaction. [19,20]

Generally speaking, resilience is how people understand, manage, and control themselves after a significant stressful or traumatic event. [21] It can be inferred from the transaction model of coping and stress[22] that resilience, as a psychological resource, can quadratically affect the evaluation process of stressors and thus moderate the relationship between stressors and their outcomes. Still, there are few studies related to the possible role of resilience in the relationship between ADHD symptoms and life satisfaction.

Given the increasing socioeconomic burden of ADHD, existing studies on ADHD symptoms are significantly underrepresented in dental students. We conducted this study to 1) analyze the prevalence of ADHD symptoms among dental students in China; 2) explore the relationship between inattention and hyperactivity dimensions of ADHD symptoms and life satisfaction in dental students; 3) explore the mediating role of resilience in the connection between ADHD symptoms and life satisfaction.

Methods

Study design and subjects

This cross-sectional study was conducted at China Medical University in May 2023. "Wenjuanxing" (www.wjx.cn), a famous professional online survey platform, was

used to conduct questionnaires and collect data. Authors had access to information that could identify individual participants during or after data collection. A stratified cluster sampling method was used to recruit dental students based on academic years. All students participated voluntarily and were informed of the study's purpose before answering the questionnaire. At the beginning of the questionnaire, we designed an informed consent link. The subjects who answered the questionnaire were considered to have signed a written informed consent agreement before answering the questionnaire. Medical Ethics Committee of China Medical University authorized the study.

Measurement of current ADHD symptoms

The Chinese arrangement of the Adult ADHD Self-Report Scale(ASRS) was adopted to assess the current ADHD symptoms of the subjects. [23] ASRS is composed of two sub-scales named inattention and hyperactivity. Each has nine items. Sample items include "How often do you fidget or squirm with your hands or feet when you have to sit down for a long time?" and "How often do you misplace or have difficulty finding things at home or work?". Based on their affairs over the past six months, the respondents were requested to rate each article on a 5-point Likert scale from 0 (never) to 4 (very often). Subjects who scored more than 24 were highly likely to have ADHD, those who scored 17-23 were classified as possibly having ADHD, and those who scored 0-16 were almost not likely to have ADHD. [24] There are good reliability and validity in the Chinese version of ASRS owing to previous studies. [23–25] Cronbach's alpha coefficients for the inattention and hyperactivity dimensions

were 0.78 and 0.85 in this study.

Measurement of nonage ADHD symptoms

Wender Utah Rating Scale (WURS) was used to assess nonage ADHD symptoms among the students. The 25-item WURS was used primarily to review individual ADHD symptoms during adolescence (e.g., "As a child, I had trouble seeing things from someone else's point of view" and "As a child, I was sad or blue, depressed, unhappy"). Each article is ranked on a Likert scale with 5 points from 0 (not at all or very slightly) to 4 (very much). An aggregate score of 46 was used as the demarcation point for having nonage ADHD. [26] Scholars believe the Chinese version of WURS has decent credibility and validity. Cronbach's alpha was 0.94 in this research.

Classification criteria of ADHD symptomatic group and asymptomatic group

In clinical practice, only individuals with both overt childhood ADHD symptoms and current ADHD symptoms are diagnosed as adult ADHD. Therefore, subjects with WURS scores ≥46 and at least one dimension of ASRS scores ≥17 were classified as a symptomatic group in this study, and the remaining were classified as asymptomatic.

Measurement of life satisfaction

The Satisfaction With Life Scale (SWLS) estimates the worldwide level of life satisfaction. [27] The scale owns five items (e.g., "So far, I have gotten the influential things I want in life" and "I am satisfied with my life"). Each article is answered on a Likert scale with 7 points from 1 (strongly disagree) to 7 (strongly agree). The higher

the ranking scored, the more satisfied the subjects were with their lives. Previous studies have demonstrated convincing reliability and predictive validity among diverse age groups. [27–29] In past studies, Cronbach's alpha for SWLS was 0.91.

Measurement of resilience

Connor‑Davidson Resilience Scale (CD-RISC) was developed to estimate the capacity to cope with pressure and adversity. [30] CD-RISC, possessing 25 items, has a higher psychometric property rating than most scales that measure resilience. [31]Sample items include "When things look hopeless, I do not give up condition" and "Past success gives me confidence for new challenges." Each article is ranked on a Likert scale with 5 points from 0 (not true at all) to 4 (accurate nearly all the time). The higher score on the full scale has a higher level of resilience they own. Owing to the scale's satisfying reliability and validity, the scale has been widely used in China. [30,31] Cronbach's alpha for CD-RISC in the past research was 0.94.

Demographic characteristics

Demographic information for the study incorporated gender, age, race, place of residence, academic year, whether the only child and parental educational level.

Statistical analysis

SPSS 26.0 was used to analyze all the data. All statistical examinations were two-sided, and results with P < 0.05 were marked as significant. T-tests and chi-squared tests were used to test the differences in psychological and categorical variables between ADHD symptomatic group and the asymptomatic group. Pearson's correlation was operated to examine correlations among inattention,

hyperactivity, resilience, and life satisfaction. Hierarchical regression analysis was conducted to investigate the influence of groups of independent variables on life satisfaction. Standardized estimates (β), F, R2, and R2-changes ($\triangle$R2) for each step were supplied. Asymptotic and resampling strategies, developed by Preacher and Hayes[29], were used to examine the mediation of resilience (a*b product) on the correlations of inattention and hyperactivity with life satisfaction. The bootstrap estimate was on the basis of 5000 bootstrap samples. The bias-corrected and accelerated 95% confidence interval (BCa 95%CI) for each a*b product was worked out, and a BCa 95%CI excluding 0 indicated a significant mediating role. All the continuous variables were standardized to avoid multicollinearity before the regression analyses.

Results

Demographic characteristics of the objects

In total,297 questionnaires were distributed, and 291 copies were collected. The demographic features of the issues and the allocations of life satisfaction are shown in Table 1. There were 120 female dental students and 171 male students. The age of the problems ranged from 17 to 23 (M =19.72, SD = 1.35)

Table 1　Demographic characteristics of the objects in life satisfaction(*N*=291)

Variables		N	%	life satisfaction(Mean±SD)
Gender	Male	120	41.2	24.46±5.97
	Female	171	58.8	24.02±5.62

Age group				
	17-20	196	67.4	24.17±5.64
	21-23	95	32.6	24.25±6.04
Race				
	Han	251	86.3	24.35±5.76
	Hui	6	2.1	22.17±4.07
	Man	25	8.6	23.92±5.98
	Other	9	3.0	22.22±6.24
Place of residence				
	Cities	192	66.0	24.96±6.12
	Towns	59	20.3	22.81±4.57
	Villages	40	13.7	22.58±4.89
Academic year				
	First year	72	27.7	25.88±6.24
	Second year	75	25.8	23.75±5.01
	Third year	69	23.7	23.04±5.43
	Fourth year	75	25.8	24.11±6.04
The only child				
	Yes	184	63.2	24.65±5.74
	No	107	36.8	23.43±5.75
Paternal educational level				
	Primary school and below	20	6.9	22.40±7.96

Secondary school	106	36.4	24.52±5.43
College and above	165	56.7	24.21±5.66
Maternal education level			
Primary school and below	32	11.0	22.31±6.61
Secondary school	106	36.4	25.34±5.86
College and above	153	52.6	23.80±5.38

Life satisfaction: Satisfaction With Life Scale

Characteristics of the objects in the ADHD symptomatic group and asymptomatic group

Among 291 dental students, 91 people scored ≥46 in WURS. Then, the 91 people were statistically analyzed by ASRS scale, and the screening method was as follows: if the scores of 1-3 questions in Part A (namely the first six questions) were ≥3, and the scores of 4-5 questions were ≥4, and more than four of the six questions met the above standards. A total of 20 people correspond to the above conditions. These 20 students are classified as ADHD symptomatic class. The comparison of the features of the subjects in the ADHD symptomatic and asymptomatic group is presented in Table 2. The male group had a higher rate of ADHD symptoms than the female group (p = 0.025). There are meaningful distinctions between the three groups in Paternal educational level (p = 0.046). There were no significant distinctions between the two groups in age(p = 0.794), Race(p = 0.216), Place of residence(p = 0.830), Academic year(p = 0.652), The only child(p = 0.865), and Maternal education level(p = 0.520).

The ADHD symptomatic group had remarkably lower ranks of resilience(p < 0.05)

and life satisfaction (p < 0.05) comparable to the asymptomatic group.

Table 2 Characteristics of the objects in the ADHD symptomatic group and

asymptomatic group(*N*=291)

Variables		ADHD symptomatic N (%)	ADHD asymptomatic N (%)	P
Gender	Male	13(10.8)	107(89.2)	0.025
	Female	7(4.1)	164(95.9)	
Age group				
	17-20	14(7.1)	182(92.9)	0.794
	21-23	6(6.3)	89(93.7)	
Race				
	Han	16(6.4)	235(93.6)	0.216
	Hui	0	6(100.0)	
	Man	4(16.0)	21(84.0)	
	Other	0	9(100.0)	
Place of residence				
	Cities	14(7.3)	178(92.7)	0.830
	Towns	3(5.1)	56(94.9)	
	Villages	3(7.5)	37(92.5)	
Academic year				

First year	4(5.6)	68(94.4)	0.652	
Second year	4(5.3)	71(94.7)		
Third year	7(10.1)	62(89.9)		
Fourth year	5(6.7)	70(93.3)		
The only child				
Yes	13(7.1)	171(92.9)	0.865	
No	7(6.5)	100(93.5)		
Paternal educational level				
Primary school and below	4(20.0)	16(80.0)	0.046	
Secondary school	5(4.7)	101(95.3)		
College and above	11(6.7)	154(93.3)		
Maternal education level				
Primary school and below	3(9.4)	29(90.6)	0.520	
Secondary school	5(4.7)	101(95.3)		
College and above	12(7.8)	141(92.2)		
	Mean ± SD	Mean ± SD		
Resilience	70.85±18.57	66.45±17.21	0.005	
Life Satisfaction	25.75±6.94	24.08±5.67	0.045	

Resilience: Connor–Davidson Resilience Scale; Life satisfaction: Satisfaction With Life Scale

Pearson correlations among resilience, ADHD symptoms, and life satisfaction

The means, standard deviations, and correlations among inattention, hyperactivity, resilience, and life satisfaction are demonstrated in Table 3. As indicated in the table, both inattention and hyperactivity were negatively linked to resilience (inattention: r = − 0.177, p < 0.05; hyperactivity: r = − 0.258, p < 0.05) and life satisfaction (inattention: r = − 0.121, p < 0.05; hyperactivity: r = − 0.190, p < 0.05). Resilience was observably positively related to life satisfaction (r = 0.752, p < 0.05).

Table 3 Correlations among ADHD symptoms, resilience, and Life Satisfaction

Variables	Mean	SD	1	2	3	4
1. inattention	15.22	6.44	1			
2. Hyperactivity	11.78	7.03	0.800**	1		
3. Resilience	66.75	17.31	-0.177**	-0.258**	1	
4. Life Satisfaction	24.20	5.77	-0.121**	-0.190**	0.752**	1

**:P<0.05(two-tailed）

The results of Hierarchical regression

The consequences of the hierarchical regression of life satisfaction are indicated in Table 4. The two proportions of ADHD symptoms, inattention, and hyperactivity, accounted for 4% of the variance in life satisfaction. After controlling for age and gender, hyperactivity was significantly negatively related to life satisfaction (β = -0.26, p < 0.05). The consequence of resilience on life satisfaction was incredibly positive (β = 0.75, p < 0.05), explaining up to 53% of the variance in life satisfaction.

Table 4 Hierarchical linear regression analyses results

Variables	Step 1 β（95%CI）	Step 2 β（95%CI）	Step 3 β（95%CI）
Step 1			
Age	-0.42(-0.68~0.32)	-0.03(-0.65~0.33)	-0.01(-0.40~0.25)
Gender	-0.04(-1.89~0.85)	-0.55(-1.97~0.73)	-0.06(-1.69~0.12)
Step 2			
inattention		0.08(-0.09~0.25)	0.02(-0.08~0.14)
Hyperactivity		-0.26**(-0.37~-0.05)	-0.02(-0.12~0.08)
Step 3			
Resilience			0.75**(0.22-0.27)
F	0.457	3.163**	75.61**
R^2	0.003	0.042	0.570
△R^2	0.003	0.039	0.528

**:$P<0.05$（two-tailed)

The mediating role of resilience in the connections between ADHD symptoms and life satisfaction

The path coefficients, the magnitude of the mediating effect (a*b products), and BCa 95%CI of the products are revealed in Table 5. Resilience has a significant mediating effect between inattention and the life satisfaction of students.(a*b=-0.199,$P<0.05$) Resilience significantly moderated the association between hyperactivity and life satisfaction in the subjects.(a*b=-0.159,$P<0.05$)

Table 5 Mediation of resilience in the correlation between ADHD symptoms and

Life Satisfaction

Predictors	Path coefficient				a*b(Bca 95%CI)
	c	a	b	C′	
inattention	-0.108**	-0.476**	0.251**	0.011	-0.199**(-0.028~-0.002)
Hyperactivity	-0.156**	-0.635**	0.251**	0.003	-0.159**(-0.266~-0.057)

**:P<0.05;

c: associations of ADHD symptoms with life satisfaction ;

a: associations of ADHD symptoms with resilience ;

b: associations of resilience with life satisfaction after controlling for the predictor variables ;

c': associations of ADHD symptoms with life satisfaction after adding resilience as the mediator

Discussion

UP till now, there has been little investigation into ADHD among dental students. Existing studies have explored the awareness and knowledge of dental students about ADHD but have not focused on ADHD in the group itself. [32] This study focuses on ADHD symptoms among dental students. It is also the first to explore the correlation between ADHD symptoms and the life satisfaction of dental students, as well as to verify the mediating effect of mental resilience in these relationships. On the one hand, the outcomes of this study indicate that 6.87% of dental students have symptomatic ADHD. The relevant data of Chinese dental students was lower than the morbidity rate of ADHD (8.45%) among Chinese medical students. [33] Several

studies have demonstrated a correlation between ADHD and learning disabilities, particularly in verbal reasoning, visual-auditory memory, and cognitive traits. [34,35] It can be inferred that some individuals with ADHD will be slightly worse than ordinary people in learning ability. Chinese undergraduates must take a national exam before being admitted and choose their prospective colleges and majors based on their scores. In most Chinese medical universities, the average score of students admitted to the stomatology program is higher than that of undergraduates majoring in clinical medicine in the same institution. Therefore, among high school graduates with ADHD symptoms who want to apply for stomatology, compared with graduates who want to use it for clinical medicine, most of them cannot study oral medication due to the entrance examination scores. On the other hand, according to the data from the World Health Organization, the result is higher than the worldwide prevalence of ADHD (2.8%) among adults. [5] This data may indicate that, unlike mental disorders such as anxiety and depression, which are easily recognized and detected, dental students facing heavy academic and life pressure often underestimate or ignore ADHD symptoms. Early detection and diagnosis of dental students with ADHD symptoms are necessary to treat them as early as possible. This study shows that the prevalence of ADHD significantly differs among students whose fathers have different levels of education. The probable reason is that Chinese fathers teach their children more harshly, while fathers with less education often cannot control their emotions and often criticize their children. These may have implications for the recovery of ADHD symptoms in children. This suggests that the

relevant department should improve the education level of residents. A review of three decades of research shows that, in the long run, people with untreated ADHD are impaired in social functioning, self-esteem, and occupation compared to ordinary people. [36] The resilience level of the asymptomatic group is significantly higher than that of the symptomatic group, which is consistent with the study of medical students. [33] The rational interpretation is that people with ADHD often have other mental disorders, including mood disorders, learning disabilities, and so on [34,37], and many have experienced significant adverse events. [38]These are all inversely related to resilience. [33]

One of the principal findings of this research was that hyperactivity was significantly associated with life satisfaction among dental students. In contrast, there was no correlation between inattention and life satisfaction after controlling the distracting factors. This is different from existing research. In the past literature, some show that inattention is correlated with life satisfaction[33], while others believe that there is no apparent difference between diverse subtypes of ADHD[39]. There are two reasons for this result: First, ADHD is more general in men than women, and it is more common for men to show hyperactivity than inattention. Second, the survey uses recall scales. Hyperactivity is more likely to attract the attention of others, such as teachers and parents. These individuals mentioned hyperactivity around subjects more often than inattention, which may make issues memorize hyperactivity more clearly than inattention.

Another important finding of the research is that resilience significantly mediates

between the two ADHD symptoms and life satisfaction in dental students. In addition to being directly related to life satisfaction, the two symptoms of ADHD are also indirectly associated with life satisfaction through the mediating effect of resilience. Students with more obvious ADHD symptoms had lower levels of resilience, which in turn was associated with lower life satisfaction. And vice versa. Unlike traditional psychology, which focuses on negative and pathological psychology, positive psychology uses human motivation as its basic tenet. Since positive psychology was proposed, it has been increasingly widely used to prevent and treat psychological disorders in the past 20 years. Resilience is an essential component of positive psychology, which refers to the excellent adaptability of individuals in the face of adversity, trauma and other significant life pressures. [40] A meta-analysis in different populations showed that multiple training programs and interventions had a specific effect. [41] Previous studies have shown that appropriate interventions to improve resilience are feasible among university faculty and military personnel. [42,43] Interventions based on mindfulness and cognitive-behavioural have been used more frequently in the psychological treatment of ADHD. This suggests that such interventions are effective in improving ADHD symptoms in adults. [44,45] Therefore, on an evidence-based basis, the School of Stomatology can design a targeted resilience intervention program to enhance the life satisfaction of students with ADHD. This may improve the enthusiasm of these students for study and life, which is conducive to cultivating more excellent future dentists. The variables in this study accounted for 57% of the variance in students'

life satisfaction. Some potential variables, such as other social and psychological factors, may affect students' satisfaction with life, which were not included in the model.

From the view of positive psychology, this study selected dental students as the subject group and obtained a large sample size and a highly effective response rate. This paper explored the correlation between the two sub-symptoms of ADHD and life satisfaction and analyzed the related mechanism between the two sub-symptoms and resilience. Admittedly, this study has some limitations. First, the studies all used self-report questionnaires, and data collected on these questionnaires can introduce errors. However, self-report questionnaires are often used in psychiatric clinical practice and large-scale epidemiological studies. [46] Second, as a cross-sectional study, it is impossible to explore the causality between the variables, and it is difficult to follow up on the subjects. Relevant prospective cohort studies can be carried out to supplement it. Third, limited by resource allocation, multi-centre research among multi-national, multi-regional, and multi-level medical schools should be carried out in the future

Conclusions

Studies have shown that the prevalence of ADHD in Chinese dental students is higher than in general adults. It was revealed that paternal education level was relevant to the morbidity. Of the two ADHD symptoms, only hyperactivity was significantly

associated with life satisfaction. Distinct mediation of resilience was found between life satisfaction and ADHD symptoms. The results of this study can ensure the early detection of ADHD symptoms in dental students to enhance the mental health level of future dentists to a certain extent. Resilience intervention programs can improve dental students' life satisfaction, especially for students with ADHD symptoms.

Acknowledgements

The authors give expressions of gratitude to all the students participating in this experiment and the faculty of stomatology who helped to collect the data.

References

1. GBD 2019 Mental Disorders Collaborators. Global, regional, and national burden of 12 mental disorders in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. Lancet Psychiatry. 2022 Feb;9(2):137–50.

2. Chhibber A, Watanabe AH, Chaisai C, Veettil SK, Chaiyakunapruk N. Global Economic Burden of Attention-Deficit/Hyperactivity Disorder: A Systematic Review. PharmacoEconomics. 2021 Apr;39(4):399–420.

3. Association AP. Diagnostic and Statistical Manual of Mental Disorders, 5th Edition: DSM-5. 5th edition. Washington, D.C: American Psychiatric Publishing; 2013. 991 p.

4. Association AP, Kennedy PJ. Understanding Mental Disorders: Your Guide to DSM-5. 1st edition. Washington, DC: American Psychiatric Pubublishing; 2015. 388 p.

5. Fayyad J, Sampson NA, Hwang I, Adamowski T, Aguilar-Gaxiola S, Al-Hamzawi A, et al. The descriptive epidemiology of DSM-IV Adult ADHD in the World Health Organization World Mental Health Surveys. Atten Deficit Hyperact Disord. 2017 Mar;9(1):47–65.

6. Harpin VA. The effect of ADHD on the life of an individual, their family, and community from preschool to adult life. Arch Dis Child. 2005 Feb;90 Suppl 1(Suppl 1):i2-7.

7. Schwanz KA, Palm LJ, Brallier SA. Attention problems and hyperactivity as

predictors of college grade point average. J Atten Disord. 2007 Nov;11(3):368–73.

8. Murphy K. Psychosocial treatments for ADHD in teens and adults: a practice-friendly review. J Clin Psychol. 2005 May;61(5):607–19.

9. Roberts W, Peters JR, Adams ZW, Lynam DR, Milich R. Identifying the facets of impulsivity that explain the relation between ADHD symptoms and substance use in a nonclinical sample. Addict Behav. 2014 Aug;39(8):1272–7.

10. Tong L, Shi HJ, Zhang Z, Yuan Y, Xia ZJ, Jiang XX, et al. Mediating effect of anxiety and depression on the relationship between Attention-deficit/hyperactivity disorder symptoms and smoking/drinking. Sci Rep. 2016 Feb 29;6:21609.

11. Shen Y, Zhang Y, Chan BSM, Meng F, Yang T, Luo X, et al. Association of ADHD symptoms, depression and suicidal behaviors with anxiety in Chinese medical college students. BMC Psychiatry. 2020 Apr 22;20(1):180.

12. Shen Y, Chan BSM, Huang C, Cui X, Liu J, Lu J, et al. Suicidal behaviors and attention deficit hyperactivity disorder (ADHD): a cross-sectional study among Chinese medical college students. BMC Psychiatry. 2021 May 18;21(1):258.

13. Shi M, Du TJ. Associations of personality traits with internet addiction in Chinese medical students: the mediating role of attention-deficit/hyperactivity disorder symptoms. BMC Psychiatry. 2019 Jun 17;19(1):183.

14. Hyman SE. The diagnosis of mental disorders: the problem of reification. Annu Rev Clin Psychol. 2010;6:155–79.

15. Das D, Cherbuin N, Butterworth P, Anstey KJ, Easteal S. A Population-Based Study of Attention Deficit/Hyperactivity Disorder Symptoms and Associated Impairment in Middle-Aged Adults. PLoS ONE. 2012 Feb 8;7(2):e31500.

16. Benoit V, Gabola P. Effects of Positive Psychology Interventions on the Well-Being of Young Children: A Systematic Literature Review. Int J Environ Res Public Health. 2021 Nov 17;18(22):12065.

17. Bazargan-Hejazi S, Shirazi A, Wang A, Shlobin NA, Karunungan K, Shulman J, et al. Contribution of a positive psychology-based conceptual framework in reducing physician burnout and improving well-being: a systematic review. BMC Med Educ. 2021 Nov 25;21(1):593.

18. Jarrett T. Warrior Resilience Training in Operation Iraqi Freedom: combining rational emotive behavior therapy, resiliency, and positive psychology. US Army Med Dep J. 2008;32–8.

19. Greven CU, Buitelaar JK, Salum GA. From positive psychology to psychopathology: the continuum of attention-deficit hyperactivity disorder. J Child Psychol Psychiatry. 2018 Mar;59(3):203–12.

20. Sedgwick JA, Merwood A, Asherson P. The positive aspects of attention deficit hyperactivity disorder: a qualitative investigation of successful adults with ADHD. Atten Deficit Hyperact Disord. 2019 Sep;11(3):241–53.

21. Carr A. Positive Psychology: The Science of Happiness and Human Strengths. Psychology Press; 2004. 412 p.

22. Spaccarelli S. Stress, appraisal, and coping in child sexual abuse: a theoretical and empirical review. Psychol Bull. 1994 Sep;116(2):340–62.

23. Yeh CB, Gau SSF, Kessler RC, Wu YY. Psychometric properties of the Chinese

version of the adult ADHD Self-report Scale. Int J Methods Psychiatr Res. 2008;17(1):45–54.

24. Kessler RC, Ustün TB. The World Mental Health (WMH) Survey Initiative Version of the World Health Organization (WHO) Composite International Diagnostic Interview (CIDI). Int J Methods Psychiatr Res. 2004;13(2):93–121.

25. Ni HC, Gau SSF. Co-occurrence of attention-deficit hyperactivity disorder symptoms with other psychopathology in young adults: parenting style as a moderator. Compr Psychiatry. 2015 Feb 1;57:85–96.

26. Ward MF, Wender PH, Reimherr FW. The Wender Utah Rating Scale: an aid in the retrospective diagnosis of childhood attention deficit hyperactivity disorder. Am J Psychiatry. 1993 Jun;150(6):885–90.

27. Diener E, Emmons RA, Larsen RJ, Griffin S. The Satisfaction With Life Scale. J Pers Assess. 1985 Feb;49(1):71–5.

28. Shi M, Wang X, Bian Y, Wang L. The mediating role of resilience in the relationship between stress and life satisfaction among Chinese medical students: a cross-sectional study. BMC Med Educ. 2015 Feb 13;15(1):16.

29. Pavot W, Diener E, Colvin R, Sandvik E. Further Validation of the Satisfaction With Life Scale: Evidence for the Cross-Method Convergence of Well-Being Measures. J Pers Assess. 1991 Sep 1;57:149–61.

30. Connor K, Davidson J. Development of a new resilience scale: The Connor-Davidson Resilience Scale (CD-RISC). Depress Anxiety. 2003 Sep 1;18:76–82.

31. Windle G, Bennett K, Noyes J. A methodological review of resilience measurement scales. Health Qual Life Outcomes. 2011 Jan 1;9:1–8.

32. Chopra A, Vishnupriya V, Gayathri R, Kavitha S. Knowledge and awareness of dental students on attention deficit hyperactivity disorder. J Adv Pharm Technol Res. 2022 Nov;13(Suppl 1):S308–13.

33. Shi M, Liu L, Sun X, Wang L. Associations between symptoms of attention-deficit/ hyperactivity disorder and life satisfaction in medical students: the mediating effect of resilience. BMC Med Educ. 2018 Jul 13;18:164.

34. Talero-Gutiérrez C, Velez-van-Meerbeke A, Reyes R. A Clinical Study of ADHD Symptoms With Relation to Symptoms of Learning Disorders in Schoolchildren in Bogota, Colombia. J Atten Disord. 2010 Oct 1;16:157–63.

35. Toffalini E, Buono S, Cornoldi C. The structure, profile, and diagnostic significance of intelligence in children with ADHD are impressively similar to those of children with a specific learning disorder. Res Dev Disabil. 2022 Oct 1;129:104306.

36. Shaw M, Hodgkins P, Caci H, Young S, Kahle J, Woods AG, et al. A systematic review and analysis of long-term outcomes in attention deficit hyperactivity disorder: effects of treatment and non-treatment. BMC Med. 2012 Sep 4;10:99.

37. Cumyn L, French L, Hechtman L. Comorbidity in Adults with Attention-Deficit Hyperactivity Disorder. Can J Psychiatry. 2009 Oct 1;54(10):673–83.

38. Dvorsky MR, Langberg JM. A Review of Factors that Promote Resilience in Youth with ADHD and ADHD Symptoms. Clin Child Fam Psychol Rev. 2016 Dec 1;19(4):368–91.

39. O'Brien JW, Dowell LR, Mostofsky SH, Denckla MB, Mahone EM.

Neuropsychological profile of executive function in girls with attention-deficit/hyperactivity disorder. Arch Clin Neuropsychol Off J Natl Acad Neuropsychol. 2010 Nov;25(7):656–70.

40. Newman R. APA's resilience initiative. Prof Psychol Res Pract. 2005;36:227–9.

41. Joyce S, Shand F, Tighe J, Laurent SJ, Bryant RA, Harvey SB. Road to resilience: a systematic review and meta-analysis of resilience training programmes and interventions. BMJ Open. 2018 Jun 14;8(6):e017858.

42. Johnson DC, Thom NJ, Stanley EA, Haase L, Simmons AN, Shih P an B, et al. Modifying Resilience Mechanisms in At-Risk Individuals: A Controlled Study of Mindfulness Training in Marines Preparing for Deployment. Am J Psychiatry. 2014 Aug;171(8):844–53.

43. Sood A, Prasad K, Schroeder D, Varkey P. Stress Management and Resilience Training Among Department of Medicine Faculty: A Pilot Randomized Clinical Trial. J Gen Intern Med. 2011 Aug;26(8):858–61.

44. Geffen J, Forster K. Treatment of adult ADHD: a clinical perspective. Ther Adv Psychopharmacol. 2018 Jan;8(1):25–32.

45. Kolar D, Keller A, Golfinopoulos M, Cumyn L, Syer C, Hechtman L. Treatment of adults with attention-deficit/hyperactivity disorder. Neuropsychiatr Dis Treat. 2008 Feb;4(1):107–21.

46. Magnússon P, Smári J, Sigurdardóttir D, Baldursson G, Sigmundsson J, Kristjánsson K, et al. Validity of self-report and informant rating scales of adult ADHD symptoms in comparison with a semistructured diagnostic interview. J Atten Disord. 2006 Feb;9(3):494–503.