Full title: **Psychiatric hospital admissions and linkages to ambulatory services in the Western Cape**

Province of South Africa (2015-2022): trends, risk factors, the impact of the COVID-19 pandemic and

possible opportunities for intervention

(197 characters)

Short title: **Western Cape psychiatric admission trends and links to ambulatory services**

(65 characters)

Hannah Hussey[1,2], Timothy Mountford[1], Alexa Heekes[1], Carol Dean[3], Marinda Roelofse[3], Lynne Hendricks[3],

Qhama Cossie[4,5], Liezel Koen[6,7], Warren Caesar[8], Vanessa Lomas[9,10], David Pienaar[11], Giovanni Perez[9],

Andrew Boulle[1,2], Katherine Sorsdahl[12], Hassan Mahomed[9,13]

1. Provincial Health Data Centre, Health Intelligence, Western Cape Department of Health and

Wellness

2. School of Public Health, University of Cape Town

3. Specialised Hospitals, Metro Health Services, Western Cape Department of Health and Wellness

4. Valkenberg Psychiatric Hospital, Western Cape Department of Health and Wellness

5. Department of Mental Health & Psychiatry, University of Cape Town

6. Stikland Psychiatric Hospital, Western Cape Department of Health and Wellness

7. Department of Psychiatry, Faculty of Medicine and Health Sciences, University of

Stellenbosch

8. Lentegeur Psychiatric Hospital, Western Cape Department of Health and Wellness

9. Metro Health Services, Western Cape Department of Health and Wellness

10. Department of Family and Emergency Medicine, University of Stellenbosch

11. Rural Health Services, Western Cape Department of Health and Wellness

12. Alan J. Flisher Centre for Public Mental Health, Department of Psychiatry & Mental Health,

University of Cape Town

13. Division of Health Systems and Public Health, Department of Global Health, Faculty of Medicine

and Health Sciences, Stellenbosch University

Contributing author: Hannah Hussey (hannah.hussey@westerncape.gov.za)

Author contributions

HH conceptualized the research, performed the formal analysis and visualization, and wrote the original

draft

TM & AH were responsible for data curation

CD, LH, MR, GP assisted with the conceptualization of the research

QC, LK, WC, VL, DP and AB assisted with reviewing and editing

KS and HM assisted with reviewing and editing, as well as supervision

Abstract

In recent years, hospital managers have reported increasing numbers of psychiatric hospital admissions

in the Western Cape Province of South Africa, driven by poverty and substance use. We aimed to examine

this trend, and the impact of the COVID-19 pandemic, as well as assess factors associated with linkage to

ambulatory services following hospital discharge and repeat psychiatric admissions.

Using electronic health data from the Provincial Health Data Centre, a consolidated routine service

database, all psychiatric hospital admissions in the Western Cape public sector from 2015 to 2022 were

analyzed, stratified by hospital level. Mixed effects logistic regression was used in this cohort study to

determine the factors associated with successful linkage to ambulatory services within 30 days following

hospital discharge, and repeat psychiatric admission within 30 and 90 days.

We found that psychiatric hospital admissions, particularly at the district/acute level, were increasing

prior to 2020 and an increasing proportion of diagnoses were substance related. 40% of admissions at the

district level had not been seen at a primary health care facility in the year before the admission. Male

patients and those with substance use disorders were less likely to be successfully linked to outpatient

services following discharge. Successful linkage was one of the most protective factors against

readmission within 90 days with an adjusted odds ratio of 0.76 (95%CI 0.73-0.79) and 0.45 (95%CI 0.42-

0.49) at district/acute and specialized hospitals respectively.

Improving linkage to ambulatory services by mental health patients post-discharge is likely to avert

hospital readmissions.

Manuscript word count: 3 920

Background

Prior to the emergence of SARS-CoV-2, and consistent with international trends, there was already a high and increasing prevalence and burden of mental health conditions in the Western Cape Province of South Africa (1). Recent surveys in the Western Cape have found a high prevalence of mental health conditions: 31.8% of respondents had probable depression in 2021, and that the lifetime prevalence for any or substance use disorder was 39.4% and 20.6% respectively (2,3). In South Africa neuropsychiatric disorders are the third highest contributors to disability-adjusted life-years, after HIV/AIDS and other infectious diseases, surpassing other non-communicable diseases (4). Social determinants such as poverty, food insecurity, experiences of trauma and violence, high rates of substance use and comorbid illnesses are all known drivers of this mental health burden (1,3,5–7).

Into this context, the COVID-19 pandemic occurred, resulting in a further increase in the burden of psychiatric symptoms and diseases, while simultaneously leaving the health system less able to adequately address them. (8). The economic crises and resultant job losses precipitated by COVID-19 also resulted in an increase in depressive symptoms in South Africa (9). High rates of depression, anxiety and post-traumatic stress disorder were also documented in front-line healthcare workers in South Africa during the pandemic (10). In addition, SARS-CoV-2 infection itself can have neuropsychiatric manifestations, including index-episode psychosis (11). At the same time, diversion of resources to COVID-19 services hampered access to mental health care for individuals known to have serious mental illnesses (12). Significant disruptions to substance use disorder treatment services in South Africa were also noted during the pandemic (13).

While the majority of patients with mental health conditions do not require inpatient care, inpatient expenditure represented 86% of total mental health expenditure in the public health services in South Africa, with specialized psychiatric hospital admissions alone requiring 46% of total mental health

expenditure (14). In an increasingly financially constrained environmental, it would be important to

understand how admissions can be reduced and what factors influence these.

This study therefore aimed to describe and interrogate trends in psychiatric hospital admissions from

2015-2022, with a particular focus on the role of substance use and the possible impact of COVID-19. This

study will also take into account ambulatory visits before and after admissions, as possible opportunities

for intervention to reduce hospital admissions.

Methods

Setting

The Western Cape Province of South Africa has a population of 7.1 million individuals, 74.9% of whom are

reliant on public health services (15). The adult population is serviced by 33 district hospitals, four regional

hospitals and two central hospitals (which we grouped together as "district/acute hospitals" for this

analysis), all of whom refer to the three specialised psychiatric hospitals of Lentegeur, Stikland and

Valkenberg. In the public sector in South Africa, the inpatient services treatment gap is high at 89%, with

only 0.31 psychiatrists and 0.97 psychologists available per 100 000 population (16).

Data sources

The Provincial Health Data Centre (PHDC) collates all available patient-level electronic health data

captured into routine health information systems within public health facilities in the Western Cape

province (17). This includes admissions to psychiatric wards, derived from Clinicom, a patient

administration database used at all public hospitals in the province, as well as eCCR (electronic Continuity

of Care Record), the electronic discharge summary that includes information on discharge diagnoses using

ICD-10 coding. Using a unique patient identifier, the admissions are linked to the patients' interactions

with the provincial health system at hospital and primary care levels prior to admission and following

discharge.

Study population and inclusion/exclusion criteria

Individuals of all ages with a psychiatric hospital admission start date from 1 January 2015 to 31 December

2022 in a Western Cape public health facility were included in this cohort study. As ICD-10 diagnostic

coding was not always reliable enough to identify psychiatric patients, admitting specialty was used

instead. A psychiatric hospital admission was defined as:

• An admission to a specialized psychiatric hospital OR

• An admission to any other hospital in the province, where patients were admitted to the

specialty of "Psychiatry" on Clinicom

Patients with psychiatric conditions seen only in the Emergency Centres or those admitted to other

specialties (e.g., suicidal patients admitted to the General Medicine Department following an overdose)

were not included.

Data analysis

Firstly, descriptive data looking at trends in admission numbers, diagnoses and outcomes after discharge

are presented. First contacts with the public health system, following hospital discharge are also explored.

While this is not limited to mental health ambulatory services only, as these can be difficult to distinguish

electronically at the primary health care (PHC) level, the assumption is that as all patients are given a

follow-up date after discharge, a visit directly after discharge was assumed to be most likely related to the

preceding admission. PHC visits in the year preceding a psychiatric admission to an district/acute hospital

are also assessed. PHC visits within 24 hours of a hospital admission were excluded here, i.e., those PHC

visits where the patient could have been referred on for hospital care were not included, but those PHC

visits where the patient went home after and were unrelated to the current hospital admission were.

Where appropriate trends are highlighted using Microsoft PowerBI   (Version: 2.116) graphs.

The second analytical section is restricted to adult general psychiatry patients, as they are the patient

population putting the most pressure on services, discharged home from either a district/acute hospital

or a specialized hospital, i.e., admissions that result in transfers from a district/acute hospital to a

specialized hospital are excluded. Elective therapeutic admissions to specialized hospitals were also

excluded. Any individuals who were known to be deceased (in-facility deaths of individuals with a previous

psychiatric admission) were also excluded. Stata (13.1, StataCorp. LLC, College Station, TX) was used for

the analysis. The outcomes assessed were:

• Successful linkage to care post-discharge, defined as an ambulatory visit (PHC clinic or hospital

out-patient) within 30 days of hospital discharge.

• Repeat psychiatric hospital admission, at any hospital level, within 30 and 90 days from the date

of discharge

All admissions discharged by 31 December 2022 were included in this analysis (the data set was pulled on

15 March 2023, allowing sufficient time for the outcomes to have occurred and been captured

electronically), except for the outcome of readmissions within 90 days, where only admissions discharged

by 30 November 2022 be included.

Mixed effects logistic regression is used, as individuals often have repeated admissions over time and

there is non-independence of observations. An alternative approach of using standard logistic regression,

but including only the latest admission for an individual, was also performed. All analyses are stratified by

hospital level (district/acute and specialized) and adjusted for:

• Age and sex

• Broad diagnosis category - based on primary discharge diagnosis ICD-10 coding: mood disorders,

including both uni- and bipolar depression (F30-F39); schizophrenia and other psychotic disorders

(F20-F29); substance related disorders (F10-F19); and other diagnoses

o As the coding at specialized hospitals is more detailed, having a substance related

diagnosis listed anywhere on the discharge summary (i.e., not just the primary diagnosis)

was also included

• Index psychiatric admission or a known previous psychiatric admission, according to the available

Clinicom data. Hospitals started using Clinicom at different times from 2001, with the psychiatric

hospitals starting in 2008. Index admission data may not be available electronically for patients

with longstanding psychiatric disorders

• PHC visit in year preceding admission or not

• Length of hospital stay,

• Admission year

• Admitting hospital

o The three specialized hospitals, labelled as A, B and C

o The district/acute hospitals grouped into rural regional or district hospitals, and the

metropolitan hospitals grouped according to their specialized hospital catchment area

(A, B or C)

Ethics

This study was approved by the University of Cape Town Health Research Ethics Committee (HREC 058

/2023). A waiver of informed consent was approved, as the study used routinely collected electronic

health data in an anonymised dataset where the researcher had no access to identifying information.

Results

General trends in increasing admission numbers

Psychiatric admissions in the province have increased markedly over the last eight years (Figure 1a). In

2015 there were 6 013 admissions in specialized hospitals and 10 991 admissions in district/acute

hospitals. This increased to 6 697 and 19 880 admissions respectively in 2022. Admissions in district/acute

level hospitals almost doubled during this period, while specialized hospitals only increased slightly. This

resulted in the proportion of total psychiatric admissions admitted to a specialized hospital decreasing

from 35.36% in 2015 to 25.2% in 2022, implying   that a larger proportion of mental health patients are

being managed at the district/acute hospital level only.

Figure 1: Psychiatric hospital admissions in the Western Cape 2015-2022 - annual admissions for all facilities (1a)
and 14 day moving average of psychiatric admissions at acute/district hospital level 2015-2022 (1b).

2020 was the only year where an increase in admissions was not seen. When we focus on the

district/acute admissions (Figures 1b), we can see a large decrease in admission numbers coinciding with

the first wave of COVID-19 infections in the province in mid-2020, followed by smaller decreases with the

second (December 2020 – January 2021) and third waves (mid-2021).

Figure 2b illustrates the primary discharge diagnosis in specialized hospitals for general acute psychiatry,

where schizophrenia and other psychotic disorders make up more than half the primary diagnoses, with

mood disorders contributing to less than 20% of admissions. Substance related disorders as the primary

discharge diagnosis has increased as a proportion from 15.96% in 2018 to 22.47% in 2022. When looking

at all diagnosis codes listed (i.e., not just the primary diagnosis code) (Figure 2d), substance related

diagnoses also increased proportionately with time, accounting for 45.84% of all admissions in 2022.

Figure 2: Discharge diagnosis trends for acute/district hospitals (2a) and specialized hospitals (2b), based on the
primary discharge diagnosis. The proportion of admissions that had any substance related ICD-10 code listed at
acute/district hospitals (2c) and specialized hospitals (2d).

Diagnostic coding is not as detailed or complete at district/acute hospitals, although this has improved

with time – as illustrated by the initially large number of diagnoses that fell into the "Other" category. But

a similar trend can be seen at these hospitals (Figures 2a and 2c), where substance related disorders now

make up 22.74% of all primary diagnoses in 2022, and 29.21% of all admissions have a substance related

diagnosis listed in 2022 (compared to 10.68% and 12.71% in 2015 respectively). While the proportion that

were substance related increased over time, so did the total absolute number of patients too, meaning

that the 12.71% of patients with any substance related diagnosis code in 2015 corresponded to 1 396

individuals and the 29.21% of patients in 2022 corresponded to 5 613 individuals.

The readmission rate has, however, remained relatively stable, with around 10% of admissions, resulting

in a psychiatric readmission in the next 30 days (Figure 3), and this increases to around 18-19% when you

increase the window to 90 days. But with the absolute number of psychiatric hospital admissions

increasing, so too have the absolute number of readmissions putting the limited beds available under

pressure. For instance, in 2018, 2 018 individuals were readmitted within 30 days, but in 2022 that number

increased to 2 668. Currently only 47.5% of admissions acute/district hospitals and 28.68% of admissions

to specialised hospitals are first known psychiatric admissions. Amongst those known with to have a

previous psychiatric admission, the median number of days between admissions decreased from 2018 to

2022, for both hospital levels.

Figure 3: Readmission rates for district/acute and specialized hospitals 2018-2022 - readmission within 30 days
(3a), 90 days (3b), readmission categories (3c) and median days between psychiatric admissions (3d).

Profile of admitted patients and outcomes after discharge

Male patients account for 62.15% of all psychiatric admissions in the province. Age is related to sex and

diagnosis. Male patients and those with a primary substance related disorder tended to be younger

(Figure 4).

Figure 4: Median age in years of admitted psychiatric patients by diagnosis category and sex, in all facilities in
the Western Cape 2015-2022.

In 2022, the median length of stay in district/acute hospitals was 7 days and for specialized hospitals 34

days. In acute/district hospitals, those awaiting referral required an additional week of admission (Figure

5). With increasing numbers of admissions at the acute/district level, a decreasing proportion were

referred on for specialist care. In 2015, 37.54% were referred on, while in 2022 only 27.08% were referred

on. Length of stay at specialised hospitals was related to discharge diagnosis. Schizophrenia and other

psychotic disorders had the highest lengths of stay, followed by mood disorders and then substance

related disorders. All diagnosis categories showed a decrease in length of stay for 2022.

A median of 12 and 16 days were taken to first contact with the health system after discharge from district/acute hospitals and specialized hospitals, respectively. For those discharged directly from an district/acute hospital in 2022, 53.2% of individuals had an outpatient visit within 30 days, a number which has been increasing slowly over the last few years (Figure 6a). An additional 10.43% do make contact with out-patient services but take >30 days after discharge. 8.37% of discharges only make contact with the health system again with their next admission and 28.0% have had no further contact with the health system (as of March 2023, when this dataset was extracted). A similar trend is seen at specialized hospitals, although a higher proportion are followed up at ambulatory services within 30 days – 65.4% in 2022 (Figure 6b). But around 20% of patients discharged from specialized psychiatric service have no further contact with the health system.

Figure 6: After discharge from hospital, the clinic or out-patient contact within 30 days is assessed at acute/district hospitals (6a) and specialized hospitals (6b). "Yes" is defined as an ambulatory (clinic or out-patient) visit within 30 days, while "No" is an ambulatory visit but at more than 30 days.

PHC visits before a psychiatric admission

In 2022, 39.4% of admitted patients had not been seen at a PHC facility in the year preceding their admission, and this increased to 72.35% if one looked at the 30 days preceding an admission (Figure 7). Since 2015 there has been a general trend of increasing PHC visits in the period before an admission. 2020, however, was the exception with the proportion of admissions with a PHC visit in the 30 days prior to admission decreasing in May to July. Despite this general increase, the median days between the last PHC

visit and the psychiatric admission has remained relatively stable between 35 and 42 days (37 days in

2022).

Figure 7: Primary health care visits preceding an district/acute psychiatric hospital admission: proportion of
admissions with a PHC visit in the 30 days prior to admission, in 31-365 days prior admission and those without
a PHC visit, for 2015-2022 (7a), and proportion of admissions with a PHC visit in the 30 days prior to admission in
January 2020 – December 2021 (7b).

Successful linkage to care

74 728 admissions at district/acute hospitals and 30 750 admissions at specialized hospitals were included

in these analyses, the detailed characteristics of which are described in Table 1.

Table 1: Detailed characteristics of included admissions, stratified according to hospital level

At district/acute hospitals (Table 2), those with a PHC visit in the preceding year were more likely to be

successfully linked to ambulatory services (adjusted Odds ratio [aOR] of 2.67 [95% confidence interval

{95%CI} 2.58-2.77]). Women were also more likely to be successfully linked to care, as were older patients

and those without a primary substance related disorder. At specialized hospitals (Table 3) similar trends

were seen, bearing in mind the slightly different categorization of diagnoses and lengths of stay. Those

with a PHC visit in the year prior to admission (aOR 2.07 [95%CI 1.93-2.22]) and women (aOR 1.14 [95%CI

1.05-1.23]) were more likely to be successfully linked to care, but increasing age and previous psychiatric

admissions were not associated with improved linkage to care.

Table 2: Mixed effects logistic regression for the outcome of successful linkage to care following discharge (i.e., a
clinic or out-patient visit within 30 days of hospital discharge) and psychiatric readmissions within 30 and 90
days for district/acute hospitals. Only patients discharged directly home are included in these analyses.

Table 3: Mixed effects logistic regression for the outcome of successful linkage to care following discharge (i.e., a
clinic or out-patient visit within 30 days of hospital discharge) and psychiatric readmissions within 30 and 90
301   **days for specialized hospitals.**

Risk factors for readmission

At district/acute hospitals, female sex, older ages, being admitted to a rural hospital and a primary

substance related diagnosis were associated with a lower risk of readmission. An increasing length of stay

(but only up until 28 days), previous psychiatric admissions and a PHC visit in the year preceding an

admission were risk factors for readmission. While the aORs for admission years decreased with time, a

relative increase was seen in 2021 (aOR = 0.86 [95%CI 0.79-0.95]) compared to 2020 (aOR = 0.77 [95%CI

0.70-0.85] for readmissions within 90 days) - i.e., 2021 was more strongly associated with readmissions

than previous years - although 2022 has seen a return to aORs of the pre-pandemic years. Successful

linkage to care decreased the risk of readmission by 31% (aOR = 0.69 [95%CI 0.65-0.72]) and 24% (aOR =

0.76 [0.73-0.79]) for readmissions within 30 and 90 days respectively.

At specialized hospitals, successful linkage to care within 30 days of discharge was the most protective

factor against readmission within 30 and 90 days, with an aOR of 0.37 (95%CI 0.32-0.43) and 0.49 (0.44-

0.55) respectively. Specialized hospital B had the highest aOR for successful linkage to care and the lowest

aORs for readmissions.

Sex was not associated with readmission but a primary diagnosis that was substance related had the

lowest association with readmission. Previous admissions were strongly associated with readmissions, as

well as a PHC visit in the preceding year and being a direct admission to a specialized hospital (i.e., patients

not transferred from a district/acute hospital), although the latter two factors only to a lesser degree.

Longer lengths of stay were also protective of readmission, although the aOR for 60-90 days and ≥ 90 days

similar. There was no clear trend for admission years.

The alternative approach of using standard logistic regression on only the latest hospital admission for an

individual found similar results (Supplementary Table 1 and 2). Successful linkage to care had an aOR of

0.79 (95%CI 0.74-0.84) and 0.49 (95%CI 0.44-0.55) for readmission within 90 days at district/acute and

specialised hospitals respectively.

Supplementary Table 1: Standard logistic regression for the outcomes of successful linkage to care and
psychiatric readmission within 30 and 90 days for district/acute hospitals. Only the latest admission for an
individual was included in this analysis.

Supplementary Table 2: Standard logistic regression for the outcomes of successful linkage to care and
psychiatric readmission within 30 and 90 days for specialized hospitals. Only the latest admission for an
individual was included in this analysis.

Discussion

The main finding of this study is that a high proportion individuals who require psychiatric hospital

admission, are not routinely attending at ambulatory services, both before and after the admission,

suggesting that opportunities to avert psychiatric hospital admissions as well prevent repeat hospital

admissions are being missed. This is particularly pertinent, given the marked increase in psychiatric

hospital admissions that the province is experiencing, particularly at the district/acute level. Substance

related disorders are contributing to not only the increase in admission numbers being seen, but once an

individual is discharged home, those with a substance related disorder are less likely to successfully link

to ambulatory services. Successful linkage to care post-discharge was found to be the most protective

factor against repeat psychiatric admission.

While COVID-19 and its after effects are likely contributing to the current increase in psychiatric

admissions, it must be highlighted that the trend of increasing admissions, particularly at the district/acute

level, was already established prior to 2020, implying that other factors, such as increasing poverty,

inequality, substance use and trauma/violence, are driving this increase (18). Food insecurity and

neighborhood level deprivation were associated with depression, even after adjusting for individual

covariates, suggesting that structural poverty was contributing to the burden of mental illness in the South

Africa (5,6). A study conducted in Lentegeur Hospital in 2016 found that 62% of patients, and 73% of male

patients, used substances concurrently in the period leading up to their admission, with cannabis and

methamphetamine being most commonly used (7). While the substance rates we found were lower, they

were dependent on clinician ICD-10 coding, which is often not detailed enough, especially at the

district/acute hospital level.

Increased admission numbers can also be related to increased access to in-patient health services,

particularly in settings where out-patient services are limited. The district/acute hospitals with the highest

number of psychiatric admissions in 2022 were the new large district hospitals of Mitchells Plain and

Khayelitsha, which opened in 2013 and 2012 respectively. And the increasing size of the general

population over time, related to both births and migration into the province, should also not be forgotten

(18). Admission numbers at specialized hospitals, however, only showed a muted increase in numbers –

likely limited by bed capacity. This is in line with provincial policy which preferentially funded the

expansion of beds at the district/acute level (Circular H18/2015). According to a Provincial Database, the

number of official psychiatry beds in the Cape Town Metro was 417 in 2022 (a 136% increase from 2015),

while the three specialized hospitals only increased by 11% in the same time period, to 1 527 beds in 2022.

24% of mental health inpatients nationally are being readmitted to hospital within three months of being

discharged (14). Our study found slightly lower rates of 18-19% within 3 months, and no increase in the

readmission rate, except for the relative increase in district/acute hospitals in 2021, although with rising

absolute numbers of admissions this might still change. Scarcity of in-patient psychiatric beds has resulted

in premature discharge policies, where patients are discharged home before full recovery, leading to an

increased risk of readmission (19). Our analysis showed that shorter lengths of stay were associated with

an increased risk of readmission at specialized hospitals, but up to a certain point only, with lengths of

stay above 90 days not decreasing the risk of readmission. Interestingly longer lengths of stay at

district/acute hospitals were associated with an increased risk of readmission, possibly suggesting that

these patients might benefit from specialized care. At the same time direct admissions to specialized

hospitals had higher aORs for readmission, highlighting the importance of patients being managed at the

appropriate level of care.

In 2022, 40% of admitted psychiatric patients had not been seen at PHC facility in the preceding year. And

although some of these patients might still be in care, if they are being seen at private facilities, hospital

out-patient departments, or by community health workers, it remains that a large proportion of patients

requiring psychiatric admission are not routinely accessing care at the PHC level. The assumption that if

these patients were regularly being followed up at the PHC level a hospital admission could be averted

might not always be true, but by not following these patients up, we are potentially missing an opportunity

to intervene and decrease hospital admissions. This relatively stable median days between PHC visit and

psychiatric hospital admission of 35 to 42 days, despite the increasing proportion of individuals visiting a

PHC in the year prior to admission, reiterates this. Presumably patients receive their medication for 28

days, but do not return after, resulting in an exacerbation of symptoms and readmission soon after.

And while out-patient services were impacted by COVID-19, particularly in early 2020, a finding which is

supported by data from the South African private sector (20), the overall trend was of increasing, albeit

slowly, PHC use prior to admission. However, it is important to note that digitized PHC headcount data

became mandatory for provincial PHC facilities in April 2020, and so at least some of the increase we are

seeing is related to better electronic data capture rather than an actual change in PHC usage.

With regards to linkage post-discharge, male patients and those with substance related diagnoses were

less likely be linked after discharge, at both hospital levels. This is in line with data from a study in patients

with schizophrenia in Cape Town that found that substance use disorders were most strongly associated

with non-adherence (21). Ongoing psychotic symptoms, impaired insight and involvement with the

criminal justice system are factors have been found to be barriers to engagement with ambulatory

services (22). Those with longer hospital admissions were also more likely to be successfully linked to care,

possibly reflecting that those individuals have more severe disease and/or that during the longer hospital

stay more effort was put into preparing post-discharge care. Later admission years were associated with

more successful linkage to care, although electronic PHC headcount data improving over time might be

biasing this as well.

Successful linkage to care after discharge was one of the most protective factors against readmission

within 30 and 90 days, at both hospital levels. All the other covariates in these analyses are relatively fixed,

with the health system having limited control over them, the exceptions being length of hospital stay -

although with limited bed capacity this ability is also constrained (PHC visits in the year prior to admission

were associated with an increased risk of readmission, possibly indicating the chronicity of the underlying

disease more than the subsequent health seeking behaviour). Linkage to ambulatory psychiatric services

remains a longstanding challenge globally (23). And this study further highlights the importance of linkage

to care post-hospital discharge, which is a major challenge in the province, not limited to psychiatry only.

Patients diagnosed with tuberculosis in hospital, for instance, are more likely to be lost to follow up and

have higher mortality than those diagnosed at clinics (24). Effective clinical bridging strategies proposed

have included communicating discharge plans to outpatient clinicians, family involvement during the

hospital stay and starting outpatient programs before discharge (23).

Limitations

By using routinely collected electronic data, a limitation of this study is firstly that we were restricted to

admitted patients only, who are not necessarily representative of the burden of disease at the out-patient

level, and secondly, that we were dependent on diagnostic coding that was not always accurate or

sufficiently detailed. Also because of the challenges with diagnostic coding, we could only identify

patients by the admitting specialty of psychiatry, excluding psychiatric patients in Emergency Centres and

those admitted to other specialties. While this will underestimate the total number of psychiatric patients

in hospital, particularly for suicidal patients, the trends are still important, and the absolute numbers of

patients admitted to psychiatric wards remain crucial to those services. In addition, with the expansion of

electronic data use at PHC level during the study period, it is impossible to determine how much of the

increase in PHC use that is related to better data capturing. Further detailed studies assessing the mental

health burden at PHCs and Emergency Centres are needed.

For the readmission analyses, data on out-of-facility deaths and migration out of the province is not

currently available at the PHDC. If, for instance, after hospital discharge, an individual moves back to the

family home in another province and relapses there, they will not be counted as a readmission, artificially

lowering the readmission rate. And if there is differential out-migration by hospital drainage area, this

would contribute to the varying facility readmission rates. The very low readmission rates at rural facilities

might be partially explained by this. Private patients also access psychiatric services in the public sector,

but billing information in the source data was unfortunately too incomplete to be adjusted for. While the

stratification of these analyses by hospital level is reasonable, it does not consider the fact that patients

often move between the two levels over many years, and more comprehensive longitudinal analyses

taking into account the individual patient journeys through the health system are needed.

Conclusion

Driven by poverty and substance use, psychiatric admissions have increased markedly, particularly at the

district/acute hospital level, and have been exacerbated further by the COVID-19 pandemic. While

ambulatory service use before and after a psychiatric hospital admission is slowly increasing with time,

more can possibly be done at that level to avert hospital admissions. Successful linkage to ambulatory

services following hospital discharge is one of the most protective factors against readmission, and

stronger measures need to be put in place to ensure that all individuals discharged home receive

appropriate follow-up.

Funding

Funding for this study was provided by the Western Cape Department of Health and Wellness, as well as

a grant by the International Epidemiology Databases to Evaluate AIDS (IeDEA) from the Swiss National

Science Foundation. Grant number: ID-NO 7060

References

1.   Jacob N, Coetzee D. Mental illness in the Western Cape Province, South Africa: A review of the

burden of disease and healthcare interventions. South African Med J [Internet]. 2018 Mar 1

[cited 2022 Oct 7];108(3):176–80. Available from: https://pubmed.ncbi.nlm.nih.gov/30004359/

2.   Herman AA, Stein DJ, Seedat S, Heeringa SG, Moomal H, Williams DR. The South African Stress

and Health (SASH) study: 12-month and lifetime prevalence of common mental disorders. S Afr

Med J [Internet]. 2009 [cited 2022 Oct 7];99(5 Pt 2):339. Available from:

/pmc/articles/PMC3191537/

3.   Craig A, Rochat T, Naicker SN, Mapanga W, Mtintsilana A, Dlamini SN, et al. The prevalence of

probable depression and probable anxiety, and associations with adverse childhood experiences

and socio-demographics: A national survey in South Africa. Front Public Heal. 2022 Oct

28;10:4076.

4.   Bradshaw D, Norman R, Schneider M. A clarion call for action based on refined DALY estimates

for South Africa. South African Med J [Internet]. 2007 [cited 2022 Oct 7];Jun;97(6). Available

from: https://pubmed.ncbi.nlm.nih.gov/17691472/

5.   Dowdall N, Ward CL, Lund C. The association between neighbourhood-level deprivation and

depression: evidence from the south african national income dynamics study. BMC Psychiatry

[Internet]. 2017 Dec 11 [cited 2022 Oct 7];17(1). Available from:

https://pubmed.ncbi.nlm.nih.gov/29228912/

6.   Shepherd DL. Food insecurity, depressive symptoms, and the salience of gendered family roles

during the COVID-19 pandemic in South Africa. Soc Sci Med [Internet]. 2022 May 1 [cited 2022

Oct 7];301. Available from: https://pubmed.ncbi.nlm.nih.gov/35367907/

7.   Franken H, Parker J, Allen R, Wicomb RA. A profile of adult acute admissions to Lentegeur

478   Psychiatric Hospital, South Africa. S Afr J Psychiatr [Internet]. 2019 [cited 2022 Oct 7];25.

479   Available from: https://pubmed.ncbi.nlm.nih.gov/31616578/

480   8.   The Lancet Psychiatry. COVID-19 and suicide. The Lancet Psychiatry. 2021 Jun 1;8(6):451.

481   9.   Posel D, Oyenubi A, Kollamparambil U. Job loss and mental health during the COVID-19

482   lockdown: Evidence from South Africa. PLoS One [Internet]. 2021 Mar 1 [cited 2022 Oct 6];16(3).

483   Available from: https://pubmed.ncbi.nlm.nih.gov/33784339/

484   10.   Dawood B, Tomita A, Ramlall S. "Unheard," "uncared for" and "unsupported": The mental health

485   impact of Covid -19 on healthcare workers in KwaZulu-Natal Province, South Africa. PLoS One

486   [Internet]. 2022 May 1 [cited 2022 Oct 6];17(5). Available from:

487   https://pubmed.ncbi.nlm.nih.gov/35507540/

488   11.   Fernandes S, Marques N, Goga L. The complexity of neuropsychiatric manifestations of COVID-19

489   in South Africa. S Afr Med J [Internet]. 2022 [cited 2022 Oct 6];112(5):313–6. Available from:

490   https://pubmed.ncbi.nlm.nih.gov/35587242/

491   12.   Kola L, Kohrt BA, Hanlon C, Naslund JA, Sikander S, Balaji M, et al. COVID-19 mental health impact

492   and responses in low-income and middle-income countries: reimagining global mental health.

493   The Lancet Psychiatry. 2021 Jun 1;8(6):535–50.

494   13.   Harker N, Johnson K, Erasmus J, Myers B. COVID 19—impact on substance use treatment

495   utilization and provision in South Africa. Subst Abuse Treat Prev Policy [Internet]. 2022 Dec 1

496   [cited 2023 May 8];17(1). Available from: /pmc/articles/PMC8892117/

497   14.   Docrat S, Besada D, Cleary S, Daviaud E, Lund C. Mental health system costs, resources and

498   constraints in South Africa: a national survey. Health Policy Plan [Internet]. 2019 Nov 1 [cited

2022 Oct 6];34(9):706–19. Available from:

https://academic.oup.com/heapol/article/34/9/706/5572608

15.   Western Cape Government - Health. WCGH Annual Report 2021-2022 [Internet]. 2022. Available

from: https://www.westerncape.gov.za/assets/annual_report_2021-2022.pdf

16.   Pillay AL, Barnes BR. Psychology and COVID-19: impacts, themes and way forward. South African

J Psychol. 2020(2):148–53.

17.   Boulle A, Heekes A, Tiffin N, Smith M, Mutemaringa T, Zinyakatira N, et al. Data centre profile:

The provincial health data centre of the western cape province, South Africa. Int J Popul Data Sci.

2019 Nov 20;4(2).

18.   Western Cape Government. Socio-economic profile - City of Cape Town 2021 [Internet]. 2021.

Available from: https://www.westerncape.gov.za/provincial-treasury/files/atoms/files/SEP-LG

2021 - City of Cape Town.pdf

19.   Coetzee D, Koen L, Niehaus D, Botha U. Descriptive outcomes for a cohort of high-frequency

psychiatric service users in the Western Cape, South Africa after 10 years. S Afr J Psychiatr

[Internet]. 2022 [cited 2022 Oct 7];28:5. Available from:

https://pubmed.ncbi.nlm.nih.gov/35747340/

20.   Wettstein A, Tlali M, Joska JA, Cornell M, Skrivankova VW, Seedat S, et al. The effect of the

COVID-19 lockdown on mental health care use in South Africa: an interrupted time-series

analysis. Epidemiol Psychiatr Sci [Internet]. 2022 Jun 15 [cited 2022 Oct 6];31. Available from:

https://pubmed.ncbi.nlm.nih.gov/35703078/

21.   Benelmokhtar JM, Chiliza B, Phahladira L, Emsley R, Asmal L. Factors associated with dropout at 2

years post-initiation of treatment in the first episode of schizophrenia. S Afr J Psychiatr [Internet].

2021 [cited 2023 May 8];27. Available from: https://pubmed.ncbi.nlm.nih.gov/33824760/

22.   Compton MT. Barriers to initial outpatient treatment engagement following first hospitalization

for a first episode of nonaffective psychosis: a descriptive case series. J Psychiatr Pract [Internet].

2005 Jan [cited 2023 May 8];11(1):62–9. Available from:

https://pubmed.ncbi.nlm.nih.gov/15650625/

23.   Boyer CA, McAlpine DD, Pottick KJ, Olfson M. Identifying risk factors and key strategies in linkage

to outpatient psychiatric care. Am J Psychiatry [Internet]. 2000 [cited 2023 May 8];157(10):1592–

8. Available from: https://pubmed.ncbi.nlm.nih.gov/11007712/

24.   Osman M, Meehan SA, von Delft A, Du Preez K, Dunbar R, Marx FM, et al. Early mortality in

tuberculosis patients initially lost to follow up following diagnosis in provincial hospitals and

primary health care facilities in Western Cape, South Africa. PLoS One [Internet]. 2021 Jun 1

[cited 2023 Mar 26];16(6):e0252084. Available from:

https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0252084

Number of new admissions

Hospital level ● Specialized ● Acute

14-day moving average of daily psychiatric admissions for the province - district/acute hospitals only

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

If discharged from district/acute hospital, first contact at clinic/OPD in 30 days or less

● Admission ● No ● No contact ● Yes

Figure 6

Figure 7

	Admissions	District/acute n = 74 432		Specialized n = 30 214	
		n	%	n	%
Sex	Male	45159	60.67	19971	66.10
	Female	29273	39.33	10243	33.90
Age category	18-29 years	31905	42.87	13437	44.48
	30-39 years	20481	27.52	9272	30.69
	40-49 years	11181	15.02	4555	15.08
	≥50 years	10850	14.58	2948	9.76
Diagnostic categories	Substance related	15140	20.34	6077	20.11
	Schizophrenia	21653	29.09	16688	55.23
	Mood disorders	11058	14.86	5919	19.59
	Other	26581	35.71	1530	5.06
	Primary substance related diagnosis	15140	20.34	6077	20.11
	Schizophrenia with any substance	2905	3.90	4636	15.34
	Mood/other with any substance	2007	2.70	1396	4.62
	Schizophrenia without any substance	18748	25.19	12052	39.89
	Mood/other without any substance	35632	47.87	6053	20.03
Previous admissions	Index admission	37677	50.62	8340	27.60
	≥1 previous admissions	36755	49.38	21874	72.40
PHC visit in year before admission	No	31531	42.36	11286	37.35
	Yes	42901	57.64	18928	62.65
Length of stay	1-3 days	17826	23.95		
	4-6 days	17421	23.41		
	7-13 days	18167	24.41		
	14-27 days	13995	18.80		
	≥28 days	7023	9.44		
	1-13 days			1717	5.68
	14-29 days			8575	28.38
	30-59 days			12138	40.17
	60-89 days			4161	13.77
	≥90 days			3623	11.99
Admission year	2015	5945	7.99	3500	11.58
	2016	7132	9.58	3573	11.83
	2017	8149	10.95	3321	10.99
	2018	9080	12.20	3451	11.42
	2019	9915	13.32	3824	12.66
	2020	9376	12.60	3820	12.64
	2021	11966	16.08	4383	14.51
	2022	12869	17.29	4342	14.37
Admitting hospital	Metro Catchment A	20502	27.54		
	Metro Catchment B	10674	14.34		
	Metro Catchment C	17273	23.21		
	Rural hospitals	25983	34.91		
	Specialized hospital A			8702	28.80
	Specialized hospital B			8859	29.32
	Specialized hospital C			12653	41.88
Outpatient linkage to care <30days	No	36381	48.88	9586	31.73
	Yes	38051	51.12	20628	68.27
Direct admission	No			24770	81.98
	Yes			5444	18.02

Table 1

		District/acute hospitals								
		Successful linkage to outpatient services <30 days			**Repeat psychiatric hospital admission <30 days**			**Repeat psychiatric hospital admission <90 days**		
	Admissions	n = 74 400			n = 74 400			n = 73 183		
	Individuals	n = 47 153			n = 47 153			n = 46 508		
		aOR	95% CI		aOR	95% CI		aOR	95% CI	
Sex	Male	Ref			Ref			Ref		
	Female	1.10	1.06	1.14	0.94	0.89	1.00	0.89	0.85	0.94
Age category	18-29 years	Ref			Ref			Ref		
	30-39 years	1.03	0.99	1.08	0.82	0.76	0.87	0.86	0.81	0.90
	40-49 years	1.22	1.15	1.28	0.79	0.73	0.86	0.85	0.79	0.90
	50-59 years	1.28	1.21	1.35	0.75	0.69	0.82	0.80	0.75	0.86
Primary diagnosis	Substance related	Ref			Ref			Ref		
	Schizophrenia	1.23	1.17	1.30	1.26	1.17	1.36	1.28	1.20	1.36
	Mood disorders	1.43	1.34	1.52	1.29	1.18	1.41	1.16	1.07	1.25
	Other	1.19	1.13	1.25	1.02	0.95	1.10	0.93	0.88	0.99
Previous admissions	Index admission	Ref			Ref			Ref		
	≥1 previous admissions	1.10	1.06	1.15	1.97	1.86	2.09	2.46	2.35	2.58
PHC visit in year before admission	No	Ref			Ref			Ref		
	Yes	2.67	2.58	2.77	1.11	1.05	1.18	1.18	1.13	1.24
Length of stay	1-3 days	Ref			Ref			Ref		
	4-6 days	1.36	1.29	1.43	1.33	1.23	1.43	1.26	1.18	1.35
	7-13 days	1.69	1.61	1.78	1.47	1.36	1.59	1.39	1.30	1.48
	14-27 days	2.00	1.89	2.11	1.23	1.13	1.34	1.18	1.10	1.27
	≥28 days	2.33	2.17	2.49	0.81	0.73	0.91	0.83	0.76	0.91
Admission year	2015	Ref			Ref			Ref		
	2016	1.03	0.95	1.12	0.93	0.83	1.04	0.91	0.83	1.00
	2017	1.10	1.02	1.19	0.88	0.79	0.99	0.88	0.80	0.96
	2018	1.10	1.02	1.19	0.87	0.78	0.97	0.86	0.78	0.94
	2019	1.10	1.02	1.18	0.77	0.68	0.86	0.79	0.72	0.87
	2020	1.11	1.03	1.20	0.77	0.69	0.86	0.77	0.70	0.85
	2021	1.18	1.09	1.27	0.83	0.74	0.92	0.86	0.79	0.95
	2022	1.24	1.16	1.34	0.78	0.70	0.86	0.80	0.73	0.88
Admitting hospital	Metro Catchment A	Ref			Ref			Ref		
	Metro Catchment B	1.02	0.96	1.08	0.93	0.86	1.02	0.87	0.81	0.93
	Metro Catchment C	1.00	0.95	1.05	0.91	0.85	0.98	0.91	0.86	0.97
	Rural hospitals	0.96	0.92	1.01	0.79	0.74	0.85	0.72	0.68	0.76
Outpatient linkage to care <30days	No				Ref			Ref		
	Yes				0.69	0.65	0.72	0.76	0.73	0.80

Table 2

		Specialized hospitals								
		Successful linkage to outpatient services			Repeat psychiatric hospital admission in 30 days			Repeat psychiatric hospital admission in 90 days		
	Admissions	n = 30 209			n = 30 209			n = 29 767		
	Individuals	n = 17 214			n = 17 214			n = 17 015		
		aOR	95% CI		aOR	95% CI		aOR	95% CI	
Sex	Male	Ref			Ref			Ref		
	Female	1.16	1.08	1.25	1.03	0.91	1.16	0.91	0.83	1.00
Age category	18-29 years	Ref			Ref			Ref		
	30-39 years	0.88	0.82	0.95	0.90	0.80	1.02	1.04	0.94	1.14
	40-49 years	0.93	0.85	1.03	0.76	0.65	0.90	0.99	0.87	1.12
	50-59 years	1.11	0.98	1.24	0.93	0.77	1.12	1.15	0.99	1.33
Diagnostic categories	y substance related diagnosis	Ref			Ref			Ref		
	Schizophrenia with any substance	1.05	0.95	1.16	1.38	1.15	1.65	1.42	1.23	1.63
	Mood/other with any substance	0.97	0.84	1.13	1.40	1.08	1.80	1.53	1.26	1.87
	Schizophrenia without any substance	1.12	1.03	1.22	1.28	1.10	1.49	1.37	1.21	1.54
	Mood/other without any substance	1.24	1.13	1.38	1.53	1.29	1.81	1.36	1.19	1.56
Previous admissions	Index psychiatric admission	Ref			Ref			Ref		
	≥1 previous admissions	0.89	0.83	0.95	2.04	1.78	2.35	2.39	2.15	2.67
PHC visits in year prior to admission	No	Ref			Ref			Ref		
	Yes	2.05	1.93	2.18	1.30	1.17	1.45	1.39	1.28	1.52
Length of stay	1-13 days	Ref			Ref			Ref		
	14-29 days	1.16	1.02	1.32	0.68	0.57	0.82	0.81	0.69	0.95
	30-59 days	1.27	1.12	1.44	0.51	0.43	0.62	0.63	0.54	0.73
	60-89 days	1.36	1.18	1.57	0.39	0.31	0.49	0.53	0.44	0.63
	≥90 days	1.57	1.35	1.83	0.39	0.31	0.49	0.53	0.44	0.63
Admission year	2015	Ref			Ref			Ref		
	2016	1.11	0.99	1.24	0.95	0.78	1.14	0.89	0.77	1.04
	2017	1.26	1.12	1.42	0.91	0.75	1.10	0.93	0.80	1.08
	2018	1.26	1.12	1.42	0.91	0.75	1.11	0.93	0.80	1.08
	2019	1.33	1.19	1.50	0.82	0.67	0.99	0.94	0.81	1.09
	2020	1.32	1.18	1.49	0.83	0.68	1.01	0.87	0.74	1.01
	2021	1.41	1.26	1.58	0.82	0.67	0.99	0.91	0.78	1.05
	2022	1.43	1.28	1.60	0.96	0.79	1.15	1.00	0.87	1.16
Direct admission	No	Ref			Ref			Ref		
	Yes	0.97	0.90	1.05	1.15	1.01	1.30	1.15	1.04	1.27
Admitting hospital	Specialized hospital A	Ref			Ref			Ref		
	Specialized hospital B	1.24	1.14	1.34	0.71	0.61	0.82	0.68	0.60	0.76
	Specialized hospital C	1.18	1.09	1.27	1.03	0.91	1.17	0.97	0.87	1.07
Successful linkage to care <30 days	No				Ref			Ref		
	Yes				0.37	0.33	0.40	0.45	0.42	0.49

Table 3