Title Page

Prepared according to manuscript instruction for an article in *Scientific Reports*

Version: Mar 20th, 2024

Title: HIV co-infection increases the risk of post-tuberculosis mortality among persons who

initiated treatment for drug-resistant tuberculosis

Running Title: HIV co-infection and post-TB mortality

Authors:

1. **Argita D. Salindri***, Department of Population Health Sciences, Georgia State

University School of Public Health, Atlanta, GA, USA; and Division of Infectious

Diseases and Geographic Medicine, Department of Medicine, Stanford University

School of Medicine, Stanford, CA, USA (Email: adsalind@stanford.edu)

2. **Maia Kipiani**, National Center for Tuberculosis and Lung Diseases, Tbilisi, Georgia;

David Tvildiani Medical University, Tbilisi, Georgia; and The University of Georgia,

Tbilisi, Georgia (Email: maiagegechkori@yahoo.com)

3. **Nino Lomtadze**, National Center for Tuberculosis and Lung Diseases, Tbilisi, Georgia;

David Tvildiani Medical University, Tbilisi, Georgia; and The University of Georgia,

Tbilisi, Georgia (Email: nlomtadze@gmail.com)

4. **Nestani Tukvadze**, National Center for Tuberculosis and Lung Diseases, Tbilisi,

Georgia; and Department of Medicine, Swiss Tropical and Public Health Institute,

Allschwil, Switzerland (Email: nestani.tukvadze@swisstph.ch)

5. **Zaza Avaliani**, National Center for Tuberculosis and Lung Diseases, Tbilisi, Georgia;

and European University, Tbilisi, Georgia (Email: avalianizaza@yahoo.com)

6. **Henry M. Blumberg**, Division of Infectious Diseases, Department of Medicine, Emory University School of Medicine, Atlanta, GA, USA; Hubert Department of Global Health and Department of Epidemiology, Emory University Rollins School of Public Health, Atlanta, GA, USA (Email: hblumbe@emory.edu)

7. **Katherine E. Masyn**, Department of Population Health Sciences, Georgia State University School of Public Health, Atlanta, GA, USA, Atlanta, GA, USA (Email: kmasyn@gsu.edu)

8. **Richard B. Rothenberg**, Department of Population Health Sciences, Georgia State University School of Public Health, Atlanta, GA, USA, Atlanta, GA, USA (Email: rrothenberg@gsu.edu)

9. **Russell R. Kempker**, Division of Infectious Diseases, Department of Medicine, Emory University School of Medicine, Atlanta, GA, USA (Email: rkempke@emory.edu)

10. **Matthew J. Magee**, Hubert Department of Global Health and Department of Epidemiology, Emory University Rollins School of Public Health, Atlanta, GA, USA; and Division of Infectious Diseases, Department of Medicine, Emory University School of Medicine, Atlanta, GA, USA (Email: mjmagee@emory.edu)

Title: 16/20

Abstract: 200/200

Main Article Word Count: 3805/4500

Table/Figure Counts: 6/8

References: 41/60

Keywords: tuberculosis, post-TB mortality; comorbidities; HIV co-infection; diabetes; hepatitis C

*Corresponding Author:

Argita Salindri, PhD, MPH

Division of Infectious Diseases and Geographic Medicine

Department of Medicine, Stanford University School of Medicine

Email: asalind@stanford.edu

Phone: +1 404 490 5906

Alternate Corresponding Author:

Matthew Magee, PhD, MPH

Hubert Department of Global Health

Emory University Rollins School of Public Health

Email: mjmagee@emory.edu

Phone: +1 404 727 9814

ABTSRACT

Little is known regarding the relationship between common comorbidities in persons with

tuberculosis (TB) (including human immunodeficiency virus [HIV], diabetes, and hepatitis C

virus [HCV]) with post-TB mortality. We conducted a retrospective cohort study among persons

who initiated treatment for rifampicin-resistant and multi/extensively drug-resistant (RR and

M/XDR) TB reported to the country of Georgia's TB surveillance during 2009-2017. Exposures

included HIV serologic status, diabetes, and HCV status. Our outcome was all-cause post-TB

mortality determined by cross-validating vital status with Georgia's death registry through

November 2019. We estimated adjusted hazard rate ratios (aHR) and 95% confidence intervals

(CI) of post-TB mortality among participants with and without comorbidities using cause-specific

hazard regressions. Among 1032 eligible participants, 34 (3.3%) died during treatment and 87

(8.7%) died post-TB treatment. Among those who died post-TB treatment, the median time to

death was 21 months (interquartile range 7–39) post-TB treatment. After adjusting for

confounders, the hazard rates of post-TB mortality were higher among participants with HIV co-

infection (aHR=3.74, 95%CI 1.77–7.91) compared to those without HIV co-infection. In our

cohort, post-TB mortality occurred most commonly in the first three years post-TB treatment.

Linkage to care for common TB comorbidities post-treatment may reduce post-TB mortality

rates.

INTRODUCTION

Tuberculosis (TB) remains a leading cause of infectious disease death with an estimated mortality rate of nearly 1.6 million persons worldwide annually.[1] The END TB strategy aims to reduce TB mortality from 2015 to 2035 by 95%.[2] Importantly, the reported number of TB deaths is likely underestimated because TB-attributable mortality after treatment is not typically measured or reported.[3] More importantly, WHO reported that TB related mortality consistently increased for the first time after more than a decade since 2020, likely due to the COVID-19 pandemic.[1] To meet the END TB goals, it is essential to better understand TB-related mortality including those that occur after the completion of TB treatment.

Emerging epidemiologic evidence suggests that individuals formerly treated for TB disease may have higher mortality rates compared to the general population.[3,4] Despite microbial cure and favorable clinical outcomes, individuals formerly treated for TB disease may have residual sequelae including pulmonary impairment or increased risk of non-communicable metabolic diseases.[5-8] However, to date, little is known regarding the relationship between pre-existing comorbidities (i.e., comorbidities diagnosed prior to or at the beginning of TB treatment) and post-TB mortality. To substantially reduce TB-related mortality, it is necessary to identify host risk factors, including pre-existing comorbidities, that are associated with post-TB mortality.

Previous studies have identified several demographic and behavioral characteristics associated with increased risks of post TB-mortality.[4,9-13] For example, highly drug-resistant forms of TB such as multidrug-resistant TB (MDR TB) are associated with higher rates of mortality post-TB treatment.[9,14,15] However, the effect of common pre-existing comorbidities such as human immunodeficiency virus (HIV) co-infection, hyperglycemia (i.e., elevated blood glucose with or without diabetes), diabetes, or hepatitis C virus (HCV) co-infection remains unclear. Understanding the extent to which these pre-existing comorbidities impact post-TB mortality rates will help identify prevention strategies to reduce post-TB mortality rates.

Given existing knowledge gaps, we aimed to a) estimate the rate of all-cause mortality

post-TB treatment and b) determine the association between pre-existing comorbidities and

post-TB treatment mortality rates using data of persons who initiated treatment with second line

TB drugs (SLDs) from the country of Georgia.

RESULTS

Study Population and Characteristics

During the study period, there were 5,385 RR and M/XDR TB treatment episodes

recorded in the national TB surveillance system, 31.5% (1,697/5,385) of which were classified

as newly diagnosed RR and M/XDR TB (i.e., no prior history of TB treatment) (Figure 1). Of

these, 1,416 (83.4%) unique case met study eligibility criteria and were cross-referenced with

the country of Georgia's vital registry. We excluded 384 individuals without unique national

identifier, leaving 1,032 individuals (72.9%) included in our analyses (Figure 1). Individuals

excluded due to missing vital status were similar to individuals included in the analyses in

regard to age, gender, prevalence of MDR or pre-XDR TB, cavitary disease, hyperglycemia,

and hepatitis C and HIV co-infections. Among individuals included, the majority were male

(73%), and median age was 35 years (interquartile range [IQR] 26 – 49) (Table 1). The

prevalence of MDR TB was 44.7% (461/1,032) and pre-XDR TB was 29.1% (300/1,032). HIV

co-infection was reported among 3.8% (39/1,032, 95% confidence interval [CI] 2.7 – 5.1) and

the prevalence of any hyperglycemia was 22.7% (234/1,032, 95%CI 20.2 – 25.3), 115 (49.1%

[115/234], 95%CI 42.8 – 55.5) of which were determined to have diabetes. HCV co-infection

was reported among 14.7% (153/1,032, 95%CI 12.8 – 17.1) of study participants.

Among 1,032 individuals included in the final analyses, 34 (3.3%) died during TB

treatment during 1502 PY follow-up (age-adjusted mortality rate during TB treatment=22.6 per

1,000 PY, 95%CI 17.9 – 27.4) with a median time to death of 4 months (IQR 1 – 7) after TB

treatment initiation. There were 87 (8.7%) post-TB deaths among 998 individuals alive at the

end of their TB treatment during 4,857 PY of follow up (age-adjusted post-TB mortality

rate=17.9 per 1,000 PY, 95%CI 11.9 – 23.9) (Table 2). Among those who died during TB

treatment, the median age at time of death was 48 years (IQR 34 – 52). Nearly half (38/87,

43.7% 95%CI 33.6 – 54.2) of post-TB deaths were observed during the first year after TB

treatment end, and the majority (67/87, 77.0%) occurred within the first three years with a

median time to death of 21 months (IQR 7 – 39) after TB treatment was completed and/or

stopped. Among those who died after-TB treatment, the median age at time of death was 50

years (IQR 39 – 63).

Pre-Existing Comorbidities and All-Cause Mortality Post-Tuberculosis Treatment

Among 32 persons living with HIV co-infection, there were 9 deaths post-TB treatment

during 130 person-years (post-TB mortality rate 69.23, 95%CI 36.02 – 133.10). The median age

at time of post-TB death among those with HIV co-infection was 41 years (IQR 39 – 46). The

post-TB mortality rates among those with pre-diabetes was 21.92 (95%CI 12.73 – 37.75) and

26.12 (95%CI 15.47 – 44.10) among those with diabetes. The median age at time of post-TB

death among those with pre-diabetes was 48 years (IQR 35 – 59) and 63 years (IQR 49 – 68)

among those with diabetes. Among 147 individuals with HCV co-infection, there were 18 deaths

post-TB treatment during 701 person-years (post-TB mortality rate 25.68, 95%CI 16.18 –

40.76). The median age at time of post-TB death among those with HCV co-infection was 57

years (IQR 39 – 62). Among 641 individuals without HIV, hyperglycemia, or HCV co-infection

(i.e., no primary comorbidity), there were 41 deaths post-TB treatment during 3128 person-

years (post-TB mortality rate 13.11, 95%CI 9.65 – 17.80) with the median age at time of post-TB

death of 49 years (IQR 38 – 59).

Persons living with HIV co-infection had significantly higher hazard rate of post-TB

mortality (adjusted hazard rate ratio [aHR]=3.74, 95%CI 1.77 – 7.91) compared to those without

HIV co-infection after adjusting for potential confounders (Table 3). In the same adjusted model,

the hazard rate of post-TB mortality among TB survivors with prediabetes was 1.48 (95%CI 0.78

– 2.78) and 1.06 (95%CI 0.53 – 2.12) among those with diabetes compared to those with

euglycemic. The hazard rate of all-cause post-TB mortality among TB survivors with HCV co-

infection was 1.21 (95%CI 0.68 – 2.59) times the hazard rate among those without HCV co-

infection. The adjusted hazard rate of all-cause post-TB mortality among TB survivors with any

primary comorbidity (i.e., HIV, prediabetes/diabetes, and HCV) was significantly higher

(aHR=2.03 (95%CI 1.33 – 3.10) compared to those without. Our estimates did not change

substantially in models with different covariates specification (Supplementary Table S1). We

also provided post-TB mortality rates (Supplementary Table S2) and estimated hazard rate

ratios (Supplementary Table S3) of post-TB mortality for other key risk factors (included other

comorbidities, smoking status, and baseline BMI).

Results from Subgroup, Interaction, and Sensitivity Analyses

Among individuals with known smoking status (n=591), we found that the multiplicative

effects of HIV co-infection and smoking, hyperglycemia and smoking, or HCV and smoking on

mortality post-TB treatment were non-significant (statistical interaction p>0.05) (Table 4). We

found that post-TB mortality rates were highest at 6-month post TB treatment among those with

HIV co-infection, pre-diabetes/diabetes or HCV (Figure 2). Among those with HIV co-infection,

pre-diabetes/diabetes, or HCV co-infection, the median time from the end of TB treatment to

post-TB deaths were lowest among those aged 16 – 40 years (median=13 month, IQR 5 – 22)

and ≥65 years (median=14 month, IQR 14 – 25) compared to those among age 41 – 65 years

(median=27 month, IQR 6 – 39). Among those with known final TB treatment end (n=980), the

hazard rate ratios of mortality post-TB treatment were similar among persons living with HIV or

pre-diabetes/diabetes regardless of their final TB treatment outcome (Supplementary Table S4).

However, those with HCV co-infection who were successfully treated for TB were more likely to

die post-TB treatment compared to those with HCV co-infection but had poor final TB treatment

outcomes.

In the log-binomial models to estimate the cumulative risk of post-TB mortality, the

cumulative risk of post-TB mortality among those living with HIV co-infection was 3.63 (95%CI

1.99 – 6.61) times the risk of those without HIV co-infection. The cumulative risk of post-TB

mortality, although non-significant, was modestly higher among those with hyperglycemia (cRR

1.59, 95%CI 1.00 – 2.52) and HCV co-infection (cRR 1.53, 95%CI 0.92 – 2.58) (Supplementary

Table S5).

The median of bias-adjusted RRs after accounting for any hyperglycemia

misclassification was 1.84 (2.5^{th} – 97.5^{th} percentile 1.10 – 2.92) in the non-differential model vs.

1.73 (2.5^{th} – 97.5^{th} percentile 1.21 – 2.37) in the differential models. This indicates that

misclassification of any hyperglycemia in our cohort may result in bias towards the null. The E-

values of ≥6.72, ≥2.56, and ≥2.45 could explain away the observed association for HIV co-

infection, hyperglycemia, HCV, and post-TB mortality risk.

METHODS

Study Design and Setting

We conducted a retrospective cohort study among adults with (>15 years old) newly

diagnosed and laboratory-confirmed pulmonary TB. Persons initiated treatment with SLDs

(including rifampicin-resistant and multidrug/extensively drug-resistant TB [RR and M/XDR TB])

and reported to the Georgian National Center for Tuberculosis and Lung Disease (NCTLD)

surveillance system from January 2009 – December 2017 were eligible. Persons with drug-

susceptible TB, persons with drug-resistant TB where strains were susceptible to rifampicin, and

those with a history of prior TB treatment were excluded from analyses. We limited our analyses

to those with newly diagnosed TB disease as Georgia's online TB surveillance developed

overtime and collection of prior TB information may not be uniform throughout our study period.

More importantly, we believe that the underlying TB susceptibility profile of individuals with

newly diagnosed TB may be different than those with a history of TB disease, thus, affecting the

counterfactual conditions. Briefly, per Georgian National TB guidelines, standard of care

performed at TB treatment initiation for those treated with SLDs includes complete blood count,

chest X-ray, sputum smear and culture, drug susceptibility testing (DST), fasting plasma

glucose (FPG), and rapid test of HIV infection. Additional tests to screen for other chronic

comorbidities like viral hepatitis (hepatitis C and B) infections were done since 2008.

Definitions and Study Measures

The primary exposures of this study were comorbidities diagnosed before or at TB

treatment initiation, including a) HIV co-infection, b) any hyperglycemia, and c) HCV co-

infection. HIV and HCV virus (HCV) co-infections were determined by antibody test performed

at the beginning of TB treatment and recorded in the surveillance system. Any hyperglycemia

was determined by either a self-reported previous diagnosis of type-2 diabetes (T2DM), or by

FPG. Individuals with FPG $\geq$5.6 mmol/L or a self-reported prior T2DM diagnosis were

categorized as "any hyperglycemia." Individuals with FPG level <5.6mmol/L and no history of

diabetes diagnosis were classified as "non-hyperglycemic." We then separately defined

diabetes status using three different levels: 1) no diabetes (FPG < 5.6 mmol/L with no previous

diagnosis of T2DM), 2) pre-diabetes (FPG 5.6 – 6.9 mmol/L with no previous diagnosis of

T2DM), and 3) diabetes (FPG $\geq$7.0 mmol/L or previous diagnosis of T2DM). "Any primary

comorbidity" was defined if individuals had a record of either HIV co-infection, any

hyperglycemia, or HCV co-infection. Other comorbidities recorded in the surveillance system

(including self-reported cardiovascular disease, kidney disease, ulcer, pancreatitis, chronic

obstructive pulmonary disease, silicosis, and sarcoidosis) were considered in the analyses by

summing the number of comorbidities and classified as "none", "1 – 2", and "$\geq$3".

Our primary study outcome, all-cause mortality post-TB treatment, was determined by

cross-referencing individuals' national unique identifier with mortality status recorded in the

death registry managed by the National Statistics Office of Georgia. Drug resistance was

determined according to phenotypic DST results and were classified as rifampicin-resistant

(RR), multidrug (MDR), pre-extensively (pre-XDR), and extensively drug-resistant (XDR) TB

according to WHO guidelines.[16] Other covariates included in our study (e.g., clinical TB

treatment outcomes, self-reported smoking status, anthropometric measures) were defined

according to surveillance records.

Statistical Analyses

Chi-square/Fisher's exact and Wilcoxon rank-sum tests were performed to assess

associations between individuals' comorbidities status and post-TB mortality (p-values were not

shown). Poisson regression was used to estimate rates of all-cause mortality post-TB treatment

(per 1,000 person-years [PY]) and rate differences (using identity link). We also built life tables

to independently estimate the mortality rates per 6-month follow-up period during and after TB

treatment. These mortality rates were then plotted with *ggplot2* package [17] in R Statistical

Software (v4.2.3; R Core Team 2023) [18] to observe mortality trend during and after TB

treatment. Proportional hazard models with competing risks were used to estimate the hazard

rate ratios (HR) of all-cause mortality during and post-TB treatment comparing individuals with

and without pre-existing comorbidities.[19] Survival time was calculated from TB treatment

initiation to a) time of death as recorded in the online TB surveillance system (for those who

died during TB treatment), b) date of death recorded in the national death registry (for those

who died after TB treatment), or c) the end of follow-up time for those who did not have the

study events (i.e., alive by November 13, 2019, when mortality queries were cross-referenced

with the vital registry's office). In the cause-specific models estimating the HR of mortality post-

TB treatment, individuals were censored if they a) died during TB treatment (i.e., competing risk)

or b) had no death indication on the date mortality status was verified. Proportional hazard

assumptions were assessed using 1) Kolmogorov-type supremum and 2) Schoenfeld's

residuals tests.[20] Covariates included in the final multivariable models were selected considering

established risk factors identified in previous studies, observed bivariate associations, and

directed acyclic graphs.[21]

Stratified analyses were performed to determine whether the effect of pre-existing

comorbidities on all-cause mortality post-TB treatment varied among those who had favorable

(n=655) vs. poor TB treatment outcomes (i.e., including those who refused to continue

treatment or were lost to follow-up) (n=325). Additionally, we performed a subgroup analysis to

assess the interactions between pre-existing comorbidities and smoking status to determine if

the association between pre-existing comorbidities and all-cause mortality post-TB treatment

varied by smoking status. We assessed statistical interaction by including the cross-product

terms (e.g., diabetes X smoking, HIV X smoking) in multivariable models. Statistical analyses

were performed using SAS version 9.4 (Cary, North Carolina).

Sensitivity Analyses

Sensitivity analyses were performed to quantify systematic errors (i.e., bias) due to a)

distribution specification in the regression analyses, b) misclassification of any hyperglycemia

status and c) confounding effects from unmeasured confounders. To assess distribution

specification, log-binomial models were used to estimate cumulative risks of post-TB mortality

comparing patients with comorbidities to those without comorbidities. Bias due to

misclassification of hyperglycemia was of concern as we did not have a way to determine

whether patients followed doctors' instruction to fast at least 8 hours before test. To quantify

bias due to hyperglycemia misclassification, we performed probabilistic bias analyses using

beta distribution. To quantify systematic errors due to unmeasured/unknown confounders, we

calculated E-values. E-value is an estimate of the minimum strength of association between

unmeasured/unknown confounders with study exposures as well as study outcome to explain

away the observed association between pre-existing comorbidities and all-cause mortality post-

TB treatment.[22]

Ethics Declaration and Approval

This study was submitted to, reviewed, and approved by the Institutional Review Boards (IRBs)

at Georgia State University, Emory University, Atlanta, USA and the ethics committee at

National Center for Tuberculosis and Lung Diseases, Tbilisi, Georgia (FWA00020831). Waiver

of informed consent was obtained as data collected for this study were collected for TB

surveillance purposes.

DISCUSSION

In our large cohort of persons who initiated treatment with SLDs (n=1,032 with a median

of follow-up time of 5 years), nearly 1 among 11 died post-TB treatment. We reported a higher

rate of all-cause mortality rate in our cohort of persons treated for RR and M/XDR when

compared to mortality rates of Georgia's general population in 2019 (17.3 vs. 12.5/1,000

person-years).[23] This increased rate of post-TB mortality is consistent with previously published

studies in different settings.[3,4,10,14,24-26] Furthermore, we found that post-TB mortality occurred

mostly in the first three years after TB treatment ended. Additionally, we observed a significantly

increased post-TB mortality risk among persons living with HIV co-infection compared to those

without HIV co-infection. Understanding the effect of pre-existing comorbidities (especially HIV

co-infection) on post-TB mortality could help determine whether additional care post-TB

treatment completion is needed to reduce premature deaths associated with TB and

comorbidities.

In our cohort, we reported nearly 4-fold increased hazard rates of all-cause mortality

post-TB treatment among persons with RR and M/XDR and HIV co-infection compared to those

without. However, this finding needs to be interpreted with caution as the number of persons

living with HIV in our cohort was small. Furthermore, we did not have any clinical HIV (e.g., CD4

count) to describe the severity of the disease manifestation. Similar to our findings, a

prospective cohort study conducted in Vietnam reported that the hazard rate of post-TB deaths

among persons with TB and HIV was 5.1 (95%CI 4.2 – 6.3) times the hazard rate among

individuals with TB but without HIV.[25] However, this hazard estimation combined individuals who

died before and after TB treatment. Another study using data from the HIV Epidemiology in

South American networks reported a higher hazard rate of death at five years after TB treatment

completion among person living with HIV and diagnosed with TB within 30 days of enrollment to

the HIV clinic vs. those without TB (aHR 1.57, 95%CI 1.25 – 1.99).[27] In the same study, the

authors reported that among persons with TB and HIV, a lower CD4 count was associated with

higher rates of deaths post-TB treatment (aHR 1.57, 95%CI 1.41 – 1.76). Unlike this South

America's study, which only included persons with drug-sensitive TB (DSTB), our study only

included individuals with RR and M/XDR TB, and this may partially explain the higher effect of

HIV on post-TB mortality reported in our study. Furthermore, our study also had longer follow-up

time (up to eight years of follow-up time) and we utilized competing risks model to distinctly

modeled the hazard rates of deaths observed during and after TB treatment.

Hyperglycemia has been associated with increased risk of mortality during TB

treatment.[28-30] However, the association between hyperglycemia and the risk of post-TB

mortality is not well characterized. We reported a non-significantly higher crude rates as well as

the relative effect for post-TB treatment mortality among TB survivors with diabetes compared to

those with pre-diabetes or euglycemic. A retrospective cohort study using data from the vital

statistics conducted in Israel found that there were less deaths due to diabetes among

individuals successfully treated for TB disease compared to the general population.[4] In contrast,

a prospective study conducted in Mexico reported a higher proportion of non-TB related deaths

post-TB treatment among individuals with diabetes (4.9%) compared to those without diabetes

(2.5%).[31] The inconsistent findings of the association between pre-existing diabetes and the risk

of post-TB deaths may be affected by the use of blood glucose-lowering agents and blood

glucose management during and post-TB treatment. For instance, metformin use was

associated with reduced mortality risk among individuals with TB in a study conducted in

Taiwan.[32] Prospective clinical studies aiming to assess the impact of diabetes clinical

characteristics (including the use of blood glucose-lowering agents during TB treatment) on the

risk of mortality post-TB deaths are still warranted.

In our cohort, TB survivors with HCV co-infection had 25% higher rate of post-TB

mortality compared to those without HCV co-infection. It is unclear why individuals with HCV

who were successfully treated for TB were more likely to die after TB treatment is ended when

compared to those with poor TB treatment outcomes. More clinical data (including information

whether individuals had acute vs. chronic infections) are needed to understand this finding. To

our knowledge, to date, there are limited studies looking at the long-term outcomes of

individuals treated for TB with HCV co-infection. It is well known that anti-tuberculosis therapy,

especially SLDs, may cause hepatotoxicity and could lead to adverse TB treatment

outcomes.[33,34] Hepatotoxicity could also lead to permanent damage to the liver and may

increase the risk of deaths.[35] Alternatively, post-TB deaths we were observing among

individuals with HCV co-infection in this cohort may be due to HCV-related liver disease.

However, since information on cause of death as well as time of HCV infection are not known in

this cohort, more epidemiologic evidence is needed to link HCV to mortality post-TB treatment.

There are several plausible biological pathways to explain how pre-existing

comorbidities could increase the risk of post-TB mortality. First, individuals with pre-existing

comorbidities may experience prolonged chronic and systematic inflammation during TB

treatment, which could lead to severe disease manifestations/complications (e.g., opportunistic

infections among persons living with HIV co-infection, renal or heart disease among individuals

with diabetes, and cirrhosis among individuals with HCV co-infection) [36,37] which, in turn, will

increase the risk of mortality post-TB treatment. Second, individuals successfully treated for TB

with pre-existing comorbidities like T2DM may have carry over or residual complications such as

proatherogenic and abnormal lipid plasma profile, an established risk factor for cardiovascular

diseases.[38] Third, individuals with pre-existing comorbidities may also have increased risks of

TB relapse (i.e., re-infection or recurrent after treatment) due to their impaired immune

system.[39-41] Further clinical studies are needed to establish which of these pathways is more

common among TB survivors with pre-existing comorbidities.

Our study subjects to several limitations. First, we performed our analyses among

individuals who initiated treatment from 2009 – 2017. Thus, cohort effects associated with

changes in national TB treatment guidelines and/or care for individuals with TB and pre-existing

comorbidities are likely affected our reported estimates. Second, we did not have data on the

cause of death. Thus, we were not able to assess whether mortalities post-TB treatment

observed in this study are directly associated with pre-existing comorbidities. Third, we did not

have clinical information on key measures for pre-existing comorbidities such as CD4 counts,

viral load, blood glucose level during TB treatment, or whether or not individuals were on

medications to manage their comorbidities. These important variables may explain how

comorbidity factors drove post-TB deaths in this cohort. Thus, future clinical studies including

clinical information on the severity of comorbidity factors are still warranted. Fourth, we identified

several plausible sources of systematic errors, including bias due to misclassification of

hyperglycemia status. However, after running the sensitivity analyses, we did not see any

significant difference in our estimates that would change the study's overall findings. Lastly, our

cohort did not include non-TB individuals as a control group. Thus, we were not able to draw

direct comparison on mortality rates between individuals with TB vs. non-TB individuals in

Georgia. Additionally, our results may not be generalizable to other settings with different TB,

HIV, diabetes, and hepatitis C burdens.

In conclusion, we reported an overall high rate of post-TB deaths among individuals who

initiated treatment for RR and M/XDR TB, especially among those with pre-existing

comorbidities. More importantly, most post-TB deaths were observed within three years after TB

treatment was stopped/completed. Aggressive approaches to diagnose comorbidities and

integrating TB and non-communicable diseases care to reduce the risk of mortality during TB

treatment are needed to help prevent premature deaths. Additionally, continuous care (e.g.,

linkage to care) for pre-existing comorbidities (especially HIV co-infection) after TB treatment

may reduce the increased risks of post-TB deaths.

REFERENCES

World Health Organization. Global Tuberculosis Report 2022. (World Health Organization, Geneva, 2022).

World Health Organization. The END TB strategy. (World Health Organization, Geneva, 2015).

Romanowski, K. *et al.* Long-term all-cause mortality in people treated for tuberculosis: a systematic review and meta-analysis. *Lancet Infect Dis*, doi:10.1016/S1473-3099(19)30309-3 (2019).

Shuldiner, J., Leventhal, A., Chemtob, D. & Mor, Z. Mortality after anti-tuberculosis treatment completion: results of long-term follow-up. *Int J Tuberc Lung Dis* **20**, 43-48, doi:10.5588/ijtld.14.0427 (2016).

Harries, A. D. *et al.* Successfully treated but not fit for purpose: paying attention to chronic lung impairment after TB treatment. *Int J Tuberc Lung Dis* **20**, 1010-1014, doi:10.5588/ijtld.16.0277 (2016).

Pearson, F. *et al.* Tuberculosis and diabetes: bidirectional association in a UK primary care data set. *J Epidemiol Community Health* **73**, 142-147, doi:10.1136/jech-2018-211231 (2019).

Salindri, A. D., Wang, J. Y., Lin, H. H. & Magee, M. J. Post-tuberculosis incidence of diabetes, myocardial infarction, and stroke: Retrospective cohort analysis of patients formerly treated for tuberculosis in Taiwan, 2002 - 2013. *Int J Infect Dis*, doi:10.1016/j.ijid.2019.05.015 (2019).

Magee, M. J. *et al.* Convergence of non-communicable diseases and tuberculosis: a two-way street? *Int J Tuberc Lung Dis* **22**, 1258-1268, doi:10.5588/ijtld.18.0045 (2018).

Blondal, K., Rahu, K., Altraja, A., Viiklepp, P. & Rahu, M. Overall and cause-specific mortality among patients with tuberculosis and multidrug-resistant tuberculosis. *Int J Tuberc Lung Dis* **17**, 961-968, doi:10.5588/ijtld.12.0946 (2013).

Christensen, A. S., Roed, C., Andersen, P. H., Andersen, A. B. & Obel, N. Long-term mortality in patients with pulmonary and extrapulmonary tuberculosis: a Danish nationwide cohort study. *Clin Epidemiol* **6**, 405-421, doi:10.2147/clep.S65331 (2014).

Dangisso, M. H., Woldesemayat, E. M., Datiko, D. G. & Lindtjorn, B. Correction: Long-term outcome of smear-positive tuberculosis patients after initiation and completion of treatment: A ten-year retrospective cohort study. *PLoS One* **13**, e0196432, doi:10.1371/journal.pone.0196432 (2018).

Wang, X. H. *et al.* Survival and associated mortality risk factors among post-treatment pulmonary tubercolosis patients in the northwest of China. *Eur Rev Med Pharmacol Sci* **19**, 2016-2025 (2015).

Kolappan, C., Subramani, R., Kumaraswami, V., Santha, T. & Narayanan, P. R. Excess mortality and risk factors for mortality among a cohort of TB patients from rural south India. *Int J Tuberc Lung Dis* **12**, 81-86 (2008).

Miller, T. L. *et al.* Mortality hazard and survival after tuberculosis treatment. *Am J Public Health* **105**, 930-937, doi:10.2105/AJPH.2014.302431 (2015).

Cox, H. *et al.* Tuberculosis recurrence and mortality after successful treatment: impact of drug resistance. *PLoS Med* **3**, e384, doi:10.1371/journal.pmed.0030384 (2006).

World Health Organization. Definitions and Reporting Framework for Tuberculosis-2013 Revision. (World Health Organization, Geneva, 2013).

Wickham, H. ggplot2: Elegant Graphics for Data Analysis (Springer-Verlag, New York, 2016).

R Core Team (2023). R: A language and environment for statistical computing v. 4.2.3 (R Foundation for Statistical Computing, Vienna, Austria, 2023).

Austin, P. C., Lee, D. S. & Fine, J. P. Introduction to the Analysis of Survival Data in the Presence of Competing Risks. *Circulation* **133**, 601-609, doi:10.1161/CIRCULATIONAHA.115.017719 (2016).

Kleinbaum, D. G. & Klein, M. *Survival Analysis: A Self-Learning*. Third Edition edn, (Springer, 2011).

Greenland, S., Pearl, J. & Robins, J. M. Causal diagrams for epidemiologic research. *Epidemiology* **10**, 37-48 (1999).

VanderWeele, T. J. & Ding, P. Sensitivity Analysis in Observational Research: Introducing the E-Value. *Ann Intern Med* **167**, 268-274, doi:10.7326/M16-2607 (2017).

Georgia, N. S. O. o. *Deaths*, <https://www.geostat.ge/en/modules/categories/320/deaths> (2020).

Basham, C. A., Karim, M. E., Cook, V. J., Patrick, D. M. & Johnston, J. C. Post-tuberculosis mortality risk among immigrants to British Columbia, Canada, 1985-2015: a time-dependent Cox regression analysis of linked immigration, public health, and vital statistics data. *Can J Public Health*, doi:10.17269/s41997-020-00345-y (2020).

Fox, G. J. *et al.* Post-treatment Mortality Among Patients With Tuberculosis: A Prospective Cohort Study of 10 964 Patients in Vietnam. *Clin Infect Dis* **68**, 1359-1366, doi:10.1093/cid/ciy665 (2019).

Tocque, K., Convrey, R. P., Bellis, M. A., Beeching, N. J. & Davies, P. D. Elevated mortality following diagnosis with a treatable disease: tuberculosis. *Int J Tuberc Lung Dis* **9**, 797-802 (2005).

Koenig, S. P. *et al.* Increased Mortality After Tuberculosis Treatment Completion in Persons Living With Human Immunodeficiency Virus in Latin America. *Clin Infect Dis* **71**, 215-217, doi:10.1093/cid/ciz1032 (2020).

Baker, M. A. *et al.* The impact of diabetes on tuberculosis treatment outcomes: a systematic review. *BMC Med* **9**, 81, doi:10.1186/1741-7015-9-81 (2011).

Faurholt-Jepsen, D. *et al.* Diabetes is a strong predictor of mortality during tuberculosis treatment: a prospective cohort study among tuberculosis patients from Mwanza, Tanzania. *Trop. Med. Int. Health* **18**, 822-829, doi:10.1111/tmi.12120 (2013).

Boillat-Blanco, N. *et al.* Transient Hyperglycemia in Patients With Tuberculosis in Tanzania: Implications for Diabetes Screening Algorithms. *J Infect Dis* **213**, 1163-1172, doi:10.1093/infdis/jiv568 (2016).

Jimenez-Corona, M. E. *et al.* Association of diabetes and tuberculosis: impact on treatment and post-treatment outcomes. *Thorax* **68**, 214-220, doi:10.1136/thoraxjnl-2012-201756 (2013).

Degner, N. R., Wang, J. Y., Golub, J. E. & Karakousis, P. C. Metformin Use Reverses the Increased Mortality Associated With Diabetes Mellitus During Tuberculosis Treatment. *Clin Infect Dis.* **66**, 198-205, doi:10.1093/cid/cix819 (2018).

Ungo, J. R. *et al.* Antituberculosis drug-induced hepatotoxicity. The role of hepatitis C virus and the human immunodeficiency virus. *Am J Respir Crit Care Med* **157**, 1871-1876, doi:10.1164/ajrccm.157.6.9711039 (1998).

Kempker, R. R. *et al.* Acquired Drug Resistance in Mycobacterium tuberculosis and Poor Outcomes among Patients with Multidrug-Resistant Tuberculosis. *Emerg Infect Dis* **21**, 992-1001, doi:10.3201/eid2106.141873 (2015).

Zhao, H., Wang, Y., Zhang, T., Wang, Q. & Xie, W. Drug-Induced Liver Injury from Anti-Tuberculosis Treatment: A Retrospective Cohort Study. *Med Sci Monit* **26**, e920350, doi:10.12659/MSM.920350 (2020).

Kornfeld, H. *et al.* High Prevalence and Heterogeneity of Diabetes in Patients With TB in South India: A Report from the Effects of Diabetes on Tuberculosis Severity (EDOTS) Study. *Chest* **149**, 1501-1508, doi:10.1016/j.chest.2016.02.675 (2016).

Samuels, J. P., Sood, A., Campbell, J. R., Ahmad Khan, F. & Johnston, J. C. Comorbidities and treatment outcomes in multidrug resistant tuberculosis: a systematic review and meta-analysis. *Sci Rep* **8**, 4980, doi:10.1038/s41598-018-23344-z (2018).

Vrieling, F. *et al.* Patients with Concurrent Tuberculosis and Diabetes Have a Pro-Atherogenic Plasma Lipid Profile. *EBioMedicine* **32**, 192-200, doi:10.1016/j.ebiom.2018.05.011 (2018).

Dooley, K. E. & Chaisson, R. E. Tuberculosis and diabetes mellitus: convergence of two epidemics. *The Lancet Infectious Diseases* **9**, 737-746, doi:10.1016/S1473-3099(09)70282-8 (2009).

Getahun, H., Gunneberg, C., Granich, R. & Nunn, P. HIV infection-associated tuberculosis: the epidemiology and the response. *Clin Infect Dis* **50 Suppl 3**, S201-207, doi:10.1086/651492 (2010).

Wu, P.-H., Lin, Y.-T., Hsieh, K.-P., Chuang, H.-Y. & Sheu, C.-C. Hepatitis C Virus Infection Is Associated With an Increased Risk of Active Tuberculosis Disease: A Nationwide Population-Based Study. *Medicine* **94**, e1328-e1328, doi:10.1097/MD.0000000000001328 (2015).

ACKNOWLEDGEMENTS

We would like to thank Mamuka Chincharauli for his assistance in pooling data from the Georgia's TB online surveillance system.

AUTHOR CONTRIBUTIONS

ADS, MJM, and RRK conceived the study design. ADS and MK obtained and had access to the full data. ADS performed the analyses. ADS, MJM, HMB, RRK, MK provided interpretation of data for the work. ADS wrote the first draft of the manuscript. MJM, HMB, RRK revised the manuscript critically for important intellectual content. All authors reviewed and approved the final version of the manuscript.

DATA SHARING STATEMENT

Data that support the findings were collected for surveillance purposes and only accessible to investigators upon request to Georgia's National TB Program and Statistics Office. Individuals included in this cohort did not give written consent for their data to be shared publicly, and since the full data contain sensitive information including individuals' private health information (e.g., name and date of birth for matching purposes), the full dataset are not made publicly available. However, deidentified individual data, will be made available upon request to corresponding authors (ADS).

CONFLICT OF INTEREST

We have no conflict of interest to declare.

STUDY FUNDING

This work was supported in part by grants from the National Institutes of Health (NIH) including the National Institute of Allergy and Infectious Diseases (NIAID) [R03AI133172 to MJM, R03AI139871 to RRK, R01AI153152 to MJM] and the Fogarty International Center (FIC) [D43TW007124 to HMB for the "Emory-Georgia TB Research Training Program", R21TW011157 to MK and MJM]. ADS was supported by a Vanderbilt Emory Cornell Duke (VECD) Global Health Fellowship, funded by the NIH FIC (D43TW009337). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

FIGURES LEGENDS

Figure 1   **Participants selection according to inclusion criteria**

This flowchart depicts the participants' selection according to inclusion criteria (i.e., non-pediatric TB individuals with no prior history of TB treatment). Individuals were excluded if they had no pulmonary involvement, culture confirmation at the beginning of TB treatment, or unknown vital status during and after TB treatment.

Figure 2   **Trend of mortality rates reported during and after TB treatment stratified by comorbidity status among persons who initiated treatment for RR or M/XDR TB in the country of Georgia from 2009 – 2017**

Mortality rates during and post-TB treatment were estimated by 6-month period by building independent life tables. Mortality rates during and post-TB treatment were expressed per 1,000 populations. Mortality rates were then plotted continuously with a single interruption point to mark the end of TB treatment (red dashed line). The beginning of survival time of 'TB treatment phase' ($T_{0\text{-treatment}}$) marks the TB treatment initiation date. Thus, the point at T_6 in the 'TB treatment' phase indicates the mortality rate at 6-month after TB treatment initiation. The beginning of survival time of 'Post-TB treatment' phase ($T_{0\text{-post-TB treatment}}$) marks the end of TB treatment (i.e., could be treatment completion or lost to follow-up dates). Thus, the point at T_6 in the 'Post-TB treatment' phase indicates the mortality rate at 6-month after TB treatment ended.

TABLES

Table 1 Demographic and clinical characteristics of persons who initiated treatment for RR and M/XDR TB according to pre-existing comorbidity status in Georgia, 2009 – 2017 (N=1,032)

Characteristics	Total N=1032 N (%)	HIV positive[*] N (%) =39 (3.8) N (%)	Hyperglycemia[^] N (%) =234 (22.7) N (%)	HCV positive[#] N (%) =153 (14.7) N (%)
Age, median (IQR)	35 (26 – 49)	40 (35 - 47)	42.5 (32 – 54)	42 (34 – 50)
Male Gender	753 (73.0)	33 (84.6)	189 (80.8)	148 (96.7)
Ethnicity, Georgian	942 (91.3)	33 (84.6)	212 (90.6)	142 (92.8)
BMI kg/m^2, median (IQR)	20.8 (19.1 – 22.9)	21.2 (18.4 – 22.0)	21.6 (19.9 – 24.8)	21.1 (19.4 – 23.2)
Smear Positive (n=983)	655 (66.6)	13 (36.1)	173 (75.9)	98 (68.1)
Culture Positive (n=883)	876 (99.2)	37 (100.0)	199 (99.0)	128 (99.2)
RR TB	195 (18.9)	4 (10.3)	42 (18.0)	28 (18.3)
MDR/pre-XDR TB	761 (73.7)	32 (82.1)	170 (72.7)	118 (77.1)
XDR TB	76 (7.4)	3 (7.7)	22 (9.4)	7 (4.6)
Abnormal Chest X-ray Findings (n=1014)	975 (96.2)	34 (89.5)	215 (93.1)	149 (98.7)
Cavitary Disease	285 (27.6)	5 (12.8)	64 (27.4)	44 (28.8)
Infiltrate	755 (73.2)	22 (56.4)	169 (72.2)	113 (73.9)
Disseminated TB disease	251 (24.3)	10 (25.6)	60 (25.6)	45 (29.4)
History of internally displaced (n=874)	36 (4.1)	2 (6.1)	5 (2.4)	6 (5.1)
Tobacco Use (n=591)	455 (44.1)	18 (90.0)	104 (75.9)	97 (91.5)
Excessive Alcohol Intake (n=998)	59 (5.9)	1 (2.7)	16 (7.1)	16 (11.0)
Treatment Outcome (n=980)				
Completed/cured	655 (65.4)	10 (26.3)	157 (68.0)	73 (51.4)
Poor treatment outcome	325 (33.2)	26 (8.0)	74 (32.1)	69 (48.6)
Failed the treatment	44/325 (13.5)	5/26 (19.2)	13/74 (17.8)	10/69 (14.5)
Lost to follow-up	247/325 (76.0)	14/26 (53.9)	58/74 (78.4)	53/69 (76.8)
Died during TB treatment	34/325 (10.5)	7/26 (26.9)	3/74 (4.1)	6/69 (8.7)
Return to care	95 (9.2)	6 (15.4)	32 (13.7)	18 (11.8)
Survival Status				
Alive	911 (88.3)	23 (59.0)	204 (87.2)	129 (84.3)
Mortality during treatment	34 (3.3)	7 (18.0)	3 (1.3)	6 (3.9)
Mortality post-TB treatment	87 (8.4)	9 (23.0)	27 (11.5)	18 (11.8)

*There were 96 individuals with missing/unknown HIV co-infection status

Characteristics	Total N=1032 N (%)	HIV positive[*] N (%) =39 (3.8) N (%)	Hyperglycemia[^] N (%) =234 (22.7) N (%)	HCV positive[#] N (%) =153 (14.7) N (%)

[^]There were 220 individuals with missing/unknown prediabetes/diabetes status
[#]There were 282 individuals with missing/unknown HCV co-infection status

Abbreviations:
BMI – body mass index; CVD – cardiovascular diseases; DRTB – drug-resistant TB; LFU – lost to follow-up; HCV – hepatitis C virus; HIV – human immunodeficiency virus; IQR – interquartile range; MDR – multidrug resistant; RR – rifampicin resistant; TB – tuberculosis; XDR – extensively drug-resistant

Table 2 All-cause post-tuberculosis mortality rates among persons who initiated treatment for RR and M/XDR TB with and without pre-existing comorbidities in Georgia, 2009 – 2017 (N=998)*

Characteristics	Death/Total (%)	Person time (Years)	Crude Rates (95%CI) per 1,000 person-years	Rate Difference (95%CI) per 1,000 person years	Median time to death post-TB treatment, months (IQR)
All cohort	**87/998 (8.7)**	**4857**	**17.91 (14.43 – 21.99)**		**21 (7 – 39)**
HIV co-infection					
No	68/878 (7.7)	4195	16.12 (12.78 – 20.56)	Reference	22 (7 – 39)
Yes	9/32 (28.1)	130	69.23 (36.02 – 133.10)	**53.02 (7.63, 98.41)**	7 (4 – 9)
Missing	10/88 (11.4)	532	18.80 (10.11 – 34.94)	2.59 (-9.68, 14.86)	23 (13 – 46)
Hyperglycemia†					
No	41/558 (7.4)	2712	15.12 (11.13 – 20.53)	Reference	28 (9 – 43)
Yes	27/231 (11.7)	1129	23.91 (16.40 – 34.87)	8.80 (-1.34, 18.93)	19 (7 – 34)
Missing	19/209 (9.1)	1016	18.70 (11.93 – 29.32)	3.58 (-6.02, 13.18)	21 (6 – 40)
Diabetes Status					
No diabetes	41/558 (7.4)	2712	15.12 (11.13 – 20.53)	Reference	28 (9 – 43)
Pre-diabetes	13/119 (10.9)	593	21.92 (12.73 – 37.75)	6.80 (-5.98, 19.59)	17 (7 – 31)
Diabetes	14/112 (12.5)	536	26.12 (15.47 – 44.10)	11.00 (-3.44, 25.44)	22 (10 – 38)
Missing	19/209 (9.1)	1016	18.70 (11.93 – 29.32)	3.58 (-6.02, 13.18)	21 (6 – 40)
HCV co-infection					
No	46/580 (7.9)	2872	16.02 (12.00 – 21.38)	Reference	20 (9 – 39)
Yes	18/147 (12.2)	701	25.68 (16.18 – 40.76)	9.66 (-3.07, 2.24)	20 (8 – 38)
Missing	23/271 (8.5)	1284	17.91 (11.90 – 26.96)	1.90 (-6.77, 10.56)	22 (6 – 45)

*34 individuals who died during treatment were excluded from the post-TB mortality analyses
†Including individuals with pre-diabetes and diabetes

Abbreviations:
BMI – body mass index; CI – confidence interval; HCV – hepatitis C virus; HIV – human immunodeficiency virus; IQR – interquartile range; TB - tuberculosis

Table 3 Crude and adjusted hazard rate ratios of mortality during and after tuberculosis treatment among persons who initiated treatment for RR and M/XDR TB with and without pre-existing comorbidities in Georgia, 2009 – 2017 (N=1,032)

Comorbidities	Study Outcomes			
	Tuberculosis mortality during treatment (n=34)		Tuberculosis mortality after treatment (n=87)	
	cHR (95%CI)	aHR[*] (95%CI)	cHR (95%CI)	aHR[*] (95%CI)
HIV co-infection				
No	Reference	Reference	Reference	Reference
Yes	**9.33 (3.92 – 22.23)**	**11.52 (4.57 – 29.06)**	**4.56 (2.27 – 9.15)**	**3.74 (1.77 – 7.91)**
Missing	**4.10 (1.80 – 9.37)**	**3.70 (1.48 – 9.27)**	1.11 (0.57 – 2.17)	1.05 (0.53 – 2.12)
Hyperglycemia				
No	Reference		Reference	
Yes	0.37 (0.11 – 1.24)		1.60 (0.98 – 2.60)	
Missing	1.47 (0.70 – 3.06)		1.24 (0.72 – 2.14)	
Diabetes Status				
No diabetes	Reference	Reference	Reference	Reference
Pre-diabetes	NA	NA	1.47 (0.79 – 2.75)	1.48 (0.78 – 2.78)
Diabetes	0.76 (0.23 – 2.55)	0.75 (0.19 – 3.00)	1.73 (0.94 – 3.18)	1.06 (0.53 – 2.12)
Missing	1.47 (0.70 – 3.06)	1.72 (0.79 – 3.75)	1.24 (0.72 – 2.14)	1.45 (0.82 – 2.59)
HCV co-infection				
No	Reference	Reference	Reference	Reference
Yes	1.40 (0.55 – 3.56)	0.93 (0.33 – 2.66)	1.58 (0.92 – 2.73)	1.21 (0.68 – 2.15)
Missing	1.39 (0.65 – 2.96)	1.26 (0.55 – 2.87)	1.12 (0.68 – 1.85)	0.90 (0.53 – 1.55)

[*]Model adjusted for HIV-coinfection, diabetes status, HCV co-infection, age, gender, cavitary disease, type of drug resistance, smoking, and BMI at TB treatment initiation (fully-adjusted model)

Abbreviations:
aHR – adjusted hazard rate ratios; BMI – body mass index; cHR – crude hazard rate ratios; CI – confidence interval; HCV – hepatitis C virus; HIV – human immunodeficiency virus

Bold indicates that the finding is statistically significant at α=0.05

Areas shaded in grey indicate that variables were not included in the final models, and thus, adjusted hazard rate ratios were not obtained.

Table 4 Subgroup analyses to assess statistical interaction between pre-existing comorbidities and smoking on mortality post-tuberculosis among persons who initiated treatment for RR and M/XDR TB in Georgia, 2009 – 2017 (N=591)

Smoking Status	Comorbidities Status	Post-TB mortality (%)	cHR (95%CI)	Adjusted Model	
				aHR* (95%CI)	p-values for the interaction term
HIV Co-infection					
Non-smokers	HIV Negative	9/117 (7.7)	Reference	Reference	
	HIV Positive	1/2 (50.0)	**13.68 (1.63 – 114.64)**	**12.72 (1.41 – 114.87)**	
Smokers	HIV Negative	32/399 (8.0)	Reference	Reference	0.8903
	HIV Positive	4/18 (22.2)	**4.72 (1.65 – 13.49)**	**4.87 (1.70 – 13.94)**	
Hyperglycemia (i.e., pre-diabetes and diabetes combined)					
Non-smokers	No hyperglycemia	7/74 (9.5)	Reference	Reference	
	Hyperglycemia	2/33 (6.1)	0.66 (0.14 – 3.17)	0.35 (0.07 – 1.76)	
Smokers	No hyperglycemia	19/256 (7.4)	Reference	Reference	0.5909
	Hyperglycemia	8/104 (7.7)	1.00 (0.44 – 2.29)	0.81 (0.35 – 1.88)	
Diabetes					
Non-smokers	No diabetes	7/74 (9.5)	Reference	Reference	
	Pre-diabetes	2/17 (11.8)	1.29 (0.27 – 6.21)	0.85 (0.15 – 4.86)	
	Diabetes	0/16 (0.0)	NA	NA	
Smokers	No diabetes	19/256 (7.4)	Reference	Reference	0.3569
	Pre-diabetes	4/51 (7.8)	1.02 (0.35 – 3.01)	1.08 (0.36 – 3.20)	
	Diabetes	4/53 (7.6)	0.98 (0.33 – 2.88)	0.64 (0.21 – 1.94)	
Hepatitis C					
Non-smokers	HCV Negative	4/84 (4.8)	Reference	Reference	
	HCV Positive	1/9 (11.1)	2.55 (0.28 – 22.86)	3.43 (0.35 – 33.69)	
Smokers	HCV Negative	15/232 (6.5)	Reference	Reference	0.8911
	HCV Positive	11/97 (11.3)	1.96 (0.90 – 4.26)	1.72 (0.79 – 3.75)	

Abbreviations: aHR – adjusted hazard rate ratios; cHR – crude hazard rate ratios; CI – confidence interval; HCV – hepatitis C virus; HIV – human immunodeficiency virus; TB – tuberculosis

[*]Model adjusted for age and gender
[†]Wald chi-square p-value for the interaction term between pre-diabetes and smoking
[‡]Wald chi-square p-value for the interaction term between diabetes and smoking
Bold indicates that the finding is statistically significant at α=0.05

Abbreviations:
EPTB – extra-pulmonary tuberculosis, IQR – inter-quartile range, TB - tuberculosis

Figure 1. Participants selection according to inclusion criteria. This flowchart depicts participants' selection according to inclusion criteria (i.e., non-pediatric TB individuals with no prior history of TB treatment). Individuals were excluded if they had no pulmonary involvement, culture confirmation at the beginning of TB treatment, or unknown vital status during and after TB treatment

Figure 2. Trend of mortality rates reported during and after TB treatment stratified by comorbidity status among persons who initiated treatment for RR or M/XDR TB in the country of Georgia from 2009 – 2017. Mortality rates during and post-TB treatment were estimated by 6-month period by building independent life tables. Mortality rates during and post-TB treatment were expressed per 1,000 populations. Mortality rates were then plotted continuously with a single interruption point to mark the end of TB treatment (red dashed line). The beginning of survival time of 'TB treatment phase' ($T_{0\text{-treatment}}$) marks the TB treatment initiation date. Thus, the point at T_6 in the 'TB treatment' phase indicates the mortality rate at 6-month after TB treatment initiation. The beginning of survival time of 'Post-TB treatment' phase ($T_{0\text{-post-TB treatment}}$) marks the end of TB treatment (i.e., could be treatment completion or lost to follow-up dates). Thus, the point at T_6 in the 'Post-TB treatment' phase indicates the mortality rate at 6-month after TB treatment ended.