Title: Bidirectional two-sample mendelian randomization study of differential white blood cell count and schizophrenia

Perry BM Leung[1,2], Zipeng Liu, Yuanxin Zhong[1], Marta Di Forti[3], Robin M Murray[2], Hon Cheong So[4,5]*, Pak Chung Sham[1,6,7]*, Simon SY Lui[1]*

[1]Department of Psychiatry, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Pok Fu Lam, Hong Kong SAR

[2]Department of Psychosis Studies, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK

[3]Social, Genetics and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK

[4]School of Biomedical Sciences, The Chinese University of Hong Kong, Shatin, Hong Kong SAR

[5]Department of Psychiatry, The Chinese University of Hong Kong, Shatin, Hong Kong SAR

[6]Centre for PanorOmic Sciences, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong SAR

[7]State Key Laboratory of Brain and Cognitive Sciences, The University of Hong Kong, Hong Kong SAR

*co-corresponding authors

Word Count (Abstract): 267
Word Count (Text): 3786
Number of Tables: 4
Number of Reference: 74

Abstract

Background: Schizophrenia and white blood cell count (WBC) are both complex and polygenic disease/traits. Previous evidence suggested WBC can lead to higher all-cause mortality, and some other evidence found elevated WBC in first-episode psychosis and chronic schizophrenia patients. However, prior observational findings may be confounded by antipsychotic exposures and their effects on WBC. Mendelian randomization (MR) is a useful method to examine the directional causal relationship between schizophrenia and WBC.

Methods: We performed a two-sample MR using summary statistics of the Psychiatric Genomics Consortium Schizophrenia Workgroup (N=130,644) and the Blood Cell Consortium (N=563,085). The MR methods included inverse variance weighted, ME Egger, weighted median, and MR-PRESSO, contamination mixture, and a novel approach called mixture model reciprocal causal inference (MRCI). False discovery rate was employed to correct for multiple testing.

Results: After correcting for horizontal pleiotropy, the MRCI method demonstrated that schizophrenia can cause by elevated lymphocyte count (OR=1.080; FDR adjusted p-value=0.026) and eosinophil count (OR=1.050; FDR adjusted p-value=0.026). The contamination mixture method showed that schizophrenia may cause elevated neutrophil counts (Beta=0.011 in unit of standard deviation of mean absolute neutrophil count; FDR adjusted p-value=0.045) and reduction of eosinophil count (Beta=-0.013 in unit of standard deviation of mean absolute eosinophil count; FDR adjusted p-value=0.045). Some further significant findings had been identified by conventional MR approaches and MR-PRESSO, but we interpreted those with cautious due to substantial heterogeneity and plausible pleiotropic effects identified.

Conclusion: This MR study provided evidence that schizophrenia has causal relationships with altered differential WBC. Our findings support the role of WBC in influencing schizophrenia risk, and may concur with the hypothesis of neuroinflammation in schizophrenia.

1. Introduction

Schizophrenia had an estimated the age standardised prevalence to be 287.4 per 100,000 people and affecting 23.6 million people globally. It leads to high disability in patients and substantial caregiver burden (GBD 2019 Mental Disorders Collaborators, 2022). The aetiology is largely unknown, but many hypotheses had been made in literature which had highlighted the important roles of neurotransmitters as well as the immune system (Vidal & Pacheco, 2020). Ermakov et al. had summarized multiple hypotheses suggesting the role of immune system in schizophrenia, including the general inflammation hypothesis, macrophage-T-cell hypothesis, autoantibody hypothesis, and microglia hypothesis (Ermakov et al., 2022).

Notably, above one-third of people with schizophrenia show abnormalities of immune biomarkers, such as abnormal levels of C-reactive protein, and dysregulations of cytokines, chemokines and microbiota (Gibney & Drexhage, 2013; Ermakov et al., 2022; Hartwig et al., 2017; Van Kestere et al., 2017). Elevated circulating level of proinflammatory cytokines were observed prior to onset of schizophrenia (Williams et al., 2022). Epidemiological evidence also revealed strong associations between prenatal and childhood infections and schizophrenia (Davies et al., 2020). In addition, the Psychiatric Genomics Consortium reported genome-wide genetic correlation between schizophrenia and various immune-related disorders, such as Crohn's disease, primary biliary cirrhosis, systemic lupus erythematosus, and ulcerative colitis (Tylee et al., 2018).

Lymphoid-derived lymphocytes, as well as myeloid-derived basophils, eosinophils, monocytes, and neutrophils, are are indispensable to innate and adaptive immunity. These five types of white blood cells have lots of interactions and crosstalk to function holistically in the immune system. Neutrophils are the most abundant in human bodies, which serve as front line of innate immunity in the immune system. Functions of neutrophils include damage pathogens, produce cytokines, and facilitate activation of macrophages. In addition, it regulates signals for proliferation and activations of B and T cells to activate adaptive immunity. It is essential to balance between inflammatory and anti-inflammatory responses (Li et al., 2019). Lymphocytes, with subclassification into B cells, T cells, natural killer cells, and innate lymphoid cells, are centre in initiate immunity and regulate adaptive immune responses (Gasteiger & Rudensky, 2014). B cells can recognize antigens and release antigen-specific antibody to eliminate foreign substances. Monocytes derive macrophages and dendritic cells at inflammatory sites. Macrophages can be both pro-and anti-inflammatory depending on the microenvironments, which is known as macrophage plasticity. Proinflmmatory monocytes release multiple chemokines with aim to remove the invading pathgens. On the other hand, anti-inflammatory monocytes negatively regulate proinflammatory cytokines and facilitate resolution of inflammation (Austermann et al., 2022). Eosinophils are also part of the innate immune system which mediate and bridge the innate and adaptive immunity. Eosinophils can affect the immune microenvironment, cellular recruitment as well as tissue homeostasis, repair and remodeling (Shamri et al., 2011). Lastly, basophils have the least amount in blood. It has been suggested as mediator in allergic reactions, and also with critical roles in immunomodulation and associations with infectious, allergic, and autoimmune diseases (Zhang et al., 2021).

Given the interconnection between levels of peripheral white blood cell counts (WBC) and immune response, they are the commonly-used laboratory-indicators for inflammatory, immune, allergic and hematologic diseases (Jain et al., 2017), and elevated WBC counts are associated with higher all-cause mortality, in particular related to respiratory, cardiac and metabolic disorders (Chan et al., 2022; Kim et al, 2013; Shim et al., 2006). Although WBC fluctuate with time to adapt to environmental pathogen exposure (Hung et al., 2021), these laboratory-markers also show 'trait-like' properties, because physically-active people tend to show persistently elevated WBC (Farhangi et al, 2013). The levels of WBC are heritable complex traits, following a polygenic model (Whitfield et al., 1985; Evans et al., 1999). The heritability of lymphocyte, monocyte, and neutrophil count were estimated at 58%, 58%, and 41% in European samples (Lin et al., 2016; Lin et al., 2017). The Blood Cell Consortium combined GWAS data of 27 cohorts and 29 blood cell phenotypes, to reveal variants responsible for regulating hematopoiesis (Vuckovic et al., 2020).

Empirical evidence supports an association between schizophrenia and elevated WBC. For instance, several meta-analyses found people with first-episode psychosis and chronic schizophrenia having elevated WBC (Miller et al., 2013; Jackson & Miller, 2020; Mazza et al., 2020). Elevated neutrophil/lymphocyte ratio (NLR), which were suggested as cost effective measurement of systemic inflammation (Imtiaz et al., 2012), was found to be consistently elevated in people with first-episode psychosis, chronic schizophrenia (Karageorgiou et al., 2019). A longitudinal study which followed a cohort of first-episode psychosis reported that NLR was higher in non-remitted schizophrenia than in remitted schizophrenia (Bioque et al., 2022). Clozapine is the only antipsychotic medication indicated for treatment-resistant schizophrenia, a severe schizophrenia subtype having failed to respond to at least two different antipsychotic medications with adequate dose and duration, and this drug has well-known effects in altering WBC in people with schizophrenia, including rare and potentially fatal clozapine-induced neutropenia, as well as more common and benign leucocytosis (Paribello et al., 2021). Non-clozapine antipsychotic medications, such as olanzapine and risperidone, have also been reported to alter WBC (Alvarez-Herrera et al., 2020; Rettenbacher et al., 2010). Findings regarding whether antipsychotic medications can alter NLR remain inconsistent (Zhou et al., 2020; Dawidowski et al., 2022; Bioque et al., 2022).

One recent sophisticated method to examine the causal relationship between traits is Mendelian Randomization (MR). MR is a statistical method to mimic randomization by using generic variants as 'instrumental variables' to estimate the causal effect of an exposure on an outcome of interest. This approach substantially reduces the risk of confounding bias typically encountered in case-control and cohort studies, based on the assumption that genetic variants are randomly allocated within a population. A recent study had attempted to explore association between schizophrenia and WBC (Gao et al., 2023). Nevertheless, there were a few shortcomings. First, their data were not most up-to-date, as Psychiatric Genomics Consortium had published their third wave data in 2022 (Trubetskoy et al., 2022), and blood cell consortium which had further combined UK Biobank with other cohorts in their meta-analysis (Vuckovic et al., 2020, Chen et al., 2021; Lettre, 2023). Moreover, several novel MR approaches had developed that try to address the horizontal pleiotropy and provide outliers-, and pleiotropic- robust estimates. We used contamination mixture developed in 2020 (Burgess et al., 2020), and MRCI which was developed most recently (Liu et al., 2023).

This MR study aimed to explore the causal relationship between schizophrenia and WBC. Furthermore, we employed two-sample (rather than one-sample) MR to minimize risk of biases caused by winners' curse and weak instrument bias (Lawlor, 2016). To our knowledge, this bidirectional two-sample MR study is the first to utilize the most updated GWAS data provided by two large scale consortiums, namely the Psychiatric Genomics Consortium wave 3 and the Blood Cell Consortium phase 2.

2. Methods

Our two-sample MR used two independent datasets as the exposure and outcome, both sensitive to causal effects of mediators and corrected for measurement errors (Carter et al., 2019). The three essential assumptions to uphold unbiased estimates of MR are (1) the genetic instruments are associated with exposure; (2) the genetic instruments are associated with the outcome only through the chosen exposure (i.e. no pleiotropic effects); and (3) the genetic instruments are independent of other factors which could influence the outcome (Boef et al., 2015).

2.1 Data source

2.1.1. Psychiatric Genomics Consortium (PGC) schizophrenia genome-wide association study (GWAS)

We used the summary statistics gathered from the most updated third wave PGC schizophrenia GWAS core dataset, which contained 53,386 schizophrenia cases and 77,258 controls from the European ancestry. Details of this dataset has been described elsewhere (Trubetskoy et al., 2022).

2.1.2. Blood Cell Consortium

We used the summary statistics of differential WBC gathered from the European working groups Blood Cell Consortium Phase 2 (BCX2), through the IEU Open GWAS Project, which contained 26 discovery cohorts of European ancestry additional to UK Biobank, which total 563,946 subjects. The log10-transformed absolute counts, in units of ($\times 10^9$ cells/L), were reported after multiplying relative counts with the total WBC count from the original cohorts. Detailed descriptions of each cohort and data management have been described elsewhere (Vuckovic et al., 2020, Chen et al., 2021; Lettre, 2023).

2.2. Statistical analysis

We first retrieved the single nucleotide polymorphisms (SNPs) from the abovementioned two GWAS meta-analysis study data, and selected those SNPs which showed a p-value smaller than the GWAS significant threshold (i.e. p-value < 5e-8). SNPs were then clumped based on the 'TwoSampleMR' package in statistical software R, with the default linkage disequilibrium threshold $r^2 < 0.001$, distance > 10000kb, and 1000 genomes European

reference panel. After harmonising the data between exposure and outcome datasets, 201 SNPs remained. Furthermore, we removed 7 SNPs which were both palindromic and ambiguous (i.e. the default option in TwoSampleMR package). The final data comprised 194 SNPs F-statistic was calculated to assess instrument strength, and F statistics >10 is a general rule of thumb that lacks evidence of weak instruments which could cause biases towards the null (Wootton et al., 2018).

We comprehensively employed different MR methods in this study, because each has its own assumptions, strengths and limitations (see Table 1 for a summary). In brief, random-effect inverse variance weighted (IVW) provides unbiased estimates when all three MR assumptions are clearly satisfied. Weighted median is robust to outliers when the majority of genetic variants are valid instruments (Slob & Burgess, 2020). Horizontal pleiotropy refers to existence of pleiotropic SNPs which have influences on the outcome other than through the exposure; this will violate the exclusion restriction assumption of MR and undermine the validity of the estimates. When horizontal pleiotropies are present, other MR methods should be considered (Slob & Burgess, 2020). For instance, MR-Egger relaxes the assumption of exclusion restriction by allowing the strength of instruments to be independent from the direct effect (InSIDE) assumption. InSIDE states that genetic variants-exposure pathways should be independent from the genetic variants-outcome pathways (Bowden et al., 2015). Furthermore, MR-PRESSO is an extended approach of IVW as it performs IVW analysis after removing the outliers. MR-PRESSO detects horizontal pleiotropy using the MR-PRESSO global test, and determines outliers using the MR-PRESSO distortion test, and then yield 'outlier-adjusted causal estimates' (Wootton et al., 2018). Nonetheless, MR-PRESSO is prone to Type 1 error, if a few invalid instruments are retained in the sample after removing the outliers (Slob & Burgess, 2020). Contamination mixture MR is a more conservative method, compared with MR-PRESSO and other outliers-removed approaches, because it assumes plurality valid and can give close to nominal rates of Type 1 error up to 40% invalid genetic variants, while maintaining reasonable power to detect causal effect (Burgess et al., 2020).

We also employed a novel MR method, namely the Mixture model Reciprocal Causation Inference (MRCI) (Liu et al., 2023). When dealing with both independent and correlated pleiotropy, MRCI utilizes all the SNPs gathered from GWAS summary statistics to provide unbiased estimates with non-inflated Type 1 error. Independent pleiotropy occurs when a SNP contribute to at least two traits with independent mechanisms. Correlated pleiotropy, on the other hand, refers to SNPs that contribute to multiple traits with shared mechanisms. The MRCI method is unique in utilizing all the available SNPs and modeling the pleiotropy explicitly by classifying SNPs into phenotype-specific, pleiotropic, or not contributing towards the specified phenotypes, based on marginal effect of each SNP. As such, MRCI provides robust estimates even when the majority of the causal variants are pleiotropic. Moreover, MRCI can operation under different (sometimes contradicting) assumptions (e.g., none of the variants are pleiotropic; all of the variants are pleiotropic; as well as situations where phenotype-specific variants could not be identified). The only required parameters to use MRCI were prevalence of schizophrenia which was set to 0.35% referencing from WHO data (World Health Organization, 2022), and the case proportion in our data which was

40.86%. MRCI outputs are the log odds effect sizes, which we then converted to OR by taking exponential of them.

Table 1 Summary of assumptions, strengths, and weaknesses of selected MR methods (Bowden et al., 2016; Burgess & Thompson, 2017; Verbanck et al., 2018; Slob & Burgess, 2020; Liu et al., 2023)

Analysis method	Consistency assumption	Strengths	Weaknesses
Random effect inverse variance weighted (IVW)	Average pleiotropic effect must be zero	Consistent and efficient estimate when MR assumptions are satisfied	All variants have to be valid IVs which is also known as 0% breakdown point
Weighted median	Majority (>50%) of genetic variants are valid IVs	Robust to outliers as the method focus on the median values only	Less efficient than IVW
MR-Egger	InSIDE (Instrument Strength Independent of Direct Effect)	The InSIDE assumption is less stringent compared with IVW	The InSIDE assumption is sometimes not plausible. This method is also sensitive to outliers
MR-PRESSO	Majority (>50%) of genetic variants are valid IVs	Outlier robust and efficient with valid IVs	High false positive rate with several invalid IVs
Contamination mixture	Plurality valid	Robust to outliers	Sensitive to variance parameters and addition/removal of genetic variants
Mixture model Reciprocal Causation Inference (MRCI)	Bivariate normality of standardized marginal effects in SNPs	Provide robust estimates even if most SNPs were pleiotropic	The robust sandwich standard errors are conservative

Two-sided p-value was used for each analysis. False discovery rate (FDR) adjusted p-value were used as the conservative estimates to address the issue of multiple hypothesis testing. False Discovery rate (FDR) correction p-value controlled the expected proportion of falsely rejected hypotheses in multiple testing (Benjamini & Hochberg, 1995). In this study, an FDR < 0.05 would be deemed as the statistically significant threshold to reject null hypothesis, and FDR < 0.1 as suggestive associations. Besides, Cochran's Q statistic and heterogeneity statistic I^2 were computed in different harmonized datasets to test for the validity of the MR assumptions. Scatter plot, forest plot, funnel plot, and leave-one-out analysis were generated to aid the detection of outliers (see Supplementary Figures 1-30). Furthermore, power calculation was performed using an online power calculator designed for MR studies, and suggested that 563,085 participants from the Blood Cell Consortium exceeded the minimum sample size to achieve 80% power (i.e. 436,100) to detect exposure R^2 and causal effect size as low as 0.02 and 0.03 standard deviation change respectively. On the other hand, 161,405 individuals from PGC with 80% statistical power can detect as little as 0.033 exposure R^2 and 1.08 odds ratio per standard deviation change in outcomes (Burgess, 2023).

All the MR analyses conducted in this study was performed using 'TwoSampleMR' packages in the R version 4.1.0. MRCI algorithm is publicly available on GitHub (Liu et al., 2023).

3. Results

We found 20,457 SNPs from PGC-SZ smaller than the GWAS threshold and retained 155 of them after clumping. Results from all the conventional MR methods were outlined in Table 2 and Table 3. Using random effect IVW analysis, our results showed schizophrenia can cause elevated peripheral lymphocyte count (beta=0.026; FDR adjusted p-val=0.008), and elevated monocyte count (beta=0.016; FDR adjusted p-val=0.067) and neutrophil count (beta=0.014; FDR adjusted p-val=0.099) as suggestive associations. Moreover, the more conservative estimate from weighted median method further illustrated associations between schizophrenia and lymphocyte count (beta=0.013; FDR adjusted p-val=0.049). On the other hand, IVW (OR=1.100; FDR adjusted p-val=0.021) and weighted median method (OR=1.136; FDR adjusted p-val=0.006) showed that lymphocyte count are risk factors of schizophrenia, suggesting a bidirectional relationship between schizophrenia and lymphocyte count.

Table 2: MR results of schizophrenia on differential WBC count; results are interpreted in units of standard deviation increase of absolute differential white blood cell count

Outcomes	MR methods	No. of SNP	Beta	P-value	FDR adjusted p-value
Basophil	IVW	146	0.008	0.152	0.190
Basophil	MR Egger	146	0.078	0.0002**	0.0003**
Basophil	Weighted median	146	0.0009	0.858	0.895
Basophil	MR-PRESSO	139 (95.2%)	0.003	0.512	0.640
Basophil	Contamination mixture	110 (75.3%)	-0.007	0.187	0.312
Eosinophil	IVW	146	0.010	0.206	0.206
Eosinophil	MR Egger	146	0.086	0.006**	0.006**
Eosinophil	Weighted median	146	-0.008	0.108	0.180
Eosinophil	MR-PRESSO	129 (88.4%)	-0.0006	0.885	0.885
Eosinophil	Contamination mixture	102 (69.9%)	-0.013	0.016**	0.045**
Lymphocyte	IVW	146	0.026	0.002**	0.008**
Lymphocyte	MR Egger	146	0.161	4.7e-7**	2.3e-6**
Lymphocyte	Weighted median	146	0.013	0.010**	0.049**
Lymphocyte	MR-PRESSO	131 (89.7%)	0.014	0.002**	0.010**
Lymphocyte	Contamination mixture	93 (63.7%)	0.006	0.299	0.374
Monocyte	IVW	146	0.016	0.027**	0.067*
Monocyte	MR Egger	146	0.116	3.7e-5**	9.3e-5**
Monocyte	Weighted median	146	-0.0006	0.895	0.895
Monocyte	MR-PRESSO	129 (88.4%)	0.009	0.027**	0.045**

Monocyte	Contamination mixture	99 (67.8%)	0.001	0.819	1.000
Neutrophil	IVW	146	0.014	0.059*	0.099*
Neutrophil	MR Egger	146	0.100	0.0007**	0.0008**
Neutrophil	Weighted median	146	0.010	0.041**	0.103
Neutrophil	MR-PRESSO	122 (83.6%)	0.011	0.011**	0.028**
Neutrophil	Contamination mixture	99 (67.8%)	0.011	0.018**	0.045**

*p-value<0.1 ,**p-value<0.05

Table 3: MR results of differential WBC count on schizophrenia; results are interpreted in units of log odds

Exposures	MR methods	No. of SNP	Beta	Odds ratio (OR)	P-value	FDR adjusted p-value
Basophil	IVW	193	0.107	1.113	0.047**	0.117
Basophil	MR Egger	193	0.065	1.067	0.545	0.545
Basophil	Weighted median	193	0.082	1.086	0.159	0.264
Basophil	MR-PRESSO	186 (96.4%)	0.052	1.053	0.255	0.638
Basophil	Contamination mixture	137 (71.0%)	0.009	1.009	0.797	0.797
Eosinophil	IVW	428	0.040	1.041	0.191	0.319
Eosinophil	MR Egger	428	0.078	1.081	0.188	0.235
Eosinophil	Weighted median	428	0.064	1.066	0.070*	0.174
Eosinophil	MR-PRESSO	417 (97.4%)	0.005	1.005	0.865	0.912
Eosinophil	Contamination mixture	306 (71.5%)	0.012	1.012	0.685	0.797
Lymphocyte	IVW	484	0.095	1.100	0.004**	0.021**
Lymphocyte	MR Egger	484	0.172	1.188	0.013**	0.065*
Lymphocyte	Weighted median	484	0.127	1.136	0.001**	0.006**
Lymphocyte	MR-PRESSO	462 (95.5%)	0.045	1.046	0.125	0.625
Lymphocyte	Contamination mixture	336 (69.4%)	-0.009	0.991	0.773	0.797
Monocyte	IVW	492	0.017	1.017	0.535	0.535
Monocyte	MR Egger	492	0.081	1.084	0.083*	0.204
Monocyte	Weighted median	492	0.003	1.003	0.922	0.997
Monocyte	MR-PRESSO	470 (95.5%)	0.002	1.002	0.912	0.912
Monocyte	Contamination mixture	351 (71.3%)	0.025	1.025	0.194	0.797

Neutrophil	IVW	412	0.032	1.033	0.410	0.512
Neutrophil	MR Egger	412	0.125	1.133	0.122	0.204
Neutrophil	Weighted median	412	-0.0002	1.000	0.997	0.997
Neutrophil	MR-PRESSO	399 (96.8%)	0.011	1.011	0.742	0.912
Neutrophil	Contamination mixture	282 (68.4%)	0.017	1.017	0.620	0.797

*p-value<0.1 ,**p-value<0.05

Nevertheless, Cochran's Q and I^2 staitstics (refer to supplementary table 1 & 2) indicate that there is substantial heterogeneity of the included genetic instruments, which could bias the results. Scatter plots and leave-one-out analysis had further confirmed the existence of a dominant SNP rs13195636. Rs13195636 was the SNP with the smallest p-value in the schizophrenia GWAS after clumping, and it also contributed heavily to the significant results on various differential WBC count. This suggested that the loci where rs13195636 was located influenced both schizophrenia and differential WBC counts more heavily than other SNPs, which could be due to horizontal pleiotropy. The MR-Egger intercept test and the MR-PRESSO global test also showed significant evidence of horizontal pleiotropy. As a result, estimates from conventional MR approaches are biased. Therefore, further MR methods were used to obtain outlier-robust estimates.

MR-PRESSO and contamination mixture methods both derived outlier-robust estimates by removing the outliers. Using MR-PRESSO method, we removed outlier and pleiotropic SNPs such as rs13195636, and the results suggested that schizophrenia might increase lymphocyte count (beta=0.014; FDR adjusted p-val=0.010), monocyte count (beta=0.009; FDR adjusted p-val=0.045) and neutrophil count (beta=0.011; FDR adjusted p-val=0.028). Nevertheless, only a handful of most significant SNPs were excluded using MR-PRESSO, and the Cochrane Q's values indicated considerable heterogeneity remained (Supplementary Table 1 & 2). Therefore, these results should be interpreted with caution. Cochrane Q's values showed that contamination mixture method reduced the heterogeneity of included SNPs substantially (Supplementary Table 1), and the results suggested that schizophrenia might increase neutrophil count (beta=0.011; FDR adjusted p-val=0.045). Surprisingly, contamination mixture method suggested that schizophrenia reduces eosinophil count (beta=-0.013, FDR adjusted p-val=0.045), which did not observe from other methods. Conversly, MR-PRESSO and contamination cixture method suggest that WBC could not causally cause schizophrenia. Nevertheless, Table 4 provides summary of results from novel MRCI method, which suggested that both eosinophil count (OR=1.050; FDR adjusted p-val=0.010) and lymphocyte count (OR=1.080; FDR adjusted p-val=0.010) may casually link to developing schizophrenia after modelling the SNPs that contribute to pleiotropic effects to both schizophrenia and WBC.

Table 4: Bidirectional estimated effect of differential WBC on schizophrenia estimated by MRCI method; results are interpreted in units of standard deviation increase of absolute differential WBC count and log odds

Exposure	Outcome	Beta	OR	P-value	FDR adjusted p-value
Schizophrenia	Basophil count	-5.1e-3	0.995	0.985	1.000
Schizophrenia	Eosinophil count	-7.1e-4	0.999	0.976	1.000
Schizophrenia	Lymphocyte count	-2.2e-5	1.000	0.999	1.000
Schizophrenia	Monocyte count	-1.2e-6	1.000	1.000	1.000
Schizophrenia	Neutrophil count	-1.6e-5	1.000	0.999	1.000
Basophil count	Schizophrenia	0.021	1.021	0.642	0.642
Eosinophil count	Schizophrenia	0.048	1.050	0.010**	0.026**

Lymphocyte count	Schizophrenia	0.077	1.080	0.010**	0.026**
Monocyte count	Schizophrenia	0.032	1.033	0.061*	0.102
Neutrophil count	Schizophrenia	0.027	1.027	0.318	0.398

*p-value<0.1 ,**p-value<0.05

4. Discussion

We conducted bidirectional two-sample MR analyses using the two largest genomic datasets to-date on schizophrenia and WBC, and investigated the possible causal relationships between the two traits, as well as putative causal mechanisms for linking them. This study used hundreds of SNPs to examine the bidirectional relationships, and the sample size was sufficient to detect true effects as low as 0.05 standard deviation away from the mean of differential WBC. In light of the substantial heterogeneity and outliers in the datasets, we carried out a few outlier-robust MR methods as well as the MRCI which explicitly modelled the situation of horizontal pleiotropy.

Our findings generally supported a causal relationship between raised lymphocyte count and schizophrenia. Based on the unbiased results generated from MRCI approach and the results corrected for multiple testing, our analysis also indicated that excess eosinophil count, which may disrupt the immune microenvironment, could also contribute to cause of schizophrenia. Furthermore, despite the possibilities of spurious estimates caused by pleiotropy from conventional MR approaches based on the substantial heterogeneity (table S2), contamination mixture method suggested schizophrenia could possibly lead to reduction of eosinophil count and elevated neutrophil count. Schizophrenia reduce eosinophil count could suggest the altered innate immune system in schizophrenia patients, which might partially explain the vulnerability of getting infection in schizophrenia patients (Pankiewicz-Dulacz et al., 2018). Moreover, elevated neutrophil count in schizophrenia had been reported by numerous studies looking into NLR, but causality could not be confirmed in their study design. Our smaller effect sizes than the reported effect sizes in observational studies indicate that likely both environmental factors and genetic factors have roles in influencing the differential WBC, as the SNPs included in this study couldn't explain all the differences.

Our findings are broadly consistent with the extant literature. For example, our finding supported the macrophage-T-lymphocyte theory of schizophrenia, i.e., activated T-lymphocytes play an important role in the neurotransmitter abnormalities and neuro-inflammation model of schizophrenia (Smith & Maes, 1995). Dopamine receptors were found to be expressed in different T cells and B cells, which could activate T cell functions in the human body (Levite, 2015). Activated T cells may affect the dopamine activity in the brain, and therefore increase the risk of developing psychosis (Kesby et al., 2018). Besides, blood-borne monocytes were associated with the immune system in central nervous system. Impaired and dysregulated monocyte activation were reported in schizophrenia (Uranova at el., 2017). Neurotoxic pro-inflammatory macrophages (M1) and anti-inflammatory macrophages (M2) can both be produced, depending on influences of different environments that can trigger the selective evolution of infiltrating monocytes (Garré & Yang, 2018). Last but not least, hypereosinophilia has long been linked to various neurological and psychiatric symptoms (Sin & Lee; 2014). As a result, understanding the relationship between

schizophrenia and different WBC can bring further insights in the risk of developing infection and autoimmune diseases.

Another notable finding from our study was that a single SNP, rs13195636, appeared to be highly influential and pleiotropic in conventional MR methods. This SNP is located at the major histocompatibility complex on chromosome 6. The population minor allele frequencies range from 1.8% in African to 10% in Non-Finnish North-West European (Open Targets, 2023). It is the strongest SNP reported by the PGC-schizophrenia and also highly significant in differential WBC reported by the Blood Cell Consortium. Genes located at MHC were believed to have critical roles in neurodevelopment and immune regulatory (Gibney & Drexhage, 2013). Its closest protein coding gene is ZNF184 gene, which is found to be associated with bipolar disorder (Huang et al., 2022), depression (Wu et al., 2021), and Parkinson's disease (Li et al., 2021). We speculated two possible explanations for this SNP. First, it could be owing to horizontal pleiotropy, which implies that this SNP has independent effects on multiple traits, including schizophrenia and different white blood cell subtypes. Another plausible explanation is that the effect size could possibly be inflated because of linkage disequilibrium with other genetic variants, which refers to non-random associations of alleles at multiple loci.

This study has several strengths. First, two largest publicly available genomic datasets of schizophrenia and differential WBC were used, providing hundreds of SNPs that had reached GWAS significance threshold, and thus conferring sufficient power to detect small effects. Although entering many instruments in the model is prone to weak instrument bias, this bias tends towards the null in two-sample MR (Yang et al., 2022). Therefore, significant results in this study were unlikely to be false positive due to weak instrument bias. Second, MR examined the causal relationships that typically could not be assessed using observational study design. Confounders, such as antipsychotic exposures, may bias the outcome measurements in observational research (Zhou et al., 2020; Dawidowski et al., 2022). Our two-sample MR study minimized the influence of confounders by including one sample with schizophrenia only and another sample with minimal subjects with schizophrenia. Third, we employed different MR methods, and carefully identified any violation against the MR assumptions. The novel MRCI method enabled us to obtain unbiased estimates with conservative standard errors. Lastly, we corrected the potential inflation of Type I error due to multiple testing using false discovery rate adjusted p-values.

Several limitations should be borne in mind. Although WBC to some extent shows trait-like properties, these indices are known to vary with time, even within a single day (Hilderink et al., 2017). This temporal variation can induce error and undermine the variance accounted for by the SNPs included in this study. Second, only autosome GWAS data of the Blood Cell Consortium were publicly available. Therefore, we only used autosome GWAS data from Psychiatric Genomics Consortium, and the potential influences of sex chromosome (i.e. sex differences in schizophrenia and WBC) were not captured in our results. In addition, using GWAS results from large meta-analysis studies may lead to higher risk of sample overlap between the exposure and outcome, which can bias the results to the null or the observational estimate (Hemani et al., 2018). Our results could be biased by a small portion of schizophrenia patients were recruited into UK biobank, which could not be identified and

excluded from the Blood Cell Consortium. Lastly, due to the different population structure of PGC and Blood Cell Consortium, our analysis only used the European ancestry for fair comparison.

5. Conclusion

To conclude, the relationship between schizophrenia and immunological system is complex. Our two-sample MR on two large datasets provided preliminary evidence that elevated lymphocyte count, a potential inflammatory biomarker, can increase the risk of schizophrenia; whereas elevated monocyte count may be protective against developing schizophrenia. Future research should explore the immune-related biological mechanisms for schizophrenia, as well as the generalizability of our results in different ancestry and broader populations.

6. Acknowledgements

This study was funded by HKU Seed Fund for Basic Research for New Staff (202009185071) and the HKU Enhanced Start-up Fund for New Staff granted to Simon S.Y. Lui.

The authors declare no competing interests.

References

Alvarez-Herrera, S., Escamilla, R., Medina-Contreras, O., Saracco, R., Flores, Y., Hurtado-Alvarado, G., … & Pavón, L. (2020). Immunoendocrine peripheral effects induced by atypical antipsychotics. *Frontiers in endocrinology*, *11*, 195.

Austermann, J., Roth, J., & Barczyk-Kahlert, K. (2022). The good and the bad: Monocytes' and macrophages' diverse functions in inflammation. *Cells*, *11*(12), 1979.

Bioque, M., Catarina Matias-Martins, A., Llorca-Bofí, V., Mezquida, G., Cuesta, M. J., Vieta, E., ... & Balanzá, V. (2022). Neutrophil to Lymphocyte Ratio in Patients With a First Episode of Psychosis: A Two-Year Longitudinal Follow-up Study. *Schizophrenia Bulletin*, *48*(6), 1327-1335.

Boef, A. G., Dekkers, O. M., & Le Cessie, S. (2015). Mendelian randomization studies: a review of the approaches used and the quality of reporting. *International journal of epidemiology*, *44*(2), 496-511.

Bowden, J., Davey Smith, G., & Burgess, S. (2015). Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *International journal of epidemiology*, *44*(2), 512-525.

Bowden, J., Davey Smith, G., Haycock, P. C., & Burgess, S. (2016). Consistent estimation in Mendelian randomization with some invalid instruments using a weighted median estimator. *Genetic epidemiology*, *40*(4), 304-314.

Burgess, S., Foley, C. N., Allara, E., Staley, J. R., & Howson, J. M. (2020). A robust and efficient method for Mendelian randomization with hundreds of genetic variants. *Nature communications*, *11*(1), 376.

Burgess, S., & Thompson, S. G. (2017). Interpreting findings from Mendelian randomization using the MR-Egger method. *European journal of epidemiology*, *32*, 377-389.

Burgess, S. (2023). Online sample size and power calculator for Mendelian randomization with a binary outcome. https://sb452.shinyapps.io/power/

Carter, A. R., Gill, D., Davies, N. M., Taylor, A. E., Tillmann, T., Vaucher, J., ... & Dehghan, A. (2019). Understanding the consequences of education inequality on cardiovascular disease: mendelian randomisation study. *bmj*, *365*.

Chan, S. H. T., Yu, T., Zhang, Z., Chang, L. Y., Guo, C., Bo, Y., ... & Lao, X. Q. (2022). Total and differential white blood cell count and cause-specific mortality in 436 750 Taiwanese adults. *Nutrition, Metabolism and Cardiovascular Diseases*, *32*(4), 937-947.

Chen, M. H., Raffield, L. M., Mousas, A., Sakaue, S., Huffman, J. E., Moscati, A., ... & Lettre, G. (2020). Trans-ethnic and ancestry-specific blood-cell genetics in 746,667 individuals from 5 global populations. *Cell*, *182*(5), 1198-1213.

Davies, C., Segre, G., Estradé, A., Radua, J., De Micheli, A., Provenzani, U., ... & Fusar-Poli, P. (2020). Prenatal and perinatal risk and protective factors for psychosis: a systematic review and meta-analysis. *The Lancet Psychiatry*, *7*(5), 399-410.

Dawidowski, B., Grelecki, G., Biłgorajski, A., Podwalski, P., Misiak, B., & Samochowiec, J. (2021). Effect of Antipsychotic Treatment on Neutrophil-to-Lymphocyte Ratio during Hospitalization for Acute Psychosis in the Course of Schizophrenia—A Cross-Sectional Retrospective Study. *Journal of Clinical Medicine*, *11*(1), 232.Dionisie, V., Filip, G. A., Manea, M. C., Movileanu, R. C., Moisa, E., Manea, M., ... & Ciobanu, A. M. (2021). Neutrophil-to-lymphocyte ratio, a novel inflammatory marker, as a predictor of bipolar type in depressed patients: a quest for biological markers. *Journal of clinical medicine*, *10*(9), 1924.

Evans, D. M., Frazer, I. H., & Martin, N. G. (1999). Genetic and environmental causes of variation in basal levels of blood cells. *Twin Research and Human Genetics*, *2*(4), 250-257.

Ermakov, E. A., Melamud, M. M., Buneva, V. N., & Ivanova, S. A. (2022). Immune system abnormalities in schizophrenia: an integrative view and translational perspectives. *Frontiers in Psychiatry*, *13*.

Farhangi, M. A., Keshavarz, S. A., Eshraghian, M., Ostadrahimi, A., & Saboor-Yaraghi, A. A. (2013). White blood cell count in women: relation to inflammatory biomarkers, haematological profiles, visceral adiposity, and other cardiovascular risk factors. *Journal of health, population, and nutrition*, *31*(1), 58.

Gao, Z., Li, B., Guo, X., Bai, W., & Kou, C. (2023). The association between schizophrenia and white blood cells count: a bidirectional two-sample Mendelian randomization study. *BMC Psychiatry*, *23*(1), 1-7.

Garré, J. M., & Yang, G. (2018). Contributions of monocytes to nervous system disorders. *Journal of Molecular Medicine*, *96*(9), 873-883

Gasteiger, G., & Rudensky, A. Y. (2014). Interactions between innate and adaptive lymphocytes. *Nature Reviews Immunology*, *14*(9), 631-639.

GBD 2019 Mental Disorders Collaborators. (2022). Global, regional, and national burden of 12 mental disorders in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *The Lancet Psychiatry*, *9*(2), 137-150.

Gibney, S. M., & Drexhage, H. A. (2013). Evidence for a dysregulated immune system in the etiology of psychiatric disorders. *Journal of Neuroimmune Pharmacology*, *8*(4), 900-920.

Grassadonia, A., Graziano, V., Iezzi, L., Vici, P., Barba, M., Pizzuti, L., ... & Tinari, N. (2021). Prognostic relevance of neutrophil to lymphocyte ratio (NLR) in luminal breast cancer: a retrospective analysis in the neoadjuvant setting. *Cells*, *10*(7), 1685.

Hannon, E., Dempster, E. L., Mansell, G., Burrage, J., Bass, N., Bohlken, M. M., ... & Mill, J. (2021). DNA methylation meta-analysis reveals cellular alterations in psychosis and markers of treatment-resistant schizophrenia. *Elife*, *10*, e58430.

Hartwig, F. P., Borges, M. C., Horta, B. L., Bowden, J., & Smith, G. D. (2017). Inflammatory biomarkers and risk of schizophrenia: a 2-sample Mendelian randomization study. *JAMA psychiatry*, *74*(12), 1226-1233.

Hemani, G., Bowden, J., & Davey Smith, G. (2018). Evaluating the potential role of pleiotropy in Mendelian randomization studies. *Human molecular genetics*, *27*(R2), R195-R208.

Hilderink, J. M., Klinkenberg, L. J., Aakre, K. M., de Wit, N. C., Henskens, Y. M., van der Linden, N., ... & Meex, S. J. (2017). Within-day biological variation and hour-to-hour reference change values for hematological parameters. *Clinical Chemistry and Laboratory Medicine (CCLM)*, *55*(7), 1013-1024.

Huang, Y., Liu, Y., Wu, Y., Tang, Y., Zhang, M., Liu, S., ... & Wang, Q. (2022). Patterns of convergence and divergence between bipolar disorder type I and type II: evidence from integrative genomic analyses. *Frontiers in Cell and Developmental Biology*, *10*.

Hung, S. C., Cheng, H. Y., Yang, C. C., Lin, C. I., Ho, C. K., Lee, W. H., ... & Chuang, H. Y. (2021). The association of white blood cells and air pollutants—A population-based study. *International Journal of Environmental Research and Public Health*, *18*(5), 2370.

Imtiaz, F., Shafique, K., Mirza, S. S., Ayoob, Z., Vart, P., & Rao, S. (2012). Neutrophil lymphocyte ratio as a measure of systemic inflammation in prevalent chronic diseases in Asian population. *International archives of medicine*, *5*(1), 1-6.

Jackson, A. J., & Miller, B. J. (2020). Meta-analysis of total and differential white blood cell counts in schizophrenia. *Acta Psychiatrica Scandinavica*, *142*(1), 18-26.

Jain, D., Hodonsky, C. J., Schick, U. M., Morrison, J. V., Minnerath, S., Brown, L., ... & Reiner, A. P. (2017). Genome-wide association of white blood cell counts in Hispanic/Latino Americans: the Hispanic Community Health Study/Study of Latinos. *Human molecular genetics*, *26*(6), 1193-1204.

Kar, N., Barreto, S., & Chandavarkar, R. (2016). Clozapine monitoring in clinical practice: beyond the mandatory requirement. *Clinical Psychopharmacology and Neuroscience*, *14*(4), 323.Karageorgiou, V., Milas, G. P., & Michopoulos, I. (2019). Neutrophil-to-lymphocyte ratio in schizophrenia: A systematic review and meta-analysis. *Schizophrenia research*, *206*, 4-12.

Kesby, J. P., Eyles, D. W., McGrath, J. J., & Scott, J. G. (2018). Dopamine, psychosis and schizophrenia: the widening gap between basic and clinical neuroscience. *Translational psychiatry*, *8*(1), 30.

Kim, K. I., Lee, J., Heo, N. J., Kim, S., Chin, H. J., Na, K. Y., ... & Kim, S. (2013). Differential white blood cell count and all-cause mortality in the Korean elderly. *Experimental Gerontology*, *48*(2), 103-108.

Lawlor, D. A. (2016). Commentary: Two-sample Mendelian randomization: opportunities and challenges. *International journal of epidemiology*, *45*(3), 908-915.

Lettre, G. (2023). Blood-cell genetics. The Lettre lab. http://www.mhi-humangenetics.org/en/welcome/#anc_projects

Levite, M. (2016). Dopamine and T cells: dopamine receptors and potent effects on T cells, dopamine production in T cells, and abnormalities in the dopaminergic system in T cells in autoimmune, neurological and psychiatric diseases. *Acta physiologica*, *216*(1), 42-89.

Li, C. Y., Ou, R. W., Chen, Y. P., Gu, X. J., Wei, Q. Q., Cao, B., ... & Shang, H. F. (2021). Genetic analysis of ZNF protein family members for early-onset Parkinson's disease in Chinese population. *Molecular Neurobiology*, *58*, 3435-3442.

Li, Y., Wang, W., Yang, F., Xu, Y., Feng, C., & Zhao, Y. (2019). The regulatory roles of neutrophils in adaptive immunity. *Cell Communication and Signaling*, *17*(1), 1-11.

Lin, B. D., Hottenga, J. J., Abdellaoui, A., Dolan, C. V., de Geus, E. J., Kluft, C., ... & Willemsen, G. (2016). Causes of variation in the neutrophil–lymphocyte and platelet–lymphocyte ratios: a twin-family study. *Biomarkers in medicine*, *10*(10), 1061-1072.

Lin, B. D., Willemsen, G., Fedko, I. O., Jansen, R., Penninx, B., De Geus, E., ... & Boomsma, D. I. (2017). Heritability and gwas studies for monocyte–lymphocyte ratio. *Twin Research and Human Genetics*, *20*(2), 97-107.

Liu, Y., Du, X., Chen, J., Jin, Y., Peng, L., Wang, H. H., ... & Zhao, Y. (2020). Neutrophil-to-lymphocyte ratio as an independent risk factor for mortality in hospitalized patients with COVID-19. *Journal of Infection*, *81*(1), e6-e12.

Liu, Z., Qin, Y., Wu, T. et al. Reciprocal causation mixture model for robust Mendelian randomization analysis using genome-scale summary data. *Nat Commun* 14, 1131 (2023).

Llorca-Bofí, V., Palacios-Garrán, R., Routo, D. R., Buil-Reiné, E., Adrados-Pérez, M., Gich, I., ... & Mur, M. (2021). High neutrophil-lymphocyte ratio upon admission is associated with better response in psychotic depression. *Journal of Psychiatric Research*, *143*, 38-42.

Mazza, M. G., Capellazzi, M., Lucchi, S., Tagliabue, I., Rossetti, A., & Clerici, M. (2020). Monocyte count in schizophrenia and related disorders: a systematic review and meta-analysis. *Acta neuropsychiatrica*, *32*(5), 229-236.

Miller, B. J., Gassama, B., Sebastian, D., Buckley, P., & Mellor, A. (2013). Meta-analysis of lymphocytes in schizophrenia: clinical status and antipsychotic effects. *Biological psychiatry*, *73*(10), 993-999.

Nora, I., Shridhar, R., Huston, J., & Meredith, K. (2018). The accuracy of neutrophil to lymphocyte ratio and platelet to lymphocyte ratio as a marker for gastrointestinal malignancies. Journal of gastrointestinal oncology, 9(5), 972.

Open Targets. (2023). 6_27541714_A_C. Open Targets Genetics. https://https://genetics.opentargets.org/variant/6_27541714_A_C

Pankiewicz-Dulacz, M., Stenager, E., Chen, M., & Stenager, E. (2018). Incidence rates and risk of hospital registered infections among schizophrenia patients before and after onset of illness: a population-based nationwide register study. *Journal of clinical medicine*, *7*(12), 485.

Paribello, P., Manchia, M., Zedda, M., Pinna, F., & Carpiniello, B. (2021). Leukocytosis associated with clozapine treatment: A case series and systematic review of the literature. *Medicina*, *57*(8), 816.

Ren, H., Meng, Y., Zhang, Y., Wang, Q., Deng, W., Ma, X., ... & Li, T. (2021). Spatial expression pattern of ZNF391 gene in the brains of patients with schizophrenia, bipolar

disorders or major depressive disorder identifies new cross-disorder biotypes: A trans-diagnostic, top-down approach. *Schizophrenia Bulletin*, *47*(5), 1351-1363.

Shah, S., Henry, A., Roselli, C., Lin, H., Sveinbjörnsson, G., Fatemifar, G., ... & Lumbers, R. T. (2020). Genome-wide association and Mendelian randomisation analysis provide insights into the pathogenesis of heart failure. *Nature communications*, *11*(1), 163.

Shamri, R., Xenakis, J. J., & Spencer, L. A. (2011). Eosinophils in innate immunity: an evolving story. *Cell and tissue research*, *343*(1), 57-83.

Shim, W. S., Kim, H. J., Kang, E. S., Ahn, C. W., Lim, S. K., Lee, H. C., & Cha, B. S. (2006). The association of total and differential white blood cell count with metabolic syndrome in type 2 diabetic patients. *Diabetes research and clinical practice*, *73*(3), 284-291.

Sin, G. L., & Lee, T. S. (2008). Cognitive and psychotic symptoms in idiopathic hypereosinophilic syndrome. *Acta Neuropsychiatrica*, *20*(1), 53-54.

Slob, E. A., & Burgess, S. (2020). A comparison of robust Mendelian randomization methods using summary data. *Genetic epidemiology*, *44*(4), 313-329.

Smith, R. S., & Maes, M. (1995). The macrophage-T-lymphocyte theory of schizophrenia: additional evidence. *Medical hypotheses*, *45*(2), 135-141.

Trubetskoy, V., Pardiñas, A. F., Qi, T., Panagiotaropoulou, G., Awasthi, S., Bigdeli, T. B., ... & Lazzeroni, L. C. (2022). Mapping genomic loci implicates genes and synaptic biology in schizophrenia. *Nature*, *604*(7906), 502-508.

Tylee, D. S., Sun, J., Hess, J. L., Tahir, M. A., Sharma, E., Malik, R., ... & Glatt, S. J. (2018). Genetic correlations among psychiatric and immune-related phenotypes based on genome-wide association data. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, *177*(7), 641-657.

Uranova, N. A., Bonartsev, P. D., Androsova, L. V., Rakhmanova, V. I., & Kaleda, V. G. (2017). Impaired monocyte activation in schizophrenia: ultrastructural abnormalities and increased IL-1β production. *European Archives of Psychiatry and Clinical Neuroscience*, *267*(5), 417-426.

Van Kesteren, C. F. M. G., Gremmels, H., De Witte, L. D., Hol, E. M., Van Gool, A. R., Falkai, P. G., ... & Sommer, I. E. C. (2017). Immune involvement in the pathogenesis of schizophrenia: a meta-analysis on postmortem brain studies. *Translational psychiatry*, *7*(3), e1075-e1075.

Verbanck, M., Chen, C. Y., Neale, B., & Do, R. (2018). Detection of widespread horizontal pleiotropy in causal relationships inferred from Mendelian randomization between complex traits and diseases. *Nature genetics*, *50*(5), 693-698.

Vidal, P. M., & Pacheco, R. (2020). The cross-talk between the dopaminergic and the immune system involved in schizophrenia. *Frontiers in Pharmacology*, *11*, 394.

Vuckovic, D., Bao, E. L., Akbari, P., Lareau, C. A., Mousas, A., Jiang, T., ... & Soranzo, N. (2020). The polygenic and monogenic basis of blood traits and diseases. *Cell*, *182*(5), 1214-1231.

Whitfield, J. B., Martin, N. G., & Rao, D. C. (1985). Genetic and environmental influences on the size and number of cells in the blood. *Genetic epidemiology*, *2*(2), 133-144.

Wildgust, H. J., Hodgson, R., & Beary, M. (2010). The paradox of premature mortality in schizophrenia: new research questions. *Journal of Psychopharmacology*, *24*(4_suppl), 9-15.

Williams, J. A., Burgess, S., Suckling, J., Lalousis, P. A., Batool, F., Griffiths, S. L., ... & PIMS Collaboration. (2022). Inflammation and brain structure in schizophrenia and other neuropsychiatric disorders: a Mendelian randomization study. *JAMA psychiatry*, *79*(5), 498-507.

Wootton, R. E., Lawn, R. B., Millard, L. A., Davies, N. M., Taylor, A. E., Munafò, M. R., ... & Haworth, C. M. (2018). Evaluation of the causal effects between subjective wellbeing and cardiometabolic health: mendelian randomisation study. *bmj*, *362*.

World Health Organization. (2022). Schizophrenia. https://www.who.int/news-room/fact-sheets/detail/schizophrenia

Wu, W., Howard, D., Sibille, E., & French, L. (2021). Differential and spatial expression meta-analysis of genes identified in genome-wide association studies of depression. *Translational psychiatry*, *11*(1), 8.

Yang, Q., Sanderson, E., Tilling, K., Borges, M. C., & Lawlor, D. A. (2022). Exploring and mitigating potential bias when genetic instrumental variables are associated with multiple non-exposure traits in Mendelian randomization. *European Journal of Epidemiology*, *37*(7), 683-700.

Zhang, N., Zhang, Z. M., & Wang, X. F. (2021). The roles of basophils in mediating the immune responses. *European Journal of Inflammation*, *19*, 20587392211047644.

Zhou, X., Wang, X., Li, R., Yan, J., Xiao, Y., Li, W., & Shen, H. (2020). Neutrophil-to-lymphocyte ratio is independently associated with severe psychopathology in schizophrenia and is changed by antipsychotic administration: a large-scale cross-sectional retrospective study. *Frontiers in psychiatry*, *11*, 581061.