Food & You: A Digital Cohort on Personalized Nutrition

Harris Héritier[1], Chloé Allémann[1], Oleksandr Balakiriev[1], Victor Boulanger[1], Sean F. Carroll[1], Noé Froidevaux[1], Germain Hugon[1], Yannis Jaquet[1], Djilani Kebaili[1], Sandra Riccardi[1], Geneviève Rousseau-Leupin[1], Rahel M. Salathé[1], Talia Salzmann[1], Rohan Singh[1], Laura Symul[1,2], Elif Ugurlu-Baud[1], Peter de Verteuil[1], Marcel Salathé[1]

[1] Digital Epidemiology Lab, School of Life Sciences, School of Computer and Communication Sciences, EPFL, Switzerland

[2] Department of Statistics, Stanford University, USA

Nutrition is a key contributor to health. Recently, several studies have identified associations between factors such as microbiota composition and health-related responses to dietary intake, raising the potential of personalized nutritional recommendations. To further our understanding of personalized nutrition, detailed individual data must be collected from participants in their day-to-day lives. However, this is challenging in conventional studies that require clinical measurements and site visits. So-called digital or remote cohorts allow *in situ* data collection on a daily basis through mobile applications, online services, and wearable sensors, but they raise questions about study retention and data quality. "Food & You" is a personalized nutrition study implemented as a fully digital cohort in which participants track food intake, physical activity, gut microbiota, glycemia, and other data for two to four weeks. Here, we describe the study protocol, report on study completion rates, and describe the collected data, focusing on assessing their quality and reliability. Overall, the study collected data from over 1000 participants, including high-resolution data of nutritional intake of more than 46 million kcal collected from 315,126 dishes over 23,335 participant days, 1,470,030 blood glucose measurements, 49,110 survey responses, and 1,024 stool samples for gut microbiota analysis. Retention was high,

32 **with over 60% of the enrolled participants completing the study. Various data quality**

33 **assessment efforts suggest the captured high-resolution nutritional data accurately**

34 **reflect individual diet patterns, paving the way for digital cohorts as a typical study**

35 **design for personalized nutrition.**

37 **Keywords** personalized nutrition · digital cohort · gut microbiota · glycemia

Introduction

Nutrition plays a significant role in moderating the risk and/or severity of several

diseases, such as type 2 diabetes[1], cardiovascular diseases,[2,3] or cancer[4]. Findings from

nutritional epidemiology studies have led to dietary guidelines and public health

campaigns designed to support healthy diets. However, while these recommendations

are generally based on results aggregated at the population level, a more individualized

approach to health[5] has led to the concept of personalized nutrition. For example, a

randomized study showed that personalized recommendations improved diet quality as

measured by the Healthy Eating Index (HEI) compared to a control group[6]. Zeevi and

colleagues showed how personalized nutrition algorithms could be used to design diets

that lower postprandial glucose responses[7]. Further studies showed the importance of

personal features such as gut microbiota compositions on glycemic responses[8,9]. Another

intervention study showed significant improvement in the food categories consumed

when receiving personalized diet advice, compared to generic or no advice[10]. These

findings highlight the need for a more holistic and individual approach to nutritional

epidemiology, encompassing diet, gut microbiota, physical activity, and lifestyle.

Blood glucose response is a particularly interesting outcome measure for nutritional

studies due to its association with insulin resistance, metabolic syndrome, and diseases

such as stroke, type 2 diabetes, and heart disease[11]. Reducing blood glucose levels is thus

a recommendation of public health authorities around the globe. Identified risk factors

for elevated blood glucose levels include a carbohydrate-rich diet, lack of physical

activity, and poor sleep, among others. In addition, studies have begun to investigate the

role of the gut microbiome in modulating the blood glucose response to food intake.
Nutritional studies trying to understand postprandial glucose response (PPGR) are thus
faced with the challenge of obtaining data on relevant factors all at once, ideally
continuously and *in situ*, that is, in the regular environment in which participants' lives
unfold.
68
In digital health studies - also called remote or siteless studies - all interactions with
participants, as well as data collection, are digital or digitally coordinated. These studies
leverage an array of digital devices, wearable sensors, and online services. Digital cohorts
and trials have been heralded as a new major development for epidemiological and clinical
studies. However, since digital cohorts are a relatively new study approach, open
questions regarding selection bias, retention, and data quality remain. Indeed, access to
devices connected to the internet and digital literacy may lead to selection bias which in
turn may lead to a lack of representativity of the study population compared to the
general population. Furthermore, the time burden generated by following the study
protocol and collecting the data might create response fatigue, which could in turn
translate into lower study adherence, or data quality. Finally, novel data collection
methods may not have been thoroughly validated.
81
Here, we address some of these questions by evaluating the completion rates, adherence,
and data quality of the "Food and You" digital cohort on personalized nutrition. The "Food
& You" study started in early 2019 and consisted of two distinct digital sub-cohorts; the
sub-cohort "Basic" (cohort B) restricted to non-diabetic participants, and the sub-cohort
"Cycle" (cohort C) restricted to non-diabetic women of reproductive age who did not use
hormonal contraceptive or medication (see Methods for inclusion / exclusion criteria).
The study duration for cohort B was 14 days, whereas cohort C participants were enrolled
for 28 days, matching the length of a typical menstrual cycle. The study was performed
in Swizterland, and all participant were required to have a postal address in Switzerland.
Throughout the study, participants were requested to report i) food consumption using
an AI-assisted food tracking app (MyFoodRepo), ii) continuous blood sugar levels using a
continuous glucose monitor, and iii) physical activity and sleep using either activity
trackers or daily surveys. They were furthermore asked to follow a protocol that included

a one-time stool sample collection for gut microbiota analysis, and the consumption of
standardized breakfasts.

The present paper details the study protocol, and reports study engagement data by
looking at the individual characteristics of participants on their journey from enrollment
to completion. Further, we provide an overview of the data collected in the "Food & You"
cohort from October 2018 to March 2023, and describe our efforts to assess data quality,
including the comparison of nutritional and microbiota data collected in "Food & You"
with data collected in traditional (on-site) studies. We also discuss the challenges of
running a complex digital cohort, and how we addressed them. Overall, retention rates
were relatively high, with more than 60% of enrolled participants completing the study.
Despite certain fatigue over time, adherence very high, especially for glucose response
data, nutritional data, microbiota data, and data from daily surveys. While the study
population shows some demographic differences compared to the overall population, the
nutrition patterns are in very good agreement with data obtained in another study from
a representative sample of the general Swiss population.

Methods

STUDY DESIGN AND SETTING

The Swiss "Food & You" study is a digital cohort study collecting data on glycemia,
nutrition, gut microbiota, lifestyle, and physical activity as well as demographic data
(Figure 1). The cohort was open to anyone fulfilling the inclusion criteria listed in
Supplementary Table 1. The study consisted of four sequential phases: enrollment phase,
preparatory phase, tracking days phase, and follow-up phase (Figure 2). In the *enrollment*
phase, interested participants were first required to perform a self-check of their
eligibility, and fill out a consent form and a short screening questionnaire. Following this,
and upon acceptance by a member of the study team (based on the available capacity to
accommodate new participants), the participants were considered enrolled. During the
subsequent *preparatory phase*, enrolled participants were given instructions on the "Food
& You" website and asked to fill out a series of questionnaires. Next, they were instructed

to download the AI-assisted nutrition tracking app MyFoodRepo[1] (MFR) to track their
food intake for a trial period of at least three days. After the successful completion of the
trial period, participants would order the study material which included, among other
things, a continuous glucose monitoring (CGM) sensor for each period of 14 days and
stool-sample self-collection material. Upon receipt, they were asked to choose the
starting date of their tracking days phase. In the *tracking days phase*, participants were
required to log all their food and drink intake via the MyFoodRepo app, wear the CGM
sensor, and answer two daily surveys for 14 (cohort B) and 28 (cohort C) days. In addition,
participants were instructed to take standardized breakfasts in accordance with dietary
restrictions from the 2nd to the 7th day during the first week. They were asked to avoid
altering standardized meals, as well as to refrain from eating or engaging in physical
activity for the subsequent two hours. Cohort C participants repeated these instructions
on their third tracking week (Supplementary Table 2). On days 6 and 7 (additionally on
days 21 and 22 for cohort C) they were asked to perform an oral glucose tolerance test by
drinking 50g of glucose. In addition, study participants were asked to provide one stool
sample collected anytime during the tracking days phase. At the end of the tracking days
phase, participants were asked to upload their physical activity and CGM data on the
"Food & You" website. In the *follow-up phase*, they were requested to fill out a feedback
questionnaire regarding their experience. Participants were also provided with
interactive visualizations of their data (Supplementary Figure 1). Cohort C participants
were followed for two additional menstrual cycles during which they continued to track
their menstrual cycle and fertility-related body signs such as morning temperature and
cervical mucus characteristics.
148
149   The Geneva ethics commission has reviewed and authorized the project (Ethical Approval
150   Number: 2017-02124). The study is registered on the website of the Federal Office of
151   Public Health (SNCTP000002833) and the platform clinicaltrials.gov (NCT03848299).
152
153
154
155

---

[1] https://www.myfoodrepo.org/

DATA COLLECTION

Questionnaires: During the enrollment phase, interested subjects were requested to fill out a screening questionnaire with items regarding age, gender, height, weight, type of mobile phone and dietary restrictions. During the preparatory phase, enrolled participants had to fill out a lifestyle and health-related questionnaire. Participants were asked about their general health (smoking, diet, food supplement intake, general hunger levels, health state, past diagnoses, antibiotic intake, and menstrual health), physical activity (exercise frequency, duration, and intensity), sleep (bedtime, wake-up time), sociodemographic variables (nationality, socioeconomic status, job status, and household description), and requested to provide self-measured anthropometric measurements (waist and hip circumference, height, and weight). In the tracking days phase, participants had to fill out a short form each evening to validate their adherence to protocol and document medication intake. Cohort C participants had to answer additional questions on menstrual blood, cervical mucus and provide self-reported temperature measurements on a daily basis.

Dietary Intake: Participants of the "Food & You" study were asked to log any dietary intake in real time on the MyFoodRepo app (MFR) using one the following options: taking pictures of the food/drink, scanning the product's barcode (if available), or describing the food item with text. A logged entry is defined as a "dish", and can contain multiple food items. For example, tuna, steamed potatoes, and green beans are all single food items, and together compose a dish. The pictures were automatically segmented and classified by an image recognition algorithm[12]. Portions, segmentations, and food classes were subsequently verified or edited by a team of trained annotators. The MyFoodRepo app also allows annotators to communicate with the participant for clarifications about the food, and participants were able to leave comments through the app. This process ensured that every single dish in the nutritional data was reviewed by a member of the study team. Each food item was linked to a nutritional value database containing 2'129 items built on the Swiss Food Composition Database[13], MenuCH data[14], and Ciqual[15]. When food intake was logged through barcode scanning, nutritional values of the food items

were fetched from the Open FoodRepo database API[16]. Manual entries were matched to

food items by the annotators.

As the aforementioned nutritional values data sources did not provide standard portion

sizes, these were manually extracted from the portion list of the WHO MONICA study[17],

and the Mean Single Unit Weights of Fruit and Vegetables report[18] by the German Federal

Office of Consumer Protection and Food Safety. When a standard portion was not

available for a particular food item, we assigned the standard portion of a similar food

item.

Each dish logged through the MyFoodRepo app carries a timestamp, which enables

dietary analysis at a high temporal resolution. Food items were classified into categories

based on the menuCH study[14] (Chatelan, Marques-Vidal, et al. 2017). Barcoded food items

from the Open FoodRepo database were categorized based on the food product

description. When such a description was not available, we assigned the category

extracted from the Open Food Facts database[2]. While "Food & You" is the first large

cohort to use MyFoodRepo, the app annotation quality had previously been validated[19].

Glycemia: Glycemic data was collected by using the Flash Glucose Monitor Freestyle

Libre (Abbott Diabetes Care). The system, which has been validated in numerous

studies,[20–22] consists of a disposable sensor applied to the back of a participants' upper

arm, and a reader device or a smartphone app allowing to collect data from the sensor

via NFC technology. It measures glycemia every 15 minutes via a subcutaneous filament

carrying enzyme glucose sensors[23]. To encourage high adherence to protocol, we chose

a non-blinded glucose monitoring system to allow the participants to see their glycemia

in real time. Participants self-applied the sensor at home following explanations provided

in writing and video. Cohort B participants wore a single sensor for 14 days, whereas

cohort C participants wore two sensors consecutively for a period of 28 days to cover the

length of a typical menstrual cycle. Notably, when participants scan the sensor, the data

from the previous eight hours is collected. Thus, unless participants scan the sensor at

least every 8 hours, some data may remain unretrievable.

---

[2] https://world.openfoodfacts.org/

Gut Microbiota: Participants were requested to collect a stool sample following detailed written and video instructions. They could collect and ship their sample anytime during the tracking days phase. Samples were collected with stool nucleic acid collection and preservation tubes from Norgen Biotek, stored at room temperature and shipped in batches of 100 to 192 samples to Microsynth AG (Balgach, Switzerland) for sequencing and bioinformatics analysis. V4 region of the bacterial 16S rRNA gene was sequenced via creation of two-step Nextera PCR libraries using the primer pair 515F (NNNNNGTGYCAGCMGCCGCGGTAA) and 806R (NNNNNGGACTACNVGGGTWTCTAAT). The primers use 5 bases at their 5′ end to increase diversity of the bases during the first five sequencing cycles. Subsequently, the Illumina MiSeq platform and a v2 500 cycles kit were used to sequence the PCR libraries. The produced paired-end reads which passed Illumina's chastity filter were subject to de-multiplexing and trimming of Illumina adaptor residuals using Illumina's real time analysis software included in the MiSeq reporter software v2.6 (no further refinement or selection). The quality of the reads was checked with the software FastQC version 0.11.8. The locus specific V4 primers were trimmed from the sequencing reads with the software cutadapt v2.8. Paired-end reads were discarded if the primer could not be trimmed. Trimmed forward and reverse reads of each paired-end read were merged to in-silico reform the sequenced molecule considering a minimum overlap of 15 bases using the software USEARCH version 11.0.667. Merged sequences were then quality filtered allowing a maximum of one expected error per merged read. Reads that contained ambiguous bases or were considered outliers regarding the amplicon size distribution were also discarded. Samples that resulted in less than 5000 merged reads were discarded, to not distort the statistical analysis. The remaining reads were denoised using the UNOISE algorithm[24] implemented in USEARCH to form amplicon sequence variants (ASVs) discarding singletons and chimeras in the process. The resulting ASV abundance table was then filtered for possible bleed-in contaminations using the UNCROSS[25] algorithm, and abundances were adjusted for 16S copy numbers using the UNBIAS[26] algorithm. ASVs were compared against the reference sequences of the RDP 16S database, and taxonomies were predicted considering a minimum confidence threshold of 0.5 using the SINTAX algorithm[27] implemented in USEARCH.

Physical Activity and Sleep: Study participant's physical activity (PA) and sleep data were collected using one of two methods: objectively via Apple Health, Google Fit, or smart-watches, or subjectively, i.e. self-reported on the study website via the morning and/or evening questionnaire. The different formats of the objective PA and sleep data were harmonized and stored in a single database, and comprised daily step count, daily calories burned, bedtime, and wake-up time. In addition, for PA measured with smart devices, the type of physical activity, the start and end times, the amount of burned calories, the average heart rate, and the maximum heart rate were collected. Participants self-reporting their sleep and PA on the website had to report the times at which they fell asleep and woke up, if they did any physical activity, and if so, when they started and finished, as well as the perceived intensity of their effort.

STATISTICAL ANALYSIS

Data Preproccessing: Sociodemographic questionnaires variables and anthropometric measurements were re-coded as follows; Monthly household income was classified in 4 categories: <6000, 6000 to 8999, 9000 to 12999 and >=13000 Swiss Francs (CHF). Note that the median income in Switzerland was 6,665 CHF per month in 2020. Household type categories were defined as either "alone" for single-person households, "couple w. children", "couple w.out children", and "other". Age at study start was transformed into the following categories; 18 to 34, 35 to 49, 50 to 64, and older than 65 years. Citizenship was categorized as "Swiss", "Binational", or "Foreigner". Education level was categorized as "low" (mandatory, primary school), "intermediate" (high school and professional diploma) or "high" (university). Smoking status was categorized as "non smoker" (never smoked), "former smoker" or "current smoker" (occasionally or daily). The self-rated health questions were re-coded as binary variables "Not good to average" (not good, average, somewhat good) or "Good to very good" (good, very good). Residential addresses were geocoded, and municipality number (Spatial Development ARE) was extracted for each address. The municipality number was then joined with a urban/rural municipality typology and linguistic region dataset from the Swiss Federal Statistical Office so that each address was categorized as "urban" or "rural," whereas linguistic region was categorized as "german" or "latin," the latter encompassing french, italian, and romansh-speaking regions.

Missing values in income and education (<0.2% and less than 8%, respectively) were

imputed via multiple imputations with chained equations using the mice package[28]. Self-

reported general physical activity levels were assessed based on the weekly frequency

and average duration in minutes. The data was then coded into "active" and "inactive"

based on the WHO cutoff of 75 min per week[29]. Body-mass index (BMI) was calculated

based on self-reported height and weight and classified into "Underweight," "Normal,"

"Overweight," "Obese" based on the WHO cutoffs for BMI. For food intake, we removed

days for which energy intake was below 1'000 kcal and aggregated by study subject to

calculate the individual weighted mean to account for day-of-week and seasonal

variations of intake over the observation period.

To assess the extent of glycemic excursions for each participant, we calculated the

proportion of readings below, in, and above the target range based on the cut-off values

of 3.9 and 10 mmol/L[30]. We also computed the mean amplitude of glycemic excursions

(MAGE) for each subject using the R package gluvarpro (4.0).

Completion rate and study population analysis: Study completion rate was defined as

the ratio of the number of participants having completed the study over the total number

of participants enrolled in the study. The completion rate was then calculated for each

cohort (B and C) and each of the following strata: gender, age group, BMI, phone type,

and dietary restriction.

For the study population analysis, we only included the participants who have completed

the study, and calculated their proportion in each cohort for the following strata: gender,

household income, household type, age group, citizenship, education, smoking status,

health status, urbanity, linguistic region, physical activity, and BMI.

Analysis of adherence to study protocol was conducted by cohort given that their

tracking phase duration differed. For each participant, we reported the number of days

with (i) CGM measurements, (ii) reported food (distinguishing between days with a total

intake above 1000 kcal or days with any food logged), (iii) sleep data, (iv) PA data, and (v)

questionnaire data. We also reported the number of standardized breakfasts taken by participants, of glucose oral tolerance tests performed, and stool samples collected.

Data quality analysis: Given the multimodal nature and multi-dimensionality of the collected data, assessment of data quality may take several forms. Here we performed a series of analyses to evaluate if each type of collected data followed expected patterns.

Specifically, we performed three analyses assessing the food data quality with respect to timing, adherence, and composition. First, we assessed the distributions of food intake times by calculating the count of the logged food intakes by day of the week and hour of the day. Second, we hypothesized that most CGM peaks should be preceded by a food intake. We thus conducted an experiment using CGM data as ground truth. For each CGM peak (*i.e.*, a local maximum above the 90th percentile of the participant's CGM distribution) we reported whether a food intake of at least 50 kcal had been logged within the 90 min interval preceding the peak, or within the 90 min preceding the peak minus 2 hours (control case). We then reported, for each participant, the proportions of CGM matching a food intake in both experiment and control situations. To test our hypothesis that the proportion of peaks with logged food intake within 90 minutes of the peak would be higher than the same proportion for the control time-window, we used a non-parametric Wilcoxon rank sum test. Third, to assess the quality of the composition of reported food intake, we compared food intake data obtained in the "Food & You" cohort to data reported by the Swiss nutritional survey menuCH which was obtained from a demographically representative sample. We considered amounts of energy, meat, dairy, water intake as well as the proportion of the study population that consumed more than five portions of fruits and vegetables a day. For each of the following strata: sex, age group and linguistic region, we calculated the weighted means of the intake to account for weekday variations.

For sleep, the data were aggregated at the participant level to calculate the weekday and weekend individual mean values for bedtime, wake-up time, and sleep duration. Sleep durations of less than 4 hours or more than 12 hours were excluded.

For gut microbiota data comparison, we used relative abundances for the microbiome
samples of other studies available from the R package CuratedMetagenomicsData
(version 3.7). These samples were selected based on the filtration criteria that all samples
were of gut origin and from healthy adults. Four studies with most of such samples were
selected for comparison with the "Food & You" microbiome[7,31,3233]. Relative abundances
from these studies and "Food & You" were aggregated at the genus taxonomic level. Bray-
Curtis distances were calculated between the samples using the vegdist function of the
vegan package in R, which were then transformed into two-dimensional principal
coordinates using the pcoa tool of ape package in R.
358
359

360 # Results

361
362 **STUDY COMPLETION RATE**
363
Overall study completion rate - defined as the ratio of the number of participants having
completed the study over the total number of participants enrolled in the study - was
relatively stable over the years, with 69.5%, 56.4%, 65.7%, and 57.3% in 2019, 2020, 2021,
and 2022, respectively. Detailed completion rates are shown in Table 1. Within the
investigated groups - gender, BMI, age, phone type, and diet - we did not see any major
differences, with the exception of diet. Participants who reported dietary restrictions had
higher completion rates (82.5% and 81.5% in cohorts B and C, respectively) than their
counterparts without restrictions (62.7% in cohort B and 56.7% in cohort C).
372
373 **COHORT CHARACTERISTICS**
374
1,014 study subjects have completed the "Food & You" study, of which 870 in cohort B and
144 in cohort C (Table 2). In cohort B, both sexes were equally distributed. The proportion
of healthy and educated subjects was found to be higher in "Food & You" as compared to
the Swiss population. A direct comparison is difficult, because representative statistics
for the Swiss subpopulation with study inclusion and exclusion criteria applied are not

available. For example, in cohort B, the proportions of overweight and obese cohort
participants was 24% and 6%, while these proportions are slightly higher in the general
Swiss population (31% and 11% respectively). However, the combination of diabetes as an
exclusion criterion and the well-known association between diabetes and overweight /
obesity[34] may explain the difference. Because representative data from the Swiss
population filtered by our exclusion and inclusion criteria is not available, we did not test
whether the observed differences could be due to chance.. Some of the large differences
are unlikely to be explained by exclusion and inclusion criteria alone. Most strikingly, the
"Food & You" study population is much more highly educated, with almost three quarters
of participants having a university degree (compared to less than one third in
Switzerland). In addition, the 18-34 and 45- to 49 age classes are substantially over-
represented, while the elderly (ages 65 and over) are severly underrepresented (less than
3% of participants).
393
394   **ADHERENCE**
395
Of the participants who completed the study, adherence to protocol was generally high,
with a large majority of participants able to collect the requested amount of data for most
modalities. In cohort B, 93.9% provided at least 13 days of food tracking data with daily
energy intake above 1000 kcal, whereas this figure was slightly lower (84.7%) for cohort
C participants, albeit for at least 27 days. Cohort C participants were asked to provide
self-reported mucus quality and temperature measurements on a daily basis. 97% and
76% reported bleeding data on any day, and during the last week, respectively. The 1,014
participants who completed the study logged a total of 297,626 dishes amounting to 43.6
million kcalk. Participants mostly logged their food intake by taking pictures (76.1%),
whereas a smaller proportion of entries were logged by barcode scanning (13.3%) or
manually (10.6%). Sleep data was available for 64.8% of the participants who completed
the study. In total, participants ate 6,944 standardized breakfasts including 2,158 glucose
drinks.
409
In total, 6'460 subjective (i.e. self-reported) physical activities were reported. A large
proportion (33% in cohort B, 42.4% in cohort C) of users in both cohorts did not report
any activity at all, but the proportion that did report activities was higher in cohort C

(Figure 3). Around 4/5 of the participants who provided physical activity data reported
only subjective activity via website questionnaire. In terms of objective activities collected
through activity trackers such as smartwatches, the most reported activity types (66.5%)
were walking, running, and cycling.
417
418   In total, 997 participants who completed the study provided a stool sample for microbiota
419   composition quantification. The distribution of relative abundances at phylum and
420   species levels are depicted in supplementary figures 2 and 3.
421
422   **DATA QUALITY**
423
Breakfasts were observed from 05:00 onwards, whereas the hourly distribution of this
meal was shifted later during the weekend as compared to weekdays (Figure 4a). Logged
lunches were found to be consistent with work schedules during the week, with a peak
at noon, whereas the number of entries at those hours was much lower during weekends.
The number of entries after 20:00 tended to be higher Fridays and Saturdays, likely
reflecting social dinners. In general, and as expected, a shift in the later hours was
observed for all meals taken during the weekend as compared to weekdays (difference in
average meal times of 33 minutes, Wilcoxon rank sum test p-value < 0.001). The exception
to this pattern was Sunday dinners: the timing of those was more similar to Monday
dinners' than to Saturday dinners'.
434
The glucose data displayed a similar pattern as the food data with a clear shift between
weekdays and weekend, reflecting the later wake-up and eating times (Figure 4b, maximal
cross-correlation between weekend and weekdays mean glucose levels is found with a
30-minute shift). Consistently, wake up and bedtimes were also shifted toward later
hours during weekends. Participants slept 7.9 hours a night on average (SD: 1.19, Figure
4f). On average, participants slept 16 minutes longer on weekends than on weekdays (n =
591, paired t-test p-value < 0.001). On average, participants' bedtime was 23:41 (SD:
1.37,Figure 4e), 23:37 on weekdays, and 23:51 on weekends. Participants woke up on
average at 07:35 (SD: 1.43, Figure 4e), and woke up significantly later on weekends, at 07:57
vs 07:27 (average difference of 30 minutes, paired t-test p-value < 0.001).
445

446   The proportion of glucose peaks matching a food item was significantly higher (median
447   proportion: 68%) within the experimental group compared to the control group (median
448   proportion: 27%, one-sided Wilcoxon rank sum test p-value < 0.001), indicating a good
449   agreement between the food intake and CGM data. However, as we have no objective
450   ground truth data on nutrition, we do not know what the expected proportions would be.
451
452   In general, participants had a healthy glycemic profile with mean amplitude of glucose
453   excursion (MAGE) values of 1.41 mmol/L and 1.21 mmol/L for cohorts B and C, while the
454   proportion of readings above 10 mmol/L (hyperglycemia) was below 0.3%.
455
456   **COMPARISON WITH OTHER STUDIES**
457
For participants who completed the study, daily mean energy, meat, dairy and water
intake were 2,205.1 kcal, 92 g, 124.9 g, and 964.9 ml, respectively. The proportion of
participants more than 5 portions of fruits and vegetables per day was 6.71%. These values
were in good agreement with the mean values reported by the Swiss nutritional survey
"menuCH" were obtained from a demographically representative sample. The only
exception was for water, for which intake was much lower in menuCH than in our study.
Means and standard deviations for energy, meat, dairy, water intake as well as proportion
of participants eating more than 5 portions of fruits and vegetables per day are reported
in supplementary table 3 for the whole population and for various strata (sex, language,
age group, and combination of sex and age group).
468
The PCoA plot of the gut microbiota profile (Figure 5) shows the Bray-Curtis distance
dissimilarities of gut microbial community samples found in healthy individuals of "Food
& You" and four other gut-related studies. To allow for comparability, bacterial
taxonomies from these microbiomes were aggregated at the genus level. The plot shows
that most participants from the "Food & You" cohort and the LifeLinesDeep study tend
to form their own cluster distinct from other studies.

Discussion

The "Food & You" study is a fully digital nutrition cohort collecting diverse multimodal data remotely, without any physical contact between the participants and any member of the study team. This study has gathered a wealth of nutritional intake data, blood glucose measurements, survey responses, and gut microbiota samples from 1,014 participants. Here, we described the study protocol, reported on study completion rates, and described the collected data, focusing on assessing their quality and reliability.

The overall completion rate was high, with over 60% of enrolled participants completing the study. In comparison with other digital health studies[35], the retention rate for 14 and 28 days (cohort B and C, respectively) was rather high. Several factors may have contributed to this outcome. Perhaps most importantly, we approached the study's design from a participant's perspective, which led to the decision to develop a new food-tracking app (MyFoodRepo) from scratch, emphasizing ease of use. We also developed the study website and data collection system from scratch in order to directly integrate data collected from sensors or apps on the study website. For example, we combined nutritional data from the app and glucose data from the CGM system to generate interactive charts on the study website. Completion rates in younger and older age groups were comparable, indicating no major hurdle related to the use of digital tools for data collection for older subjects. Subjects with dietary restrictions were particularly committed to the study in both cohorts (completion rates over 80%), in line with previous findings from disease-based digital health studies showing that participants affected by the disease showed higher study retention[35].

With respect to adherence, data availability was high for most indicators in both cohorts, with the exception of physical activity and sleep. The limited provision of physical activity and sleep data by participants possibly indicates that these aspects, unlike diet, were not viewed as central to the study, despite encouragement to include this information. The minimal study duration was 14 days (cohort B), which may represent a significant time investment for participants. Besides participant's fatigue, other parameters may have impacted adherence. For example, technical issues such as faulty glucose sensors or omission to scan the sensor and temporary technical issues in the early versions of the

food tracking app or the "Food & You" website may at times have contributed to a lower
amount of data delivered. However, across both cohorts, data availability was high for
most indicators.

Because the "Food & You" study obtained high-resolution dietary data from a mobile app
(MyFoodRepo) that was originally created specifically for the study, care needed to be
taken to assess the quality of the nutritional data. As other studies have also begun to use
the app, the first independent validation study indicated strong data quality[19]. In addition,
the fact that every submitted data point on nutrition was reviewed by a study annotator
provides additional confidence in the data quality. We observed expected patterns
related to weekdays and weekends in terms of timing of food intake, glucose curves, and
wake-up and bedtimes at the population level. The high proportion of CGM peaks
matching a food intake further suggests that participants logged their food intake
appropriately and that the number of missing intakes is expected to be low. In addition,
overall intake, and intake of main food groups is in agreement with results reported from
a representative sample of the Swiss population.

"Food & You" is the first study that collected food intake via the AI-assisted nutrition
tracking app MyFoodRepo, generating an unparalleled dietary high temporal resolution
dataset of over 300'000 food dishes with a total of over 45 million kcal. We therefore have
a very precise picture of dietary patterns over at least two weeks from over 1000
participants. The data provided by the MyFoodRepo app is primarily based on time-
stamped pictures, and thus allows for objective temporal assessment of food intake. The
high study completion rates after participants completed the test of the app, combined
with the overall positive feedback from post-study surveys indicate that MyFoodRepo
was well accepted by participants.

Self-selection bias may have led to a non-representative study population with regard to
some particular sociodemographic variables. Based on the exclusion criteria targeting the
non pre-diabetic and diabetic population, we expected our study population to lead a
healthy lifestyle. Indeed, the high proportion of physically active people who self-report
a good health status among the participants points in that direction. Similarly, blood
glucose-related indicators such as MAGE and proportion of hypo- and hyperglycemic

excursion show no indication of subjects affected by pre-diabetes. Recruitment via social
media may have increased the proportion of younger participants with scientific affiinity.
Further, the digital nature of the project could have increased the selection of digitally
savvy participants. The lack of socio-demographic representation of the "Food & You"
study can be overcome through appropriate weighting of the imbalanced strata as
conducted in other studies[36]. This adjustment procedure will be crucial for further
publications, since it has been shown that socio-demographic factors greatly influence
dietary factors[37]. Finally, by excluding pre-diabetic people from the study, we may have
filtered out participants with high BMI, since the former is known to be associated with
the latter. A Swiss study[38] on self-reported anthropometric measurements has reported
that BMI measurements based on self-reported height and weight are underestimated by
a factor of 1.6. The true distribution of BMI observed in the "Food & You" project may
therefore be shifted towards higher BMI ranges.

The use of a non-blinded CGM system may be considered a limitation of this study as
participants may have adjusted their dietary habits with the intention to control their
glucose levels. The decision to use a non-blinded CGM system was motivated by two
factors. First, we believed that participants would remain more engaged than with a
blinded system. Second, as self-tracking health sensors become more commonly
available, future personalized nutrition systems deployed at scale must be designed in
the context of full data visibility to participants. Whether such potential self-adjustments
in digital cohorts are limited to the short term and initial use is an important question for
further research.

Microbiome samples in "Food & You" were sequenced using 16S rRNA, whereas the
samples from the other studies to which the microbiome data was compared to used
shotgun sequencing. The distinct clustering of "Food & You" and LifeLinesDeep samples,
originating from Switzerland and the Netherlands respectively, suggests a potential
geographical influence. These geographical differences are likely contributing factors to
the observed variations in gut microbiota between the two cohorts. More detailed
analyses of the microbiome and its associations with other data collected in the study are
ongoing.

Taken together, our results show that collecting a large amount of high-quality data with
high study protocol adherence is feasible in the framework of a digital nutrition cohort,
opening the path toward large-scale and detailed personalized nutrition studies.

576 Acknowledgments

We are deeply grateful to the participants of the "Food & You" study. This work was
supported by the Kristian Gerhard Jebsen Foundation, the Seerave Foundation, and the
Fondation Leenaards. The funders had no role in the design or execution of this study
and will have no role in the analyses, interpretation of the data, or decision to submit
results. We thank Stéphanie Bonnewyn, Sandra Rüegsegger-Tanner, and Bennan Tong
for help with the annotations, as well as Dr Jardena Puder for her clinical guidance. We
further thank everyone who helped with various parts of the project, especially Chloé
Greub, Leila Munaretto, and Jil Zanolin.

585 Author Contributions

Study Design: CA, MS

Technological Development: AB, SC, NF, GH, YJ, DK, PdV

Nutritional Analysis: CA, GR-L, RMS, EU-B

Project Management: CA, SR, TS, HH, MS

Data analysis: HH, VB, DK, RS, LS, MS

Writing: HH, RS, LS, MS

597 Competing Interests

598 The authors declare that there are no competing interests.

599 # Data Availability

600   The data to reproduce the results presented will be made available in a public repository
601   upon publication. We're happy to share the repository with reviewers during the review
602   process.

Figures

Figure 1

Data collection. a) Schematic illustrating the data collection process. (Left) Participants track the study variables *in situ* (from home, work, etc.). (Center) Data or samples are collected via web platforms and apps, or shipped to the lab by mail. (Right) Data is processed in the Food & You database. b and c) Example of data collected by one participant over 5 days. Top panel shows blood glucose levels (orange line), physical activity (turquoise spikes), and sleep (translucent turquoise rectangles). Bottom panel shows time and micronutrient composition (colors) of reported food intake. Like in the top panels, translucent turquoise rectangles show when the participant is asleep.

619 **Figure 2**

623 Study phases with participants per phase, and exit numbers.

Figure 3

Distribution of collected data.

(a–d) and (f–i): Distribution of the total number of days (x-axis) with collected data by participants (y-axis) in cohort B (a–d) and cohort C (f–i) for each of the study data modality. Light gray vertical bar indicates the duration of the study for both cohorts (i.e., the number of days for which participants were instructed to report data). Top panels (a,f) show that distribution for the blood glucose data (CGM: continuous glucose monitoring). Darker shades show the distribution in the case where days are included if there is at least 1 data point collected that day, while lighter shades show the distribution in the case that days are included only if there are data points for at least 90% of the day. 2nd row panels (b, g) show the distribution for the food intake data. In a similar fashion to the top panels, darker shades show the distribution for days with any food intake, while lighter shades only include days during which total caloric intake was above 1000 kcal.

3rd row panels (c, h) show the distribution for the sleep and physical activity data. 4th

row panels (d, i) show the distribution for daily questionnaires. Days were included if

participants filled the morning (lighter shade) or evening (darker shade) surveys. 5th row

panels (e, j) show the number of glucose drink (left), standardized breakfast (center), and

stool samples (right) reported or sent by participants. Cohort B (resp. C) participants were

instructed to eat two (four) glucose drinks and 6 (12) standardized breakfasts. For panel

(i) and (h), note that participants in Cohort C were allowed to keep answering survey for

150 days after start.

Figure 4

Temporal patterns of measured or self-reported data

a. Total number across all participants of logged dishes (any food or drink intake) per hour (x-axis) and weekday.

b. Mean (solid line) and 50% interquartile range (shaded areas) of the blood glucose levels measured during weekdays (black) or weekends (blue).

c. Schematic illustrating the experiment to assess if glucose peaks are more likely to be directly preceded by food intake. The time-window of interest (*i.e.*, in which food intake is expected) is displayed in purple, while the control time-window is displayed in gray.

d. Distribution of the percentage of glucose peaks directly (purple) or distantly (control - gray) preceded by food intake. Percentages are computed per participant: for each participant, glucose peaks are identified, food intake in the two time-window is reported as a binary variable, and percentages are computed as the fraction of time-windows with reported food intake.

e. Distribution of the times at which participants woke up (upper panel) or fell asleep (lower panel) during weekdays (gray) or weekend (blue). These times are either

670   self-reported by participants when filling the morning questionnaires or obtained

671   from participants' connected devices (smartwatches) data.

672   f.   Distribution of average sleep durations (x-axis, in hours) during weekdays (gray)

673   or weekends (blue).

Figure 5

Comparison with other studies

(Left panels): Comparison with the menuCH results. Distribution of the daily intake in energy (top row), meat (2nd row), dairy (third row), and water (bottom row). In each panel, the colored violins and dots represent the distribution and mean of the Food & You data while the smaller black dot is the mean for the corresponding population stratum reported by the menuCH study. (Top right panel): Proportion of participants that, on average, eat more than 5 portions of fruits and vegetables daily. For menuCH, the averages are over 2 days of data collection. For "Food & You", the average is over 14 (cohort B) or 28 (cohort C) days of data collection. Bar heights show the proportion of participants, black whiskers show the 95% CI. (Bottom right panel): Microbiota composition of F&Y participants compared to that of other cohorts. Each dot is a sample, and samples are colored by cohort. The coordinates of each sample are the first two coordinates resulting from a principal coordinate analysis on the Bray-Curtis distances between samples (Methods).

Tables

706 **Table 1**

		Cohort B	Cohort C
Age Group	18–34	63.27 (348)	61.76 (84)
	35–49	64.23 (316)	59.18 (58)
	50–65	60.4 (183)	50.0 (2)
	65+	57.5 (23)	0.0 (0)
BMI	Normal	63.64 (574)	62.15 (110)
	Underweight	61.54 (16)	63.64 (7)
	Overweight	63.4 (220)	47.62 (14)
	Obese	57.14 (60)	58.33 (7)
Gender	female	63.24 (425)	59.5 (144)
	male	61.38 (445)	0.0 (0)
Dietary Restriction	No	62.73 (771)	56.74 (122)
	Yes	82.5 (99)	81.48 (22)
Phone Type	Android	66.44 (396)	56.14 (64)
	iPhone	59.18 (474)	62.5 (80)

710 **Cohort completion rates**. For each section, the table shows percentages per section and
711 cohort, and absolute numbers in parentheses. Percentages are calculated as the number
712 of participants having completed the study over the number of participants enrolled in
713 the study.

715   **Table 2**

		Cohort B (n)	Cohort C (n)
Gender	Female	48.85 (425)	100 (144)
	Male	51.15 (445)	0 (0)
Household Income	<6000	17.82 (155)	22.92 (33)
	6000-8999	20.11 (175)	24.31 (35)
	9000-12999	29.31 (255)	28.47 (41)
	>=13000	32.76 (285)	24.31 (35)
Household Type	Alone	18.28 (159)	21.53 (31)
	Couple w. Children	30.69 (267)	33.33 (48)
	Couple w.out Children	35.06 (305)	29.17 (42)
	Other	15.98 (139)	15.97 (23)
Age Group	18-34	37.82 (329)	57.64 (83)
	35-49	35.86 (312)	40.97 (59)
	50-64	23.56 (205)	1.39 (2)
	65+	2.76 (24)	0 (0)
Citizenship	Swiss	52.64 (458)	45.14 (65)
	Binational	10.69 (93)	13.19 (19)
	Foreigner	36.67 (319)	41.67 (60)
Education	Low	1.72 (15)	2.78 (4)
	Intermediate	24.71 (215)	15.28 (22)
	High	73.56 (640)	81.94 (118)
Smoking	Non Smoker	48.05 (418)	51.39 (74)
	Former Smoker	42.18 (367)	34.03 (49)
	Smoker	9.77 (85)	14.58 (21)
Health Status	Not Good to Average	8.85 (77)	6.25 (9)
	Good to very Good	91.15 (793)	93.75 (135)
Urbanity	Urban	76.55 (666)	78.47 (113)
	Rural	23.45 (204)	21.53 (31)
Linguistic Region	German	47.93 (417)	53.47 (77)
	Latin	52.07 (453)	46.53 (67)
Physical Activity	Active	66.55 (579)	65.97 (95)
	Not active	33.45 (291)	34.03 (49)
BMI	Underweighted	1.84 (16)	4.86 (7)
	Normal	65.98 (574)	76.39 (110)
	Overweighted	25.29 (220)	13.89 (20)
	Obese	6.9 (60)	4.86 (7)

717   **Cohort characteristics.** For each section, the table shows percentages per section and
718   cohort, and absolute numbers in parentheses. Household income is monthly in Swiss
719   Francs (CHF), which is roughly in parity with USD.

References

1. Papamichou, D., Panagiotakos, D. B. & Itsiopoulos, C. Dietary patterns and
management of type 2 diabetes: A systematic review of randomised clinical trials.
Nutrition Metabolism Cardiovasc Dis **29**, 531–543 (2019).

2. Rodríguez-Monforte, M., Flores-Mateo, G. & Sánchez, E. Dietary patterns and CVD: a
systematic review and meta-analysis of observational studies. *Brit J Nutr* **114**, 1341–1359
(2015).

3. Grosso, G. *et al.* Mediterranean Diet and Cardiovascular Risk Factors: A Systematic
Review. *Crit Rev Food Sci* **54**, 593–610 (2014).

4. Key, T. J. et al. Diet, nutrition, and cancer risk: what do we know and what is the way
forward? *Bmj* **368**, m511 (2020).

5. Hood, L. & Friend, S. H. Predictive, personalized, preventive, participatory (P4) cancer
medicine. *Nat Rev Clin Oncol* **8**, 184–187 (2011).

6. Celis-Morales, C. et al. Effect of personalized nutrition on health-related behaviour
change: evidence from the Food4Me European randomized controlled trial. *Int J*
Epidemiol 578–588 (2016) doi:10.1093/ije/dyw186.

7. Zeevi, D. et al. Personalized Nutrition by Prediction of Glycemic Responses. *Cell* **163**,
1079–1094 (2015).

8. Mendes-Soares, H. et al. Assessment of a Personalized Approach to Predicting
Postprandial Glycemic Responses to Food Among Individuals Without Diabetes. *Jama*
Netw Open **2**, e188102 (2019).

9. Berry, S. E. et al. Human postprandial responses to food and potential for precision
nutrition. *Nat Med* **26**, 964–973 (2020).

10. Hoevenaars, F. P. M. et al. Evaluation of Food-Intake Behavior in a Healthy
Population: Personalized vs. One-Size-Fits-All. *Nutrients* **12**, 2819 (2020).

11. Collaboration, E. R. F. et al. Diabetes mellitus, fasting blood glucose concentration,
and risk of vascular disease: a collaborative meta-analysis of 102 prospective studies.
Lancet **375**, 2215–2222 (2010).

12. Mohanty, S. P. et al. The Food Recognition Benchmark: Using Deep Learning to
Recognize Food in Images. *Frontiers Nutrition* **9**, 875143 (2022).

13. Federal Food Safety and Veterinary Office. Swiss Food Composition Database.
https://valeursnutritives.ch/en.

14. Chatelan, A. et al. Lessons Learnt About Conducting a Multilingual Nutrition Survey
in Switzerland: Results from menuCH Pilot Survey. *Int J Vitam Nutr Res* **87**, 25–36 (2017).

15. French Agency for Food Environmental and Occupational Health and Safety. ANSES-
CIQUAL food composition table. https://ciqual.anses.fr/.

16. Lazzari, G., Jaquet, Y., Kebaili, D. J., Symul, L. & Salathe, M. FoodRepo: An Open Food
Repository of Barcoded Food Products. *Frontiers in nutrition* **5**, 57 (2018).

17. Dinauer, M., Winkler, A. & Winkler, G. Moncia Mengenliste. https://www.ble-
medienservice.de/0654/monica-mengenliste (1991).

18. Prüße, U., Hüther, L. & Hohgardt, K. Mittlere Gewichte einzelner Obst- und
Gemüseerzeugnisse.
https://www.bvl.bund.de/SharedDocs/Downloads/04_Pflanzenschutzmittel/rueckst
_gew_obst_gem%C3%BCde_pdf.pdf?__blob=publicationFile (2002).
19. Zuppinger, C. et al. Performance of the Digital Dietary Assessment Tool
MyFoodRepo. Nutrients 14, 635 (2022).
20. Ólafsdóttir, A. F. et al. A Clinical Trial of the Accuracy and Treatment Experience of
the Flash Glucose Monitor FreeStyle Libre in Adults with Type 1 Diabetes. *Diabetes*
Technol The **19**, 164–172 (**20**17).
21. Scott, E. M., Bilous, R. W. & Kautzky-Willer, A. Accuracy, User Acceptability, and
Safety Evaluation for the FreeStyle Libre Flash Glucose Monitoring System When Used
by Pregnant Women with Diabetes. Diabetes Technol The 20, 180–188 (2018).
22. Tsoukas, M. et al. Accuracy of FreeStyle Libre in Adults with Type 1 Diabetes: The
Effect of Sensor Age. Diabetes Technol The 22, 203–207 (2020).
23. Blum, A. Freestyle Libre Glucose Monitoring System. *Clin Diabetes Publ Am Diabetes*
Assoc **36**, 203–204 (2018).
24. Edgar, R. C. & Flyvbjerg, H. Error filtering, pair assembly and error correction for
next-generation sequencing reads. *Bioinformatics* **31**, 3476–3482 (2015).
25. Edgar, R. C. Accuracy of microbial community diversity estimated by closed- and
open-reference OTUs. *Peerj* **5**, e3889 (2017).
26. Edgar, R. C. UNBIAS: An attempt to correct abundance bias in 16S sequencing, with
limited success. *Biorxiv* 124149 (2017) doi:10.1101/124149.
27. Edgar, R. C. SINTAX: a simple non-Bayesian taxonomy classifier for 16S and ITS
sequences. Biorxiv 074161 (2016) doi:10.1101/074161.
28. Buuren, S. V. & Groothuis-Oudshoorn, K. mice: Multivariate imputation by chained
equations in R. *Journal of statistical software* **45**, 1–67 (2011).
29. Bull, F. C. et al. World Health Organization 2020 guidelines on physical activity and
sedentary behaviour. *Brit J Sport Med* **54**, 1451–1462 (2020).
30. Battelino, T. et al. Clinical Targets for Continuous Glucose Monitoring Data
Interpretation: Recommendations From the International Consensus on Time in Range.
Diabetes Care **42**, 1593–1603 (2019).
31. Asnicar, F. et al. Microbiome connections with host metabolism and habitual diet
from 1,098 deeply phenotyped individuals. Nat Med 27, 321–332 (2021).
32. Zhernakova, A. et al. Population-based metagenomics analysis reveals markers for
gut microbiome composition and diversity. *Science* **352**, 565–569 (2016).
33. Mehta, R. S. et al. Stability of the human faecal microbiome in a cohort of adult men.
Nat Microbiol **3**, 347–355 (2018).
34. Abdullah, A., Peeters, A., Courten, M. de & Stoelwinder, J. The magnitude of
association between overweight and obesity and the risk of diabetes: A meta-analysis of
prospective cohort studies. *Diabetes Res Clin Pr* **89**, 309–319 (2010).
35. Pratap, A. et al. Indicators of retention in remote digital health studies: a cross-study
evaluation of 100,000 participants. *Npj Digital Medicine* **3**, 21 (2020).

36. Chatelan, A. et al. Major Differences in Diet across Three Linguistic Regions of
Switzerland: Results from the First National Nutrition Survey menuCH. Nutrients 9, 1163
(2017).
37. Krieger, J.-P. et al. Dietary Patterns and Their Sociodemographic and Lifestyle
Determinants in Switzerland: Results from the National Nutrition Survey menuCH.
Nutrients 11, 62 (2018).
38. Faeh, D., Marques-Vidal, P., Chiolero, A. & Bopp, M. Obesity in Switzerland: do
estimates depend on how body mass index has been assessed? *Swiss Med Wkly* **138**,
204–210 (2008).