Full title:

Association of neurotransmitter pathway polygenic risk with specific symptom profiles in psychosis

Tracy L. Warren*[1], Justin D. Tubbs, Ph.D.*[2,3,4], Tyler A. Lesh, Ph.D[5], Mylena B. Corona[1], Sarvenaz Pakzad[5], Marina Albuquerque[5], Praveena Singh[5], Vanessa Zarubin[5], Sarah Morse, M.S.[6], Pak Chung Sham, Ph.D[†2,7,8], Cameron S. Carter, M.D[†5], Alex S. Nord, Ph.D[†1]

* co-first authors

† co-senior authors

[1] UC Davis Center for Neuroscience

[2] Department of Psychiatry, The University of Hong Kong

[3] Psychiatric and Neurodevelopmental Genetics Unit, Center for Genomic Medicine, Massachusetts General Hospital

[4] Department of Psychiatry, Harvard Medical School

[5] UC Davis Imaging Research Center

[6] Washington University in St. Louis

[7] Centre for PanorOmic Sciences, The University of Hong Kong

[8] State Key Laboratory of Brain and Cognitive Sciences, The University of Hong Kong

Abstract:

Merging genetic risk, neurological phenotypes, and clinical presentation is a primary goal for psychiatry. Pursuing this goal, we tested association between phenotypes and overall and pathway-specific polygenic risk in patients with early-stage psychosis. Subjects included 206 demographically diverse cases with a psychotic disorder and 115 matched controls with comprehensive psychiatric and neurological phenotyping. DNA was

extracted from blood and genotyped. We calculated polygenic scores (PGSs) for schizophrenia (SZ) and bipolar

disorder (BP) using Psychiatric Genomics Consortium GWAS summary statistics. To dissect convergent

mechanisms of symptoms, we calculated pathway PGSs (pPGSs) for SZ risk affecting each of four major

neurotransmitter systems: glutamate, GABA, dopamine, and serotonin. Psychosis subjects had elevated SZ and

BP PGS versus controls; cases with SZ or BP diagnoses had stronger SZ or BP risk, respectively. There was no

significant association between individual symptom measures and overall PGS. However, neurotransmitter-

specific pPGSs were significantly associated with specific symptoms; most notably, increased glutamatergic pPGS

was associated with deficits in cognitive control and altered cortical activation during cognitive control task-

based fMRI. Finally, unbiased symptom-driven clustering identified three diagnostically mixed case groups with

distinct symptom profiles that separated on primary deficits of positive symptoms, negative symptoms, global

functioning, and cognitive control. These clusters had specific genetic risk profiles and differential response to

treatment, and outperformed diagnosis in predicting glutamate and GABA pPGS. Our findings suggest pathway-

based PGS analysis may be a powerful path forward for identifying convergent mechanisms driving psychotic

disorders and linking genetic risk with endophenotypes.

Introduction

Psychosis-spectrum disorders, including schizophrenia (SZ), schizoaffective disorder (SA), and bipolar disorder

(BP), affect 0.5-2.3% of people worldwide(1–5). Evidence suggests shared etiology for these disorders,

supported by family studies and genome-wide association studies (GWASs) showing high interheritability and

shared genomic risk(6–8). Given symptom heterogeneity across the psychosis spectrum, a deeper understanding

of the biology underlying specific symptoms may drive psychiatry towards improved patient outcomes using

stratified medicine.

One approach toward revealing shared neurobiology is identifying transdiagnostic endophenotypes. Recently,

the Bipolar & Schizophrenia Network for Intermediate Phenotypes (B-SNIP) Consortium identified three biotypes

of psychotic disorders using neuropsychiatric markers, agnostic to diagnoses(9), that were primarily characterized by differences in cognitive control and sensorimotor reactivity. As knowledge of biology underlying symptom heterogeneity expands, reclassifying psychosis patients using biologically grounded phenotypes may allow for more effective, targeted interventions.

Identification of causal and clinically informative genetic components of psychotic disorders has been aided by large-scale GWASs and estimation of overall genetic risk using polygenic scores (PGSs). While large cohorts were initially required to develop PGSs, once defined, research leveraging PGSs in cohorts with phenotyping that extends beyond diagnosis is revealing how genetic burden is associated with specific symptomatology. For example, recent efforts have found associations between SZ PGS and treatment response(10) as well as neurological and cognitive measures(11,12). Such uses of PGS enable study of genetic burden in small cohorts that have been characterized at a level not feasible at the scale required for GWASs focused on the discovery of new risk loci.

While PGS is a useful metric of genetic risk, it fails to assign burden into relevant biological pathways and elucidate mechanisms underlying symptoms. A novel approach to overcome this is pathway-specific PGS (pPGS), which partitions variants into those contained only in genes of a given function or pathway(13). By assigning burden into pathways, it is possible to test how specific processes contribute to phenotype. This has been used to associate miR-137 pPGS with SZ risk(14) and neurological markers(15,16). Another recent implementation identified SZ and BP subjects with risk in pathways targetable by available pharmaceuticals, showing the utility of pPGS in targeted medicine(17). Among pathways relevant to psychosis, altered glutamate(18–22), GABA(21–23), dopamine(24–27), and serotonin(27–29) signaling have been strongly associated with psychotic disorders and symptoms.

There is a major opportunity to use pPGS to identify pathway contributions to psychosis symptoms. Here, we investigated the relationship of overall and neurotransmitter-associated PGS to psychotic disorder presentation at the diagnostic and endophenotypic level. We estimated overall and pathway-level PGS for a diverse cohort

recruited from California following a first psychotic episode, and tested association between genetic burden, diagnosis, and clinical and neuroimaging endophenotypes. We found that overall and pathway PGSs were elevated in cases, pPGS was associated with specific endophenotypes, and unbiased clustering on phenotypes outperformed diagnosis in associations with pPGS and treatment response. Our results demonstrate the power of pPGS to link genetic and neurobiological underpinnings of psychotic disorders.

Methods

Study participants

Psychosis subjects were all outpatients within two years of their first psychotic episode. Subjects were selected from an ongoing psychosis research cohort, which includes 196 first-episode SZ-spectrum patients, 53 patients with first-episode BP with psychotic features, and 135 controls aged 12-38. After all quality control (QC), 119 SZ, 39 SA, 48 BP, and 115 controls were included for this study from an ongoing early psychosis research cohort. The study was approved by the University of California, Davis, Institutional Review Board and all subjects gave written consent and were paid for their participation.

Psychiatric and neuroimaging phenotyping

All participants were assessed using the Structured Clinical Interview for the DSM-IV-TR (SCID I/P)(30). Clinical interviews were conducted by clinicians with masters or doctoral degrees trained to high reliability (kappa > .70; range = .70-1.0). All patients were followed longitudinally and diagnoses were confirmed 6 months after ascertainment. Exclusion criteria for all groups included: Wechsler Abbreviated Scale of Intelligence (WASI) IQ score below 70, alcohol or drug dependence or abuse within 3 months before testing, positive urine toxicology screen for illicit drugs, prior head trauma worse than a Grade I concussion, or contraindication to MRI scanning. Control subjects were excluded for the following additional criteria: any lifetime diagnosis of an Axis I or Axis II disorder or any first-degree relatives with a psychotic disorder. Before testing, a detailed description of the study was provided and written informed consent obtained.

Subjects were evaluated on the Global Assessment of Functioning (GAF)(31), Global Social Functioning scale (GSF)(32), Global Role Functioning scale (GRF)(33), Young Mania Rating Scale (YMRS)(34), Scale for the Assessment of Positive Symptoms (SAPS)(35), Scale for the Assessment of Negative Symptoms (SANS)(35), and Brief Psychiatric Rating Scale (BPRS)(36). Reality distortion, poverty symptoms, and disorganization scores were defined from the BPRS, SAPS, and SANS(37). Treatment response was defined as > 20% decrease in BPRS from baseline(38).

The GSF and GRF were measured at multiple time points; for each scale, we computed an average of the highest and lowest values measured for each subject during the past year and proceeded with these values for subsequent analyses. Finally, we removed item #8 ("Content") from scores for the YMRS, as this question asks specifically about positive psychotic symptoms and can skew YMRS scores for subjects with schizophrenia. As such, scores on the YMRS used in these analyses more directly represent mania-specific symptoms. At baseline, all patients had BPRS scores >=5 to offer sufficient resolution to detect a 20% improvement in score at follow-up. For treatment response calculation, BPRS was rescaled to a lowest score of zero (i.e. score of 24=score of 0)(39).

Behavioral and neuroimaging methods are described previously(40). Consequently, we present these methods in a condensed form. The AX-Continuous Performance Task (AX-CPT)(41) was performed during fMRI. In short, the task requires participants to respond to a series of cue and probe letters and correctly identify the target pair ("AX" trials) while correctly rejecting other pairs. The frequency manipulation of trial types creates a prepotent tendency to make a target response when the "X" probe letter is presented. Consequently when a non-A cue is presented and followed by an X (i.e. "BX" trials) the participant must engage proactive control to both retain the goal, keep the incorrect cue in mind, and correctly reject the trial at the probe phase. Participants were excluded if performance did not meet a minimal threshold(42). The primary behavioral measure used for this study is d'-context, which represents a contrast of AX hits versus BX false alarms.

Functional Blood Oxygenation Level Dependent (BOLD) data were acquired using a 1.5T GE Signa and 3.0T Siemens TimTrio. Two regions of interest were defined *a priori* and comprised CueB versus CueA contrast, reflecting high versus low cognitive control-related activity. Specifically, bilateral dorsolateral prefrontal cortex (DLPFC) and bilateral superior parietal cortex (SPC) were defined as 5mm radius spheres based on coordinates from two independent datasets(43,44). All fMRI data were preprocessed using SPM8 (Wellcome Dept. of Imaging Neuroscience, London) and included slice timing correction, realignment, normalization to the Montreal Neurological Institute (MNI) template, and smoothing with an 8mm FWHM Gaussian kernel. Individual runs were excluded when framewise displacement measures of movement exceeded 0.45mm (calculated with https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/FSLMotionOutliers) and whole subjects were excluded if more than half of their data exceeded this threshold. All trial types were modeled (CueA, CueB, AX, AY, BX, BY) and correct responses were used to create first-level images.

Based on the exclusion criteria mentioned above, five controls, nine patients with schizophrenia, and two patients with bipolar disorder were excluded due to excess motion. Two controls, five patients with schizophrenia, and two patients with bipolar disorder were excluded for poor behavioral performance. Finally, three controls and six patients with schizophrenia were excluded for other reasons, including scanning artifacts, scanner failure, or button pad failure.

Genotyping, Quality Control, and Imputation

Subjects underwent blood draws using PAXgene blood DNA tubes, which were subsequently stored at -80 °C until DNA extraction. Tubes were thawed at 37 °C for 15 minutes before extracting DNA using the Qiagen QIAamp DNA Blood Mini Kit. The protocol was followed as written with minor modifications: volumes of blood, protease, lysis buffer, and ethanol were tripled prior to binding the DNA to the spin column; the final elution incubation was carried out at 50 °C for five minutes; and DNA was eluted in 100 µl of nuclease-free water. After DNA extraction, samples were cleaned on the Zymo Research Genomic DNA Clean & Concentrator-10 kit with minor modifications: an extra two-minute spin in a clean collection tube was added after completing the wash

steps; samples were incubated at 50 °C for five minutes at the elution step; and DNA was eluted in 100 μl of nuclease-free water. After cleaning, all samples were analyzed on a spectrophotometer and verified to have concentrations ≥ 50 ng/μl and 260/280 and 260/230 ratios ≥ 1.70.

DNA was genotyped using the Illumina (San Diego, California) Infinium PsychArray-24 Kit at the UC Davis DNA Technologies Core. Initial QC was performed using Illumina GenomeStudio following published guidance(45,46). Additional quality control was applied using PLINK. First, variants with greater than 5% missingness were removed. We confirmed that no individuals had more than 1% of SNPs missing, nor did we observe any mismatch between the genetic sex inferred by PLINK and subjects' self-reported sex. We removed variants which had significantly different missingness rates between cases and controls (p < 0.001) or variants which indicated significant deviation from Hardy-Weinberg equilibrium (p < 1e-6). No severe heterozygosity outliers were observed. A small number of pairs of individuals appeared to be related; in these cases one subject from each pair was randomly excluded from subsequent analyses. Following imputation, the QC filters outlined above were performed again. Additionally, variants with imputation quality INFO scores less than 0.7 or empirical INFO scores less than 0.8 were removed.

PLINK (v1.9) was used to calculate genetic principal components (PCs) in the 1000 Genomes(47) Phase 3 dataset, which were projected onto our sample. Samples were submitted to the Michigan Imputation Server for genotype imputation, using the full 1000 Genomes(47) Version 3 dataset as the reference panel. Following all QC, 7,608,150 SNPs were available across 338 unrelated subjects. Following additional review, 14 subjects with schizophreniform disorder and 2 subjects with schizotypal disorder that had initially been included were excluded from subsequent analyses due to insufficient numbers, such that the final number of subjects was 321 as described in "Study Participants."

We additionally checked for presence of known copy number variants (CNVs) of high penetrance for schizophrenia risk in our samples using iPsychCNV5 and PennCNV(48–50), using default parameters. CNVs called by both software were retained as consensus calls for further analysis. This resulted in a total of 333 CNVs across

all subjects. We cross-referenced these results with genome-wide significant CNV loci associated with SZ(51) and filtered for CNVs in our sample that overlapped with at least 50% of one of these previously identified SZ risk CNV loci. One subject with SZ appeared to have a previously described SZ risk CNV: 15q11.2 deletion. As the role of deletion at 15q11.2 is currently of uncertain significance in schizophrenia risk(52,53), we retained this individual for all analyses.

Polygenic Score Calculation

Using Psychiatric Genomics Consortium GWAS summary statistics for SZ (2021)(54) and BP (2022)(55), we employed PRS-CS(56) to calculate SZ and BP PGS, with the phi parameter set to 0.01 as recommended without a validation sample. PRS-CS applies shrinkage to optimize PGS prediction and may not be applicable when restricting PGS to genes from specific pathways. Therefore, we used the PRSet function from PRSice(57) to calculate pPGS for the four neurotransmitter pathways using SZ GWAS summary statistics(54), as our subjects were weighted toward SZ diagnosis and showed high correlation between SZ and BP PGS (Fig. 1). As recommended by the authors of PRSet, in order to avoid eliminating SNPs in certain genes, no p-value filter was applied when calculating pPGS. However, in order to choose an optimal R^2 cutoff for clumping, we tested five potential values (0.1, 0.3, 0.5, 0.7, 0.9). In the subset of European ancestry samples, we found that using all SNPs as input, a clumping R^2 of 0.7 maximized prediction of case-control status of the overall SZ PGS. Thus, this value was used when calculating pPGS for each of the four pathways. Otherwise, default parameters of PRSet were retained, with the 1000 Genomes(47) European subset used as the LD reference panel.

To reduce bias from population stratification, we employed an ancestry-specific standardization process to model PGS in a mixed-ancestry sample. First, we assigned each subject to their nearest 1000 Genomes(47) population by minimizing Euclidean distance to population centroid in 20-dimensional PC space, as recommended by Prive et al.(58) Then, we calculated overall PGSs and pPGSs for all samples in the 1000 Genomes(47) dataset using the same SNPs and weights used in our target sample. Finally, we z-score normalized PGS for each subject relative to their matching 1000 Genomes(47) ancestry. Thus, PGSs are standardized within

ancestry-matched population such that a subject with CEU ancestry and a standardized PGS of 1 has a raw PGS one standard deviation above the average of the 1000 Genomes(47) CEU samples.

pPGS pathways

Pathways include genes relevant in glutamate, GABA, dopamine, and serotonin. Genes were sourced from KEGG(59), REACTOME(60), and AmiGO(61) by searching for pathways and ontologies that include these neurotransmitters or variations on them (e.g. "glutamatergic"). The complete gene list is in Table S1.

Unbiased clustering

Phenotypes were z-score normalized within cases. Subjects were clustered on all phenotypes. We used all subjects with complete phenotype data (n = 90) to calculate the optimal number of clusters using the NbClust R package(62). Estimates converged on three clusters. Subsequently, we used all subjects with greater than 50% of phenotype variables available (n = 167) for k-means clustering using the flipCluster R package(63).

Statistical analysis

All analyses were completed in R(64). One BP subject was excluded as an outlier from analyses based on Cook's distance near 1 in regression models. Correlation analyses and plots were produced using the psych(65) and corrplot(66) packages. Violin plots were produced using the vioplot package(67).

Nagelkerke's R^2 was calculated for variance explained in logistic regression models of disease status by PGS using the RMS R package(68). All phenotypes were tested against PGS and the following covariates in regression models: chromosomal sex, age, self-reported race, self-reported ethnicity, and the first four genetic PCs. Protocol/scanner type was controlled for in cognitive control analyses. Overall SZ PGS was included as a covariate for endophenotype analyses (Fig. 3). Models of continuous variable phenotypes against PGS are represented with partial regression plots.

For cluster versus diagnosis model comparison (Fig. 4E), we regressed PGSs separately against clusters, plus

standard covariates, and against diagnoses, plus covariates. In separate analyses (Fig. 4F), we regressed PGS

against both cluster and diagnosis in the same model, as well as covariates. BP and Cluster 3 were the reference

levels for, respectively, diagnosis and cluster.

Multiple testing corrections

Multiple testing corrections were applied for analyses of phenotype, including diagnosis and case status, by

pathway PGS (Figs. 2, 3). Given the correlated structure of phenotypes and of pPGS in our subjects, we used the

poolr R package(69) to calculate the effective number of statistical tests following Galwey (2009)(70). This

methodology identified effectively three (of initially four) independent pPGS variables and eight (of initially ten)

independent endophenotype variables. As differential diagnoses of SZ, SA, or BP were mutually exclusive, they

were not adjusted for correlated structure. Total tested dependent variables were multiplied by total tested

independent variables such that findings reported in Figure 2 were corrected for either three (case status by

three pPGS) or nine (three diagnoses by three pPGS) total tests, and findings reported in Figure 3 were corrected

for 24 total tests (eight endophenotypes by three pPGS) using the Benjamini-Hochberg(71) false discovery rates

reported in Results..

Permuted null pathway calculation (Fig. S2)

For each of the pathways in our main analyses, we constructed 10,000 matched null gene sets with an

equivalent number of genes, chosen such that the probability of a gene's inclusion is proportional to its length.

We then calculated pPGS for these 40,000 permuted null gene sets in the same way as described above, i.e.

using PRSet to calculate pPGS for our target sample and the 1000 genomes sample with weights from the most

recent SZ GWAS and standardizing the target sample pPGS within each ancestry relative to the matched-

ancestry 1000 genomes samples.

Results

Demographics

Subjects were demographically heterogenous and well matched across cases and controls (Table 1).

	Cases (N)	(%)	Controls (N)	(%)
Total	206	64%	115	36%
Diagnosis				
Schizophrenia	119	58%	-	-
Bipolar Disorder	48	23%	-	-
Schizoaffective Disorder	39	19%	-	-
Race				
African American/Black	28	14%	7	6%
American Indian/Alaskan Native	2	1%	0	0%
Asian	15	7%	27	23%
Caucasian/White	131	64%	61	53%
Native Hawaiian/Pacific Islander	3	1%	1	1%
Multiple	23	11%	17	15%
Unknown	4	2%	2	2%
Ethnicity				
Hispanic/Latino	50	24%	17	15%
Non-Hispanic/Latino	151	73%	97	84%
Unknown	5	2%	1	1%
Inferred 1000 Genomes superpopulation				
African (AFR)	58	28%	19	17%
American (AMR)	20	10%	8	7%
East Asian (EAS)	14	7%	26	23%
European (EUR)	111	54%	53	46%
South Asian (SAS)	3	1%	9	8%
Sex				
Male	158	77%	67	58%
Female	48	23%	48	42%
Mean age (s.d.)	19.7 (4.1)		19.7 (4.2)	

Table 1: Demographics of study subjects. Study subjects were racially and ethnically diverse across both cases and controls. Potential differences in demographics were considered when controlling for covariates in subsequent analyses. (s.d. = standard deviation)

Overall and pathway PGSs predict diagnostic status

We first tested whether overall PGS was associated with psychosis status. SZ PGS was associated with case

status (OR = 1.37 [CI: 1.15,1.63]; p = 3.7×10^{-4}) (Fig. 1A) and explained 5.2% of variance in status (p = 4.5×10^{-4}),

similar to previous reports(72–76) (Fig. S1). This association was stronger in SZ (OR = 1.50 [CI: 1.22,1.84]; p =

1.3×10^{-4}) and SA (OR = 1.59 [CI: 1.11,2.26]; p = 0.01) (Fig. 1B). While BP PGS was not significant in cases overall, it

was elevated in BP subjects (OR = 1.47 [CI: 1.11,1.93]; p < 0.01) (Figs.1A,B). SZ and BP PGS were moderately

correlated in cases (Fig. 1C). Thus, overall PGS captured both transdiagnostic and diagnosis-specific genetic risk

for psychotic disorders, consistent with a strong literature on partial genetic overlap between SZ, SA, and

BP(6,7,77–79).

Figure 1: SZ and BP PGS are associated with psychotic disorder diagnoses. (A) SZ PGS (top) is significantly associated with psychosis case status, while BP PGS (bottom) does not reach significance. (B) SZ PGS (top) PGS is associated with SZ and SA status versus controls, but not BP status. Conversely, BP PGS (bottom) is associated with BP status versus controls, but not SZ or SA status. For each (A) and (B), the top row represents a

comparison of SZ PGS across diagnostic groups, while the bottom row represents a comparison of BP PGS across diagnostic groups. (C) Pearson's correlations of SZ and BP PGS in psychosis cases show moderate-to-high correlations between PGS for all psychosis subjects, as well as for people in individual diagnostic groups. This correlation is somewhat lower for BP subjects.

Association of pPGS with phenotypes

To dissect genetic contributions of biological pathways to clinical and neurobiological phenotypes, we calculated pPGS for four neurotransmitter systems relevant to psychosis: glutamate(18–22), GABA(21–23), dopamine(24–27), and serotonin(27–29) (Fig. 2A, Table S1). These showed moderate intercorrelations and correlations with overall PGS (Fig. 2B).

We first tested association with psychosis status. Glutamate and GABA pPGS explained a substantial portion of variance in disease status (R^2 = 0.02-0.04, $p < 0.05$) (Fig. S2). Glutamate and GABA pPGS were significantly elevated in cases (both: OR = 1.3 [CI: 1.1-1.6]; p = 0.04) (Fig. 2C), specifically SZ (both: OR = 1.4 [CI: 1.1-1.8]; p = 0.01) (Fig. 2D); these findings passed multiple testing corrections at a false discovery rate (FDR) < 0.10. This implies disease status in our cohort is partially explained by glutamatergic and GABAergic genetic risk.

Figure 2: Pathway polygenic scores (pPGS) based on four main neurotransmitter systems. (A) KEGG(59), AmiGO(61), and REACTOME(60) were searched for genes associated with glutamatergic, GABAergic, dopaminergic, and serotonergic neurotransmission. (B) pPGS developed from these gene sets showed substantial correlation in psychosis subjects when regressing against the first four genetic principal components. All correlations were significant with p-value < 0.05 except for the correlations between serotonin pPGS and the overall SZ and BP PGS. (C) Glutamate and GABA pPGS were able to separate psychosis cases from controls and passed FDR < 0.10. (D) Glutamate and GABA pPGS much more strongly separated SZ cases from controls (FDR < 0.10) than either SA or BP subjects from controls.

<u>pPGSs are associated with endophenotypes in cases</u>

We next tested if PGS was associated with symptom variation across 10 phenotypes within subjects with a psychotic disorder (Fig. 3A). Overall PGSs were not significantly associated with measured phenotypes (Fig. S2). However, we found associations of pathway-specific risk to six phenotypes in social functioning, global functioning, cognitive control, and treatment response. While no endophenotype associations passed multiple testing corrections at FDR < 0.10, all but social functioning and treatment response passed FDR < 0.15.

Dopamine and GABA pPGS were associated with, respectively, poorer global (β = -1.18 [CI: -3.32,-0.29]; p = 0.02) and social (β = -0.26 [CI: -0.05,-0.01]; p = 0.04) functioning (Fig. 3B), though the social functioning association did not pass FDR < 0.15. Perhaps the most robust finding was a strong association between glutamate pPGS and cognitive control in cases (β = -0.16 [CI: -0.30,-0.02] to -0.34 [CI: -0.62,-0.06]; p = 0.01-0.02) (Fig. 3C). This implies that perturbations in glutamate may be a determinant of poor cognitive control in psychosis subjects, recapitulating a robust literature associating glutamate with cognitive control(80–83).

Of 63 subjects with available treatment response data, 35 were responders, consistent with reported efficacy rates(84). We hypothesized subjects with higher dopaminergic risk would be more likely to respond to treatment, as they might be enriched for the kinds of biological differences that dopamine-targeting antipsychotics can address. While power to identify an association was reduced due to subsetting of our cohort, treatment response showed an association with higher dopamine pPGS (OR = 1.88 [CI: 0.98,3.59]; p = 0.057) (Fig. 3D), though this did not pass FDR < 0.15. This adds onto a strong literature associating dopamine with antipsychotic efficacy, providing evidence that an accumulation of dopaminergic risk may be associated with treatment efficacy.

Figure 3: pPGSs identify mechanisms for endophenotypes of psychotic disorders. (A) Subjects were phenotyped on a range of psychological, clinical, and neurological measures. (B) Dopamine and GABA pPGS were associated with poorer global functioning in psychosis subjects when controlling for overall SZ PGS. Only the association with between dopamine and global functioning passed FDR < 0.15. (C) Glutamate pPGS showed strong and significant associations with poorer performance and reduced cortical activation in psychosis subjects during the AX-CPT cognitive control task when controlling for overall SZ PGS. These relationships passed FDR < 0.15. (D) Treatment response showed a slight association with increased dopamine pPGS, though this did not pass FDR < 0.15. (Resp. = responder to treatment, non-resp. = non-responder to treatment)

Unbiased phenotype clustering of psychosis cases and genetic burden

Biotype-level grouping of psychosis cases, such as by B-SNIP(9), may advance the use of biological data to better model psychopathology. We hypothesized a biotype-approach might yield stronger relationships with PGS compared to diagnosis. Toward this goal, we used k-means clustering to group endophenotype presentation among cases. We clustered subjects on z-score normalized phenotype data into three groups following cluster optimization using the NbClust R package(62).

Clusters showed distinct symptom profiles (Fig. 4A, Table S2). Cluster 1 was distinguished by high mania, disorganization, and reality distortion, and moderate impairments in cognitive control. Cluster 2 had deficits in

negative symptoms, showed poor outcomes in role and social functioning, and had the poorest measures of cognitive control. Cluster 3 had low pathology. Consistent with its high degree of positive symptoms, Cluster 1 showed enrichment for treatment responders relative to Clusters 2 and 3, though this did not reach significance (p = 0.057) (Fig. 4B). In contrast, treatment response showed no separation by diagnosis (p = 0.845). Clusters were demographically and diagnostically heterogenous (Fig. 4C, Table S2), though Cluster 3 captured most BP subjects. This may indicate our clusters better distinguished transdiagnostic variation in SZ and SA subjects, though the presence of BP subjects in Clusters 1 and 2 suggests there is merit to psychosis-spectrum endophenotypic stratification for these subjects as well.

We next tested associations between cluster and PGS (Fig. 4D, Table S2). SZ PGS was associated with all clusters (OR = 1.28 [CI: 1.01-1.62] to 1.94 [CI: 1.31-2.89]; p = 0.001-0.039). BP PGS was strongly associated with Cluster 3 (OR = 1.42 [CI: 1.11-1.81]; p = 0.006), consistent with its preponderance of BP subjects, though also with Cluster 1 (OR = 1.40 [CI: 1.01-1.96]; p = 0.046). Cluster 3 was not associated with any pPGS, suggesting genetic risk may lie outside of neurotransmitter-linked loci. In contrast, GABA pPGS was elevated in Clusters 1 and 2 (OR = 1.47 [CI: 1.09-1.98] to 1.59 [CI: 1.02-2.49]; p = 0.011-0.040) while glutamate pPGS was only elevated in Cluster 2 (OR = 1.71 [CI: 1.24-2.37]; p = 0.001). This may have implications for the symptom profiles of these clusters. Cluster 2 showed strong negative symptoms, which are thought to largely arise from glutamatergic dysfunction(85,86). Cluster 2 also showed deficits in social and role functioning, which are predicted by negative symptoms(87), and strong cognitive control deficits. pPGS findings suggest Cluster 2's symptoms may be more specific to GABAergic and glutamatergic risk, consistent with the hypothesis of an imbalance between these neurotransmitters in psychosis(22).

We next compared explanatory power of our clusters versus diagnoses to model genetic risk in our cohort. For this, we defined models regressing PGSs against either cluster or diagnosis. Per Bayesian information criteria and R^2 (Fig. 4E), models using clusters performed better than models using diagnosis to predict glutamatergic, GABAergic, and overall SZ PGS. In contrast, overall BP PGS and serotonin pPGS were somewhat better modeled by diagnosis, and there was no difference for dopamine pPGS. We next regressed PGS against both cluster and

diagnosis as covariates to test independence between diagnosis and cluster (Fig. 4F). Again, glutamate and GABA pPGS were significantly associated with Cluster 2 when including diagnosis in the model (β = 0.46 [CI: 0.05,0.88] to 0.49 [CI: 0.05,0.94]; both p = 0.03). Neither cluster nor diagnosis was independently associated with other PGSs. These results provide evidence for the biological validity of these clusters overall and within the context of molecular pathways. In particular, glutamate and GABA pPGS seem to be more closely associated with cluster than with diagnosis.

A Phenotypes by cluster

Cognitive control *Lower indicates more pathology.* Psychotic symptoms *Higher indicates more pathology.* Global functioning *Lower indicates more pathology.*

Z-score in all psychosis cases

d'-Context, Bilat. DLPFC β-val., Bilat. SPC β-val., Poverty symptoms, Reality distortion, Disorganization, YMRS, GRF, GSF, GAF

● Cluster 1 (n = 34) ● Cluster 2 (n = 72) ● Cluster 3 (n = 61)

B Treatment response separates better by cluster than by diagnosis

By cluster
X^2, p = 0.057

By diagnosis
X^2, p = 0.845

Resp. Non-resp.

C
Diagnosis by clust.
Clust. by diagnosis

SZ SA BP
C. 1 C. 2 C. 3

D PGS by cluster

SZ PGS: p < 0.01 v. control, p < 0.01 v. control, p = 0.04 v. control
Glut. PGS: p = 0.17 v. control, p < 0.01 v. control, p = 1.00 v. control
Dop. PGS: p = 0.39 v. control, p = 0.31 v. control, p = 0.73 v. control
BP PGS: p = 0.05 v. control, p = 0.46 v. control, p < 0.01 v. control
GABA PGS: p = 0.04 v. control, p = 0.01 v. control, p = 0.80 v. control
Ser. PGS: p = 0.85 v. control, p = 0.62 v. control, p = 0.27 v. control

Control (n = 115) Clust. 1 (n = 34) Clust. 2 (n = 72) Clust. 3 (n = 61)

E Model performance of PGS by cluster v. PGS by diagnosis

	BIC for PGS by:		R^2 for PGS by:	
	Cluster	Diagnosis	Cluster	Diagnosis
SZ PGS	**676**	678	**0.11**	0.09
BP PGS	670	**669**	0.16	0.16
Glut. pPGS	**566**	570	**0.12**	0.10
GABA pPGS	**588**	594	**0.12**	0.09
Dop. pPGS	554	554	0.17	0.17
Ser. pPGS	534	**532**	0.09	**0.10**

Better values indicated in **bold**.

F Significance of cluster v. diagnosis as covariates in regression model for PGS

-log(p)

p = 0.01
p = 0.05
p = 0.10

C1 (v. C3) C2 (v. C3) SZ (v. BP) SA (v. BP)

SZ PGS BP PGS Glut. pPGS GABA pPGS Dop. pPGS Ser. pPGS

Figure 4: Psychosis subjects separate into clusters based on symptom profiles. (A) Psychosis subjects separate into three clusters that are characterized by (Cluster 1) mania, disorganization, reality distortion, and moderate impairments in cognitive control; (Cluster 2) high poverty symptoms, poor social functioning, and strong deficits in cognitive control; and (Cluster 3) overall low levels of pathology. (B) Cluster 1 showed enrichment for treatment responders. In contrast, treatment efficacy did not separate by diagnosis. (C) Diagnoses are mixed across clusters. (D) SZ and GABA PGS are elevated in Clusters 1 and 2. BP PGS is elevated in Clusters 1 and 3. Glutamate pPGS is elevated in Cluster 2. (E) Models regressing PGS against cluster perform better for glutamate and GABA PGS than models regressing PGS against diagnosis. Models using cluster also perform better in predicting SZ PGS, though models using diagnosis better predict BP and serotonin PGS. (F) Glutamate and GABA PGS are significantly associated with cluster membership when controlling for diagnosis. (C. 1 = Cluster 1, C. 2 = Cluster 2, C. 3 = Cluster 3, resp. = responder to treatment, non-resp. = non-responder to treatment)

Discussion

We tested how overall and pathway-specific PGS are related to diagnosis and variation in psychosis symptoms in a diverse patient sample. We curated gene sets representing four major neurotransmitter systems(59–61) and calculated pPGS for SZ variants in these pathways. We also calculated overall SZ and BP PGS. Overall PGS, as well as glutamate and GABA pPGS, were associated with case status. pPGSs outperformed overall PGS in explaining endophenotypes: dopamine was associated with treatment response and global functioning, GABA was associated with social functioning, and glutamate was associated with cognitive control. Unsupervised clustering identified three phenotypically distinct groups of psychosis subjects that differed predominantly on positive symptoms, negative symptoms, cognitive control, and global functioning. Though our data largely represent different phenotypic information than that used by B-SNIP, we reproduced those findings which our measures can capture and generated new insights on the relationship between biotypes and genetic burden. Like B-SNIP, we identified three diagnostically mixed endophenotypes that separate on general impairment and cognitive control. Stratification was supported by distinct PGS profiles. Notably, Cluster 2, which showed primary deficits in cognitive control, negative symptoms, and social functioning, had elevated glutamatergic and GABAergic risk. Treatment efficacy separated by cluster and model comparisons showed that GABA and glutamate pPGS were better modeled by cluster than diagnosis. Our results link overall and pathway-specific partitions of genomic risk to transdiagnostic endophenotypes of psychotic disorders.

Our study features limitations. Notably, our cohort is smaller than many genomics studies, though our deep level of phenotyping is a strength and generally not possible for the larger or combined cohorts needed for GWAS. Focusing on PGS rather than SNP discovery mitigates these limitations, though limits our study to integration of previously-defined associations. We tested *a priori*-defined and hypothesis-driven pathways in the present study, however, there are certain to be other contributing pathways not investigated here. Our demographic heterogeneity, while a strength, presents methodological challenges. Thus, we controlled for race, ancestry, and genetic PCs in analyses. We further z-score normalized PGSs to subjects' genetically matched(58) 1000 Genomes(47) ancestries. Finally, k-means clustering will identify clusters even when the true nature of the data is continuous or cluster differences are not biologically relevant. The identification of unique PGS profiles for our clusters, as well as by-cluster differences in treatment response, provide auxiliary evidence supporting separate mechanisms driving distinct clusters of primary symptoms. Comparisons of clusters versus diagnoses show that clusters better predict glutamate and GABA pPGS in our cohort. Additionally, we replicate key findings from previous clustering attempts on psychosis subjects(9).

Our results show the utility of pPGS to interrogate convergent neurobiological mechanisms underlying disease phenotypes. The neurotransmitter systems tested here have been robustly associated with psychotic disorders(18–29). However, their relationship to specific symptoms have not been elucidated. Mechanisms underlying cognitive and negative symptoms in psychosis are particularly poorly understood, despite being some of the strongest predictors of poor outcomes in psychotic disorders(88–91). Our results robustly support a role for glutamate dysfunction in cognitive, as well as negative and social, symptoms in psychosis. This provides evidence that the biological basis of these symptoms may be partly explained by genetic risk driving glutamatergic abnormalities, which has been widely hypothesized. Our study and findings represent a model for going beyond overall genetic burden in investigation of the genetic components of psychosis, and, in particular, of maximizing power in smaller but deeply characterized cohorts. While our findings will need to be replicated, they offer a promising look into how genetic burden can be partitioned and can explain specific psychosis diagnosis and endophenotypes. Our work is a step towards a more precise understanding of convergent

molecular mechanisms underlying psychotic symptoms, which is critical for realizing the promise of stratified medicine and targeted treatments for those suffering from psychotic disorders.

Acknowledgements

Subject recruitment, blood draws, and phenotyping were supported by NIMH grant R01MH059883. DNA extraction and genotyping were supported by the UC Davis-HKU Collaboration Grant. Analyses and writing were supported by the UC Davis LaMP T32 (NIMH T32MH112507) and the NIMH NRSA F31MH129135.

Conflict of interest statement

The authors declare no conflicts of interest or competing financial interests.

References

1. Moreno-Küstner B, Martín C, Pastor L. Prevalence of psychotic disorders and its association with methodological issues. A systematic review and meta-analyses. McKenna PJ, editor. PLOS ONE. 2018 Apr 12;13(4):e0195687.

2. van Os J, Hanssen M, Bijl RV, Vollebergh W. Prevalence of Psychotic Disorder and Community Level of Psychotic Symptoms: An Urban-Rural Comparison. Arch Gen Psychiatry. 2001 Jul 1;58(7):663.

3. Subramaniam M, Abdin E, Vaingankar JA, Sambasivam R, Zhang YJ, Shafie S, et al. Lifetime Prevalence and Correlates of Schizophrenia and Other Psychotic Disorders in Singapore. Front Psychiatry. 2021 Mar 11;12:650674.

4. Sharifi V, Amin-Esmaeili M, Hajebi A, Motevalian A, Radgoodarzi R, Hefazi M, et al. Twelve-month prevalence and correlates of psychiatric disorders in Iran: the Iranian Mental Health Survey, 2011. Arch Iran Med. 2015 Feb;18(2):76–84.

5. Phillips MR, Zhang J, Shi Q, Song Z, Ding Z, Pang S, et al. Prevalence, treatment, and associated disability of mental disorders in four provinces in China during 2001-05: an epidemiological survey. Lancet Lond Engl. 2009 Jun 13;373(9680):2041–53.

6. Lee PH, Anttila V, Won H, Feng YCA, Rosenthal J, Zhu Z, et al. Genomic Relationships, Novel Loci, and Pleiotropic Mechanisms across Eight Psychiatric Disorders. Cell. 2019 Dec;179(7):1469-1482.e11.

7. Lee PH, Feng YCA, Smoller JW. Pleiotropy and Cross-Disorder Genetics Among Psychiatric Disorders. Biol Psychiatry. 2021 Jan;89(1):20–31.

8. Light G, Greenwood TA, Swerdlow NR, Calkins ME, Freedman R, Green MF, et al. Comparison of the Heritability of Schizophrenia and Endophenotypes in the COGS-1 Family Study. Schizophr Bull. 2014 Nov 1;40(6):1404–11.

9. Clementz BA, Sweeney JA, Hamm JP, Ivleva EI, Ethridge LE, Pearlson GD, et al. Identification of Distinct Psychosis Biotypes Using Brain-Based Biomarkers. Am J Psychiatry. 2016 Apr;173(4):373–84.

10. Zhang JP, Robinson D, Yu J, Gallego J, Fleischhacker WW, Kahn RS, et al. Schizophrenia Polygenic Risk Score

as a Predictor of Antipsychotic Efficacy in First-Episode Psychosis. Am J Psychiatry. 2019 Jan;176(1):21–8.

11. Shafee R, Nanda P, Padmanabhan JL, Tandon N, Alliey-Rodriguez N, Kalapurakkel S, et al. Polygenic risk for schizophrenia and measured domains of cognition in individuals with psychosis and controls. Transl Psychiatry. 2018 Apr 12;8(1):78.

12. He Q, Jantac Mam-Lam-Fook C, Chaignaud J, Danset-Alexandre C, Iftimovici A, Gradels Hauguel J, et al. Influence of polygenic risk scores for schizophrenia and resilience on the cognition of individuals at-risk for psychosis. Transl Psychiatry. 2021 Oct 9;11(1):518.

13. Choi SW, Garcia-Gonzalez J, Ruan Y, Wu HM, Johnson J, Hoggart C, et al. The power of pathway-based polygenic risk scores [Internet]. In Review; 2021 Jun [cited 2023 Jan 7]. Available from: https://www.researchsquare.com/article/rs-643696/v1

14. Yao Y, Guo W, Zhang S, Yu H, Yan H, Zhang H, et al. Cell type-specific and cross-population polygenic risk score analyses of MIR137 gene pathway in schizophrenia. iScience. 2021 Jul;24(7):102785.

15. Cosgrove D, Harold D, Mothersill O, Anney R, Hill MJ, Bray NJ, et al. MiR-137-derived polygenic risk: effects on cognitive performance in patients with schizophrenia and controls. Transl Psychiatry. 2017 Jan 24;7(1):e1012–e1012.

16. Cosgrove D, Mothersill DO, Whitton L, Harold D, Kelly S, Holleran L, et al. Effects of MiR-137 genetic risk score on brain volume and cortical measures in patients with schizophrenia and controls. Am J Med Genet B Neuropsychiatr Genet. 2018 Apr;177(3):369–76.

17. Reay WR, Geaghan MP, Atkins JR, Carr VJ, Green MJ, Cairns MJ. Genetics-informed precision treatment formulation in schizophrenia and bipolar disorder. Am J Hum Genet. 2022 Sep;109(9):1620–37.

18. Moghaddam B. Targeting metabotropic glutamate receptors for treatment of the cognitive symptoms of schizophrenia. Psychopharmacology (Berl) [Internet]. 2004 Jun [cited 2023 Jan 16];174(1). Available from: http://link.springer.com/10.1007/s00213-004-1792-z

19. Javitt DC. Glutamate as a therapeutic target in psychiatric disorders. Mol Psychiatry. 2004 Nov 1;9(11):984–97.

20. Namkung H, Yukitake H, Fukudome D, Lee BJ, Ursini G, Lam S, et al. The miR-124-AMPAR pathway connects polygenic risks with behavioral changes shared between schizophrenia and bipolar disorder [Internet]. Neuroscience; 2021 Oct [cited 2022 May 4]. Available from: http://biorxiv.org/lookup/doi/10.1101/2021.10.19.465053

21. Fromer M, Roussos P, Sieberts SK, Johnson JS, Kavanagh DH, Perumal TM, et al. Gene expression elucidates functional impact of polygenic risk for schizophrenia. Nat Neurosci. 2016 Nov;19(11):1442–53.

22. Koshiyama D, Kirihara K, Tada M, Nagai T, Fujioka M, Ichikawa E, et al. Electrophysiological evidence for abnormal glutamate-GABA association following psychosis onset. Transl Psychiatry. 2018 Oct 8;8(1):211.

23. Karolinska Schizophrenia Project (KaSP) Consortium, Orhan F, Fatouros-Bergman H, Goiny M, Malmqvist A, Piehl F, et al. CSF GABA is reduced in first-episode psychosis and associates to symptom severity. Mol Psychiatry. 2018 May;23(5):1244–50.

24. Seeman P, Chau-Wong M, Tedesco J, Wong K. Brain receptors for antipsychotic drugs and dopamine: direct binding assays. Proc Natl Acad Sci. 1975 Nov;72(11):4376–80.

25. Creese I, Burt DR, Snyder SH. Dopamine Receptor Binding Predicts Clinical and Pharmacological Potencies of Antischizophrenic Drugs. Science. 1976 Apr 30;192(4238):481–3.

26. Seeman P, Schwarz J, Chen JF, Szechtman H, Perreault M, McKnight GS, et al. Psychosis pathways converge via D2High dopamine receptors. Synapse. 2006 Sep 15;60(4):319–46.

27. Richtand N, Welge J, Logue A, Keckjr P, Strakowski S, Mcnamara R. Role of serotonin and dopamine receptor binding in antipsychotic efficacy. In: Progress in Brain Research [Internet]. Elsevier; 2008 [cited 2023 Jan 16]. p. 155–75. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0079612308009084

28. Coyle N, Jones I, Robertson E, Lendon C, Craddock N. Variation at the serotonin transporter gene influences susceptibility to bipolar affective puerperal psychosis. The Lancet. 2000 Oct;356(9240):1490–1.

29. Malhotra AK, Goldman D, Mazzanti C, Clifton A, Breier A, Pickar D. A functional serotonin transporter (5-HTT) polymorphism is associated with psychosis in neuroleptic-free schizophrenics. Mol Psychiatry. 1998 Jul 1;3(4):328–32.

30. First M, Spitzer R, Gibbon M, Williams J. Structured Clinical Interview for DSM-IV-TR Axis I Disorders, Research Version, Patient Edition. (SCID-I/P). Biometrics Research, New York State Psychiatric Institute; 2002.

31. American Psychiatric Association. Global assessment of functioning (GAF). In: Diagnostic and Statistical Manual on Mental Disorders, third ed text rev. Washington, D.C.: American Psychiatric Association; 1987.

32. Auther A, Smith C, Cornblatt B. Global Functioning: Social Scale (GF: Social). In Glen Oaks, New York: Zucker-Hillside Hospital; 2006.

33. Niendam T, Bearden C, Johnson J, Cannon T. Global Functioning: Role Scale (GF: Role). In Los Angeles: University of California, Los Angeles; 2006.

34. Young RC, Biggs JT, Ziegler VE, Meyer DA. A Rating Scale for Mania: Reliability, Validity and Sensitivity. Br J Psychiatry. 1978 Nov;133(5):429–35.

35. Andreasen NC. The Scales of Negative (SANS) and Positive (SAPS) Symptoms. Iowa City: University of Iowa; 1984.

36. Overall JE, Gorham DR. The Brief Psychiatric Rating Scale. Psychol Rep. 1962 Jun;10(3):799–812.

37. Barch DM, Carter CS, MacDonald AW, Braver TS, Cohen JD. Context-processing deficits in schizophrenia: diagnostic specificity, 4-week course, and relationships to clinical symptoms. J Abnorm Psychol. 2003 Feb;112(1):132–43.

38. Howes OD, Bose SK, Turkheimer F, Valli I, Egerton A, Valmaggia LR, et al. Dopamine Synthesis Capacity Before Onset of Psychosis: A Prospective [^{18} F]-DOPA PET Imaging Study. Am J Psychiatry. 2011 Dec;168(12):1311–7.

39. Howes OD, McCutcheon R, Agid O, de Bartolomeis A, van Beveren NJM, Birnbaum ML, et al. Treatment-Resistant Schizophrenia: Treatment Response and Resistance in Psychosis (TRRIP) Working Group Consensus Guidelines on Diagnosis and Terminology. Am J Psychiatry. 2017 Mar;174(3):216–29.

40. Smucny J, Lesh TA, Carter CS. Baseline Frontoparietal Task-Related BOLD Activity as a Predictor of Improvement in Clinical Symptoms at 1-Year Follow-Up in Recent-Onset Psychosis. Am J Psychiatry. 2019

Oct 1;176(10):839–45.

41. Rosvold HE, Mirsky AF, Sarason I, Bransome ED, Beck LH. A continuous performance test of brain damage. J Consult Psychol. 1956 Oct;20(5):343–50.

42. Henderson D, Poppe AB, Barch DM, Carter CS, Gold JM, Ragland JD, et al. Optimization of a Goal Maintenance Task for Use in Clinical Applications. Schizophr Bull. 2012 Jan 1;38(1):104–13.

43. Minzenberg MJ, Laird AR, Thelen S, Carter CS, Glahn DC. Meta-analysis of 41 Functional Neuroimaging Studies of Executive Function in Schizophrenia. Arch Gen Psychiatry. 2009 Aug 1;66(8):811.

44. MacDonald AW, Cohen JD, Stenger VA, Carter CS. Dissociating the Role of the Dorsolateral Prefrontal and Anterior Cingulate Cortex in Cognitive Control. Science. 2000 Jun 9;288(5472):1835–8.

45. Infinium® Genotyping Data Analysis: A guide for analyzing Infinium genotyping data using the GenomeStudio® Genotyping Module [Internet]. Available from: https://pre.illumina.com.cn/content/dam/illumina-marketing/documents/products/technotes/technote_infinium_genotyping_data_analysis.pdf

46. Patel H, Hyuck Sang L. GenomeStudio Genotyping QC SOP v.1.6 [Internet]. Available from: https://khp-informatics.github.io/COPILOT/GenomeStudio_genotyping_SOP.html

47. Siva N. 1000 Genomes project. Nat Biotechnol. 2008 Mar;26(3):256–256.

48. Wang K, Li M, Hadley D, Liu R, Glessner J, Grant SFA, et al. PennCNV: An integrated hidden Markov model designed for high-resolution copy number variation detection in whole-genome SNP genotyping data. Genome Res. 2007 Nov;17(11):1665–74.

49. Diskin SJ, Li M, Hou C, Yang S, Glessner J, Hakonarson H, et al. Adjustment of genomic waves in signal intensities from whole-genome SNP genotyping platforms. Nucleic Acids Res. 2008 Nov;36(19):e126–e126.

50. Wang K, Chen Z, Tadesse MG, Glessner J, Grant SFA, Hakonarson H, et al. Modeling genetic inheritance of copy number variations. Nucleic Acids Res. 2008 Dec;36(21):e138–e138.

51. Rees E, Walters JTR, Georgieva L, Isles AR, Chambert KD, Richards AL, et al. Analysis of copy number variations at 15 schizophrenia-associated loci. Br J Psychiatry. 2014 Feb;204(2):108–14.

52. Grozeva D, Conrad DF, Barnes CP, Hurles M, Owen MJ, O'Donovan MC, et al. Independent estimation of the frequency of rare CNVs in the UK population confirms their role in schizophrenia. Schizophr Res. 2012 Mar;135(1–3):1–7.

53. Levinson DF, Duan J, Oh S, Wang K, Sanders AR, Shi J, et al. Copy Number Variants in Schizophrenia: Confirmation of Five Previous Findings and New Evidence for 3q29 Microdeletions and VIPR2 Duplications. Am J Psychiatry. 2011 Mar;168(3):302–16.

54. Trubetskoy V, Pardiñas AF, Qi T, Panagiotaropoulou G, Awasthi S, Bigdeli TB, et al. Mapping genomic loci implicates genes and synaptic biology in schizophrenia. Nature. 2022 Apr;604(7906):502–8.

55. Mullins N, Forstner AJ, O'Connell KS, Coombes B, Coleman JRI, Qiao Z, et al. Genome-wide association study of more than 40,000 bipolar disorder cases provides new insights into the underlying biology. Nat Genet 2021 536. 2021 May 17;53(6):817–29.

56. Ge T, Chen CY, Ni Y, Feng YCA, Smoller JW. Polygenic prediction via Bayesian regression and continuous shrinkage priors. Nat Commun. 2019 Apr 16;10(1):1776.

57. Choi SW, O'Reilly PF. PRSice-2: Polygenic Risk Score software for biobank-scale data. GigaScience. 2019 Jul 1;8(7):giz082.

58. Privé F, Aschard H, Carmi S, Folkersen L, Hoggart C, O'Reilly PF, et al. Portability of 245 polygenic scores when derived from the UK Biobank and applied to 9 ancestry groups from the same cohort. Am J Hum Genet. 2022 Jan 6;109(1):12–23.

59. Ogata H, Goto S, Sato K, Fujibuchi W, Bono H, Kanehisa M. KEGG: Kyoto Encyclopedia of Genes and Genomes. Nucleic Acids Res. 1999 Jan 1;27(1):29–34.

60. Joshi-Tope G. Reactome: a knowledgebase of biological pathways. Nucleic Acids Res. 2004 Dec 17;33(Database issue):D428–32.

61. Carbon S, Ireland A, Mungall CJ, Shu S, Marshall B, Lewis S, et al. AmiGO: online access to ontology and annotation data. Bioinformatics. 2009 Jan 15;25(2):288–9.

62. Charrad M, Ghazzali N, Boiteau V, Niknafs A. **NbClust** : An *R* Package for Determining the Relevant Number

of Clusters in a Data Set. J Stat Softw [Internet]. 2014 [cited 2023 Jan 10];61(6). Available from:

http://www.jstatsoft.org/v61/i06/

63. DisplayR. flipCluster: Cluster analysis. 2023.

64. R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna, Austria: R

Foundation for Statistical Computing; 2022. Available from: https://www.R-project.org/

65. Revelle W. psych: Procedures for Psychological, Psychometric, and Personality Research [Internet].

Evanston, Illinois: Northwestern University; 2022. Available from: https://CRAN.R-

project.org/package=psych

66. Wei T, Simko V. R package "corrplot": Visualization of a Correlation Matrix [Internet]. 2021. Available from:

https://github.com/taiyun/corrplot

67. Adler D, Kelly ST, Elliot T, Adamson J. vioplot: violin plot [Internet]. 202AD. Available from:

https://github.com/TomKellyGenetics/vioplot

68. Harrell FE. rms: Regression Modeling Strategies [Internet]. 2023. Available from: https://CRAN.R-

project.org/package=rms

69. Cinar O, Viechtbauer W. The **poolr** Package for Combining Independent and Dependent p Values. J Stat

Softw [Internet]. 2022 [cited 2023 Mar 28];101(1). Available from: https://www.jstatsoft.org/v101/i01/

70. Galwey NW. A new measure of the effective number of tests, a practical tool for comparing families of non-

independent significance tests. Genet Epidemiol. 2009 Nov;33(7):559–68.

71. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to

Multiple Testing. J R Stat Soc Ser B Methodol. 1995 Jan;57(1):289–300.

72. Perkins DO, Olde Loohuis L, Barbee J, Ford J, Jeffries CD, Addington J, et al. Polygenic Risk Score Contribution

to Psychosis Prediction in a Target Population of Persons at Clinical High Risk. Am J Psychiatry. 2020 Feb

1;177(2):155–63.

73. Calafato MS, Thygesen JH, Ranlund S, Zartaloudi E, Cahn W, Crespo-Facorro B, et al. Use of schizophrenia

and bipolar disorder polygenic risk scores to identify psychotic disorders. Br J Psychiatry. 2018

Sep;213(3):535–41.

74. Smigielski L, Papiol S, Theodoridou A, Heekeren K, Gerstenberg M, Wotruba D, et al. Polygenic risk scores across the extended psychosis spectrum. Transl Psychiatry. 2021 Nov 26;11(1):600.

75. Vassos E, Di Forti M, Coleman J, Iyegbe C, Prata D, Euesden J, et al. An Examination of Polygenic Score Risk Prediction in Individuals With First-Episode Psychosis. Biol Psychiatry. 2017 Mar;81(6):470–7.

76. Ripke S, Neale BM, Corvin A, Walters JTR, Farh KH, Holmans PA, et al. Biological insights from 108 schizophrenia-associated genetic loci. Nature. 2014;

77. Cardno AG, Owen MJ. Genetic Relationships Between Schizophrenia, Bipolar Disorder, and Schizoaffective Disorder. Schizophr Bull. 2014 May 1;40(3):504–15.

78. ReproGen Consortium, Psychiatric Genomics Consortium, Genetic Consortium for Anorexia Nervosa of the Wellcome Trust Case Control Consortium 3, Bulik-Sullivan B, Finucane HK, Anttila V, et al. An atlas of genetic correlations across human diseases and traits. Nat Genet. 2015 Nov;47(11):1236–41.

79. Ruderfer DM, Ripke S, McQuillin A, Boocock J, Stahl EA, Pavlides JMW, et al. Genomic Dissection of Bipolar Disorder and Schizophrenia, Including 28 Subphenotypes. Cell. 2018;

80. Morgenroth E, Orlov N, Lythgoe DJ, Stone JM, Barker H, Munro J, et al. Altered relationship between prefrontal glutamate and activation during cognitive control in people with high trait anxiety. Cortex J Devoted Study Nerv Syst Behav. 2019 Aug;117:53–63.

81. Falkenberg LE, Westerhausen R, Craven AR, Johnsen E, Kroken RA, L Berg EM, et al. Impact of glutamate levels on neuronal response and cognitive abilities in schizophrenia. NeuroImage Clin. 2014;4:576–84.

82. Falkenberg LE, Westerhausen R, Specht K, Hugdahl K. Resting-state glutamate level in the anterior cingulate predicts blood-oxygen level-dependent response to cognitive control. Proc Natl Acad Sci. 2012 Mar 27;109(13):5069–73.

83. Bustillo JR, Chen H, Gasparovic C, Mullins P, Caprihan A, Qualls C, et al. Glutamate as a Marker of Cognitive Function in Schizophrenia: A Proton Spectroscopic Imaging Study at 4 Tesla. Biol Psychiatry. 2011 Jan;69(1):19–27.

84. Leucht S, Leucht C, Huhn M, Chaimani A, Mavridis D, Helfer B, et al. Sixty Years of Placebo-Controlled Antipsychotic Drug Trials in Acute Schizophrenia: Systematic Review, Bayesian Meta-Analysis, and Meta-Regression of Efficacy Predictors. Am J Psychiatry. 2017 Oct;174(10):927–42.

85. Goff DC, Coyle JT. The Emerging Role of Glutamate in the Pathophysiology and Treatment of Schizophrenia. Am J Psychiatry. 2001 Sep;158(9):1367–77.

86. Gruber O, Chadha Santuccione A, Aach H. Magnetic Resonance Imaging in Studying Schizophrenia, Negative Symptoms, and the Glutamate System. Front Psychiatry [Internet]. 2014 Apr 3 [cited 2023 Feb 10];5. Available from: http://journal.frontiersin.org/article/10.3389/fpsyt.2014.00032/abstract

87. Ereshefsky S, Zarubin V, Tryon V, Lesh T, Carter C, Niendam T. T76. RETROSPECTIVE ADHD SYMPTOMS IN EARLY PSYCHOSIS: RELATION TO CURRENT CLINICAL SYMPTOMS AND PSYCHOSOCIAL FUNCTIONING. Schizophr Bull. 2020 May 18;46(Supplement_1):S260–1.

88. Bow-Thomas CC, Velligan DI, Miller AL, Olsen J. Predicting quality of life from symptomatology in schizophrenia at exacerbation and stabilization. Psychiatry Res. 1999 May;86(2):131–42.

89. Dickerson F. Social functioning and neurocognitive deficits in outpatients with schizophrenia: a 2-year follow-up. Schizophr Res. 1999 May 4;37(1):13–20.

90. Bowie CR, Reichenberg A, Patterson TL, Heaton RK, Harvey PD. Determinants of Real-World Functional Performance in Schizophrenia Subjects: Correlations With Cognition, Functional Capacity, and Symptoms. Am J Psychiatry. 2006 Mar;163(3):418–25.

91. Green MF, Kern RS, Braff DL, Mintz J. Neurocognitive Deficits and Functional Outcome in Schizophrenia: Are We Measuring the "Right Stuff"? Schizophr Bull. 2000 Jan 1;26(1):119–36.

Correlations of SZ and BP PGS in cases	All psychosis	SZ only	SA only	BP only
Pearson's r	0.64	0.75	0.64	0.37
p-value	< 0.001	< 0.001	< 0.001	0.01

A Gene counts per pPGS

B Pearson's correlations of PGS in psychosis subjects

C Psychosis v. control by pPGS

D SZ, SA, and BP v. control by pPGS

A Phenotypes measured

Measure	Phenotype
12-month BPRS change	Treatment response
GAF, GSF, GRF	Global, social, and role functioning
AX-CPT during fMRI	Cognitive control
SANS, SAPS, BPRS	Disorganization
SANS, SAPS, BPRS	Poverty symptoms
SANS, SAPS, BPRS	Reality distortion
YMRS	Mania

B Dop. and GABA are associated with poorer global functioning in psychosis

Glut., p = 0.223
GABA, p = 0.118
Dop., p = 0.020
Ser., p = 0.134

Glut., p = 0.744
GABA, p = 0.038
Dop., p = 0.405
Ser., p = 0.383

C Glut. pPGS is associated with altered cognitive control in psychosis

Glut., p = 0.022
GABA, p = 0.932
Dop., p = 0.653
Ser. p = 0.713

Glut., p = 0.007
GABA, p = 0.335
Dop., p = 0.829
Ser., p = 0.852

Glut., p = 0.017
GABA, p = 0.572
Dop., p = 0.806
Ser., p = 0.426

D Dop. pPGS is marginally associated with treatment response

p = 0.057

A

Phenotypes by cluster

Cognitive control
Lower indicates more pathology.

Psychotic symptoms
Higher indicates more pathology.

Global functioning
Lower indicates more pathology.

Z-score in all psychosis cases

1.5
1
0.5
0
-0.5
-1

d'-Context | Bilat. DLPFC β-val. | Bilat. SPC β-val. | Poverty symptoms | Reality distortion | Disorganization | YMRS | GRF | GSF | GAF

● Cluster 1 (n = 34)   ● Cluster 2 (n = 72)   ● Cluster 3 (n = 61)

B

Treatment response separates better by cluster than by diagnosis

By cluster

100%
80%
60%
40%
20%
0%

C. 1: 3 (Non-resp.), 13 (Resp.)
C. 2: 15, 13
C. 3: 10, 9

■ Resp. □ Non-resp.

X^2, p = 0.057

By diagnosis

100%
80%
60%
40%
20%
0%

SZ: 18, 23
SA: 3, 5
BP: 7, 7

■ Resp. □ Non-resp.

X^2, p = 0.845

C

Diagnosis by clust.

100%
80%
60%
40%
20%
0%

C. 1: 4, 11, 19
C. 2: 6, 13, 53
C. 3: 27, 8, 26

■ SZ ■ SA ■ BP

Clust. by diagnosis

SZ: 26, 53, 19
SA: 8, 13, 11
BP: 27, 6, 4

■ C. 1 ■ C. 2 ■ C. 3

D

PGS by cluster

SZ PGS
p < 0.01 v. control | p < 0.01 v. control | p = 0.04 v. control
Control | Clust. 1 | Clust. 2 | Clust. 3

Glut. PGS
p = 0.17 v. control | p < 0.01 v. control | p = 1.00 v. control
Control | Clust. 1 | Clust. 2 | Clust. 3

Dop. PGS
p = 0.39 v. control | p = 0.31 v. control | p = 0.73 v. control
Control | Clust. 1 | Clust. 2 | Clust. 3

BP PGS
p = 0.05 v. control | p = 0.46 v. control | p < 0.01 v. control
Control (n = 115) | Clust. 1 (n = 34) | Clust. 2 (n = 72) | Clust. 3 (n = 61)

GABA PGS
p = 0.04 v. control | p = 0.01 v. control | p = 0.80 v. control
Control (n = 115) | Clust. 1 (n = 34) | Clust. 2 (n = 72) | Clust. 3 (n = 61)

Ser. PGS
p = 0.85 v. control | p = 0.62 v. control | p = 0.27 v. control
Control (n = 115) | Clust. 1 (n = 34) | Clust. 2 (n = 72) | Clust. 3 (n = 61)

E

Model performance of PGS by cluster v. PGS by diagnosis

	BIC for PGS by:		R^2 for PGS by:	
	Cluster	Diagnosis	Cluster	Diagnosis
SZ PGS	**676**	678	**0.11**	0.09
BP PGS	670	**669**	0.16	0.16
Glut. pPGS	**566**	570	**0.12**	0.10
GABA pPGS	**588**	594	**0.12**	0.09
Dop. pPGS	554	554	0.17	0.17
Ser. pPGS	534	**532**	0.09	**0.10**

Better values indicated in **bold**.

F

Significance of cluster v. diagnosis as covariates in regression model for PGS

-log(p)

2.0
1.5
1.0
0.0

p = 0.01
p = 0.05
p = 0.10

■ C1 (v. C3) ■ C2 (v. C3) ■ SZ (v. BP) ■ SA (v. BP)

SZ PGS | BP PGS | Glut. pPGS | GABA pPGS | Dop. pPGS | Ser. pPGS