Full title: Assessment of sewer connectivity in the United States and its

implications for equity in wastewater-based epidemiology

Short title: Sewer connectivity and equity in wastewater-based epidemiology

Authors: QinQin Yu[1], Scott W. Olesen[2], Claire Duvallet[2], Yonatan H. Grad[1*]

Affiliations:

1. Department of Immunology and Infectious Diseases, Harvard T. H. Chan School of

Public Health, Boston, Massachusetts, USA

2. Biobot Analytics, Inc., Cambridge, Massachusetts, USA

[*]Corresponding author

Email: ygrad@hsph.harvard.edu (YG)

Abstract

Wastewater-based epidemiology is a promising public health tool that can yield a more representative view of the population than case reporting. However, only about 80% of the U.S. population is connected to public sewers, and the characteristics of populations missed by wastewater-based epidemiology are unclear. To address this gap, we used publicly available datasets to assess sewer connectivity in the U.S. by location, demographic groups, and economic groups. Data from the U.S. Census' American Housing Survey revealed that sewer connectivity was lower than average when the head of household was American Indian and Alaskan Native, White, non-Hispanic, older, and for larger households and those with higher income, but smaller geographic scales revealed local variations from this national connectivity pattern. For example, data from the U.S. Environmental Protection Agency showed that sewer connectivity was positively correlated with income in Minnesota, Florida, and California. Data from the U.S. Census' American Community Survey and Environmental Protection Agency also revealed geographic areas with low sewer connectivity, such as Alaska, the Navajo Nation, Minnesota, Michigan, and Florida. However, with the exception of the U.S. Census data, there were inconsistencies across datasets. Using mathematical modeling to assess the impact of wastewater sampling inequities on inferences about epidemic trajectory at a local scale, we found that in some situations, even weak connections between communities may allow wastewater monitoring in one community to serve as a reliable proxy for an interacting community with no wastewater monitoring, when cases are widespread. A systematic, rigorous assessment of sewer connectivity will be

37   important for ensuring an equitable and informed implementation of wastewater-based

38   epidemiology as a public health monitoring system.

Introduction

Wastewater-based epidemiology (WBE) plays an important role in surveillance of SARS-CoV-2 [1,2], polio virus [3,4] and other pathogens [5,6] and has applications to monitoring a variety of other public health concerns [7], including opioid usage [8]. One proposed benefit of wastewater-based epidemiology is that wastewater data is more representative of the population than case reporting, which can be biased towards those with health-seeking behavior or access to healthcare [9]. For example, the populations served by the Health and Human Services SARS-CoV-2 National Wastewater Surveillance System were more representative of the entire US's age distribution and Black and Hispanic populations than the vaccinated population [10].

While WBE offers convenient sampling of populations served by public sewers, about 20% of individuals in the US live in homes not connected to public sewers [11,12]. This includes those on decentralized wastewater systems and those with no wastewater treatment systems. The most common decentralized wastewater system is septic tanks [13], which collect and treat wastewater onsite, typically the yard of the home. Households without wastewater treatment systems may have outhouses or privies, chemical toilets, or no plumbing. Variability in sewer connectivity exists across the U.S. [14,15], with more connectivity in urban areas than rural areas [11,16–18] and disparities driven by structural inequalities [19].

As with any emerging public health tools, it is important to ask to what extent the tool exacerbates or alleviates inequities. A few studies have evaluated equity of sewer

connectivity on broad geographic scales or at the sewershed level. A 2017 study from

Environmental Protection Agency Office of Water showed that households in the U.S.

that earned less than the national median household income (MHI, $61,000) were

almost 10% more likely to have a decentralized wastewater system or no wastewater

treatment system compared to households that earned more than the MHI [20].

Additionally, this study found that as household income decreased, decentralized

wastewater system usage increased in Florida, Hawaii, and Delaware, but not in Rhode

Island. Based on the 2019 U.S. Census Bureau's American Housing Survey, rural areas

are less connected to sewers, and while the income in rural areas was lower than that in

metropolitan areas, households connected to septic tanks were wealthier than those

connected to sewers within both rural and metropolitan areas [11]. In sewersheds in

North Carolina with comprehensive sewershed maps, the sewered population had

higher social vulnerability, more minorities, lower income, and lower educational

attainment than the unsewered population [21].

However, we have an incomplete understanding of the factors associated with sewer

connectivity across the US and the implications for the interpretation of wastewater

data. In this study, we sought to address the following questions, focusing our analyses

on the U.S.: (1) To what extent is there demographic and economic inequity in sewer

connectivity? (2) Which geographic areas have low sewer connectivity? (3) What is the

applicability of WBE data to neighboring unsampled communities? To address the first

question, we analyzed household-level data of sewer connectivity stratified by

geographic, demographic, and economic variables. For the second question, we

evaluated datasets aggregated at the county or county subdivision levels to qualitatively

identify geographic areas that have low sewer connectivity. To address the third

question, we used a mathematical model to simulate WBE in two interacting

populations.

Methods

Dataset compilation

We assembled publicly available datasets on sewer connectivity, septic connectivity, or

no plumbing at the household level across the U.S. from federal and state agencies

informed by (1) discussions with experts from federal and state agencies and (2) web

searches using terms including "Number of people connected with treatment plants in

United States", "Number of households on septic", and "U.S. census sewer". The types

of data included household-level data, locations of sewer systems and population

served, and population served by or lacking wastewater systems aggregated by county

or county subdivision. All datasets are summarized in **Table 1** and described in more

detail in the Supplementary Information.

Table 1: Datasets used in this study.

Dataset	Location	Year(s)	Description	Sampling method	Dataset completeness and potential biases	Ref.
American Housing Survey	US excluding Island areas	2013, 2019, 2021	This was a longitudinal survey of housing, demographic, and economic	The sampling method was a representative sample of all U.S. households at the time of sample selection (most	The survey excludes group quarters, businesses, hotels, and motels. Additionally, the survey was more likely to classify rural seasonal homes as vacant units, counted	[12,22–26]

			characteristics of approximately 60,000-80,000 representative households.	recently in 2015 and before that, in 1985). New housing units were added each survey cycle.	large cluster septic systems as public sewer, and was thought to lose new households built in rural areas due to its longitudinal design.	
American Community Survey 5-year estimates	US excluding Island areas	2021	This was a monthly survey of characteristics of populations and households of a representative subsample of addresses in the US. The survey reports the number of households lacking complete plumbing facilities by geographic subdivision.	The survey was sent to a random sample of addresses in the U.S. every month. Each address was selected no more than once every 5 years.	The surveys were completed by self-response (although a subset of non-responses were followed up with personal visits). Responses were reviewed for completeness and questionnaires needing clarification were followed up by phone calls. The data was averaged over 5 years to reduce statistical fluctuations.	[27]
Florida Department of Health septic tank inspections	Florida	Up to 2012	This dataset includes the onsite sewage treatment and disposal systems locations inspected by the Florida Department of Health reported in June 2012.	The sample included onsite sewage treatment and disposal system permits recorded in the Florida Department of Health system.	As of 2022, a permit was required for construction and inspection was recommended every 3-5 years after. From this dataset, an estimated 7% of households were connected to septic tanks across the state. This was substantially less than the Florida Department of Environmental Protection's estimate that approximately one third of Florida's population used septic tanks [28].	[29]
EPA Clean Watersheds Needs Survey	US	2012	This survey included voluntary submissions of the locations of publicly owned wastewater collection and treatment facilities	The responses were coordinated by states.	This survey had voluntary responses and variable effort and resources that each state put into the survey. States that had the most comprehensive responses were New York, California, Florida, New Jersey, Maryland, Iowa, Minnesota and Michigan. Missing	[30,31]

			and their estimated population served for the purposes to assess funding needs for treatment works projects.		facilities included those in South Carolina, the Northern Mariana Islands, and American Samoa; facilities whose projects did not have documented solutions or cost estimates; privately owned wastewater facilities; facilities on tribal lands and Alaskan Native Villages; and facilities whose projects received funding from other sources. Small community facilities are thought to be underrepresented due to having less resources for completing the survey, but this was not quantified.	
US Census Island Areas Decennial Survey	US Island Areas	2020	This was a survey of housing, social, and economic information of all housing units.	The survey was conducted in person and via phone interviews.	The COVID-19 pandemic impacted collection of data on group quarters [32] and the survey had higher nonresponse rates than previous years' surveys, particularly for Guam. Quality control checks were performed on responses.	[33]

103 The datasets are further described in the S1 Appendix and the dataset completeness

104 and potential biases are further assessed in the S2 Appendix. Additional datasets that

105 are briefly described in the main text but primarily analyzed in the Supplementary

106 information are summarized in **Table S 4**.

Calculation of weights, error bars, and summary statistics in American Housing Survey (AHS) data

All summary statistics were calculated by applying the weights reported in the AHS Public Use

File [26]. Because not every household was sampled and there was uneven sampling of

households across the U.S., the weight estimates the number of similar households that each

surveyed household represented. Error bars on summary statistics represent the middle 95% of

values calculated using the reported replicate weights (160 replicate weights per household).

Each summary statistic was calculated for each replicate separately before taking the middle

95% of values. Only categories with at least 5 households were used.

Analysis of sewer connectivity by urban or rural areas in AHS data

We used the most recent AHS National Sample with publicly available data on the

urban or rural location of the household (2013) which categorized households into being

in a central city of a Metropolitan Statistical Area (MSA), in an urban area of an MSA but

not in a central city, in a rural area of an MSA but not in a central city, in an urban area

outside of an MSA, or in a rural area outside of an MSA [34]. Central cities are defined

as having either (1) ≥250,000 population or at least 100,000 people working within

corporate limits, (2) ≥25,000 population, at least 75 jobs for each 100 residents who

were employed, and 60% or fewer of the city's resident workers commuted to jobs

outside, or (3) 15,000-25,000 population, at least one third the size of the metropolitan

statistical area's largest city, and met the two commuting requirements in (2).

Metropolitan Statistical Areas were defined as whole counties that have significant

levels of community and contiguous urban areas in common and may include rural

areas. Urban areas were defined as having ≥2500 people, with at least 1500 residing

outside of institutional group quarters. Rural areas were defined as those not in urban

areas.

Analyses of geographic areas with low connectivity to sewer

Data of individual locations of septic tanks or sewer collection systems, when they

existed, were aggregated based on latitude and longitude into county or county

subdivision boundaries as defined by the U.S. Census' TIGER shapefiles of the

corresponding year [35]. To calculate the fraction of a geographic area connected to

septic or sewer, the total population size or total number of households in that

geographic area was taken from the corresponding year in the American Community

Survey (ACS) 5-year estimate [27]. For county subdivision and census designated place

analyses, only areas with at least 5 households and 20 population size were used for

the analyses. This thresholding removed at most 1% of the total population and at most

0.03% of the total number of households, making it unlikely to bias our results. Detailed

data analyses of the state and island area datasets are described in the Supplementary

information.

Counties and county subdivisions were classified into Metropolitan statistical areas (at

least one urbanized area of 50,000 or more population), Micropolitan statistical areas (at

least one urban cluster of 10,000-50,000 population), or Rural areas (all other areas)

using the U.S. Census classification of the encompassing Core Based Statistical Area

(core area containing a substantial population nucleus together with adjacent territory

that has a high degree of social and economic integration with the core as measured by

commuting ties) [36].

Demographic and economic variable correlations with sewer

or septic connectivity

To assess correlations of demographic and economic variables with sewer and septic

connectivity when aggregating at the county, county subdivision, or census designated place

levels, we used the ACS 5-year estimates from the corresponding year and geographic scale.

Only geographic subdivisions with at least 5 households and 20 population size were used. The

Pearson correlation coefficient between demographic or economic variables of interest and the

percentage of households in a geographic subdivision connected to septic tanks or sewer was

calculated using the stats.pearsonr function in the scipy package (version 1.10.1) in Python

(version 3.11.0). A Bonferroni correction was applied to correct for multiple hypothesis testing to

calculate the q values.

169   Simulations of interacting populations and wastewater

170   sampling

Two interacting populations (A and B) were modeled in a deterministic compartmental model

with susceptible, infected, and recovered (SIR) compartments for each population and a

fraction, ε, of cross-population contacts. The base parameters were set at population size

$N_A=N_B=5000$, recovery rate $\gamma_I=0.18$ inverse days (corresponding to an infectious period of 5.6

days), basic reproduction number $R_0^A = R_0^B = \frac{\beta_A}{\gamma_I} = \frac{\beta_B}{\gamma_I} = 1.5$ and were varied in sensitivity

analyses. The differential equations were solved using the integrate.odeint function in the scipy

package (version 1.6.2) in Python (version 3.11.0) using a timestep of 0.1 days.

Wastewater sampling was modeled by the number of copies of pathogen genetic material shed

per day by each population that was sampled by wastewater divided by the volume of sampled

wastewater produced per day by each population. Both numerator and denominator depended

on the fraction of waste produced (including shed pathogen genetic material) by each

population that is sampled by wastewater, f_A and f_B. A detailed description of the simulations is

provided in S1 Appendix.

Code availability

187   All code and data to reproduce the analyses and simulations are available at

188   https://github.com/gradlab/wastewater_equity.

Results

Is there inequity in who could be sampled by wastewater-

based epidemiology?

To address this question, we used the American Housing Survey (AHS), a biannual

longitudinal survey of representative housing units within the U.S that reports

household-level data of sewer connectivity, demographic factors, and economic factors

(**Table 1**). We focused our analyses on geographic scales with a reliable sample size for

statistical analyses (census division, urban/rural at the national level, metropolitan

statistical areas) [37].

According to the 2021 AHS, 83% of US households were connected to public sewers

and 16% to standard septic tanks (**Fig S 1**). Each of the other forms of sewage disposal

(non-standard septic tanks, chemical toilets, outhouses or privies, other, none, or not

reported) was more than an order of magnitude less common (**Fig S 1**). Across census

divisions in 2021, the western US was overall better connected to sewers than the

eastern US, with the highest levels of sewer connection in the Pacific (91.3% of

households, CI: 90.8%-91.7%) and Mountain (87.6% of households, CI: 86.8%-88.4%)

divisions and the lowest levels of sewer connection in the New England (72.9% of

households, CI:72.1%-74.0%) and East South Central (74.9% of households, CI:

73.7%-76.3%) divisions (**Fig 1**).

Fig 1*: The percentage of households connected to sewers in each census division. Data

are from the 2021 American Housing Survey National Survey [12]. The number of sampled

households for each census division are 3890 (New England), 7018 (Middle Atlantic), 8703

(East North Central), 2862 (West North Central), 13643 (South Atlantic), 2903 (East South

Central), 8051 (West South Central), 4570 (Mountain), and 12501 (Pacific).

Across census divisions, households with an Asian, Black or African American, or Native

Hawaiian and Other Pacific Islander householder (the owner or renter of the unit) were on

average 14.0%, 13.1%, and 12.5% more connected to sewers, respectively, than the overall

census division; however, specifically in the Pacific census division, households with a Native

Hawaiian and Other Pacific Islander householder were 5.3% less connected to sewers than the

overall census division (**Fig 2**). Households with an American Indian and Alaska Native or White

householder were both on average 2.6% less connected to sewers than the overall census

division. Across census divisions, households with a Hispanic householder were on average

11.7% more connected to sewers whereas households with a non-Hispanic householder were

1.1% less connected to sewers than the overall census division.

Connection to sewers decreased with the age of the householder (**Fig 2**), with the exception

that households with a householder aged 75+ were better connected in most census divisions

than those with a householder aged 65-74 (79.7% vs 77.9% of households). Connection to

sewers decreased with household income, as measured as a percentage of the federal poverty

threshold determined by the Census [38], with the exception of the Mountain region where

households with an income at 50-99% of the poverty level were less connected than those with

income at ≥200% of the poverty level (84.5% vs 88.2% of households). Additionally, households

with 1 person were on average 4.2% more connected whereas households with 2 people were

on average 3.6% less connected to sewers than the overall census division. There were no

consistent trends across census divisions for households with more than 2 people.

Fig 2*:* **The percentage of households connected to sewers in each census division by**

demographic and economic characteristics. Data are from the 2021 American Housing

Survey National Survey [12]. The number of sampled households for each category is shown in

Fig S 2.

We sought to describe how the differences in sewer connection between demographic and

economic groups observed at the census division level was affected by household location in

urban or rural areas (see Methods). The qualitative trends observed at the census division level

for race, ethnicity, age, household size, and income were largely preserved across national

urban and rural areas (**Fig 3**). One notable exception was that lower sewer connectivity for

households with an American Indian and Alaska Native or Hawaiian and Other Pacific Islander

householder compared to those with a Black or Asian householder was only observed in

locations inside of an Metropolitan Statistical Area (MSA) but outside of a central city (both

urban and rural). In central cities and locations outside of an MSA (both urban and rural),

households with an American Indian and Alaska Native or Hawaiian and Other Pacific Islander

householder had similar levels of sewer connectivity as households with a Black or Asian

householder.

By central city/suburban status (2013)

**Fig 3*: The percentage of households connected to sewers by urban and rural

categorizations and by demographic and economic characteristics.** See Methods for

definitions of Metropolitan Statistical Area (MSA), central city, urban, and rural. Data from the

2013 American Housing Survey National Survey [39]. The number of sampled households for

each category is shown in **Fig S 3**. MSA: metropolitan statistical area.

Next, we assessed sewer connectivity in large metropolitan areas, which was the

smallest geographic scale resolved by the AHS due to sample size. In 35 of the largest

metropolitan areas in 2019 and 2021, household connectivity to sewers ranges from

69.2% (Birmingham-Hoover, AL) to 99.4% (LA-Long Beach-Anaheim, CA) (**Fig S 4**). In

each metropolitan area, households with a White or American Indian and Alaska Native

householder, with an older householder, with higher income, with a larger household

size, and with a non-Hispanic householder were less connected to sewers than the

average household, which was similar to what was seen when stratifying by census

division. Exceptions were lower connectivity for households with a Hispanic

householder in San Jose, CA (Hispanic householder: 96.1% of households, CI: 95.1%-

97.2%; non-Hispanic householder: 97.8% of households, CI: 97.5%-98.2%) and lower

connectivity for households with lower income in San Francisco, CA; Dallas, TX;

Chicago, IL; and Tampa, FL (**Fig S 4**).

In summary, at the census division level, broad trends of sewer connectivity by racial

group revealed higher connectivity for households with an Asian, Black or African

American, Native Hawaiian and Other Pacific Islander, or Hispanic householder and

lower connectivity for households with an American Indian and Alaska Native or White

householder. Sewer connectivity decreased with age, decreased with household

income, and was highest in households with one person. These trends were similar

when stratifying by urban, rural, central city, and MSA status, except that the broad

racial disparities described above were only observed in urban and rural areas that are

inside of an MSA but outside of a central city. Broad trends were also similar for

individual large metropolitan areas, with some exceptions for households with a

Hispanic householder and trends by household income in particular MSAs.

Which geographic areas have low connectivity to sewers and

what are their demographics?

Next, we sought to assess which geographic areas in the U.S. have low connectivity to

sewer systems. These are areas where wastewater-based epidemiology may provide

little direct information on disease dynamics. Additionally, we assessed correlations of

sewer connectivity at the county or county subdivision level with demographic and

economic factors. In the absence of comprehensive national data on sewer connectivity,

we assembled publicly available state-based datasets, which included California,

Florida, Iowa, Maryland, Michigan, Minnesota, New Jersey, and New York, as well as

the U.S. Island Areas. The compilation of the datasets and an assessment of data

completeness are described in the S2 Appendix, **Table *1***, and **Table S 4**.

Using data from the 2021 American Community Survey on the percentage of occupied

housing units lacking complete plumbing facilities by county, we defined the lower

bound on the percentage of the population not connected to sewers. In 14 counties,

≥10% of occupied housing units lacked complete plumbing facilities; of these counties,

12 were in Alaska and 2 were in the Navajo Nation. In 5 counties, ≥20% of occupied

housing units lacked complete plumbing facilities, all in Alaska. The Yukon-Koyukuk

Census Area in Alaska, had the highest percentage of occupied housing units lacking

complete plumbing facilities at 36% (**Fig S 5**, **Table S *1***).

The Florida Department of Health onsite sewage treatment and disposal systems

inspection data (**Fig *4***) revealed that the percentage of households not on septic tanks

(expected to correlate with the percentage of households connected to sewer) was

higher in metropolitan compared to micropolitan county subdivisions and higher in

micropolitan compared to rural county subdivisions, but the effect was not significant (p

= 0.052 and p = 0.37, respectively) (**Fig S 6**a-b). The percentage of households not on

septic tanks was significantly lower in county subdivisions in the panhandle (74.0% vs

92.0%, $p<10^{-9}$) and in county subdivisions that did not border the coast (86.6% vs

95.7%, $p<10^{-9}$) (**Fig S 6**). By combining this dataset with the 2012 American Community

Survey (ACS) describing demographic and economic characteristics of county

subdivisions, we found that county subdivisions with less septic usage (suggesting more

sewer connectivity) were significantly more Hispanic, more Asian, and had a smaller

household size (**Table *2*** and **Fig S 7**), consistent with the nation-wide results from the

AHS. These trends were driven by metropolitan county subdivisions (**Fig S 8**), and an

additional trend in metropolitan county subdivisions not observed for all county

subdivisions was that county subdivisions with less septic usage had a significantly

lower percentage of American Indian and Alaska Natives. There were too few

micropolitan and rural county subdivisions to observe significant trends (**Fig S 9** and **Fig**

S 10).

Fig 4*: **Map of the percentage of households in each county subdivision of Florida *not***

connected to septic tanks. Data are from the Florida Department of Health septic tank

inspection data from June 2012. County subdivisions with fewer than 5 households or 20

population size (<0.01% of total households and <0.01% of total population size in the state) are

displayed in white.

Table 2: Correlation of the percentage of a Florida county subdivision *not* connected to

septic tanks with different demographic or economic variables.

Demographic or economic variable	Correlation of variable with % of county subdivision *not* on septic (Pearson correlation coefficient)	Significance (q-value)

Percent one race and Asian	0.31	**<0.001**
Percent Hispanic	0.21	**<0.01**
Average household size	-0.19	**<0.01**
Percent one race and American Indian and Alaska native	-0.15	0.07
Median age	0.12	0.36
Percent one race and White	-0.12	0.40
Percent one race and Black or African American	0.09	1.11
Median income	0.07	2.27
Percent one race and some other race	0.07	2.33
Percent one race and Native Hawaiian and other Pacific Islander	-0.04	5.76
Percent uninsured (health insurance)	-0.02	8.24

Correlations significant to 5% are shown in bold. Data are from the Florida Department

of Health septic tank inspection permits reported in 2012.

In the 8 states with robust data in the EPA Clean Watersheds Needs Survey, multiple

neighboring counties in Florida, Michigan, and Minnesota had less than 20% of

residents receiving sewage collection, whereas this was not the case in California,

Florida, Iowa, Maryland, New Jersey, and New York (**Fig S 11**). Across multiple states,

sewer connectivity by county was positively correlated with percent Asian and percent

Black or African American and negatively correlated with percent White (**Table 3**, full

table shown in **Table S 2**), consistent with the nationwide trends from the AHS data.

While counties more connected to sewers appeared to have a significantly lower

percent of American Indian and Alaskan Natives in California, this is affected by missing

data from Indian reservations. Age and percent without health insurance were

negatively correlated with sewer connectivity. Counties more connected to sewers had

significantly higher income in Minnesota, Florida, and California, in contrast to the

national results from the AHS data. Significant trends of sewer connectivity with

demographic and economic factors were predominantly driven by metropolitan counties

(**Table S 3**).

Table 3: Correlation of the percentage of households in a county connected to

public sewers with different demographic or economic factors by state.

Demographic or economic variable	State	Correlation of variable with % of households by county connected to public sewers (Pearson correlation coefficient)	Significance (q-value)
Median age	MN	-0.43	<0.01
	MI	-0.41	0.01
Median income	MN	0.44	<0.01
	FL	0.48	<0.01
	CA	0.49	<0.01

Percent one race and American Indian and Alaska Native	CA	-0.50	<0.01
	MN	0.58	<0.001
	MI	0.51	<0.001
	NY	0.55	<0.001
Percent one race and Asian	CA	0.56	<0.001
	FL	0.53	<0.001
	MN	0.62	<0.001
	MI	0.59	<0.001
Percent one race and Black or African American	NY	0.55	<0.001
	IA	0.38	<0.01
	CA	0.46	0.02
	MI	-0.57	<0.001
	CA	-0.54	<0.001
Percent one race and White	NY	-0.51	<0.01
	IA	-0.35	0.03
Percent uninsured (health insurance)	MI	-0.38	0.03

Only correlations that are significant (q-value ≤ 0.05) are shown. Data are from the 2012 EPA Clean Watersheds Needs Survey.

363 Analysis of the Island Areas of Guam, the Northern Mariana Islands, the Virgin Islands,

364 and American Samoa revealed lower levels of sewer connectivity than in the states,

365 with considerable spatial variability in connectivity (see S2 Appendix).

Factors influencing applicability of wastewater-based

epidemiology to communities lacking sewer connection

The inequities in sewer connections among communities raise the question of the

applicability of inferences drawn from wastewater data in one community to

communities lacking sewer connections. To explore this question, we used a

deterministic compartmental model of two interacting populations with tunable levels of

interaction, each with susceptible-infected-recovered (SIR) dynamics, and with

wastewater sampling in each population into a common sample (**Fig 5**a). We set

population A to be entirely connected to sewers and population B to have a tunable

level of sampling by wastewater.

A common application of wastewater data is to aid in determining when an outbreak has

peaked, which can inform policy decisions on when to ease restrictions. We first asked

how well the wastewater data could predict the time of peak infections in population B

when it is completely unconnected to sewer systems and thus the only sampling is from

population A. We found that when the two populations have similar sizes, the

wastewater concentration peak and the infection peak in population B was within one

generation time except when the interactions between the two population is weak (**Fig**

5b). The generation time is the average time between an individual's infection and

transmission, which in an SIR model is the same as the infectious period. For example,

in two populations with 5000 individuals each, a disease basic reproduction number of

1.5, and infectious period of 5.6 days, the percentage of contacts that occur across

populations must drop below 4% before the peak in wastewater concentration occurred

more than one generation time apart from the peak in infections in the unconnected

population (**Fig 5**c).

Fig 5: Deterministic compartmental model of two interacting populations with

susceptible, infected, and recovered compartments and sampling by wastewater. (a)

Schematic of model. Population A is entirely sampled by wastewater and a fraction of

population B, f_B, is sampled by wastewater. (b) Number of infected individuals (black curves) in

population A, population B, and population A and B combined and wastewater concentration

(blue curves) over time for different interaction strengths or fractions of cross-population

contacts (ε). The vertical lines indicate the peak time for the curves with matching line style and

color. The time between the population B peak in infections and the wastewater concentration

peak is indicated by Δt. (c) The time between the population B peak in infections and the

wastewater concentration peak (defined in (b)) when varying the fraction of cross-population

contacts (ε) and the fraction of population B that is sampled (f_B). Light blue dots indicate

parameter regimes in which Δt is less than 1 generation time. In both (b) and (c), the simulation

starts with a single infected individual in population A at time 0 and no infected individuals in

population B. The parameters are set to population size $N_A = N_B = 5000$, basic reproduction

number $R^A_0 = R^B_0 = R_0 = \frac{\beta_A}{\gamma_I} = \frac{\beta_B}{\gamma_I} = 1.5$, rate of recovery $\gamma_I = 0.18$, and the overall contact rate times

the probability of infection given contact $\beta_A = \beta_A = \gamma_I R_0$.

We then weakened the assumption that population B had no contribution to the

sampled wastewater and allowed a fraction of population B to be sampled by

wastewater. For example, individuals in population B may commute to a workplace

sampled by wastewater, or a fraction of population B may be connected at home. As the

fraction of population B that is sampled by wastewater increases, the time between

infection and wastewater peaks decreases; however, the fraction of B that is sampled

has a much weaker effect on the wastewater peak time than the interaction parameter

between the communities. For example, with the above parameters and equal

population sizes, with 10% sampling of population B, the cross-population contacts

must drop below 4%, and with 50% below 3% and with 100% below 2% before the

wastewater concentration peak and the peak in the number of infections in population B

drops below one generation time.

Varying the population size of both populations (with an equal population size in the two

populations), the recovery rate or generation time, and the R_0 of both populations (with

an equal R_0 in the two populations) has only a weak effect on the discrepancy between

wastewater concentration and population B infection peaks (**Fig S 12**).

When the unconnected population has a smaller size than the connected population,

then even substantial interactions between the two populations can lead the wastewater

concentration to peak before the infections in the unconnected population peaks (**Fig S

12**). Both very weak and very strong interactions cause the wastewater peak to coincide

with the peak number of infections in the unconnected population.

Additionally, when the unconnected population has a smaller R_0 than the connected

population (which could occur for instance if the unconnected population was more

spread out geographically and thus had a lower contact rate), the outbreak in population

B peaks later than the wastewater peak leading to a discrepancy in the peak times (**Fig

S 12**).

Discussion

Equity of sewer connectivity

Our analysis of available datasets revealed considerable variability in sewer connectivity

within and between communities, across locations, demographics, and economic

statuses in the United States. The western US had higher levels of sewer connectivity

than the eastern US. Across the US, lower connectivity to sewers was observed for

American Indian and Alaska Native householders; White householders; non-Hispanic

householders; older populations; and larger households in most census divisions. In the

Pacific census division, lower connectivity to sewers was observed for Native Hawaiian

and Other Pacific Islander householders specifically in areas inside of a metropolitan

statistical area but outside of a central city. Across census divisions (with the exception

of the Mountain census division), households with higher income were less connected

to sewers, consistent with nation-wide observations [11]. The decrease in sewer

connectivity by income could not be explained by collinearity between age and income

as age and income exhibited a weak negative correlation (**Fig S 13**). These results are

also consistent with the observation that in North Carolina, minorities and those with

social vulnerabilities are overrepresented in the sewered population [21]. Aggregating

by large metropolitan areas yielded similar trends as when aggregating by census

divisions, except for lower connection for Hispanic householders in San Jose area and

for households with lower income in San Francisco, Dallas, Chicago, and Tampa

metropolitan areas, whereas these groups were better connected to sewers at the

census division level. The differences in the MSA-level data suggested heterogeneity in

connectivity across locations.

We have not attempted to determine why the observed differences in sewer connectivity

exist. Many of the variables that we analyzed are likely to be correlated with one

another.

Geographic variation in sewer connectivity

Large parts of Alaska and the Navajo Nation lacked plumbing. This is consistent with

the finding from the Annual Report to Congress on Sanitation Deficiency Levels for

Indian Homes and Communities in 2019 that approximately 20% of Indian homes in

Alaska do not have sewage disposal and about 10% of Indian homes in the Navajo

Nation do not have sewage disposal [40].

State-level datasets revealed that sewer connectivity varied between and within

communities. While in some states most areas had sewer connectivity (California, Iowa,

Maryland, New Jersey, New York), in others there were large regions that lacked

connectivity (Minnesota, Michigan, Florida). Asian and Hispanic populations were less

connected to septic tanks across county subdivisions in Florida; however, there was

considerable variability across the state and potential biases in the data. In contrast to

the national data, in Minnesota, Florida, and California, counties with lower median

household incomes were less connected to sewers, suggesting variability across the

nation. This result in Florida is consistent with the observation that as income levels

rose, household decentralized system usage declined in Florida [20]. However, we did

not observe any significant correlation of income with septic usage in the Florida

Department of Health onsite sewage treatment and disposal systems inspection

permitting dataset, possibly due to not all septic tanks having a permit that is reported in

this dataset.

While these correlations may be largely driven by differences between urban and rural

areas within a county or county subdivision, they may still be very useful in assessing

inequities in sewer connectivity in large catchments that span neighboring counties or

county subdivisions.

Data limitations

Substantial data gaps and biases prevented a comprehensive analysis of disparities in

sewer system connectivity within and between communities across the US, particularly

for small communities, tribal lands and Alaska Native Villages, and state and local

geographic scales. The current design of the American Housing Survey has too small of

sample size to study state and local geographic scales [37]. Additionally, the AHS may

have underestimated the fraction of households using septic systems due to its survey

design (**Table 1**). The differences between Florida state data supports the need for

detailed and reproducible longitudinal studies of sewer connectivity (both in developing

sampling and analysis methods). Additionally, multiple datasets were over a decade old,

and changes to which households are served by sewer and septic systems will have

occurred since the datasets were collected.

Public health officials may find it beneficial to conduct local equity studies of sewer

connectivity. Data on the catchment area sizes of the wastewater treatment plants

would allow a better understanding of the geographic extent of the sampled population

and the level of geographic aggregation needed to study sewer connectivity inequities.

A recent study mapping wastewater treatment plant catchment areas and their

population sizes served in New York state provides a template for how to create these

sewer catchment maps from a combination of permitting, survey, and tax record data

[16]. If possible, it would be helpful to consolidate and standardize existing data where it

exists (for instance, sewage disposal permitting at the local or state levels). Additionally,

if the EPA's request to expand the question on the ACS about access to plumbing

facilities to additionally ask about the type of plumbing facility [20,41] is granted, then

this would be a valuable dataset with the geographic and temporal resolution to study

equity in sewer connectivity moving forward.

523 Generalizability of wastewater-based epidemiology data in

524 light of heterogeneities in sewer connectivity

Our modeling results suggest that even weak interactions between two communities

allow wastewater monitoring in one community to serve as reliable proxy for the time of

maximum infections in the other community when the population sizes and R_0 of the two

populations are comparable, but not when the unconnected population has a

substantially lower population size or R_0. In the scenario with unequal population size,

outbreaks that are seeded at the same time in the two populations in the absence of

interactions peak earlier in the unconnected population due to its smaller size. With

weak interactions, seeding of infections from population A to B occurs slower, leading to

more coincidence in peaks; with strong interactions, then the dynamics of B are

dominated by those in A, leading to more coincidence in peaks; at intermediate

interaction strengths, infections are seeded earlier and peaks early, leading to the

largest discrepancy in wastewater and infection peaks.

As the purpose of our model was to explore the impact of factors on the generalizability

of wastewater rather than to accurately capture the dynamics in all scenarios, we chose

the most simple compartmental model (SIR) and made simplifying assumptions,

including that all infected individuals shed the same amount of pathogen genetic

material into wastewater and shed only during the infected period; that all individuals

contribute equally to wastewater; and that pathogen detection in wastewater has perfect

sensitivity. We also note that compartmental models have previously been used to

analyze wastewater data [42].

Our results suggest that in assessing the generalizability of wastewater data, it would be

useful to estimate the extent of mobility between connected and unconnected

communities. Interestingly, Ref [43] found no correlation between the size of a

catchment area and the correlation of wastewater with case data for SARS-CoV-2,

consistent with the result that interactions between connected and unconnected

communities cause the disease dynamics to look similar to the wastewater data. In

regions without sewer connectivity or with little interaction between connected and

unconnected communities, wastewater data from neighboring communities will be less

informative. In areas with low sewer connectivity in households, sampling wastewater

outflow at frequently visited non-household locations (i.e., schools, offices, malls, etc.)

may capture a more representative population.

Additional considerations for equity in WBE

In addition to inequities in sewer connectivity, the sewer locations used for wastewater

sampling should be considered to promote demographic equity and ensuring the ability

to capture spatiotemporal trends [21,44–46]. While we have focused on analysis on the

US due to data availability, internationally disadvantaged populations are associated

with lower access to sewers [47–49]. The ongoing development of wastewater sampling

in non-sewered settings, for example in water channels in Las Vegas Valley [50], onsite

sanitation facilities in Bangladesh [51], a refugee camp in Lebanon [52], and various

non-sewered settings in low and middle income countries [53–55], represents a critical

area for research and development.

Conclusions

In summary, while wastewater-based epidemiology is a useful tool to monitor disease

burden and dynamics, our analyses suggest that access to this new tool varies across

the US. More comprehensive data on sewer connectivity is needed, and in combination

with assessments of mobility and population parameters, these data can help with the

design of wastewater sampling schemes and the interpretation for epidemic trends in

sampled and neighboring unsampled communities.

Acknowledgements

We thank Cara Omana from the Minnesota Pollution Control Agency for generously

sharing the Minnesota WINS data and Harry Campbell from the Utah Department of

Environment Quality, Division of Water Quality for generously sharing the Utah MWPP

survey data and to both for answering questions about the datasets. We thank

colleagues in the Grad lab and the Harvard T. H. Chan School of Public Health Center

for Communicable Disease Dynamics for helpful discussions, particularly Stephen

Kissler.

References

1. Peccia J, Zulli A, Brackney DE, Grubaugh ND, Kaplan EH, Casanovas-Massana A, et al. Measurement of SARS-CoV-2 RNA in wastewater tracks community infection dynamics. Nat Biotechnol. 2020;38: 1164–1167.

2. Medema G, Been F, Heijnen L, Petterson S. Implementation of environmental surveillance for SARS-CoV-2 virus to support public health decisions: Opportunities and challenges. Curr Opin Environ Sci Health. 2020;17: 49–71.

3. Pöyry T, Stenvik M, Hovi T. Viruses in sewage waters during and after a poliomyelitis outbreak and subsequent nationwide oral poliovirus vaccination campaign in Finland. Appl Environ Microbiol. 1988;54: 371–374.

4. Link-Gelles R, Lutterloh E, Schnabel Ruppert P, Backenson PB, St George K, Rosenberg ES, et al. Public Health Response to a Case of Paralytic Poliomyelitis in an Unvaccinated Person and Detection of Poliovirus in Wastewater - New York, June-August 2022. MMWR Morb Mortal Wkly Rep. 2022;71: 1065–1068.

5. Kilaru P, Hill D, Anderson K, Collins MB, Green H, Kmush BL, et al. Wastewater Surveillance for Infectious Disease: A Systematic Review. Am J Epidemiol. 2023;192: 305–322.

6. Boehm AB, Hughes B, Duong D, Chan-Herur V, Buchman A, Wolfe MK, et al. Wastewater concentrations of human influenza, metapneumovirus, parainfluenza, respiratory syncytial virus, rhinovirus, and seasonal coronavirus nucleic-acids during the COVID-19 pandemic: a surveillance study. Lancet Microbe. 2023. doi:10.1016/S2666-5247(22)00386-X

7.   Baz-Lomba JA, Salvatore S, Gracia-Lor E, Bade R, Castiglioni S, Castrignanò E, et

al. Comparison of pharmaceutical, illicit drug, alcohol, nicotine and caffeine levels in

wastewater with sale, seizure and consumption data for 8 European cities. BMC

Public Health. 2016;16: 1035.

8.   Gushgari AJ, Venkatesan AK, Chen J, Steele JC, Halden RU. Long-term tracking of

opioid consumption in two United States cities using wastewater-based

epidemiology approach. Water Res. 2019;161: 171–180.

9.   Holm RH, Pocock G, Severson MA, Huber VC, Smith T, McFadden LM. Using

wastewater to overcome health disparities among rural residents. Geoforum.

2023;144: 103816.

10.   Analytics B. Wastewater for equitable Covid-19 monitoring. In: Medium [Internet]. 8

Jul 2021 [cited 9 Feb 2023]. Available:

https://biobotanalytics.medium.com/wastewater-for-equitable-covid-19-monitoring-

e3b947d91e0a

11.   Olesen SW, Young CCW, Duvallet C. White paper: The effect of septic systems on

wastewater-based epidemiology. 2022. doi:10.5281/zenodo.7130726

12.   U.S. Census Bureau. AHS 2021 National Public Use File. Available:

https://www2.census.gov/programs-

surveys/ahs/2021/AHS%202021%20National%20PUF%20v1.0%20CSV.zip

13.   US EPA Decentralized Wastewater Program}. Septic Systems Fact Sheet. 2008

Oct.

14.   Leker HG, MacDonald Gibson J. Relationship between race and community water

and sewer service in North Carolina, USA. PLoS One. 2018;13: e0193225.

15. Meehan K, Jurjevich JR, Chun NMJW, Sherrill J. Geographies of insecure water

access and the housing-water nexus in US cities. Proc Natl Acad Sci U S A.

2020;117: 28700–28707.

16. Hill DT, Larsen DA. Using geographic information systems to link population

estimates to wastewater surveillance data in New York State, USA. PLOS Glob

Public Health. 2023;3: e0001062.

17. Rural Community Assistance Partnership. Still Living Without the Basics in the 21st

Century: Analyzing the Availability of Water and Sanitation Services in the United

States. 2004. Available: http://opportunitylinkmt.org/wp-

content/uploads/2015/07/Still-Living-Without-the-Basics-Water.pdf

18. Hernandez A, Pierce G. The geography and socioeconomic characteristics of U.S.

households reliant on private wells and septic systems. J Am Water Resour Assoc.

2023;59: 1397–1412.

19. Brown J, Acey CS, Anthonj C, Barrington DJ, Beal CD, Capone D, et al. The effects

of racism, social exclusion, and discrimination on achieving universal safe water

and sanitation in high-income countries. Lancet Glob Health. 2023;11: e606–e614.

20. U.S. Environmental Protection Agency, Office of Water. Report to Congress on The

Prevalence Throughout the U.S. of Low- and Moderate-Income Households Without

Access to a Treatment Works and The Use by States of Assistance under Section

603(c)(12) of the Federal Water Pollution Control Act. U.S. Environmental

Protection Agency; 2021 Jul. Available:

https://www.epa.gov/system/files/documents/2022-01/low-mod-income-without-

treatment_report-to-congress.pdf

21. Hu XC, Reckling SK, Keshaviah A. Assessing health equity in wastewater

monitoring programs: Differences in the demographics and social vulnerability of

sewered and unsewered populations across North Carolina. medRxiv. 2023. p.

2023.10.06.23296680. doi:10.1101/2023.10.06.23296680

22. U.S. Census Bureau. AHS 2019 National Public Use File. Available:

https://www2.census.gov/programs-

surveys/ahs/2019/AHS%202019%20National%20PUF%20v1.1%20CSV.zip

23. U.S. Census Bureau. AHS 2019 Metropolitan Public Use File. Available:

https://www2.census.gov/programs-

surveys/ahs/2019/AHS%202019%20Metropolitan%20PUF%20v1.0%20CSV.zip

24. U.S. Census Bureau, Department of Commerce Department of Housing and Urban

Development. 2019 AHS Integrated National Sample: Sample Design, Weighting,

and Error Estimation. U.S. Census Bureau, Department of Commerce Department

of Housing and Urban Development; 2020 Sep. Available:

https://www2.census.gov/programs-

surveys/ahs/2019/2019%20AHS%20National%20Sample%20Design,%20Weightin

g,%20and%20Error%20Estimation.pdf

25. U.S. Census Bureau, Department of Commerce Department of Housing and Urban

Development. 2019 AHS Metropolitan Sample: Sample Design, Weighting, and

Error Estimation. U.S. Census Bureau, Department of Commerce Department of

Housing and Urban Development; 2020 Sep. Available:

https://www2.census.gov/programs-

surveys/ahs/2019/2019%20AHS%20Metropolitan%20Sample%20Design,%20Weig

hting,%20and%20Error%20Estimation.pdf

26. U.S. Census Bureau, Department of Commerce Department of Housing and Urban

Development. 2021 AHS Integrated National Sample: Sample Design, Weighting,

and Error Estimation. U.S. Census Bureau, Department of Commerce Department

of Housing and Urban Development; 2022 Aug. Available:

https://www2.census.gov/programs-

surveys/ahs/2021/2021%20AHS%20National%20Sample%20Design,%20Weightin

g,%20and%20Error%20Estimation.pdf

27. US Census Bureau. American Community Survey Information Guide. [cited 5 Jan

2023]. Available: https://www.census.gov/programs-surveys/acs/library/information-

guide.html

28. General Facts and Statistics about Wastewater in Florida. [cited 10 Feb 2023].

Available: https://floridadep.gov/water/domestic-wastewater/content/general-facts-

and-statistics-about-wastewater-florida

29. Florida Department of Health. Onsite Sewage Locations in Florida - 2012. Available:

https://fgdl.org/zips/metadata/xml/septic_jun12.xml

30. Environmental Protection Agency. Clean Watersheds Needs Survey Detailed

Scope and Methods. Environmental Protection Agency; 2012. Available:

https://www.epa.gov/sites/default/files/2015-

10/documents/cwns_2012_detailed_scope_and_methods-508.pdf

31. Environmental Protection Agency. Clean Watersheds Needs Survey 2012 Report to

Congress. Environmental Protection Agency; 2016. Available:

699 https://www.epa.gov/sites/default/files/2015-

700 12/documents/cwns_2012_report_to_congress-508-opt.pdf

701 32. U.S. Census Bureau. Guidance on Using 2020 Island Areas Censuses Data. 20 Oct

702 2022 [cited 31 Mar 2023]. Available: https://www2.census.gov/programs-

703 surveys/decennial/2020/technical-documentation/island-areas-tech-docs/2020-iac-

704 guidance.pdf

705 33. U.S. Census Bureau. 2020 Island Areas Censuses (IAC) Demographic Profile

706 Summary File. U.S. Census Bureau; 2022 Oct. Available:

707 https://www2.census.gov/programs-surveys/decennial/2020/technical-

708 documentation/island-areas-tech-docs/demographic_profile/2020-iac-dpsf-

709 technical-documentation.pdf

710 34. U.S. Department of Housing and Urban Development and U.S. Census Bureau.

711 2013 AHS Definitions. In: 2013 AHS Definitions [Internet]. [cited 9 Feb 2023].

712 Available: https://www2.census.gov/programs-

713 surveys/ahs/2013/2013%20AHS%20Definitions.pdf

714 35. TIGER/Line® Shapefiles. [cited 25 Apr 2023]. Available:

715 https://www.census.gov/cgi-bin/geo/shapefiles/index.php

716 36. U.S. Census Bureau. Metropolitan and Micropolitan About. In: Metropolitan and

717 Micropolitan About [Internet]. 22 Nov 2021 [cited 9 Feb 2023]. Available:

718 https://www.census.gov/programs-surveys/metro-micro/about.html

719 37. Schwartz MB. 2007 American Community Survey: A Comparison to Selected

720 Housing and Financial Characteristics for the United States from the 2007 American

721 Housing Survey. Housing and Household Economic Statistics Division U.S. Census

Bureau; Available: https://www.huduser.gov/portal/datasets/ahs/acs-

ahscomparisonreport.pdf

38. U.S. Census Bureau. Poverty Thresholds. 12 Sep 2023 [cited 7 Dec 2023].

Available: https://www.census.gov/data/tables/time-series/demo/income-

poverty/historical-poverty-thresholds.html

39. U.S. Census Bureau. AHS 2013 National Public Use File. Available:

https://www2.census.gov/programs-

surveys/ahs/2013/AHS%202013%20National%20PUF%20v2.0%20CSV.zip

40. Indian Health Service; Office of Environmental Health and Engineering; Division of

Sanitation Facilities Construction. Annual Report To the Congress of the United

States On Sanitation Deficiency Levels for Indian Homes and Communities: Fiscal

Year 2019. Rockville (MD): Indian Health Service (US); 2020.

41. Environmental Protection Agency. EPA's Proposal: Justification for Requested

Revision, Addition or Deletion to Questions on the American Community Survey

(ACS). 2018. Available:

https://foiaonline.gov/foiaonline/action/public/submissionDetails?trackingNumber=E

PA-HQ-2018-010667&type=request

42. McMahan CS, Self S, Rennert L, Kalbaugh C, Kriebel D, Graves D, et al. COVID-19

wastewater epidemiology: a model to estimate infected populations. Lancet Planet

Health. 2021;5: e874–e881.

43. Duvallet C, Wu F, McElroy KA, Imakaev M, Endo N, Xiao A, et al. Nationwide

Trends in COVID-19 Cases and SARS-CoV-2 RNA Wastewater Concentrations in

the United States. ACS EST Water. 2022. doi:10.1021/acsestwater.1c00434

44. Medina CY, Kadonsky KF, Roman FA Jr, Tariqi AQ, Sinclair RG, D'Aoust PM, et al.

The need of an environmental justice approach for wastewater based epidemiology

for rural and disadvantaged communities: A review in California. Curr Opin Environ

Sci Health. 2022;27: 100348.

45. Haak L, Delic B, Li L, Guarin T, Mazurowski L, Dastjerdi NG, et al. Spatial and

temporal variability and data bias in wastewater surveillance of SARS-CoV-2 in a

sewer system. Sci Total Environ. 2022;805: 150390.

46. Naughton CC, Roman FA Jr, Alvarado AGF, Tariqi AQ, Deeming MA, Bibby K, et

al. Show us the Data: Global COVID-19 Wastewater Monitoring Efforts, Equity, and

Gaps. bioRxiv. medRxiv; 2021. doi:10.1101/2021.03.14.21253564

47. Adhikari S, Halden RU. Opportunities and limits of wastewater-based epidemiology

for tracking global health and attainment of UN sustainable development goals.

Environ Int. 2022;163: 107217.

48. Wolf J, Johnston RB, Ambelu A, Arnold BF, Bain R, Brauer M, et al. Burden of

disease attributable to unsafe drinking water, sanitation, and hygiene in domestic

settings: a global analysis for selected adverse health outcomes. Lancet. 2023;401:

2060–2071.

49. WHO/UNICEF Joint Monitoring Programme for Water Supply, Sanitation and

Hygiene. Progress on Household Drinking Water, Sanitation and Hygiene. WHO

and UNICEF; 2021.

50. Gerrity D, Papp K, Dickenson E, Ejjada M, Marti E, Quinones O, et al.

Characterizing the chemical and microbial fingerprint of unsheltered homelessness

in an urban watershed. Sci Total Environ. 2022;840: 156714.

51. Jakariya M, Ahmed F, Islam MA, Al Marzan A, Hasan MN, Hossain M, et al.

Wastewater-based epidemiological surveillance to monitor the prevalence of SARS-

CoV-2 in developing countries with onsite sanitation facilities. Environ Pollut.

2022;311: 119679.

52. Tokajian S, Moghnieh R, Salloum T, Arabaghian H, Alousi S, Moussa J, et al.

Extended-spectrum β-lactamase-producing Escherichia coli in wastewaters and

refugee camp in Lebanon. Future Microbiol. 2018;13: 81–95.

53. Gwenzi W. Wastewater, waste, and water-based epidemiology (WWW-BE): A novel

hypothesis and decision-support tool to unravel COVID-19 in low-income settings?

Sci Total Environ. 2022;806: 150680.

54. Gina Pocock, Leanne Coetzee, Janet Mans, Maureen Taylor, and Bettina Genthe.

Proof of Concept Study Application of wastewater-based surveillance to monitor

SARS-CoV-2 prevalence in South African communities. Water Research

Commission Report. 2020. Available: https://wrcwebsite.azurewebsites.net/wp-

content/uploads/mdocs/TT%20832-20%20final%20web.pdf

55. Barnes KG, Levy JI, Gauld J, Rigby J, Kanjerwa O, Uzzell CB, et al. Utilizing river

and wastewater as a SARS-CoV-2 surveillance tool in settings with limited formal

sewage systems. Nat Commun. 2023;14: 7883.

56. U.S. Department of Housing and Urban Development and U.S. Census Bureau.

Appendix B: Sample Design and Weighting: 2013. U.S. Department of Housing and

Urban Development and U.S. Census Bureau; 2013. Available:

https://www2.census.gov/programs-

790  surveys/ahs/2013/2013%20AHS%20National%20Sample%20Design%20and%20

791  Weighting.pdf

792 57. Onsite Sewage FAQ - Permitting. [cited 10 Feb 2023]. Available:

793  https://floridadep.gov/water/onsite-sewage/content/onsite-sewage-faq-permitting

794 58. Florida Department of Health. Septic System Owner's Guide. In: floridahealth.gov

795  [Internet]. [cited 10 Feb 2023]. Available:

796  https://www.floridahealth.gov/environmental-health/onsite-

797  sewage/_documents/septic-folder-8x11-link2.pdf

798 59. Utah Department of Environmental Quality Water Quality. Municipal Wastewater

799  Planning Program (MWPP). In: Municipal Wastewater Planning Program (MWPP)

800  [Internet]. 3 Feb 2023 [cited 10 Feb 2023]. Available: https://deq.utah.gov/water-

801  quality/municipal-wastewater-planning-program-mwpp

802 60. Campbell H. 2021 MWPP Survey. In: 2021 MWPP Survey [Internet]. 27 Oct 2021

803  [cited 10 Feb 2023]. Available: https://documents.deq.utah.gov/water-

804  quality/engineering/municipal-wastewater-planning-program/DWQ-2021-026722.pdf

805 61. Minnesota Pollution Control Agency. 2021 Wastewater Infrastructure Needs

806  Instructions and Frequently Asked Questions (FAQ). Apr 2021 [cited 31 Mar 2023].

807  Available: https://www.pca.state.mn.us/sites/default/files/wq-wwtp3-06.pdf

808 62. Robinson C. 2017 SSTS Annual Report Subsurface Sewage Treatment Systems in

809  Minnesota. Minnesota Pollution Control Agency; 2018 Apr. Available:

810  https://www.pca.state.mn.us/sites/default/files/wq-wwists1-58.pdf

811 63. U.S. Environmental Protection Agency. 2012 Clean Watersheds Needs Survey.

812  Available: https://ordspub.epa.gov/ords/cwns2012/f?p=cwns2012:25

813 64. Weidhaas J, Aanderud ZT, Roper DK, VanDerslice J, Gaddis EB, Ostermiller J, et

814   al. Correlation of SARS-CoV-2 RNA in wastewater with COVID-19 disease burden

815   in sewersheds. Sci Total Environ. 2021;775: 145790.

816 65. U.s E. WaterSense Statistics and Facts. 11 May 2022 [cited 3 Apr 2023]. Available:

817   https://www.epa.gov/watersense/statistics-and-facts

Supporting information

S1 Appendix. Supplementary method

S2 Appendix. Supplementary analyses. Compilation of data, assessment of data completeness, sewer connectivity in U.S. Island Areas, supplementary figures, and supplementary tables.

S1 Appendix. Supplementary methods.

Dataset descriptions and data cleaning

U.S. Census Bureau American Housing Survey 2019 and 2021

The American Housing Survey (AHS) National and Metropolitan Public Use Files were downloaded from Ref. [12,22,23]. The sample design, weighting, and error estimation survey is described in Refs. [24–26]. Briefly, the AHS is a biannual survey that assesses housing characteristics. Two samples of households are chosen every survey cycle, the Integrated National Sample and the Independent Metropolitan Sample. The Integrated National Sample consists of nationally representative households and includes an oversample of each of 15 largest metropolitan areas. The Independent Metropolitan Sample consists of representative households in 20 large metropolitan areas that are representative of the next largest 50 metropolitan areas. Ten of the 20 large metropolitan areas are sampled every other survey cycle (10 metropolitan areas surveyed every 2 years). The Integrated National Sample in 2019 included 86,257 selected representative housing units, of which about 63,500 met the AHS definition of a housing unit and were able to be interviewed, and in 2021 included 95,295 selected representative housing units, of which about 64,100 were interviewed. The Integrated

Metropolitan Samples in 2019 and 2021 included around 3000 selected representative

housing units in each metropolitan area, of which around 1000-3000 were eligible and

were able to be interviewed (same criteria as in the Integrated National Sample).

Geographic data of each household available in the Public Use File included the census

division; core based statistical area (CBSA) for the 15 largest metropolitan areas and

the 20 large metropolitan areas in the metropolitan data. All other households are

classified as in "all other metropolitan areas" or "not in a metropolitan area" (all other

locations).

The race, ethnicity, and age of the householder (the person or one of the persons who

is an owner or renter of the unit) was used to categorize the race, ethnicity, and age of

the household. Due to the low numbers of households where the householder

responded as having more than one race, we focused on households where

householders identified as having a single race.

U.S. Census Bureau American Housing Survey National Sample 2013

The American Housing Survey National (2013) Use File was downloaded from Ref. [39].

The sample design, weighting, and error estimation survey is described in Ref [56]. The

National Sample included 84,400 selected housing units representative nationally and

including a supplemental sample of 15,553 housing units in the Chicago, Detroit, New

York City, Northern New Jersey, and Philadelphia metropolitan statistical areas, of

which about 71,600 met the definition of a housing unit and were able to be interviewed.

Geographic data of each household available in the Public Use File included the census

division; census region; central city, urban, or rural status; and select metropolitan area

codes. All other housing units are classified into "not in a metropolitan area" or had their

code suppressed for confidentiality [34].

Florida Department of Health onsite sewage treatment and disposal

system inspections

The locations of onsite sewage treatment and disposal systems inspected by Florida

Department of Health reported June 2012 was downloaded from Ref. [29]. As of 2022,

an inspection permit is required for construction [57] and inspection is recommended

every 3 to 5 years after [58]. The number of systems inspected that are active at the

latest inspection (after removing repeat inspections) is 601,430. The years of inspection

are primarily 1998-2011, with some outliers as early as 1900, which may be old septic

systems or typos, and as late as 2200, which are likely typos.

Utah Municipal Wastewater Planning Program (MWPP) Survey 2021

Data from the Utah Municipal Wastewater Planning Survey was provided by the Utah

Department of Environment Quality, Division of Water Quality. The dataset includes an

estimate of the population receiving collection by each utility that owns or operates a

sanitary sewerage system and its latitude and longitude. The data gathered from

treatment plant operators and participation is reported as mandatory [59], although only

71% of contacts responded [60]. We filtered the data to include any facility that has

collection (this included collection only, treatment and collection, or small lagoons which

are often collection and treatment). Manual correction of facility longitude and latitude is

described in the supplementary methods.

Manual editing of facility longitude and latitude was done for 5 facilities' that did not fall

within the state of Utah when combining with the 2021 TIGER shapefile (BRIAN HEAD

TOWN, CANYONG LAND IMPROVEMENT DISTRICT, HILL AFB (AMERICAN

WATER), JORDANELLE SSD, PLAIN CITY) and 1 facility that did not report a latitude

and longitude (PANGUITCH LAKE S. S. D.). For these facilities, the latitude and

longitude were manually edited to be that of the location found in a Google Maps search

of the facility name. If the Google Maps search found a city, then an arbitrarily chosen

location in that city was used to obtain a latitude and longitude.

Minnesota Wastewater Infrastructure Needs Survey (WINS) 2021

Data for the Minnesota Wastewater Infrastructure Needs Survey [61] was provided by

the Minnesota Pollution Control Agency. Communities (places and townships)

responded to the question "Does your community have a collection system?" with "yes",

"no", or no response. The survey did not receive a response from every community in

Minnesota.

The Census Designated Place (CDP) name was obtained from the reported community

name with manual changes reported in **Table S 5**.

Minnesota Subsurface Sewage Treatment Systems 2017

The Minnesota Subsurface Sewage Treatment Systems (SSTS) dataset reported the

number of SSTS by local government units with a known SSTS program. Data was

taken from Ref. [62].

Two hundred and eleven out of 2018 known SSTS programs submitted an annual

report. Not all SSTS programs were able to be identified and contacts were not always

provided. Fourteen reported having zero SSTS within their jurisdiction despite permitting

SSTS in 2017. Data were reported to sometimes be estimates.

U.S. Environmental Protection Agency Clean Watersheds Needs Survey 2012

Data for the most recent Environmental Protection Agency (EPA) Clean Watersheds Needs

Survey (CWNS) survey (2012) [31] was downloaded from [63] and extracted from a Microsoft

Access database file to csv files using the mdbtools package

(https://github.com/mdbtools/mdbtools). The survey is administered by the EPA and the States

to assess funding needs for unfunded capital costs of treatment works projects that (1) address

a water quality or related public health problem existing as of January 1, 2012, or is expected to

occur within the next 20 years and (2) meet the CWNS documentation criteria. CWNS

documentation criteria includes (1) description and location of the problem, (2) site-specific

solution to the problem, and (3) detailed cost information for implementing the solution. Data is

collected about (1) publicly owned wastewater collection and treatment facilities, (2) combined

sewer overflow control facilities, (3) stormwater management activities, and (4) decentralized

wastewater treatment facilities. Data is collected on (a) estimated needs, costs and technical

information, (b) facility location and contact information, and (c) facility population served, flow,

effluent, and unit process information. Supporting documentation was required for the entered

data with some exceptions (for instance small communities could use a more simple form), and

quality control was performed by reviewers to compare documentation with data entered in the

system to ensure consistency of technical and cost data [30]. Data entry occurred from January

to December 2012.

There was large variability in the level of effort and resources that each state put into the survey.

New York, California, Florida, New Jersey, Maryland, Iowa, Minnesota, and Michigan likely

invested heavily in participating in the survey or already had similar state-level data collection

systems in place for getting comprehensive responses [29,61,62].

Facilities not included in the dataset are those in South Carolina, the Northern Mariana Islands,

and American Samoa as they did not participate in the survey; facilities whose projects did not

have documented solutions or cost estimates; privately owned wastewater facilities that serve

privately owned industrial facilities, military installations, national parks, or other federal facilities;

facilities on tribal lands and in Alaskan Native Villages, which were separately surveyed by the

Indian Health Service; and facilities with projects that had existing funding prior to January 1,

2012. Additionally, larger facilities were more likely to be captured by the survey due to the

resource requirements for obtaining the right documentation. To address this bias, the survey

had decreased documentation needs for small communities (population <=10,000); despite this,

the small community facilities were still underrepresented (pg. 2 of Ref. [31]).

We filtered the facilities to the current facilities of type "collection: combined sewers" or

"collection: separate sewers". The population receiving collection is taken from the reported total

number of residents who are connected to a sewer system which empties into a treatment plant.

This does not include non-resident populations, populations served by acceptable decentralized

wastewater treatment systems, or populations connected to sewers that do not discharge to a

treatment plant.

U.S. Census Bureau Island Areas Decennial Survey 2020

The Island Areas Decennial Survey surveys housing, social, and economic information

for all housing units in Guam, the Northern Mariana Islands, the Virgin Islands, and

American Samoa [33]. Data of general demographic characteristics (DP01), selected

economic characteristics (DP03), and selected housing characteristics (DP04) were

downloaded from https://data.census.gov/. Housing characteristics included data for

both occupied and vacant housing units.

U.S. Census Bureau American Community Survey

The American Community Survey (ACS) is a monthly survey of a representative

subsample of addresses in the US [27] on characteristics of populations and

households. The ACS 5 year estimates (average over 5 years, less subject to

fluctuations) for selected social characteristics (DP02), selected economic

characteristics (DP03), and demographic and housing estimates (DP05) stratified

county or county subdivision for select years were downloaded from

https://data.census.gov/.

Data analysis and visualization

Florida Department of Health onsite sewage treatment and disposal

system inspections dataset analysis

983 The number of septic tanks in each county subdivision was calculated as the number of septic

984 permits whose latitude and longitude occurred inside the boundaries of each county subdivision

985 according to the 2012 U.S. Census Bureau county subdivision TIGER shapefile. To calculate

986 the fraction of a county subdivision connected to septic tanks, the number of septic tanks in

987 each county subdivision was divided by the number of households in the county subdivision as

988 reported by the 2012 American Community Survey 5-year estimate. Only county subdivisions

989 with at least 5 households and 20 population size were used.

991 **Utah Municipal Wastewater Planning Program Survey analysis**

992 The population in each county receiving collection was calculated by summing the population

993 receiving collection across all reported facilities in that county. The population in each county

994 subdivision receiving collection was calculated by summing the population receiving collection

995 across all reported facilities whose latitude and longitude fell within that county subdivision's

996 boundaries from the 2021 TIGER shapefile. To calculate the fraction of a county or county

997 subdivision receiving sewer collection, the population receiving collection was divided by the

998 population size of each county or county subdivision reported by the 2021 American Community

999 Survey 5-year estimate. Only county subdivisions with at least 5 households and 20 population

1000 size were used.

1001

1002 **U.S. Environmental Protection Agency Clean Watersheds Needs**

1003 **Survey analysis**

1004 To compare the EPA CWNS and AHS data in Core Based Statistical Areas (CBSA), counties

1005 were categorized into CBSAs using the February 2013 U.S. Census Bureau file. To calculate

1006 the fraction of a county receiving collection, the population size receiving collection in that

1007 county was divided by the population size of each county reported by the 2012 American

1008   Community Survey 5-year estimate. We flagged counties that had no facilities reported, which

1009   could either be because there were no facilities in that county or because of a lack of reporting,

1010   and counties that had >100% sewer connectivity by population (potentially due to misreporting

1011   or collection occurring across multiple counties) were set to 100% connectivity.

1012

1013 U.S. Census Bureau Island Areas Decennial Survey analysis

1014   To calculate the fraction of a Census Designated Place (CDP) connected to sewers, the number

1015   of households connected to sewers was divided by the total number of households in a CDP.

1016   Only CDPs with at least 5 households and 20 population size were used.

1017

1018 Generation of maps

1019   TIGER shapefiles from the U.S. Census Bureau in the closest matching year to the dataset

1020   were used for generating maps. Maps were generated using the geopandas package (version

1021   0.12.2) in Python (version 3.11.0) and were displayed in the Mercator projection.

1022

1023 Simulations of interacting populations and wastewater

1024 sampling

1025   Two interacting populations are modeled in a compartmental model with susceptible, infected,

1026   and recovered (SIR) compartments for each population. The number of susceptible, infected,

1027   and recovered individuals in population A (S_A, I_A, and R_A, respectively) and in population B (S_B,

1028 I_B, and R_B, respectively) are described by:

1029

1030
$$\frac{dS_A}{dt} = -\beta_A(1-\varepsilon)I_A\frac{S_A}{N_A} - \beta_B\varepsilon I_B\frac{S_A}{N_A}$$

1031
$$\frac{dS_B}{dt} = -\beta_B(1-\varepsilon)I_B\frac{S_B}{N_B} - \beta_A\varepsilon I_A\frac{S_B}{N_B}$$

1032
$$\frac{dI_A}{dt} = \beta_A(1-\varepsilon)I_A\frac{S_A}{N_A} + \beta_B\varepsilon I_B\frac{S_A}{N_A} - \gamma_I I_A$$

1033
$$\frac{dI_B}{dt} = \beta_B(1-\varepsilon)I_B\frac{S_B}{N_B} + \beta_A\varepsilon I_A\frac{S_B}{N_B} - \gamma_I I_B$$

1034
$$\frac{dR_A}{dt} = \gamma_I I_A$$

1035
$$\frac{dR_B}{dt} = \gamma_I I_B$$

1036

where N_A and N_B are the number of individuals in populations A and B respectively, β_A and β_B

are the overall contact rates of populations A and B respectively times the probability of infection

given contact, γ_I is the rate of recovery. ε is the fraction of the total contacts made by individuals

in a population that occurs with individuals in the other population and it describes the

interaction strength between the two populations. ε can take on values between 0 and 1, where

$\varepsilon=0$ indicates that all interactions occur within each population and none between populations,

and $\varepsilon=1$ indicates that all interactions occur between the populations and none within

populations. The base parameters are set at $N_A=N_B=5000$, $\gamma_I=0.18$ inverse days, $R_0^A = R_0^B =$

$\frac{\beta_A}{\gamma_I} = \frac{\beta_B}{\gamma_I} = 1.5$, which represent an outbreak similar to COVID-19, but the parameters are varied

in sensitivity analyses. The initial condition was a single infected individual in population A ($I_A=1$)

and all other individuals were susceptible ($S_A=N$-1, $S_B=N$, $I_B=0$, $R_A=0$, $R_B=0$). The differential

equations are solved deterministically using the integrate.odeint function in the scipy package

(version 1.6.2) in Python using a timestep of 0.1 days.

1050

1051   The concentration C of the pathogen in wastewater is given by

1052

1053
$$C = \frac{n_A + n_B}{v_A + v_B}$$

1054

1055   where n_A and n_B are the number of copies of pathogen genetic material (DNA or RNA) shed per

1056   day by population A and B, respectively, into the sampled wastewater and v_A and v_B are the

1057   volume of wastewater produced per day by population A and B, respectively, into the sampled

1058   wastewater.

1059

1060   The number of copies of pathogen genetic material, n_A and n_B, shed per day by population A

1061   and B, respectively, into the sampled wastewater is given by

1062

1063 $$n_A = n I_A f_A$$

1064 $$n_B = n I_B f_B$$

1065

1066   Where n is the number of copies of pathogen genetic material shed per infected person per day

1067   (assumed to be the same for all infected individuals) and f_A and f_B are the fraction of total shed

1068   pathogen genetic material by population A and B respectively that is sampled by wastewater

1069   (determined by the sewer connectivity of each of the populations).

1070

1071   The volume of wastewater, v_A and v_B, produced per day by population A and B, respectively,

1072   into the sampled wastewater is given by

1073 $$v_A = v N_A f_A$$

1074 $$v_B = v N_B f_B$$

1075

1076   where v is the volume of wastewater produced per person per day (assumed to be the same for

1077   all individuals). The value used for n was $5 \times 10^{6.7}$ gene copies/person/day (estimate taken from

1078   Ref. [64] by multiplying $10^{4.7}$ gene copies/mL feces by 500 mL feces/day/person) and for v was

1079   20 liters wastewater/person/day (per person daily wastewater usage approximated from per

1080   household daily wastewater usage from Ref. [65] assuming 4 people per household).

1081

In addition to the assumptions mentioned above, the model makes the following additional

assumptions: The probability of an infected individual shedding pathogen genetic material to

wastewater is 1. Given that there is pathogen genetic material present in wastewater, the

probability of detecting it is 1. Shedding of pathogen genetic material occurs only when an

individual is infected. There are no stormwater contributions to wastewater to dilute the

pathogen concentration.

1088

1089   To calculate the time of the maximum number of infections in a population and the time of the

1090   maximum concentration of pathogen gene content in wastewater data, we numerically

1091   calculated the time derivative at the midpoints of the time points as the difference between

1092   consecutive datapoints divided by the time interval and took the first time point at which the

1093   derivative crossed 0.

S2 Appendix. Supplementary analyses.

Compilation of data and assessment of data completeness

As the AHS does not provide finer scale geographic information than the census

division for households outside of the 35 selected large metropolitan areas, we used the

American Community Survey, which reports the percentage of occupied housing units

lacking complete plumbing facilities by county and which allowed us to define the lower

bound on the percentage of the population not connected to sewers. Additionally, we

identified several state-level datasets, including locations of permitted septic tanks in

Florida, sewer locations and populations served in Utah and Minnesota, and the 2012

EPA Clean Watersheds Needs Survey of sewer locations and populations served for 49

states (South Carolina did not participate). However, many of these state-level datasets

are incomplete or potentially biased and thus only allow qualitative analyses. Finally, the

2020 US Census' Island Areas Decennial survey provided information the percentage of

households connected to sewer by Census Designated Place in the island areas of

American Samoa, Guam, the Northern Mariana Islands, and the Virgin Islands (see

Error! Reference source not found. and **Table S 4** for a summary of datasets).

1110

We checked the consistency of the Florida Department of Health onsite sewage

treatment and disposal systems inspections data reported in 2012 with other sources.

We noted that an estimated 7% of households across the state were connected to

septic tanks, but that this was substantially less than the Florida Department of

Environmental Protection's estimate that approximately one third of Florida's population

uses septic tanks [28] even after accounting for potential correlations of septic usage

with household size (approximate increase in household size of 10% for a 50% increase

in septic connectivity (**Fig S 7**) cannot explain the almost 5 time difference in septic

connectivity across the two sources), suggesting that more households may be on

septic tanks, and thus less on sewer systems. This discrepancy may exist because not

every septic tank had a permit recorded in the state system, because there were

changes since 2012, or because of uncertainty in the Florida Department of

Environmental Protection's estimate. Thus, we focused our analysis on relative

comparisons across county subdivisions (Fig 4).

1125

The 2012 EPA Clean Watersheds Needs Survey dataset includes voluntary

submissions of the locations of publicly owned wastewater collection and treatment

facilities and their estimated population served. The following states had among the

highest reported needs and survey participation level (pg. 6 and Table A-1 of Ref. [31])

and were the focus of our analyses: New York, California, Florida, New Jersey,

Maryland, Iowa, Minnesota, and Michigan. Comparison of Florida in the EPA and

Florida Department of Public Health datasets showed consistent spatial trends (less

connection to sewers and more connection to septic systems in the panhandle and

inland) (Fig 4 and Fig S *11*). Additionally, in both datasets, counties with more

connection to sewers are more Asian (EPA: Pearson r = 0.53 (p = 4.2×10^{-4}), Florida

Department of Public Health: Pearson r = 0.31 (p = 3.2×10^{-7})). Comparison of

Minnesota in the EPA dataset, Minnesota Wastewater Infrastructure Needs Survey, and

Minnesota SSTS datasets also showed consistent spatial trends (more connection to

sewers and less connection to subsurface sewage treatment systems around

Minneapolis) (**Fig S 14**). We found a significant correlation between the AHS and EPA

datasets for the largest 35 metropolitan areas in the states with good data collection

(**Fig S 15**), despite deviations from a one-to-one correspondence suggesting overall

systematic biases in the data.

1144

1145   The 2021 Utah Municipal Wastewater Planning Survey estimated the populations receiving

1146   collection by municipal utilities that owned or operated a sanitary sewerage system. The

1147   purpose of the survey is to aid in overseeing and communicating with the wastewater industry in

1148   Utah. We noticed that these data may have not be quantitatively accurate but are likely

qualitatively informative. First, large swaths of the state appeared not to receive wastewater

collection (**Fig S 16**), but this may partly reflect that only 71% of contacts responding to the

survey [60]. Second, in some county subdivisions, >100% of the population appeared

connected to sewers, likely due to errors in estimation, differences in how combined treatment

and collection facilities reported population estimates, or collection populations resided in

neighboring county subdivisions. While the 5 county subdivisions with the highest percentages

of American Indian and Alaska Natives (~20-100%) were reported to not be connected to

sewers at all in this dataset, we believe this is because Indian reservations are not included in

the dataset (as they are under federal, not state jurisdiction). In fact, all 5 of these counties

(Uintah and Ouray, Casa del Eco Mesa-White Mesa, West Juab, Oljato, Blanding) include

Indian reservations. Additionally, the county subdivision with the highest percentage of Black or

African American; Native Hawaiian and Other Pacific Islander; or Hispanic are in county

subdivisions has nobody receiving collection (**Fig S 17**); however, we cannot rule out that this is

due to underreporting from these county subdivisions.

The 2021 Minnesota Wastewater Infrastructure Needs Survey reported whether

communities have a collection system. The data from this survey was geographically

sparse, as communities (the level of aggregation of the dataset) often directly mapped

to a census designated place (CDP), and CDPs do not cover the entire state (**Fig S

14**a). Consequently, the sewer connectivity of the rural areas between census

designated places could not be determined from this dataset. Additionally, not all

communities responded to the survey. Of the 523 communities that participated in the

survey, all but 11 had a collection system. The geographic distribution of sewer

connectivity qualitatively matched that of the 2017 Subsurface Sewage Treatment

Systems (SSTS) Annual Report of the number of reported subsurface sewage

1174 treatment systems (septic tanks) by county in Minnesota (**Fig S 14**b); however, 7 out of

1175 the 218 contacted local government units did not respond, 14 reported having zero

1176 SSTS within their jurisdiction despite permitting SSTS and not all SSTS programs were

1177 able to be identified to be contacted.

1178

1179 Given the data incompleteness and biases for the Utah and Minnesota datasets, we

1180 excluded them from the analyses in the main text.

1181

Sewer connectivity in the U.S. Island Areas

From the 2020 US Census Island Areas Decennial Survey of household characteristics,

the overall levels of household sewer connection in the island areas were lower than in

the states (**Fig S 18**). In Guam, the Northern Mariana Islands, the Virgin Islands, and

American Samoa, the overall connection to sewer systems across each island was

65%, 51%, 66%, and 52% of households, respectively (compared to an overall 83%

connection across mainland US). To ask whether there was strong spatial variability

across each island, we looked at the median household connection by Census

Designated Place (CDP). A median of 64%, 32%, 70%, and 13% households by CDP in

Guam, the Northern Mariana Islands, the Virgin Islands, and American Samoa,

respectively, were connected to sewers. This suggests that sewer connectivity is

particularly spatially concentrated in the Northern Mariana Islands and American Samoa

(difference in median household connection by CDP and overall island connection),

although all island areas showed some spatial variability in connectivity when visually

assessing maps (**Fig S 19**). The only significant correlation of sewer connectivity with

demographic or economic characteristics was in American Samoa, where CDPs with a

higher percentage of Black or African Americans were more connected to sewers

(Pearson r = 0.38, q-value = 0.01).

Supplementary figures

Fig S 1: **Percentage of households connected to various wastewater processing systems.**

The number of households sampled (n) for each category are also shown. Data are from 2021

American Housing Survey.

Fig S 2: **The percentage of households connected to sewers in each census division by demographic and economic characteristics (same as Fig 2) including the number of sampled households in each category.**

1209

1210 **Fig S 3: The percentage of households connected to sewers by urban and rural**

1211 **categorizations and by demographic and economic characteristics (Fig 3) including the**

1212 **number of sampled households in each category.**

1213

1214   **Fig S 4*: *Top 35 metropolitan areas' connectivity to public sewers by demographics.** Data are from the 2019 and 2021 U.S.

1215   Census American Housing Survey [12,23]. The number of sampled households for each category can be found in the Github

1216   repository in outputs/ahs_sewer_connectivity_by_cbsa.csv in the "num_observations" column.

Percent of occupied housing units lacking complete plumbing facilities by county

1217

1218 **Fig S 5: Percent of occupied housing units lacking complete plumbing facilities by**

1219 **county.** Data are from the 2021 U.S. Census American Community Survey.

1220

Fig S 6: Comparison of percentage of households not on septic across different geographies in Florida. Data are from the Florida Department of Health septic tank inspection permits reported in 2012. (a) Map of county subdivision colored by whether it is a Metropolitan Statistical Area (blue), Micropolitan Statistical Area (orange), or Rural Area (neither metropolitan not micropolitan, green). (b) Percentage of households not on septic in county subdivisions stratified by whether the county subdivision is a Metropolitan Statistical Area, Micropolitan Statistical Area, or Rural Area. County subdivisions that are Metropolitan Statistical Areas have a higher percentage of households not on sewer than those that are Micropolitan Statistical Areas, but the effect is not significant (p = 0.052). County subdivisions that are Micropolitan Statistical Areas have a higher percentage of households not on sewer than those that are Rural Areas, but the effect is not significant (p = 0.37). (c) Map of county subdivisions colored by whether they are always included in references to the Florida panhandle. Orange: county

subdivisions in the panhandle. Blue: county subdivisions not in the panhandle. (d) Percentage of

households not on septic in county subdivisions stratified by whether the county subdivision is in

the panhandle. Each point represents a county subdivision. County subdivisions in the

panhandle have a significantly lower percentage of households not on septic ($p < 10^{-9}$). (e) Map

of county subdivisions colored by whether they have a coastline. Orange: county subdivision

with coastline. Blue: county subdivisions without coastline. (f) Percentage of households not on

septic in county subdivisions stratified by coastal and not coastal county subdivisions. Each

point represents a county subdivision. County subdivisions not on the coast have a significantly

higher fraction of households on septic ($p<10^{-9}$).

1242

1243

1244

1245   **Fig S 7***: **Correlation of percentage of households in county subdivisions of Florida** *not*

1246   **connected to septic tanks with demographic and economic factors.** Data are from the

1247   Florida Department of Health septic tank inspection permits reported in 2012. Each point

1248   represents a county subdivision. Only county subdivisions with at least 5 households and 20

1249   population size were included.

1250

County subdivisions in metropolitan statistical areas

1251

Fig S 8: Correlation of percentage of households in county subdivisions of Florida in metropolitan statistical areas not connected to septic tanks with demographic and economic factors. Data are from the Florida Department of Health septic tank inspection

1255   permits reported in 2012. Each point represents a county subdivision. Only metropolitan county

1256   subdivisions with at least 5 households and 20 population size were included.

1257

Fig S 9: Correlation of percentage of households in county subdivisions of Florida in

micropolitan statistical areas not connected to septic tanks with demographic and

economic factors. Each point represents a county subdivision. Data are from the Florida

1261   Department of Health septic tank inspection permits reported in 2012. Only micropolitan county

1262   subdivisions with at least 5 households and 20 population size were included.

County subdivisions in rural areas

1263

Fig S 10: Correlation of percentage of households in county subdivisions of Florida in rural areas not connected to septic tanks with demographic and economic factors. Data are from the Florida Department of Health septic tank inspection permits reported in 2012. Each

1267   point represents a county subdivision. Only rural county subdivisions with at least 5 households

1268   and 20 population size were included.

1269
1270

1271

Fig S 11: Fraction of present resident population receiving collection by county in 8

states. Data are from the 2012 EPA Clean Watsheds Needs Survey. (a) Fraction shown on a

continuous scale. (b) Fraction above and below 0.2. Only states that had more comprehensive

responses in the survey are shown. Counties that did not report any data in the survey are

shown in gray. Counties than report more than 100% of the present resident population

receiving collection are shown in (a) as 1 and in (b) as ≥0.2.

1278

1279

1280

1281 **Fig S 12: Sensitivity analysis of Fig 5c from main text**. The effect on the time between

1282 population B peak and wastewater peak of varying the (a) population size of the two populations

1283 (N_A, N_B), (b) recovery rate (γ_I in units of inverse days), and (c) basic reproduction number of the

1284   two populations (R_0^A, R_0^B) across a range of values. Unless where indicated, the parameters

1285   were $N_A=N_B=5000$, $\gamma_I=0.18$ days^{-1}, and R_0^A, $R_0^B=1.5$.

1286
1287   **Fig S 13: Correlation between householder age and household income in the American**

1288   **Housing Survey dataset by census division.**

1289

Fig S 14: Sewer connectivity in Minnesota. (a) Map showing whether a community in

Minnesota (mapped to Census Designated Place) reported having a collection system in the

2021 Minnesota Wastewater Infrastructure Needs Survey. Almost all communities mapped to

Census Designated Places. (b) Percentage of households without subsurface sewage treatment

systems (i.e. without septic tanks, suggesting sewered) by county reported in the 2017

Subsurface Sewage Treatment Systems in Minnesota Annual Report [62]. Counties reporting

more than 100% of households having subsurface sewage treatment systems are set at 100%

(shown as 0% without subsurface sewage treatment systems in the map).

1298

Fig S 15*: Comparison of the 2012 EPA Clean Watersheds Needs Survey (CWNS) and the

2019 and 2021 U.S. Census American Housing Survey (AHS) datasets in core based

statistical areas (CBSA). Data from the EPA CWNS are of the fraction of residents in the

CBSA receiving sewage collection. Data from the AHS are of the fraction of households in the

CBSA on sewer. Only CBSAs that were oversampled in the AHS data and were in states that

had more comprehensive responses in the EPA CWNS were included (see Methods). The

dashed black line shows y=x as a reference.

% of population receiving collection by county subdivision in Utah

1306

1307 **Fig S 16: The percentage of the population that received wastewater collection by a**

1308 **municipal utility in Utah by county subdivision of Utah.** Data are from the 2021 Utah

1309 Municipal Wastewater Planning Survey. All values less than 0 are displayed as 0. County

1310 subdivisions with fewer than 5 households or 20 population size are displayed in white. Note

1311 that Indian reservations were not surveyed in this dataset.

1312

Fig S 17: Correlation of fraction households in county subdivisions of Utah with

wastewater collection with demographic and economic factors. Data are from the 2021

Utah Municipal Wastewater Planning Survey. Note that Indian reservations were not surveyed

1316   in this dataset. Values of the fraction of the county subdivision receiving collection greater than

1317   1 were set to 1.

1318

1319

Fig S 18: Fraction of households by Census Designated Place that are connected to

sewers, septic tanks, or other forms of sewage disposal in the U.S. Island Areas. (a)

American Samoa, (b) Guam, (c) the Northern Mariana Islands, and (d) the Virgin Islands. Data

are from the 2020 U.S. Census Island Areas Decennial Survey.

1324

1325 **Fig S 19: Map of fraction of households connected to sewers by Census Designated**

1326 **Place.** (a) American Samoa, (b) Guam, (c) the Northern Mariana Islands, and (d) the Virgin

1327 Islands. Data are from the 2020 U.S. Census Island Areas Decennial Survey.

1328

1329

Supplementary tables

Table S 1: Counties in the US with >=5% of occupied housing units lacking complete plumbing facilities. Data are from the 2021 U.S. Census American Community Survey.

	County, State	Percent of occupied housing units lacking complete plumbing facilities
0	Yukon-Koyukuk Census Area, Alaska	36.1
1	Bethel Census Area, Alaska	31.1
2	Kusilvak Census Area, Alaska	25.3
3	Nome Census Area, Alaska	24.5
4	Northwest Arctic Borough, Alaska	20
5	Denali Borough, Alaska	18.1
6	Lake and Peninsula Borough, Alaska	13.8
7	Apache County, Arizona	12.8
8	Dillingham Census Area, Alaska	12
9	Southeast Fairbanks Census Area, Alaska	11.1
10	North Slope Borough, Alaska	10.8
11	Haines Borough, Alaska	10.5
12	Copper River Census Area, Alaska	10.1
13	San Juan County, Utah	10
14	Adams County, Indiana	9.7

15	McKinley County, New Mexico	9.6
16	Oglala Lakota County, South Dakota	9.2
17	Hoonah-Angoon Census Area, Alaska	7.6
18	Mora County, New Mexico	7.6
19	Navajo County, Arizona	7
20	Prince of Wales-Hyder Census Area, Alaska	6.7
21	Lawrence County, Mississippi	6.2
22	Catahoula Parish, Louisiana	6.1
23	Hudspeth County, Texas	6.1
24	Fairbanks North Star Borough, Alaska	5.6
25	Salinas Municipio, Puerto Rico	5.1
26	Yakutat City and Borough, Alaska	5.1

1333

1334 **Table S 2: Correlation of the percentage of households in a county connected to**

1335 **public sewers with different demographic or economic factors by state; same as**

1336 **Table 3**, **but includeing all state-variable pairs.**

Demographic or economic variable	State	Correlation of variable with % of households by county connected to public	Significance (q-value)

sewers (Pearson correlation

coefficient)

	MN	0.29	0.49
	IA	-0.24	1
	NJ	0.34	1
	MD	-0.32	1
Average household size	NY	0.19	1
	FL	-0.12	1
	CA	0.25	1
	MI	0.01	1
	MN	-0.43	<0.01
	MI	-0.41	0.01
	CA	-0.41	0.13
	NY	-0.37	0.26
Median age	MD	0.17	1
	IA	-0.08	1
	NJ	-0.42	1
	FL	-0.05	1
	MN	0.44	<0.01
	FL	0.48	<0.01
Median income	CA	0.49	<0.01
	NJ	-0.27	1
	IA	-0.11	1

	MD	-0.17	1
	MI	0.21	1
	NY	0.23	1
	FL	0.36	0.26
	NY	0.31	1
	NJ	0.51	1
	IA	0.15	1
Percent Hispanic	MD	-0.29	1
	MI	0.24	1
	CA	0.25	1
	MN	0.19	1
	CA	-0.50	<0.01
	FL	-0.29	1
	MN	-0.14	1
Percent one race and	NJ	0.09	1
American Indian and	NY	0.03	1
Alaska Native	MI	-0.02	1
	MD	-0.04	1
	IA	-0.01	1
	MN	0.58	<0.001
Percent one race and	MI	0.51	<0.001
Asian	NY	0.55	<0.001
	CA	0.56	<0.001

	FL	0.53	<0.001
	IA	0.22	1
	NJ	0.25	1
	MD	-0.02	1
Percent one race and Black or African American	MN	0.62	<0.001
	MI	0.59	<0.001
	NY	0.55	<0.001
	IA	0.38	<0.01
	CA	0.46	0.02
	NJ	0.39	1
	MD	-0.05	1
	FL	0.01	1
Percent one race and Native Hawaiian and other Pacific Islander	IA	0.26	0.85
	MN	-0.18	1
	MI	-0.15	1
	MD	0.28	1
	NY	0.10	1
	CA	0.07	1
	NJ	-0.07	1
	FL	-0.01	1
Percent one race and some other race	NJ	0.57	0.63
	CA	0.35	0.68
	MN	0.26	1

	NY	0.23	1
	IA	0.16	1
	MI	0.17	1
	MD	-0.28	1
	FL	0.03	1
	MI	-0.57	<0.001
	CA	-0.54	<0.001
	NY	-0.51	<0.01
Percent one race and	IA	-0.35	0.03
White	MN	-0.28	1
	NJ	-0.54	1
	FL	-0.08	1
	MD	0.08	1
	MI	-0.38	0.03
	NJ	0.51	1
	CA	-0.24	1
Percent uninsured (health	MN	-0.15	1
insurance)	NY	-0.10	1
	MD	-0.15	1
	FL	0.06	1
	IA	-0.03	1

1337   q-values above 1 are set to 1.

 Table S 3: Correlation of the fraction of households by county connected to public

 sewers and different demographic or economic factors, stratified by state and

 Metropolitan, Micropolitan, and Rural Statistical Areas. Only correlations that are significant

 (q-value<=0.05) are shown.

Demographic or economic variable	State	Metropolitan, Micropolitan, or Rural Statistical Area	Pearson r	q-value
Median income	FL	Micropolitan Statistical Area	0.98	9.8E-03
Percent Hispanic	FL	Metropolitan Statistical Area	0.53	1.8E-02
	MN	Metropolitan Statistical Area	0.72	2.3E-03
Percent one race and Asian	FL	Metropolitan Statistical Area	0.52	3.0E-02
Percent one race and Black or African American	MN	Metropolitan Statistical Area	0.73	1.1E-03
	MI	Metropolitan Statistical Area	0.67	1.4E-02
	MN	Metropolitan Statistical Area	-0.74	7.6E-04
Percent one race and White	MI	Metropolitan Statistical Area	-0.71	4.6E-03

| | Metropolitan Statistical | | |
| IA | Area | -0.69 | 5.0E-02 |

1342

1343

Table S 4: Additional datasets discussed in the Supplementary Information but excluded

from the main text due to data incompleteness and potential biases.

Dataset	Location	Year(s)	Description	Sampling method	Dataset completeness and potential biases	Ref.
Utah Municipal Wastewater Planning Survey	Utah	2021	This survey reported the population receiving collection by each municipal utility that owns or operates a sanitary sewerage system.	The survey completed was by treatment plant operators. Participation was reported in the survey information as being mandatory.	Of those contacted, 71% responded to the survey [60]. In some county subdivisions, >100% of the population appeared connected to sewers, likely due to errors in estimation, differences in how combined treatment and collection facilities reported population estimates, or collection populations residing in neighboring county subdivisions.	[60]
Minnesota Wastewater Infrastructure Needs Survey (WINS)	Minnesota	2021	Communities (places and townships) responded to the question "does your community have a collection system?" (yes/no/did not respond).	This was a voluntary survey.	Some communities (number unknown) did not respond to the survey.	[61]
Minnesota Subsurface Sewage Treatment Systems (SSTS)	Minnesota	2017	This dataset reports the number of subsurface sewage treatment systems (SSTS) in various local government units.	Surveys were sent to local government units with a known SSTS program. According to the SSTS Annual Report [62], 'some of the data is "hard" data, such as the reported number and types of permits issued. Other data is considered "soft" data, such as the reported best estimates provided to determine SSTS	Two hundred and eleven out of 218 known SSTS programs submitted an annual report. Not all SSTS programs were identified and contacts were not always provided. Fourteen reported having zero SSTS within their jurisdiction despite permitting SSTS in 2017.	[62]

				compliance rates and, sometimes, the total number of SSTS in each jurisdiction.'	

1347

1348 **Table S 5: Manual changes community names reported in the Minnesota**

1349 **Wastewater Infrastructure Needs Survey to be able to match with a U.S. Census**

1350 **Designated Place.**

Original name in dataset	Edited name for merging with Census Designated Place name	Note
St X (i.e. St Paul)	St. X (i.e. St. Paul)	
Plainview - Elgin Sanitary District	[Deleted]	There was another entry for Plainview (same zip code) with the same response for whether there was collection facility
Clear Lake - Clearwater Sewer Authority	Clear Lake	
Tower-Breitung Wastewater Board	Tower	
Birchwood	Birchwood Village	
Coleraine, Bovey, Taconite Joint Wastewater Commission	[Deleted]	Two entries for Coleraine with same zip code and similar population sizes

Glacial Lakes Sanitary Sewer and Water District	[Deleted]	Two entries for Spicer with same zip code and similar population sizes
Central Iron Range Sanitary Sewer District	[Deleted]	Two entries for Chisholm with same zip code and similar population sizes
Shorewood Park Sanitary District	[Deleted]	Two entries for Rush City with same zip code. Deleted the entry with the smaller population size.
Rich Prairie Water & Sewer District (Pierz/Genola)	Pierz	
Pine River Area Sanitary District	Pine River	
Moose Lake Windemere Area Sanitary Sewer District	[Deleted]	Two entries for Moose Lake with same zip code. Deleted the entry with no reported population
Alexandria Lakes Area Sanitary District	Alexandria	
North Long Lake Sewer District	Brainerd	

Serpent Lake Sanitary Sewer District	Crosby	
Blomkest/Svea Sewer Board	Blomkest	
GEM Sanitary District	Greenwald	
Chisago Lakes Joint Sewage Treatment Commission	Center City	
Duluth	[Deleted]	3 entries in Duluth - kept the one with the largest population size (Western Lake Superior Sanitary District (WLSSD))
Duluth – North Shore Sanitary District	[Deleted]	3 entries in Duluth - kept the one with the largest population size (Western Lake Superior Sanitary District (WLSSD))
MCES - Metro Council Env. Services	[Deleted]	Two entries in St. Paul. Deleted the entry with the smaller population size.

1351