Full title: Assessment of sewer connectivity in the United States and its implications for

equity in wastewater-based epidemiology

Short title: Sewer connectivity and equity in wastewater-based epidemiology

Authors: QinQin Yu[1], Scott W. Olesen[2], Claire Duvallet[2], Yonatan H. Grad[1*]

Affiliations:

1. Department of Immunology and Infectious Diseases, Harvard T. H. Chan School of

10   Public Health, Boston, Massachusetts, USA

11   2. Biobot Analytics, Inc., Cambridge, Massachusetts, USA

13   [*]Corresponding author

14   Email: ygrad@hsph.harvard.edu (YG)

Abstract

Wastewater-based epidemiology is a promising public health tool that can yield a more representative view of the population than case reporting. However, only about 80% of the U.S. population is connected to public sewers, and the characteristics of populations missed by wastewater-based epidemiology are unclear. To address this gap, we used publicly available datasets to assess sewer connectivity in the U.S. by location, demographic groups, and economic groups. Data from the U.S. Census' American Housing Survey revealed that sewer connectivity was lower than average for American Indian and Alaskan Native, White, non-Hispanic, older, not in poverty, and larger households, but smaller geographic scales revealed local variations from this national connectivity pattern. For example, data from the U.S. Environmental Protection Agency showed that sewer connectivity was positively correlated with income in Minnesota, Florida, and California. Data from the U.S. Census' American Community Survey and Environmental Protection Agency also revealed geographic areas with low sewer connectivity, such as Alaska, the Navajo Nation, Minnesota, Michigan, and Florida. However, with the exception of the U.S. Census data, there were inconsistencies across datasets. Using mathematical modeling to assess the impact of wastewater sampling inequities on inferences about epidemic trajectory at a local scale, we found that in some situations, even weak connections between communities may allow wastewater monitoring in one community to serve as a reliable proxy for an interacting community with no wastewater monitoring, when cases are widespread. A systematic, rigorous assessment of sewer connectivity will be important for ensuring an equitable and informed implementation of wastewater-based epidemiology as a public health monitoring system.

Introduction

Wastewater-based epidemiology (WBE) plays an important role in surveillance of SARS-CoV-2 [1,2], polio virus [3,4] and other pathogens [5,6] and has applications to monitoring a variety of other public health concerns [7], including opioid usage [8]. One proposed benefit of wastewater-based epidemiology is that wastewater data is more representative of the population than case reporting, which can be biased towards those with health-seeking behavior or access to healthcare. For example, the populations served by the Health and Human Services SARS-CoV-2 National Wastewater Surveillance System were more representative of the entire US's age distribution and Black and Hispanic populations than the vaccinated population [9].

While WBE offers convenient sampling of populations served by public sewers, about 20% of individuals in the US live in homes not connected to public sewers [10,11]. This includes those on decentralized wastewater systems and those with no wastewater treatment systems. The most common decentralized wastewater system is septic tanks [12], which collect and treat wastewater onsite, typically the yard of the home. Households without wastewater treatment systems may have outhouses or privies, chemical toilets, or no plumbing. Variability in sewer connectivity exists across the U.S., with more connectivity in urban areas than rural areas [10,13].

As with any emerging public health tools, it is important to ask to what extent the tool exacerbates or alleviates inequities. A few studies have evaluated equity of sewer connectivity on broad geographic scales. A 2017 study from Environmental Protection Agency Office of Water showed that households in the U.S. that earned less than the national median household income (MHI, $61,000) were almost 10% more likely to have a decentralized wastewater

system or no wastewater treatment system compared to households that earned more than the

MHI [14]. Additionally, this study found that as household income decreased, decentralized

wastewater system usage increased in Florida, Hawaii, and Delaware, but not in Rhode Island.

Based on the 2019 U.S. Census Bureau's American Housing Survey, rural areas are less

connected to sewers, and while the income in rural areas was lower than in metropolitan areas,

households connected to septic tanks were wealthier than those connected to sewers in both

rural and metropolitan areas [10].

However, we have an incomplete understanding of the factors associated with sewer

connectivity across the US and the implications for the interpretation of wastewater data. In this

study, we sought to address the following questions, focusing our analyses on the U.S.: (1) To

what extent is there demographic and economic inequity in sewer connectivity? (2) Which

geographic areas have low sewer connectivity? (3) What is the applicability of WBE data to

neighboring unsampled communities? To address the first question, we analyzed household-

level data of sewer connectivity stratified by geographic, demographic, and economic variables.

For the second question, we evaluated datasets aggregated at the county or county subdivision

levels to qualitatively identify geographic areas that have low sewer connectivity. To address the

third question, we used a mathematical model to simulate WBE in two interacting populations.

Methods

Dataset compilation

We used a convenience sample of publicly available datasets on sewer connectivity, septic

connectivity, or no plumbing at the household level. The types of data included household-level

data, locations of sewer systems and population served, and population served by or lacking

wastewater systems aggregated by county or county subdivision. All datasets are summarized

in **Table *1*** and described in more detail in the Supplementary Information.

Table 1: Datasets used in this study.

Dataset	Location	Year(s)	Description	Sampling method	Dataset completeness and potential biases	Ref.
American Housing Survey	US excluding Island areas	2013, 2019, 2021	This was a longitudinal survey of housing, demographic, and economic characteristics of approximately 60,000-80,000 representative households.	The sampling method was a representative sample of all U.S. households at the time of sample selection (most recently in 2015 and before that, in 1985). New housing units were added each survey cycle.	The survey excludes group quarters, businesses, hotels, and motels. Additionally, the survey was more likely to classify rural seasonal homes as vacant units, counted large cluster septic systems as public sewer, and was thought to lose new households built in rural areas due to its longitudinal design.	[11,15–19]
American Community Survey 5-year estimates	US excluding Island areas	2021	This was a monthly survey of characteristics of populations and households of a representative subsample of addresses in the US. The survey reports the number of households lacking complete plumbing facilities by geographic subdivision.	The survey was sent to a random sample of addresses in the U.S. every month. Each address was selected no more than once every 5 years.	The surveys were completed by self-response (although a subset of non-responses were followed up with personal visits). Responses were reviewed for completeness and questionnaires needing clarification were followed up by phone calls. The data was averaged over 5 years to reduce statistical fluctuations.	[20]
Florida Department of Health	Florida	Up to 2012	This dataset includes the onsite sewage treatment and disposal systems locations	The sample included onsite sewage treatment and disposal system permits recorded in the	As of 2022, a permit was required for construction and inspection was recommended every 3-5 years after. From this dataset, an estimated 7% of households	[22]

septic tank inspections			inspected by the Florida Department of Health reported in June 2012.	Florida Department of Health system.	were connected to septic tanks across the state. This was substantially less than the Florida Department of Environmental Protection's estimate that approximately one third of Florida's population used septic tanks [21].	
EPA Clean Watersheds Needs Survey	US	2012	This survey included voluntary submissions of the locations of publicly owned wastewater collection and treatment facilities and their estimated population served for the purposes to assess funding needs for treatment works projects.	The responses were coordinated by states.	This survey had voluntary responses and variable effort and resources that each state put into the survey. States that had the most comprehensive responses were New York, California, Florida, New Jersey, Maryland, Iowa, Minnesota and Michigan. Missing facilities included those in South Carolina, the Northern Mariana Islands, and American Samoa; facilities whose projects did not have documented solutions or cost estimates; privately owned wastewater facilities; facilities on tribal lands and Alaskan Native Villages; and facilities whose projects received funding from other sources. Small community facilities are thought to be underrepresented due to having less resources for completing the survey, but this was not quantified.	[23,24]
US Census Island Areas Decennial Survey	US Island Areas	2020	This was a survey of housing, social, and economic information of all housing units.	The survey was conducted via in person and phone interviews.	The COVID-19 pandemic impacted collection of data on group quarters [25] and the survey had higher nonresponse rates than previous years' surveys, particularly for Guam. Quality control checks were performed on responses.	[26]

90   The datasets are further described in the S1 Appendix and the dataset completeness and

91   potential biases are further assessed in the S2 Appendix. Additional datasets that are briefly

92 described in the main text but primarily analyzed in the Supplementary information are

93 summarized in **Table S *3***.

Calculation of weights, error bars, and summary statistics in American Housing Survey (AHS) data

All summary statistics were calculated by applying the weights reported in the AHS Public Use

File [19]. Because not every household was sampled and there was uneven sampling of

households across the U.S., the weight estimates the number of similar households that each

surveyed household represented. Error bars on summary statistics represent the middle 95% of

values calculated using the reported replicate weights (160 replicate weights per household).

Each summary statistic was calculated for each replicate separately before taking the middle

95% of values. Only categories with at least 5 households were used.

Analysis of sewer connectivity by urban or rural areas in AHS data

We used the most recent AHS National Sample with publicly available data on the urban or rural

location of the household (2013) which categorized households into being in a central city of a

Metropolitan Statistical Area (MSA), in an urban area of an MSA but not in a central city, in a

rural area of an MSA but not in a central city, in an urban outside of an MSA, or in a rural area

outside of an MSA [27]. Central cities are defined as having either (1) ≥250,000 population or at

least 100,000 people working within corporate limits, (2) ≥25,000 population, at least 75 jobs for

each 100 residents who were employed, and 60% or fewer of the city's resident workers

commuted to jobs outside, or (3) 15,000-25,000 population, at least one third the size of the

metropolitan statistical area's largest city, and met the two commuting requirements in (2).

Metropolitan Statistical Areas were defined as whole counties that have significant levels of

community and contiguous urban areas in common and includes rural areas. Urban areas were

defined as having ≥2500 people, with at least 1500 residing outside of institutional group

quarters. Rural areas were defined as those not in urban areas.

Analyses of geographic areas with low connectivity to sewer

Data of individual locations of septic tanks or sewer collection systems, when they existed, were

aggregated based on latitude and longitude into county or county subdivision boundaries as

defined by the U.S. Census' TIGER shapefiles of the corresponding year [28]. To calculate the

fraction of a geographic area connected to septic or sewer, the total population size or total

number of households in that geographic area was taken from the corresponding year in the

American Community Survey (ACS) 5-year estimate [20]. For county subdivision and census

designated place analyses, only areas with at least 5 households and 20 population size were

used for the analyses. Detailed data analyses of the state and island area datasets are

described in the Supplementary information.

Counties and county subdivisions were classified into Metropolitan statistical areas (at least one

urbanized area of 50,000 or more population), Micropolitan statistical areas (at least one urban

cluster of 10,000-50,000 population), or Rural areas (all other areas) using the U.S. Census

classification of the encompassing Core Based Statistical Area (core area containing a

substantial population nucleus together with adjacent territory that has a high degree of social

and economic integration with the core as measured by commuting ties) [29].

Demographic and economic variable correlations with sewer

or septic connectivity

To assess correlations of demographic and economic variables with sewer and septic

connectivity when aggregating at the county, county subdivision, or census designated place

levels, we used the ACS 5-year estimates from the corresponding year and geographic scale.

Only geographic subdivisions with at least 5 households and 20 population size were used. The

Pearson correlation coefficient between demographic or economic variables of interest and the

percentage of households in a geographic subdivision connected to septic tanks or sewer was

calculated using the stats.pearsonr function in the scipy package (version 1.10.1) in Python

(version 3.11.0). A Bonferroni correction was applied to correct for multiple hypothesis testing to

calculate the q values.

Simulations of interacting populations and wastewater

sampling

Two interacting populations (A and B) were modeled in a deterministic compartmental model

with susceptible, infected, and recovered (SIR) compartments for each population and a

fraction, ε, of cross-population contacts. The base parameters were set at population size

$N_A = N_B = 5000$, recovery rate $\gamma_I = 0.18$ inverse days (corresponding to an infectious period of 5.6

days), basic reproduction number $R_0^A = R_0^B = \frac{\beta_A}{\gamma_I} = \frac{\beta_B}{\gamma_I} = 1.5$ and were varied in sensitivity

analyses. The differential equations were solved using the integrate.odeint function in the scipy

package (version 1.6.2) in Python (version 3.11.0) using a timestep of 0.1 days.

Wastewater sampling was modeled by the number of copies of pathogen genetic material shed

per day by each population divided the volume of sampled wastewater produced per day by

each population. Both numerator and denominator depended on the fraction of waste produced

(including shed pathogen genetic material) by each population that is sampled by wastewater, f_A

and f_B. A detailed description of the simulations is provided in S1 Appendix.

Code availability

167 All code and data to reproduce the analyses and simulations are available at

168 https://github.com/gradlab/wastewater_equity.

Results

Is there inequity in who is sampled by wastewater?

To address this question, we used the American Housing Survey (AHS), a biannual longitudinal

survey of representative housing units within the U.S that reports household-level data of sewer

connectivity, demographic factors, and economic factors (**Table *1***). We focused our analyses on

geographic scales with a reliable sample size for statistical analyses (census division,

urban/rural at the national level, metropolitan statistical areas) [30].

According to the 2021 AHS, 83% of US households were connected to public sewers and 16%

to standard septic tanks (**Fig S *1***). Each of the other forms of sewage disposal (non-standard

septic tanks, chemical toilets, outhouses or privies, other, none, or not reported) was more than

an order of magnitude less common (**Fig S *1***). Across census divisions in 2021, the western US

was overall better connected to sewers than the eastern US, with the highest levels of sewer

connection in the Pacific (91.3% of households, CI: 90.8%-91.7%) and Mountain (87.6% of

households, CI: 86.8%-88.4%) divisions and the lowest levels of sewer connection in the New

England (72.9% of households, CI:72.1%-74.0%) and East South Central (74.9% of

households, CI: 73.7%-76.3%) divisions (**Fig *1***).

Fig 1: **The percentage of households connected to sewers in each census division.** Data

are from the 2021 American Housing Survey National Survey [11].

Across census divisions, households with an Asian, Black or African American, or Native

Hawaiian and Other Pacific Islander householder (the owner or renter of the unit) were on

average 14.0%, 13.1%, and 12.5% more connected to sewers, respectively, than the overall

census division; however, specifically in the Pacific census division, households with a Native

Hawaiian and Other Pacific Islander householder were 5.3% less connected to sewers than the

overall census division (**Fig 2**). Households with an American Indian and Alaska Native or White

householder were both on average 2.6% less connected to sewers than the overall census

division. Across census divisions, households with a Hispanic householder were on average

11.7% more connected to sewers whereas households with a non-Hispanic householder were

1.1% less connected to sewers than the overall census division.

Connection to sewers decreased with the age of the householder (**Fig 2**), with the exception

that households with a householder aged 75+ were better connected in most census divisions

than those with a householder aged 65-74 (79.7% vs 77.9% of households). Households with 1

person were on average 4.2% more connected whereas households with 2 people were on

average 3.6% less connected to sewers than the overall census division. There were no

consistent trends across census divisions for households with more than 2 people. Additionally,

households that were in poverty were better connected to sewers (average of 87.5% of

households across census divisions) than households not in poverty (82.7% of households

across census divisions; **Fig 2**). The threshold used to determine whether a household was in

poverty depended on family size and age of persons in the household; for example, in 2021 the

threshold for 1 person under 65 years is $14,097 and for 4 persons with two children is $27,479

[31].

Fig 2*:* **The percentage of households connected to sewers in each census division by**

demographic and economic characteristics. Data are from the 2021 American Housing

Survey National Survey [11].

We sought to describe how the differences in sewer connection between demographic and

economic groups observed at the census division level was affected by household location in

urban or rural areas (see Methods). The qualitative trends observed at the census division level

for race, ethnicity, age, household size, and poverty level were largely preserved across

national urban and rural areas (**Fig 3**). One notable exception was that lower sewer connectivity

for households with an American Indian and Alaska Native or Hawaiian and Other Pacific

Islander householder compared to those with a Black or Asian householder was only observed

in locations inside of an Metropolitan Statistical Area (MSA) but outside of a central city (both

urban and rural). In central cities and locations outside of an MSA (both urban and rural),

households with an American Indian and Alaska Native or Hawaiian and Other Pacific Islander

householder had similar levels of sewer connectivity as households with a Black or Asian

householder.

By central city/suburban status (2013)

Fig 3*: The percentage of households connected to sewers by urban and rural

categorizations and by demographic and economic characteristics. See Methods for

definitions of Metropolitan Statistical Area (MSA), central city, urban, and rural. Data from the

2013 American Housing Survey National Survey [32]. MSA: metropolitan statistical area.

Next, we assessed sewer connectivity in large metropolitan areas, which was the smallest

geographic scale resolved by the AHS due to sample size. In 35 of the largest metropolitan

areas in 2019 and 2021, household connectivity to sewers ranges from 69.2% (Birmingham-

Hoover, AL) to 99.4% (LA-Long Beach-Anaheim, CA) (**Fig S *2***). In each metropolitan area,

households with a White or American Indian and Alaska Native householder, with an older

householder, not in poverty, with a larger household size, and with a non-Hispanic householder

were less connected to sewers than the average household, which was similar to what was

seen when stratifying by census division. Exceptions were lower connectivity for households

with a Hispanic householder in San Jose, CA (Hispanic householder: 96.1% of households, CI:

95.1%-97.2%; non-Hispanic householder: 97.8% of households, CI: 97.5%-98.2%) and lower

connectivity for households in poverty in San Francisco, CA (in poverty: 97.7%, CI: 96.7%-

98.7%; not in poverty: 99.3%, CI:99.1%-99.5%); Dallas, TX (in poverty: 92.5%, CI: 90.5%-

94.5%; not in poverty: 95.1%, CI: 94.7%-95.6%); Chicago, IL (in poverty: 92.8%, CI:91.2%-

94.3%, not in poverty: 94.8%, CI: 94.3%-95.3%); and Tampa, FL (in poverty: 83.9%, CI: 81.2%-

87.0%; not in poverty: 88.8%, CI: 88.1%-89.6%) (**Fig S *2***).

In summary, at the census division level, broad trends of sewer connectivity by racial group

revealed higher connectivity for households with an Asian, Black or African American, Native

Hawaiian and Other Pacific Islander, or Hispanic householder and lower connectivity for

households with an American Indian and Alaska Native or White householder. Sewer

connectivity decreased with age, was highest in households with one person, and higher for

households in poverty. These trends were similar when stratifying by urban, rural, central city,

and MSA status, except that the broad racial disparities described above were only observed in

urban and rural areas that are inside of an MSA but outside of a central city. Broad trends were

also similar for individual large metropolitan areas, with some exceptions for households with a

Hispanic householders and households in poverty in particular MSAs.

Which geographic areas have low connectivity to sewers and what are their demographics?

Next, we sought to assess which geographic areas in the U.S. have low connectivity to sewer

systems. These are areas where wastewater-based epidemiology may provide little direct

information on disease dynamics. Additionally, we assessed correlations of sewer connectivity

at the county or county subdivision level with demographic and economic factors. The

compilation of the datasets and an assessment of data completeness are described in the S2

Appendix, **Table *1***, and **Table S *3***.

Using data from the 2021 American Community Survey on the percentage of occupied housing

units lacking complete plumbing facilities by county, we defined the lower bound on the

percentage of the population not connected to sewers. In 14 counties, ≥10% of occupied

housing units lacked complete plumbing facilities; of these counties, 12 were in Alaska and 2

were in the Navajo Nation. In 5 counties, ≥20% of occupied housing units lacked complete

plumbing facilities, all in Alaska. The Yukon-Koyukuk Census Area in Alaska, had the highest

percentage of occupied housing units lacking complete plumbing facilities at 36% (**Fig S 3**,

Table S *1*).

The Florida Department of Health onsite sewage treatment and disposal systems inspection

data (**Fig *4***) revealed that the percentage of households not on septic tanks (expected to

correlate with the percentage of households connected to sewer) was higher in metropolitan

compared to micropolitan county subdivisions and higher in micropolitan compared to rural

county subdivisions, but the effect was not significant (p = 0.052 and p = 0.37, respectively) (**Fig**

S 4a-b). The percentage of households not on septic tanks was significantly lower in county

subdivisions in the panhandle (74.0% vs 92.0%, $p<10^{-9}$) and in county subdivisions that did not

border the coast (86.6% vs 95.7%, $p<10^{-9}$) (**Fig S 4**). By combining this dataset with the 2012

American Community Survey (ACS) describing demographic and economic characteristics of

county subdivisions, we found that county subdivisions with less septic usage (suggesting more

sewer connectivity) were significantly more Hispanic, more Asian, and had a smaller household

size (**Table *2*** and **Fig S 5**), consistent with the nation-wide results from the AHS. These trends

were driven by metropolitan county subdivisions (**Fig S 6**), and an additional trend in

metropolitan county subdivisions not observed for all county subdivisions was that county

subdivisions with less septic usage had a significantly lower percentage of American Indian and

Alaska Natives. There were too few micropolitan and rural county subdivisions to observe

significant trends (**Fig S 7** and **Fig S *8***).

% of households not on septic
by county subdivision in Florida

Fig 4*: **Map of the percentage of households in each county subdivision of Florida** *not*

connected to septic tanks. Data are from the Florida Department of Health septic tank

inspection data from June 2012. County subdivisions with fewer than 5 households or 20

population size are displayed in white.

Table 2: Correlation of the percentage of a Florida county subdivision *not* **connected to**

septic tanks with different demographic or economic variables.

Demographic or economic variable	Correlation of variable with % of county subdivision *not* on septic (Pearson correlation coefficient)	Significance (q-value)
Percent one race and Asian	0.31	**3.2E-07**
Percent Hispanic	0.21	**1.7E-03**
Average household size	-0.19	**9.1E-3**
Percent one race and American Indian and Alaska native	-0.15	0.07

Median age	0.12	0.36
Percent one race and White	-0.12	0.40
Percent one race and Black or African American	0.09	1.11
Median income	0.07	2.27
Percent one race and some other race	0.07	2.33
Percent one race and Native Hawaiian and other Pacific Islander	-0.04	5.76
Percent uninsured (health insurance)	-0.02	8.24

Correlations significant to 5% are shown in bold. Data are from the Florida Department of Health

septic tank inspection permits reported in 2012.

In the 8 states with robust data in the EPA Clean Watersheds Needs Survey, multiple

neighboring counties in Florida, Michigan, and Minnesota had less than 20% of residents

receiving sewage collection, whereas this was not the case in California, Florida, Iowa,

Maryland, New Jersey, and New York (**Fig S 9**). Across multiple states, sewer connectivity by

county was positively correlated with percent Asian and percent Black or African American and

negatively correlated with percent White (**Table 3**), consistent with the nationwide trends from

the AHS data. While counties more connected to sewers appeared to have a significantly lower

percent of American Indian and Alaskan Natives in California, this is affected by missing data

from Indian reservations. Age and percent without health insurance were negatively correlated

with sewer connectivity. Counties more connected to sewers had significantly higher income in

Minnesota, Florida, and California, in contrast to the national results from the AHS data.

Significant trends of sewer connectivity with demographic and economic factors were

predominantly driven by metropolitan counties (**Table S 2**).

Table 3: Correlation of the percentage of households in a county connected to public sewers with different demographic or economic factors by state.

Demographic or economic variable	State	Correlation of variable with % of households by county connected to public sewers (Pearson correlation coefficient)	Significance (q-value)
Median age	MN	-0.43	2.9E-03
	MI	-0.41	1.1E-02
	MN	0.44	1.7E-03
	FL	0.48	3.4E-03
Median income	CA	0.49	7.2E-03
Percent one race and American Indian and Alaska Native	CA	-0.50	4.9E-03
	MN	0.58	2.7E-07
	MI	0.51	7.7E-05
	NY	0.55	2.8E-04
Percent one race and Asian	CA	0.56	3.9E-04
	FL	0.53	4.2E-04
	MN	0.62	1.2E-08
	MI	0.59	4.6E-07
	NY	0.55	3.0E-04
Percent one race and Black or African American	IA	0.38	7.7E-03
	CA	0.46	2.1E-02
	MI	-0.57	1.4E-06

		-0.54	9.6E-04
	CA		
Percent one race and	NY	-0.51	1.8E-03
White	IA	-0.35	2.8E-02
Percent uninsured (health			
insurance)	MI	-0.38	3.3E-02

Only correlations that are significant (q-value $\leq$ 0.05) are shown. Data are from the 2012 EPA

Clean Watersheds Needs Survey.

Analysis of the Island Areas of Guam, the Northern Mariana Islands, the Virgin Islands, and

American Samoa revealed lower levels of sewer connectivity than in the states, with

considerable spatial variability in connectivity (see S2 Appendix).

Factors influencing applicability of wastewater-based

epidemiology to communities lacking sewer connection

The inequities in sewer connections among communities raise the question of the applicability

of inferences drawn from wastewater data in one community to communities lacking sewer

connections. To explore this question, we used a deterministic compartmental model of two

interacting populations with tunable levels of interaction, each with susceptible-infected-

recovered (SIR) dynamics, and with wastewater sampling in each population into a common

sample (**Fig 5**a). We set population A to be entirely connected to sewers and population B to

have a tunable level of sampling by wastewater.

A common application of wastewater data is to aid in determining when an outbreak has

peaked, which can inform policy decisions on when to ease restrictions. We first asked how well

the wastewater data could predict the time of peak infections in population B when it is

completely unconnected to sewer systems and thus the only sampling is from population A. We

found that when the two populations have similar sizes, the wastewater concentration peak and

the infection peak in population B was within one generation time except when the interactions

between the two population is weak (**Fig 5**b). The generation time is the average time between

an individual's infection and transmission, which in an SIR model is the same as the infectious

period. For example, in two populations with 5000 individuals each, a disease basic

reproduction number of 1.5, and infectious period of 5.6 days, the percentage of contacts that

occur across populations must drop below 4% before the peak in wastewater concentration

occurred more than one generation time apart from the peak in infections in the unconnected

population (**Fig 5**c).

Fig 5: Deterministic compartmental model of two interacting populations with

susceptible, infected, and recovered compartments and sampling by wastewater. (a)

Schematic of model. Population A is entirely sampled by wastewater and a fraction of

population B, f_B, is sampled by wastewater. (b) Number of infected individuals (black curves) in

population A, population B, and population A and B combined and wastewater concentration

(blue curves) over time for different interaction strengths or fractions of cross-population

contacts (ε). The vertical lines indicate the peak time for the curves with matching line style and

color. The time between the population B peak in infections and the wastewater concentration

peak is indicated by Δt. (c) The time between the population B peak in infections and the

wastewater concentration peak (defined in (b)) when varying the fraction of cross-population

contacts (ε) and the fraction of population B that is sampled (f_B). Light blue dots indicate

parameter regimes in which Δt is less than 1 generation time. In both (b) and (c), the simulation

starts with a single infected individual in population A at time 0 and no infected individuals in

population B. The parameters are set to population size $N_A = N_B = 5000$, basic reproduction

number $R^A_0 = R^B_0 = R_0 = \frac{\beta_A}{\gamma_I} = \frac{\beta_B}{\gamma_I} = 1.5$, rate of recovery $\gamma_I = 0.18$, and the overall contact rate times

the probability of infection given contact $\beta_A = \beta_A = \gamma_I R_0$.

We then weakened the assumption that population B had no contribution to the sampled

wastewater and allowed a fraction of population B to be sampled by wastewater. For example,

individuals in population B may commute to a workplace sampled by wastewater, or a fraction of

population B may be connected at home. As the fraction of population B that is sampled by

wastewater increases, the time between infection and wastewater peaks decreases; however,

the fraction of B that is sampled has a much weaker effect on the wastewater peak time than

the interaction parameter between the communities. For example, with the above parameters

and equal population sizes, with 10% sampling of population B, the cross-population contacts

must drop below 4%, and with 50% below 3% and with 100% below 2% before the wastewater

concentration peak and the peak in the number of infections in population B drops below one

generation time.

Varying the population size of both populations (with an equal population size between the two

populations), the recovery rate or generation time, and the R_0 of both populations (with an equal

R_0 in the two populations) has only a weak effect on the discrepancy between wastewater

concentration and population B infection peaks (**Fig S 10**).

When the unconnected population has a smaller size than the connected population, then even

substantial interactions between the two populations can lead the wastewater concentration to

peak before the infections in the unconnected population peaks (**Fig S 10**). Both very weak and

very strong interactions cause the wastewater peak to coincide with the peak number of

infections in the unconnected population.

Additionally, when the unconnected population has a smaller R_0 than the connected population

(which could occur for instance if the unconnected population was more spread out

geographically and thus had a lower contact rate), the outbreak in population B peaks later than

the wastewater peak leading to a discrepancy in the peak times (**Fig S 10**).

Discussion
403

Equity of sewer connectivity
405

Our analysis of available datasets revealed considerable variability in sewer connectivity within

and between communities, across locations, demographics, and economic statuses in the

United States. The western US had higher levels of sewer connectivity than the eastern US.

Across the US, lower connectivity to sewers was observed for American Indian and Alaska

Native householders; White householders; non-Hispanic householders; older populations; and

larger households in most census divisions. In the Pacific census division, lower connectivity to

sewers was observed for Native Hawaiian and Other Pacific Islander householders specifically

in areas inside of a metropolitan statistical area but outside of a central city. Across census

divisions, households not in poverty were less connected to sewers, consistent with the

observation that, nation-wide, higher income households were less connected to sewers [10].

Aggregating by large metropolitan areas yielded similar trends as when aggregating by census

divisions, except for lower connection for Hispanic householders in San Jose area and for

households in poverty in San Francisco, Dallas, Chicago, and Tampa metropolitan areas,

whereas these groups were better connected to sewers at the census division level. The

differences in the MSA-level data suggested heterogeneity in connectivity across locations.

Geographic variation in sewer connectivity

Large parts of Alaska and the Navajo Nation lacked plumbing. This is consistent with the finding

from the Annual Report to Congress on Sanitation Deficiency Levels for Indian Homes and

Communities in 2019 that approximately 20% of Indian homes in Alaska do not have sewage

disposal and about 10% of Indian homes in the Navajo Nation do not have sewage disposal

[33].

State-level datasets revealed that sewer connectivity varied between and within communities.

While in some states most areas had sewer connectivity (California, Iowa, Maryland, New

Jersey, New York), in others there were large regions that lacked connectivity (Minnesota,

Michigan, Florida). Asian and Hispanic populations were less connected to septic tanks across

county subdivisions in Florida; however, there was considerable variability across the state and

potential biases in the data. In contrast to the national data, in Minnesota, Florida, and

California, counties with lower median household incomes were less connected to sewers,

suggesting variability across the nation. This result in Florida is consistent with the observation

that as income levels rose, household decentralized system usage declined in Florida [14].

However, we did not observe any significant correlation of income with septic usage in the

Florida Department of Health onsite sewage treatment and disposal systems inspection

permitting dataset, possibly due to not all septic tanks having a permit that is reported in this

dataset.

While these correlations may be largely driven by differences between urban and rural areas

within a county or county subdivision, they may still be very useful in assessing inequities in

sewer connectivity in large catchments that span neighboring counties or county subdivisions.

Data limitations

Substantial data gaps and biases prevented a comprehensive analysis of disparities in sewer

system connectivity within and between communities across the US, particularly for small

communities, tribal lands and Alaska Native Villages, and state and local geographic scales.

The current design of the American Housing Survey has too small of sample size to study state

and local geographic scales [30]. Additionally, the AHS may have underestimated the fraction of

households using septic systems due to its survey design (**Table 1**). The differences between

Florida state data supports the need for detailed and reproducible longitudinal studies of sewer

connectivity (both in developing sampling and analysis methods).

Public health officials may find it beneficial to conduct local equity studies of sewer connectivity.

Data on the catchment area sizes of the wastewater treatment plants would allow a better

understanding of the geographic extent of the sampled population and the level of geographic

aggregation needed to study sewer connectivity inequities. A recent study mapping wastewater

treatment plant catchment areas and their population sizes served in New York state provides a

template for how to create these sewer catchment maps from a combination of permitting,

survey, and tax record data [13]. If possible, it would be helpful to consolidate and standardize

existing data where it exists (for instance, sewage disposal permitting at the local or state

levels). Additionally, if the EPA's request to expand the question on the ACS about access to

plumbing facilities to additionally ask about the type of plumbing facility [14,34] is granted, then

this would be a valuable dataset with the geographic and temporal resolution to study equity in

sewer connectivity moving forward.

470 Generalizability of wastewater-based epidemiology data in

471 light of heterogeneities in sewer connectivity

Our modeling results suggest that even weak interactions between two communities allow

wastewater monitoring in one community to serve as reliable proxy for the time of maximum

infections in the other community when the population sizes and R_0 of the two populations are

comparable, but not when the unconnected population has a substantially lower population size

or R_0. In the scenario with unequal population size, outbreaks that are seeded at the same time

in the two populations in the absence of interactions peak earlier in the unconnected population

due to its smaller size. With weak interactions, seeding of infections from population A to B

occurs slower, leading to more coincidence in peaks; with strong interactions, then the

dynamics of B are dominated by those in A, leading to more coincidence in peaks; at

intermediate interaction strengths, infections are seeded earlier and peaks early, leading to the

largest discrepancy in wastewater and infection peaks.

As the purpose of our model was to explore the impact of factors on the generalizability of

wastewater rather than to accurately capture the dynamics in all scenarios, we made simplifying

assumptions, including that all infected individuals shed the same amount of pathogen genetic

material into wastewater and shed only during the infected period; that all individuals contribute

equally to wastewater; and that pathogen detection in wastewater has perfect sensitivity.

Our results suggest that in assessing the generalizability of wastewater data, it would be useful

to estimate the extent of mobility between connected and unconnected communities.

Interestingly, Ref [35] found no correlation between the size of a catchment area and the

correlation of wastewater with case data for SARS-CoV-2, consistent with the result that

interactions between connected and unconnected communities cause the disease dynamics to

look similar to the wastewater data. In regions without sewer connectivity or with little interaction

between connected and unconnected communities, wastewater data from neighboring

communities will be less informative. In areas with low sewer connectivity in households,

sampling wastewater outflow at frequently visited non-household locations (i.e., schools, offices,

malls, etc.) may capture a more representative population.

501 Additional considerations for equity in WBE

In addition to inequities in sewer connectivity, the sewer locations used for wastewater sampling

should be considered to promote demographic equity and ensuring the ability to capture

spatiotemporal trends [36,37]. While we have focused on analysis on the US due to data

availability, internationally disadvantaged populations are associated with lower access to

sewers [38]. The ongoing development of wastewater sampling in non-sewered settings [39], for

example in water channels in Las Vegas Valley [40], onsite sanitation facilities in Bangladesh, a

refugee camp in Lebanon [41], and various non-sewered settings in low and middle income

countries [42], represents a critical area for research and development.

511 Conclusions

In summary, while wastewater-based epidemiology is a useful tool to monitor disease burden

and dynamics, our analyses suggest that access to this new tool varies across the US. More

comprehensive data on sewer connectivity is needed, and in combination with assessments of

mobility and population parameters, these data can help with the design of wastewater sampling

schemes and the interpretation for epidemic trends in sampled and neighboring unsampled

communities.

Acknowledgements

We thank Cara Omana from the Minnesota Pollution Control Agency for generously sharing the

Minnesota WINS data and Harry Campbell from the Utah Department of Environment Quality,

Division of Water Quality for generously sharing the Utah MWPP survey data and to both for

answering questions about the datasets. We thank colleagues in the Grad lab and the Harvard

T. H. Chan School of Public Health Center for Communicable Disease Dynamics for helpful

discussions, particularly Stephen Kissler.

References

1. Peccia J, Zulli A, Brackney DE, Grubaugh ND, Kaplan EH, Casanovas-Massana A, et al. Measurement of SARS-CoV-2 RNA in wastewater tracks community infection dynamics. Nat Biotechnol. 2020;38: 1164–1167.

2. Medema G, Been F, Heijnen L, Petterson S. Implementation of environmental surveillance for SARS-CoV-2 virus to support public health decisions: Opportunities and challenges. Curr Opin Environ Sci Health. 2020;17: 49–71.

3. Pöyry T, Stenvik M, Hovi T. Viruses in sewage waters during and after a poliomyelitis outbreak and subsequent nationwide oral poliovirus vaccination campaign in Finland. Appl Environ Microbiol. 1988;54: 371–374.

4. Link-Gelles R, Lutterloh E, Schnabel Ruppert P, Backenson PB, St George K, Rosenberg ES, et al. Public Health Response to a Case of Paralytic Poliomyelitis in an Unvaccinated Person and Detection of Poliovirus in Wastewater - New York, June-August 2022. MMWR Morb Mortal Wkly Rep. 2022;71: 1065–1068.

5. Kilaru P, Hill D, Anderson K, Collins MB, Green H, Kmush BL, et al. Wastewater Surveillance for Infectious Disease: A Systematic Review. Am J Epidemiol. 2023;192: 305–322.

6. Boehm AB, Hughes B, Duong D, Chan-Herur V, Buchman A, Wolfe MK, et al. Wastewater concentrations of human influenza, metapneumovirus, parainfluenza, respiratory syncytial virus, rhinovirus, and seasonal coronavirus nucleic-acids during the COVID-19 pandemic: a surveillance study. Lancet Microbe. 2023. doi:10.1016/S2666-5247(22)00386-X

7. Baz-Lomba JA, Salvatore S, Gracia-Lor E, Bade R, Castiglioni S, Castrignanò E, et al. Comparison of pharmaceutical, illicit drug, alcohol, nicotine and caffeine levels in wastewater with sale, seizure and consumption data for 8 European cities. BMC Public Health. 2016;16: 1035.

8. Gushgari AJ, Venkatesan AK, Chen J, Steele JC, Halden RU. Long-term tracking of opioid

consumption in two United States cities using wastewater-based epidemiology approach.

Water Res. 2019;161: 171–180.

9. Analytics B. Wastewater for equitable Covid-19 monitoring. In: Medium [Internet]. 8 Jul

2021 [cited 9 Feb 2023]. Available: https://biobotanalytics.medium.com/wastewater-for-

equitable-covid-19-monitoring-e3b947d91e0a

10. Olesen SW, Young CCW, Duvallet C. White paper: The effect of septic systems on

wastewater-based epidemiology. 2022. doi:10.5281/zenodo.7130726

11. U.S. Census Bureau. AHS 2021 National Public Use File. Available:

https://www2.census.gov/programs-

surveys/ahs/2021/AHS%202021%20National%20PUF%20v1.0%20CSV.zip

12. US EPA Decentralized Wastewater Program}. Septic Systems Fact Sheet. 2008 Oct.

13. Hill DT, Larsen DA. Using geographic information systems to link population estimates to

wastewater surveillance data in New York State, USA. PLOS Glob Public Health. 2023;3:

e0001062.

14. U.S. Environmental Protection Agency, Office of Water. Report to Congress on The

Prevalence Throughout the U.S. of Low- and Moderate-Income Households Without

Access to a Treatment Works and The Use by States of Assistance under Section

603(c)(12) of the Federal Water Pollution Control Act. U.S. Environmental Protection

Agency; 2021 Jul. Available: https://www.epa.gov/system/files/documents/2022-01/low-

mod-income-without-treatment_report-to-congress.pdf

15. U.S. Census Bureau. AHS 2019 National Public Use File. Available:

https://www2.census.gov/programs-

surveys/ahs/2019/AHS%202019%20National%20PUF%20v1.1%20CSV.zip

16. U.S. Census Bureau. AHS 2019 Metropolitan Public Use File. Available:

https://www2.census.gov/programs-

surveys/ahs/2019/AHS%202019%20Metropolitan%20PUF%20v1.0%20CSV.zip

17. U.S. Census Bureau, Department of Commerce Department of Housing and Urban

Development. 2019 AHS Integrated National Sample: Sample Design, Weighting, and Error

Estimation. U.S. Census Bureau, Department of Commerce Department of Housing and

Urban Development; 2020 Sep. Available: https://www2.census.gov/programs-

surveys/ahs/2019/2019%20AHS%20National%20Sample%20Design,%20Weighting,%20a

nd%20Error%20Estimation.pdf

18. U.S. Census Bureau, Department of Commerce Department of Housing and Urban

Development. 2019 AHS Metropolitan Sample: Sample Design, Weighting, and Error

Estimation. U.S. Census Bureau, Department of Commerce Department of Housing and

Urban Development; 2020 Sep. Available: https://www2.census.gov/programs-

surveys/ahs/2019/2019%20AHS%20Metropolitan%20Sample%20Design,%20Weighting,%

20and%20Error%20Estimation.pdf

19. U.S. Census Bureau, Department of Commerce Department of Housing and Urban

Development. 2021 AHS Integrated National Sample: Sample Design, Weighting, and Error

Estimation. U.S. Census Bureau, Department of Commerce Department of Housing and

Urban Development; 2022 Aug. Available: https://www2.census.gov/programs-

surveys/ahs/2021/2021%20AHS%20National%20Sample%20Design,%20Weighting,%20a

nd%20Error%20Estimation.pdf

20. US Census Bureau. American Community Survey Information Guide. [cited 5 Jan 2023].

Available: https://www.census.gov/programs-surveys/acs/library/information-guide.html

21. General Facts and Statistics about Wastewater in Florida. [cited 10 Feb 2023]. Available:

https://floridadep.gov/water/domestic-wastewater/content/general-facts-and-statistics-

about-wastewater-florida

22. Florida Department of Health. Onsite Sewage Locations in Florida - 2012. Available:

https://fgdl.org/zips/metadata/xml/septic_jun12.xml

23. Environmental Protection Agency. Clean Watersheds Needs Survey Detailed Scope and

Methods. Environmental Protection Agency; 2012. Available:

https://www.epa.gov/sites/default/files/2015-

10/documents/cwns_2012_detailed_scope_and_methods-508.pdf

24. Agency USEP. Clean watersheds needs survey 2012 report to congress. US

Environmental Protection Agency; 2016.

25. U.S. Census Bureau. Guidance on Using 2020 Island Areas Censuses Data. 20 Oct 2022

[cited 31 Mar 2023]. Available: https://www2.census.gov/programs-

surveys/decennial/2020/technical-documentation/island-areas-tech-docs/2020-iac-

guidance.pdf

26. U.S. Census Bureau. 2020 Island Areas Censuses (IAC) Demographic Profile Summary

File. U.S. Census Bureau; 2022 Oct. Available: https://www2.census.gov/programs-

surveys/decennial/2020/technical-documentation/island-areas-tech-

docs/demographic_profile/2020-iac-dpsf-technical-documentation.pdf

27. U.S. Department of Housing and Urban Development and U.S. Census Bureau. 2013 AHS

Definitions. In: 2013 AHS Definitions [Internet]. [cited 9 Feb 2023]. Available:

https://www2.census.gov/programs-surveys/ahs/2013/2013%20AHS%20Definitions.pdf

28. TIGER/Line® Shapefiles. [cited 25 Apr 2023]. Available: https://www.census.gov/cgi-

bin/geo/shapefiles/index.php

29. U.S. Census Bureau. Metropolitan and Micropolitan About. In: Metropolitan and

Micropolitan About [Internet]. 22 Nov 2021 [cited 9 Feb 2023]. Available:

https://www.census.gov/programs-surveys/metro-micro/about.html

30. Schwartz MB. 2007 American Community Survey: A Comparison to Selected Housing and

Financial Characteristics for the United States from the 2007 American Housing Survey.

Housing and Household Economic Statistics Division U.S. Census Bureau; Available:

https://www.huduser.gov/portal/datasets/ahs/acs-ahscomparisonreport.pdf

31. U.S. Department of Housing and Urban Development and U.S. Census Bureau. 2021 AHS

Definitions. In: 2021 AHS Definitions [Internet]. [cited 22 May 2023]. Available:

https://www2.census.gov/programs-surveys/ahs/2021/2021%20AHS%20Definitions.pdf

32. U.S. Census Bureau. AHS 2013 National Public Use File. Available:

https://www2.census.gov/programs-

surveys/ahs/2013/AHS%202013%20National%20PUF%20v2.0%20CSV.zip

33. Indian Health Service; Office of Environmental Health and Engineering; Division of

Sanitation Facilities Construction. Annual Report To the Congress of the United States On

Sanitation Deficiency Levels for Indian Homes and Communities: Fiscal Year 2019.

Rockville (MD): Indian Health Service (US); 2020.

34. Environmental Protection Agency. EPA's Proposal: Justification for Requested Revision,

Addition or Deletion to Questions on the American Community Survey (ACS). 2018.

Available:

https://foiaonline.gov/foiaonline/action/public/submissionDetails?trackingNumber=EPA-HQ-

2018-010667&type=request

35. Duvallet C, Wu F, McElroy KA, Imakaev M, Endo N, Xiao A, et al. Nationwide Trends in

COVID-19 Cases and SARS-CoV-2 RNA Wastewater Concentrations in the United States.

ACS EST Water. 2022. doi:10.1021/acsestwater.1c00434

36. Medina CY, Kadonsky KF, Roman FA Jr, Tariqi AQ, Sinclair RG, D'Aoust PM, et al. The

need of an environmental justice approach for wastewater based epidemiology for rural and

disadvantaged communities: A review in California. Curr Opin Environ Sci Health. 2022;27:

100348.

37. Haak L, Delic B, Li L, Guarin T, Mazurowski L, Dastjerdi NG, et al. Spatial and temporal

variability and data bias in wastewater surveillance of SARS-CoV-2 in a sewer system. Sci

Total Environ. 2022;805: 150390.

38. Adhikari S, Halden RU. Opportunities and limits of wastewater-based epidemiology for

tracking global health and attainment of UN sustainable development goals. Environ Int.

2022;163: 107217.

39. Jakariya M, Ahmed F, Islam MA, Al Marzan A, Hasan MN, Hossain M, et al. Wastewater-

based epidemiological surveillance to monitor the prevalence of SARS-CoV-2 in developing

countries with onsite sanitation facilities. Environ Pollut. 2022;311: 119679.

40. Gerrity D, Papp K, Dickenson E, Ejjada M, Marti E, Quinones O, et al. Characterizing the

chemical and microbial fingerprint of unsheltered homelessness in an urban watershed. Sci

Total Environ. 2022;840: 156714.

41. Tokajian S, Moghnieh R, Salloum T, Arabaghian H, Alousi S, Moussa J, et al. Extended-

spectrum β-lactamase-producing Escherichia coli in wastewaters and refugee camp in

Lebanon. Future Microbiol. 2018;13: 81–95.

42. Gwenzi W. Wastewater, waste, and water-based epidemiology (WWW-BE): A novel

hypothesis and decision-support tool to unravel COVID-19 in low-income settings? Sci

Total Environ. 2022;806: 150680.

43. U.S. Department of Housing and Urban Development and U.S. Census Bureau. Appendix

B: Sample Design and Weighting: 2013. U.S. Department of Housing and Urban

Development and U.S. Census Bureau; 2013. Available:

https://www2.census.gov/programs-

surveys/ahs/2013/2013%20AHS%20National%20Sample%20Design%20and%20Weightin

g.pdf

44. Onsite Sewage FAQ - Permitting. [cited 10 Feb 2023]. Available:

https://floridadep.gov/water/onsite-sewage/content/onsite-sewage-faq-permitting

45. Florida Department of Health. Septic System Owner's Guide. In: floridahealth.gov [Internet].

[cited 10 Feb 2023]. Available: https://www.floridahealth.gov/environmental-health/onsite-

sewage/_documents/septic-folder-8x11-link2.pdf

46. Utah Department of Environmental Quality Water Quality. Municipal Wastewater Planning

Program (MWPP). In: Municipal Wastewater Planning Program (MWPP) [Internet]. 3 Feb

2023 [cited 10 Feb 2023]. Available: https://deq.utah.gov/water-quality/municipal-

wastewater-planning-program-mwpp

47. Campbell H. 2021 MWPP Survey. In: 2021 MWPP Survey [Internet]. 27 Oct 2021 [cited 10

Feb 2023]. Available: https://documents.deq.utah.gov/water-quality/engineering/municipal-

wastewater-planning-program/DWQ-2021-026722.pdf

48. Minnesota Pollution Control Agency. 2021 Wastewater Infrastructure Needs Instructions

and Frequently Asked Questions (FAQ). Apr 2021 [cited 31 Mar 2023]. Available:

https://www.pca.state.mn.us/sites/default/files/wq-wwtp3-06.pdf

49. Robinson C. 2017 SSTS Annual Report Subsurface Sewage Treatment Systems in

Minnesota. Minnesota Pollution Control Agency; 2018 Apr. Available:

https://www.pca.state.mn.us/sites/default/files/wq-wwists1-58.pdf

50. U.S. Environmental Protection Agency. 2012 Clean Watersheds Needs Survey. Available:

https://ordspub.epa.gov/ords/cwns2012/f?p=cwns2012:25

51. Weidhaas J, Aanderud ZT, Roper DK, VanDerslice J, Gaddis EB, Ostermiller J, et al.

Correlation of SARS-CoV-2 RNA in wastewater with COVID-19 disease burden in

sewersheds. Sci Total Environ. 2021;775: 145790.

52. U.s E. WaterSense Statistics and Facts. 11 May 2022 [cited 3 Apr 2023]. Available:

https://www.epa.gov/watersense/statistics-and-facts

Supporting information

S1 Appendix. Supplementary method

S2 Appendix. Supplementary analyses. Compilation of data, assessment of data

completeness, sewer connectivity in U.S. Island Areas, supplementary figures, and

supplementary tables.

S1 Appendix. Supplementary methods.

Dataset descriptions and data cleaning

U.S. Census Bureau American Housing Survey 2019 and 2021

The American Housing Survey (AHS) National and Metropolitan Public Use Files were

downloaded from Ref. [11,15,16]. The sample design, weighting, and error estimation survey is

described in Refs. [17–19]. Briefly, the AHS is a biannual survey that assesses housing

characteristics. Two samples of households are chosen every survey cycle, the Integrated

National Sample and the Independent Metropolitan Sample. The Integrated National Sample

consists of nationally representative households and includes an oversample of each of 15

largest metropolitan areas. The Independent Metropolitan Sample consists of representative

households in 20 large metropolitan areas that are representative of the next largest 50

metropolitan areas. Ten of the 20 large metropolitan areas are sampled every other survey

cycle (10 metropolitan areas surveyed every 2 years). The Integrated National Sample in 2019

included 86,257 selected representative housing units, of which about 63,500 met the AHS

definition of a housing unit and were able to be interviewed, and in 2021 included 95,295

selected representative housing units, of which about 64,100 were interviewed. The Integrated

Metropolitan Samples in 2019 and 2021 included around 3000 selected representative housing

units in each metropolitan area, of which around 1000-3000 were eligible and were able to be

interviewed (same criteria as in the Integrated National Sample).

Geographic data of each household available in the Public Use File included the census

division; core based statistical area (CBSA) for the 15 largest metropolitan areas and the 20

large metropolitan areas in the metropolitan data. All other households are classified as in "all

other metropolitan areas" or "not in a metropolitan area" (all other locations).

The race, ethnicity, and age of the householder (the person or one of the persons who is an

owner or renter of the unit) was used to categorize the race, ethnicity, and age of the household.

Due to the low numbers of households where the householder responded as having more than

one race, we focused on households where householders identified as having a single race.

U.S. Census Bureau American Housing Survey National Sample 2013

The American Housing Survey National (2013) Use File was downloaded from Ref. [32]. The

sample design, weighting, and error estimation survey is described in Ref [43]. The National

Sample included 84,400 selected housing units representative nationally and including a

supplemental sample of 15,553 housing units in the Chicago, Detroit, New York City, Northern

New Jersey, and Philadelphia metropolitan statistical areas, of which about 71,600 met the

definition of a housing unit and were able to be interviewed.

Geographic data of each household available in the Public Use File included the census

division; census region; central city, urban, or rural status; and select metropolitan area codes.

All other housing units are classified into "not in a metropolitan area" or had their code

suppressed for confidentiality [27].

Florida Department of Health onsite sewage treatment and disposal system inspections

The locations of onsite sewage treatment and disposal systems inspected by Florida Department of Health reported June 2012 was downloaded from Ref. [22]. As of 2022, an inspection permit is required for construction [44] and inspection is recommended every 3 to 5 years after [45]. The number of systems inspected that are active at the latest inspection (after removing repeat inspections) is 601,430. The years of inspection are primarily 1998-2011, with some outliers as early as 1900, which may be old septic systems or typos, and as late as 2200, which are likely typos.

Utah Municipal Wastewater Planning Program (MWPP) Survey 2021

Data from the Utah Municipal Wastewater Planning Survey was provided by the Utah Department of Environment Quality, Division of Water Quality. The dataset includes an estimate of the population receiving collection by each utility that owns or operates a sanitary sewerage system and its latitude and longitude. The data gathered from treatment plant operators and participation is reported as mandatory [46], although only 71% of contacts responded [47]. We filtered the data to include any facility that has collection (this included collection only, treatment and collection, or small lagoons which are often collection and treatment). Manual correction of facility longitude and latitude is described in the supplementary methods.

Manual editing of facility longitude and latitude was done for 5 facilities' that did not fall within the state of Utah when combining with the 2021 TIGER shapefile (BRIAN HEAD TOWN, CANYONG LAND IMPROVEMENT DISTRICT, HILL AFB (AMERICAN WATER), JORDANELLE SSD, PLAIN CITY) and 1 facility that did not report a latitude and longitude (PANGUITCH LAKE S. S. D.). For these facilities, the latitude and longitude were manually

774　edited to be that of the location found in a Google Maps search of the facility name. If the

775　Google Maps search found a city, then an arbitrarily chosen location in that city was used to

776　obtain a latitude and longitude.

Minnesota Wastewater Infrastructure Needs Survey (WINS) 2021

Data for the Minnesota Wastewater Infrastructure Needs Survey [48] was provided by the

Minnesota Pollution Control Agency. Communities (places and townships) responded to the

question "Does your community have a collection system?" with "yes", "no", or no response. The

survey did not receive a response from every community in Minnesota.

The Census Designated Place (CDP) name was obtained from the reported community name

with manual changes reported in **Table S 2**.

Minnesota Subsurface Sewage Treatment Systems 2017

The Minnesota Subsurface Sewage Treatment Systems (SSTS) dataset reported the number of

SSTS by local government units with a known SSTS program. Data was taken from Ref. [49].

Two hundred and eleven out of 2018 known SSTS programs submitted an annual report. Not all

SSTS programs were able to be identified and contacts were not always provided. Fourteen

reported having zero SSTS within their jurisdiction despite permitting SSTS in 2017. Data were

reported to sometimes be estimates.

U.S. Environmental Protection Agency Clean Watersheds Needs Survey 2012

Data for the most recent Environmental Protection Agency (EPA) Clean Watersheds Needs

Survey (CWNS) survey (2012) [24] was downloaded from [50] and extracted from a Microsoft

Access database file to csv files using the mdbtools package

(https://github.com/mdbtools/mdbtools). The survey is administered by the EPA and the States

to assess funding needs for unfunded capital costs of treatment works projects that (1) address

a water quality or related public health problem existing as of January 1, 2012, or is expected to

occur within the next 20 years and (2) meet the CWNS documentation criteria. CWNS

documentation criteria includes (1) description and location of the problem, (2) site-specific

solution to the problem, and (3) detailed cost information for implementing the solution. Data is

collected about (1) publicly owned wastewater collection and treatment facilities, (2) combined

sewer overflow control facilities, (3) stormwater management activities, and (4) decentralized

wastewater treatment facilities. Data is collected on (a) estimated needs, costs and technical

information, (b) facility location and contact information, and (c) facility population served, flow,

effluent, and unit process information. Supporting documentation was required for the entered

data with some exceptions (for instance small communities could use a more simple form), and

quality control was performed by reviewers to compare documentation with data entered in the

system to ensure consistency of technical and cost data [23]. Data entry occurred from January

to December 2012.

There was large variability in the level of effort and resources that each state put into the survey.

New York, California, Florida, New Jersey, Maryland, Iowa, Minnesota, and Michigan likely

invested heavily in participating in the survey or already had similar state-level data collection

systems in place for getting comprehensive responses [22,48,49].

Facilities not included in the dataset are those in South Carolina, the Northern Mariana Islands,

and American Samoa as they did not participate in the survey; facilities whose projects did not

have documented solutions or cost estimates; privately owned wastewater facilities that serve

privately owned industrial facilities, military installations, national parks, or other federal facilities;

facilities on tribal lands and in Alaskan Native Villages, which were separately surveyed by the

Indian Health Service; and facilities with projects that had existing funding prior to January 1,

2012. Additionally, larger facilities were more likely to be captured by the survey due to the

resource requirements for obtaining the right documentation. To address this bias, the survey

had decreased documentation needs for small communities (population <=10,000); despite this,

the small community facilities were still underrepresented (pg. 2 of Ref. [24]).

We filtered the facilities to the current facilities of type "collection: combined sewers" or

"collection: separate sewers". The population receiving collection is taken from the reported total

number of residents who are connected to a sewer system which empties into a treatment plant.

This does not include non-resident populations, populations served by acceptable decentralized

wastewater treatment systems, or populations connected to sewers that do not discharge to a

treatment plant.

U.S. Census Bureau Island Areas Decennial Survey 2020

The Island Areas Decennial Survey surveys housing, social, and economic information for all

housing units in Guam, the Northern Mariana Islands, the Virgin Islands, and American Samoa

[26]. Data of general demographic characteristics (DP01), selected economic characteristics

(DP03), and selected housing characteristics (DP04) were downloaded from

https://data.census.gov/. Housing characteristics included data for both occupied and vacant

housing units.

U.S. Census Bureau American Community Survey

The American Community Survey (ACS) is a monthly survey of a representative subsample of

addresses in the US [20] on characteristics of populations and households. The ACS 5 year

estimates (average over 5 years, less subject to fluctuations) for selected social characteristics

(DP02), selected economic characteristics (DP03), and demographic and housing estimates

(DP05) stratified county or county subdivision for select years were downloaded from

https://data.census.gov/.

Data analysis and visualization

Florida Department of Health onsite sewage treatment and disposal system inspections dataset analysis

The number of septic tanks in each county subdivision was calculated as the number of septic

permits whose latitude and longitude occurred inside the boundaries of each county subdivision

according to the 2012 U.S. Census Bureau county subdivision TIGER shapefile. To calculate

the fraction of a county subdivision connected to septic tanks, the number of septic tanks in

each county subdivision was divided by the number of households in the county subdivision as

reported by the 2012 American Community Survey 5-year estimate. Only county subdivisions

with at least 5 households and 20 population size were used.

Utah Municipal Wastewater Planning Program Survey analysis

The population in each county receiving collection was calculated by summing the population

receiving collection across all reported facilities in that county. The population in each county

subdivision receiving collection was calculated by summing the population receiving collection

across all reported facilities whose latitude and longitude fell within that county subdivision's

boundaries from the 2021 TIGER shapefile. To calculate the fraction of a county or county

subdivision receiving sewer collection, the population receiving collection was divided by the

population size of each county or county subdivision reported by the 2021 American Community

Survey 5-year estimate. Only county subdivisions with at least 5 households and 20 population

size were used.

U.S. Environmental Protection Agency Clean Watersheds Needs

Survey analysis

To compare the EPA CWNS and AHS data in Core Based Statistical Areas (CBSA), counties

were categorized into CBSAs using the February 2013 U.S. Census Bureau file. To calculate

the fraction of a county receiving collection, the population size receiving collection in that

county was divided by the population size of each county reported by the 2012 American

Community Survey 5-year estimate. We flagged counties that had no facilities reported, which

could either be because there were no facilities in that county or because of a lack of reporting,

and counties that had >100% sewer connectivity by population (potentially due to misreporting

or collection occurring across multiple counties) were set to 100% connectivity.

U.S. Census Bureau Island Areas Decennial Survey analysis

889   To calculate the fraction of a Census Designated Place (CDP) connected to sewers, the number

890   of households connected to sewers was divided by the total number of households in a CDP.

891   Only CDPs with at least 5 households and 20 population size were used.

Generation of maps

894   TIGER shapefiles from the U.S. Census Bureau in the closest matching year to the dataset

895   were used for generating maps. Maps were generated using the geopandas package (version

896   0.12.2) in Python (version 3.11.0) and were displayed in the Mercator projection.

Simulations of interacting populations and wastewater sampling

Two interacting populations are modeled in a compartmental model with susceptible, infected, and recovered (SIR) compartments for each population. The number of susceptible, infected, and recovered individuals in population A (S_A, I_A, and R_A, respectively) and in population B (S_B, I_B, and R_B, respectively) are described by:

$$\frac{dS_A}{dt} = -\beta_A(1-\varepsilon)I_A\frac{S_A}{N_A} - \beta_B\varepsilon I_B\frac{S_A}{N_A}$$

$$\frac{dS_B}{dt} = -\beta_B(1-\varepsilon)I_B\frac{S_B}{N_B} - \beta_A\varepsilon I_A\frac{S_B}{N_B}$$

$$\frac{dI_A}{dt} = \beta_{AA}(1-\varepsilon)I_A\frac{S_A}{N_A} + \beta_{BA}\varepsilon I_B\frac{S_A}{N_A} - \gamma_I I_A$$

$$\frac{dI_B}{dt} = \beta_{BB}(1-\varepsilon)I_B\frac{S_B}{N_B} + \beta_{AB}\varepsilon I_A\frac{S_B}{N_B} - \gamma_I I_B$$

$$\frac{dR_A}{dt} = \gamma_I I_A$$

$$\frac{dR_B}{dt} = \gamma_I I_B$$

where N_A and N_B are the number of individuals in populations A and B respectively, β_A and β_B are the overall contact rates of populations A and B respectively times the probability of infection given contact, γ_I is the rate of recovery. ε is the fraction of the total contacts made by individuals in a population that occurs with individuals in the other population and it describes the interaction strength between the two populations. ε can take on values between 0 and 1, where $\varepsilon=0$ indicates that all interactions occur within each population and none between populations, and $\varepsilon=1$ indicates that all interactions occur between the populations and none within

populations. The base parameters are set at $N_A = N_B = 5000$, $\gamma_I = 0.18$ inverse days, $R_0^A = R_0^B =$

$\frac{\beta_A}{\gamma_I} = \frac{\beta_B}{\gamma_I} = 1.5$, which represent an outbreak similar to COVID-19, but the parameters are varied

in sensitivity analyses. The initial condition was a single infected individual in population A ($I_A = 1$)

and all other individuals were susceptible ($S_A = N-1$, $S_B = N$, $I_B = 0$, $R_A = 0$, $R_B = 0$). The differential

equations are solved deterministically using the integrate.odeint function in the scipy package

(version 1.6.2) in Python using a timestep of 0.1 days.

The concentration C of the pathogen in wastewater is given by

928
$$C = \frac{n_A + n_B}{v_A + v_B}$$

where n_A and n_B are the number of copies of pathogen genetic material (DNA or RNA) shed per

day by population A and B, respectively, into the sampled wastewater and v_A and v_B are the

volume of wastewater produced per day by population A and B, respectively, into the sampled

wastewater.

The number of copies of pathogen genetic material, n_A and n_B, shed per day by population A

and B, respectively, into the sampled wastewater is given by

938
$$n_A = n I_A f_A$$

939
$$n_B = n I_B f_B$$

Where n is the number of copies of pathogen genetic material shed per infected person per day

(assumed to be the same for all infected individuals) and f_A and f_B are the fraction of total shed

pathogen genetic material by population A and B respectively that is sampled by wastewater

(determined by the sewer connectivity of each of the populations).

The volume of wastewater, v_A and v_B, produced per day by population A and B, respectively,

into the sampled wastewater is given by

$$v_A = v N_A f_A$$

$$v_B = v N_B f_B$$

where v is the volume of wastewater produced per person per day (assumed to be the same for

all individuals). The value used for n was $5 \times 10^{6.7}$ gene copies/person/day (estimate taken from

Ref. [51] by multiplying $10^{4.7}$ gene copies/mL feces by 500 mL feces/day/person) and for v was

20 liters wastewater/person/day (per person daily wastewater usage approximated from per

household daily wastewater usage from Ref. [52] assuming 4 people per household).

In addition to the assumptions mentioned above, the model makes the following additional

assumptions: The probability of an infected individual shedding pathogen genetic material to

wastewater is 1. Given that there is pathogen genetic material present in wastewater, the

probability of detecting it is 1. Shedding of pathogen genetic material occurs only when an

individual is infected. There are no stormwater contributions to wastewater to dilute the

pathogen concentration.

To calculate the time of the maximum number of infections in a population and the time of the

maximum concentration of pathogen gene content in wastewater data, we numerically

calculated the time derivative at the midpoints of the time points as the difference between

consecutive datapoints divided by the time interval and took the first time point at which the

derivative crossed 0.

S2 Appendix. Supplementary analyses.

Compilation of data and assessment of data completeness

As the AHS does not provide finer scale geographic information than the census division for households outside of the 35 selected large metropolitan areas, we used the American Community Survey, which reports the percentage of occupied housing units lacking complete plumbing facilities by county and which allowed us to define the lower bound on the percentage of the population not connected to sewers. Additionally, we identified several state-level datasets, including locations of permitted septic tanks in Florida, sewer locations and populations served in Utah and Minnesota, and the 2012 EPA Clean Watersheds Needs Survey of sewer locations and populations served for 49 states (South Carolina did not participate). However, many of these state-level datasets are incomplete or potentially biased and thus only allow qualitative analyses. Finally, the 2020 US Census' Island Areas Decennial survey provided information the percentage of households connected to sewer by Census Designated Place in the island areas of American Samoa, Guam, the Northern Mariana Islands, and the Virgin Islands (see **Error! Reference source not found.** and Table S 3 for a summary of datasets).

We checked the consistency of the Florida Department of Health onsite sewage treatment and disposal systems inspections data reported in 2012 with other sources. We noted that an estimated 7% of households across the state were connected to septic tanks, but that this was substantially less than the Florida Department of Environmental Protection's estimate that approximately one third of Florida's population uses septic tanks [21] even after accounting for potential correlations of septic usage with household size (approximate increase in household size of 10% for a 50% increase in septic connectivity (**Fig S 5**) cannot explain the almost 5 time

difference in septic connectivity across the two sources), suggesting that more households may

be on septic tanks, and thus less on sewer systems. This discrepancy may exist because not

every septic tank had a permit recorded in the state system, because there were changes since

2012, or because of uncertainty in the Florida Department of Environmental Protection's

estimate. Thus, we focused our analysis on relative comparisons across county subdivisions

(Fig 4).

1001

The 2012 EPA Clean Watersheds Needs Survey dataset includes voluntary submissions of the

locations of publicly owned wastewater collection and treatment facilities and their estimated

population served. The following states had among the highest reported needs and survey

participation level (pg. 6 and Table A-1 of Ref. [24]) and were the focus of our analyses: New

York, California, Florida, New Jersey, Maryland, Iowa, Minnesota, and Michigan. Comparison of

Florida in the EPA and Florida Department of Public Health datasets showed consistent spatial

trends (less connection to sewers and more connection to septic systems in the panhandle and

inland) (Fig 4 and Fig S *9*). Additionally, in both datasets, counties with more connection to

sewers are more Asian (EPA: Pearson r = 0.53 (p = 4.2×10^{-4}), Florida Department of Public

Health: Pearson r = 0.31 (p = 3.2×10^{-7})). Comparison of Minnesota in the EPA dataset,

Minnesota Wastewater Infrastructure Needs Survey, and Minnesota SSTS datasets also

showed consistent spatial trends (more connection to sewers and less connection to subsurface

sewage treatment systems around Minneapolis) (**Fig S 11**). We found a significant correlation

between the AHS and EPA datasets for the largest 35 metropolitan areas in the states with

good data collection (**Fig S 12**), despite deviations from a one-to-one correspondence

suggesting overall systematic biases in the data.

1018

The 2021 Utah Municipal Wastewater Planning Survey estimated the populations receiving

collection by municipal utilities that owned or operated a sanitary sewerage system. The

purpose of the survey is to aid in overseeing and communicating with the wastewater industry in

Utah. We noticed that these data may have not be quantitatively accurate but are likely

qualitatively informative. First, large swaths of the state appeared not to receive wastewater

collection (**Fig S 13**), but this may partly reflect that only 71% of contacts responding to the

survey [47]. Second, in some county subdivisions, >100% of the population appeared

connected to sewers, likely due to errors in estimation, differences in how combined treatment

and collection facilities reported population estimates, or collection populations resided in

neighboring county subdivisions. While the 5 county subdivisions with the highest percentages

of American Indian and Alaska Natives (~20-100%) were reported to not be connected to

sewers at all in this dataset, we believe this is because Indian reservations are not included in

the dataset (as they are under federal, not state jurisdiction). In fact, all 5 of these counties

(Uintah and Ouray, Casa del Eco Mesa-White Mesa, West Juab, Oljato, Blanding) include

Indian reservations. Additionally, the county subdivision with the highest percentage of Black or

African American; Native Hawaiian and Other Pacific Islander; or Hispanic are in county

subdivisions has nobody receiving collection (**Fig S 14**); however, we cannot rule out that this is

due to underreporting from these county subdivisions.

1037

The 2021 Minnesota Wastewater Infrastructure Needs Survey reported whether communities

have a collection system. The data from this survey was geographically sparse, as communities

(the level of aggregation of the dataset) often directly mapped to a census designated place

(CDP), and CDPs do not cover the entire state (**Fig S 11**a). Consequently, the sewer

connectivity of the rural areas between census designated places could not be determined from

this dataset. Additionally, not all communities responded to the survey. Of the 523 communities

that participated in the survey, all but 11 had a collection system. The geographic distribution of

sewer connectivity qualitatively matched that of the 2017 Subsurface Sewage Treatment

Systems (SSTS) Annual Report of the number of reported subsurface sewage treatment

systems (septic tanks) by county in Minnesota (**Fig S 11**b); however, 7 out of the 218 contacted

local government units did not respond, 14 reported having zero SSTS within their jurisdiction

despite permitting SSTS and not all SSTS programs were able to be identified to be contacted.

1050

1051 Given the data incompleteness and biases for the Utah and Minnesota datasets, we excluded

1052 them from the analyses in the main text.

1053

Sewer connectivity in the U.S. Island Areas

From the 2020 US Census Island Areas Decennial Survey of household characteristics, the

overall levels of household sewer connection in the island areas were lower than in the states

(**Fig S 15**). In Guam, the Northern Mariana Islands, the Virgin Islands, and American Samoa,

the overall connection to sewer systems across each island was 65%, 51%, 66%, and 52% of

households, respectively (compared to an overall 83% connection across mainland US). To ask

whether there was strong spatial variability across each island, we looked at the median

household connection by Census Designated Place (CDP). A median of 64%, 32%, 70%, and

13% households by CDP in Guam, the Northern Mariana Islands, the Virgin Islands, and

American Samoa, respectively, were connected to sewers. This suggests that sewer

connectivity is particularly spatially concentrated in the Northern Mariana Islands and American

Samoa (difference in median household connection by CDP and overall island connection),

although all island areas showed some spatial variability in connectivity when visually assessing

maps (**Fig S 16**). The only significant correlation of sewer connectivity with demographic or

economic characteristics was in American Samoa, where CDPs with a higher percentage of

Black or African Americans were more connected to sewers (Pearson r = 0.38, q-value = 0.01).

1070 **Supplementary figures**

1071

1072 **Fig S 1**: **Percentage of households connected to various wastewater processing systems.**

1073 Data are from 2021 American Housing Survey.

1074

1075   **Fig S 2***: **Top 35 metropolitan areas' connectivity to public sewers by demographics.** Data are from the 2019 and 2021 U.S.

1076   Census American Housing Survey [11,16].

Percent of occupied housing units lacking complete plumbing facilities by county

Fig S 3: Percent of occupied housing units lacking complete plumbing facilities by county. Data are from the 2021 U.S. Census American Community Survey.

1080

Fig S 4: Comparison of percentage of households not on septic across different

geographies in Florida. Data are from the Florida Department of Health septic tank inspection

permits reported in 2012. (a) Map of county subdivision colored by whether it is a Metropolitan

Statistical Area (blue), Micropolitan Statistical Area (orange), or Rural Area (neither metropolitan

not micropolitan, green). (b) Percentage of households not on septic in county subdivisions

stratified by whether the county subdivision is a Metropolitan Statistical Area, Micropolitan

Statistical Area, or Rural Area. County subdivisions that are Metropolitan Statistical Areas have

a higher percentage of households not on sewer than those that are Micropolitan Statistical

Areas, but the effect is not significant (p = 0.052). County subdivisions that are Micropolitan

Statistical Areas have a higher percentage of households not on sewer than those that are

Rural Areas, but the effect is not significant (p = 0.37). (c) Map of county subdivisions colored by

whether they are always included in references to the Florida panhandle. Orange: county

subdivisions in the panhandle. Blue: county subdivisions not in the panhandle. (d) Percentage of

households not on septic in county subdivisions stratified by whether the county subdivision is in

the panhandle. Each point represents a county subdivision. County subdivisions in the

panhandle have a significantly lower percentage of households not on septic ($p < 10^{-9}$). (e) Map

of county subdivisions colored by whether they have a coastline. Orange: county subdivision

with coastline. Blue: county subdivisions without coastline. (f) Percentage of households not on

septic in county subdivisions stratified by coastal and not coastal county subdivisions. Each

point represents a county subdivision. County subdivisions not on the coast have a significantly

higher fraction of households on septic ($p<10^{-9}$).

1102

1103

1104

1105   **Fig S 5***: **Correlation of percentage of households in county subdivisions of Florida** *not*

1106   **connected to septic tanks with demographic and economic factors.** Data are from the

1107   Florida Department of Health septic tank inspection permits reported in 2012. Each point

1108   represents a county subdivision. Only county subdivisions with at least 5 households and 20

1109   population size were included.

1110

1111

Fig S 6: Correlation of percentage of households in county subdivisions of Florida in metropolitan statistical areas not connected to septic tanks with demographic and economic factors. Data are from the Florida Department of Health septic tank inspection

1115   permits reported in 2012. Each point represents a county subdivision. Only metropolitan county

1116   subdivisions with at least 5 households and 20 population size were included.

1117

1118 **Fig S 7: Correlation of percentage of households in county subdivisions of Florida in**

1119 **micropolitan statistical areas not connected to septic tanks with demographic and**

1120 **economic factors. Each point represents a county subdivision.** Data are from the Florida

1121   Department of Health septic tank inspection permits reported in 2012. Only micropolitan county

1122   subdivisions with at least 5 households and 20 population size were included.

1123

Fig S 8: Correlation of percentage of households in county subdivisions of Florida in rural areas not connected to septic tanks with demographic and economic factors. Data are from the Florida Department of Health septic tank inspection permits reported in 2012. Each

1127   point represents a county subdivision. Only rural county subdivisions with at least 5 households

1128   and 20 population size were included.

1129

1130

1131

Fig S 9: Fraction of present resident population receiving collection by county in 8

states. Data are from the 2012 EPA Clean Watersheds Needs Survey. (a) Fraction shown on a

continuous scale. (b) Fraction above and below 0.2. Only states that had more comprehensive

responses in the survey are shown. Counties that did not report any data in the survey are

shown in gray. Counties than report more than 100% of the present resident population

receiving collection are shown in (a) as 1 and in (b) as ≥0.2.

1138

1139

1140

1141   **Fig S 10: Sensitivity analysis of Fig 5c from main text**. The effect on the time between

1142   population B peak and wastewater peak of varying the (a) population size of the two populations

1143 (N_A, N_B), (b) recovery rate (γ_I in units of inverse days), and (c) basic reproduction number of the

1144   two populations (R_0^A, R_0^B) across a range of values. Unless where indicated, the parameters

1145   were $N_A = N_B = 5000$, $\gamma_I = 0.18$ days^{-1}, and R_0^A, $R_0^B = 1.5$.

1146

Fig S 11: Sewer connectivity in Minnesota. (a) Map showing whether a community in

Minnesota (mapped to Census Designated Place) reported having a collection system in the

2021 Minnesota Wastewater Infrastructure Needs Survey. Almost all communities mapped to

Census Designated Places. (b) Percentage of households without subsurface sewage treatment

systems (i.e. without septic tanks, suggesting sewered) by county reported in the 2017

Subsurface Sewage Treatment Systems in Minnesota Annual Report [49]. Counties reporting

more than 100% of households having subsurface sewage treatment systems are set at 100%

(shown as 0% without subsurface sewage treatment systems in the map).

1155

Fig S 12*: Comparison of the 2012 EPA Clean Watersheds Needs Survey (CWNS) and the

2019 and 2021 U.S. Census American Housing Survey (AHS) datasets in core based

statistical areas (CBSA). Data from the EPA CWNS are of the fraction of residents in the

CBSA receiving sewage collection. Data from the AHS are of the fraction of households in the

CBSA on sewer. Only CBSAs that were oversampled in the AHS data and were in states that

had more comprehensive responses in the EPA CWNS were included (see Methods). The

dashed black line shows y=x as a reference.

% of population receiving collection by county subdivision in Utah

1163

1164 **Fig S 13: The percentage of the population that received wastewater collection by a**

1165 **municipal utility in Utah by county subdivision of Utah.** Data are from the 2021 Utah

1166 Municipal Wastewater Planning Survey. All values less than 0 are displayed as 0. County

1167 subdivisions with fewer than 5 households or 20 population size are displayed in white. Note

1168 that Indian reservations were not surveyed in this dataset.

1169

Fig S 14: Correlation of fraction households in county subdivisions of Utah with

wastewater collection with demographic and economic factors. Data are from the 2021

Utah Municipal Wastewater Planning Survey. Note that Indian reservations were not surveyed

1173   in this dataset. Values of the fraction of the county subdivision receiving collection greater than

1174   1 were set to 1.

1175

1176

1177   **Fig S 15: Fraction of households by Census Designated Place that are connected to**

1178   **sewers, septic tanks, or other forms of sewage disposal in the U.S. Island Areas.** (a)

1179   American Samoa, (b) Guam, (c) the Northern Mariana Islands, and (d) the Virgin Islands. Data

1180   are from the 2020 U.S. Census Island Areas Decennial Survey.

1181

Fig S 16: Map of fraction of households connected to sewers by Census Designated Place. (a) American Samoa, (b) Guam, (c) the Northern Mariana Islands, and (d) the Virgin Islands. Data are from the 2020 U.S. Census Island Areas Decennial Survey.

1185

1186

 # Supplementary tables

1188 **Table S 1: Counties in the US with >=5% of occupied housing units lacking complete**

1189 **plumbing facilities.** Data are from the 2021 U.S. Census American Community Survey.

	County, State	Percent of occupied housing units lacking complete plumbing facilities
0	Yukon-Koyukuk Census Area, Alaska	36.1
1	Bethel Census Area, Alaska	31.1
2	Kusilvak Census Area, Alaska	25.3
3	Nome Census Area, Alaska	24.5
4	Northwest Arctic Borough, Alaska	20
5	Denali Borough, Alaska	18.1
6	Lake and Peninsula Borough, Alaska	13.8
7	Apache County, Arizona	12.8
8	Dillingham Census Area, Alaska	12
9	Southeast Fairbanks Census Area, Alaska	11.1
10	North Slope Borough, Alaska	10.8
11	Haines Borough, Alaska	10.5
12	Copper River Census Area, Alaska	10.1
13	San Juan County, Utah	10
14	Adams County, Indiana	9.7
15	McKinley County, New Mexico	9.6

16	Oglala Lakota County, South Dakota	9.2
17	Hoonah-Angoon Census Area, Alaska	7.6
18	Mora County, New Mexico	7.6
19	Navajo County, Arizona	7
20	Prince of Wales-Hyder Census Area, Alaska	6.7
21	Lawrence County, Mississippi	6.2
22	Catahoula Parish, Louisiana	6.1
23	Hudspeth County, Texas	6.1
24	Fairbanks North Star Borough, Alaska	5.6
25	Salinas Municipio, Puerto Rico	5.1
26	Yakutat City and Borough, Alaska	5.1

1190

1191

1192 **Table S 2: Correlation of the fraction of households by county connected to public**

1193 **sewers and different demographic or economic factors, stratified by state and**

1194 **Metropolitan, Micropolitan, and Rural Statistical Areas.** Only correlations that are significant

1195 (q-value<=0.05) are shown.

Demographic or economic variable	State	Metropolitan, Micropolitan, or Rural Statistical Area	Pearson r	q-value
Median income	FL	Micropolitan Statistical Area	0.98	9.8E-03
Percent Hispanic	FL	Metropolitan Statistical Area	0.53	1.8E-02
Percent one race and Asian	MN	Metropolitan Statistical Area	0.72	2.3E-03
	FL	Metropolitan Statistical Area	0.52	3.0E-02
	MN	Metropolitan Statistical Area	0.73	1.1E-03

Percent one race and					
Black or African American	MI	Metropolitan Statistical Area		0.67	1.4E-02
	MN	Metropolitan Statistical Area		-0.74	7.6E-04
Percent one race and	MI	Metropolitan Statistical Area		-0.71	4.6E-03
White	IA	Metropolitan Statistical Area		-0.69	5.0E-02

1196

1197

Table S 3: Additional datasets discussed in the Supplementary Information but excluded

from the main text due to data incompleteness and potential biases.

Dataset	Location	Year(s)	Description	Sampling method	Dataset completeness and potential biases	Ref.
Utah Municipal Wastewater Planning Survey	Utah	2021	This survey reported the population receiving collection by each municipal utility that owns or operates a sanitary sewerage system.	The survey completed was by treatment plant operators. Participation was reported in the survey information as being mandatory.	Of those contacted, 71% responded to the survey [47]. In some county subdivisions, >100% of the population appeared connected to sewers, likely due to errors in estimation, differences in how combined treatment and collection facilities reported population estimates, or collection populations residing in neighboring county subdivisions.	[47]
Minnesota Wastewater Infrastructure Needs Survey (WINS)	Minnesota	2021	Communities (places and townships) responded to the question "does your community have a collection system?" (yes/no/did not respond).	This was a voluntary survey.	Some communities (number unknown) did not respond to the survey.	[48]
Minnesota Subsurface Sewage Treatment Systems (SSTS)	Minnesota	2017	This dataset reports the number of subsurface sewage treatment systems (SSTS) in various local government units.	Surveys were sent to local government units with a known SSTS program. According to the SSTS Annual Report [49], 'some of the data is "hard" data, such as the reported number and types of permits issued. Other data is considered "soft" data, such as the reported best estimates provided to determine SSTS	Two hundred and eleven out of 218 known SSTS programs submitted an annual report. Not all SSTS programs were identified and contacts were not always provided. Fourteen reported having zero SSTS within their jurisdiction despite permitting SSTS in 2017.	[49]

				compliance rates and, sometimes, the total number of SSTS in each jurisdiction.'	

1201

1202 **Table S 4: Manual changes community names reported in the Minnesota Wastewater**

1203 **Infrastructure Needs Survey to be able to match with a U.S. Census Designated Place.**

Original name in dataset	Edited name for merging with Census Designated Place name	Note
St X (i.e. St Paul)	St. X (i.e. St. Paul)	
Plainview - Elgin Sanitary District	[Deleted]	There was another entry for Plainview (same zip code) with the same response for whether there was collection facility
Clear Lake - Clearwater Sewer Authority	Clear Lake	
Tower-Breitung Wastewater Board	Tower	
Birchwood	Birchwood Village	
Coleraine, Bovey, Taconite Joint Wastewater Commission	[Deleted]	Two entries for Coleraine with same zip code and similar population sizes
Glacial Lakes Sanitary Sewer and Water District	[Deleted]	Two entries for Spicer with same zip code and similar population sizes

Central Iron Range Sanitary Sewer District	[Deleted]	Two entries for Chisholm with same zip code and similar population sizes
Shorewood Park Sanitary District	[Deleted]	Two entries for Rush City with same zip code. Deleted the entry with the smaller population size.
Rich Prairie Water & Sewer District (Pierz/Genola)	Pierz	
Pine River Area Sanitary District	Pine River	
Moose Lake Windemere Area Sanitary Sewer District	[Deleted]	Two entries for Moose Lake with same zip code. Deleted the entry with no reported population
Alexandria Lakes Area Sanitary District	Alexandria	
North Long Lake Sewer District	Brainerd	
Serpent Lake Sanitary Sewer District	Crosby	
Blomkest/Svea Sewer Board	Blomkest	
GEM Sanitary District	Greenwald	
Chisago Lakes Joint Sewage Treatment Commission	Center City	

Duluth	[Deleted]	3 entries in Duluth - kept the one with the largest population size (Western Lake Superior Sanitary District (WLSSD))
Duluth – North Shore Sanitary District	[Deleted]	3 entries in Duluth - kept the one with the largest population size (Western Lake Superior Sanitary District (WLSSD))
MCES - Metro Council Env. Services	[Deleted]	Two entries in St. Paul. Deleted the entry with the smaller population size.

1204