THE NEURAL CORRELATES OF AUTONOMIC INTEROCEPTION: PERCEPTUAL MISMATCH AND CLINICAL IMPLICATIONS

A PREPRINT

Poppy Z. Grimes[*]
Division of Psychiatry
Centre for Clinical Brain Sciences
University of Edinburgh, UK
p.grimes@ed.ac.uk

Christina N. Kampoureli
Department of Clinical Neuroscience
Brighton and Sussex Medical School
University of Sussex, UK

Charlotte L. Rae
School of Psychology
University of Sussex, UK

Neil A. Harrison
Cardiff University Brain Research
Imaging Centre (CUBRIC)
University of Cardiff, UK

Sarah N. Garfinkel
Institute of Cognitive Neuroscience
University College London, UK

Hugo D. Critchley[†]
Department of Clinical Neuroscience
Brighton and Sussex Medical School
University of Sussex, UK

Jessica A. Eccles[‡]
Department of Clinical Neuroscience
Brighton and Sussex Medical School
University of Sussex, UK

May 22, 2023

ABSTRACT

Interoceptive mismatch is a perceptual discrepancy between ascending bodily signals and higher-order representation of anticipated physiological state. This mismatch is implicated in emotion inference within predictive coding models. We present *autonomic perceptual mismatch* as a novel in vivo measure of the discrepancy between actual and perceived autonomic signals, for clinical application to brain-body interactions, specifically in the expression of anxiety. Joint hypermobility is disproportionately found in individuals with anxiety disorders, revealing an underexplored link between the mind and body. Atypical autonomic reactivity represents a likely mediating mechanism consequent of altered connective tissue within the vasculature and nervous system.

The present fMRI study determined neural substrates of autonomic perceptual mismatch on affective processing in the hypermobility-anxiety interaction. We compared regional brain activity during emotional face processing in participants with and without hypermobility and generalized anxiety disorder diagnosis, then tested association with perceptual mismatch. In the brain, autonomic perceptual mismatch correlated with enhanced activation in emotion processing and autonomic control regions, notably anterior cingulate cortex. Anxious individuals exhibited increased mid-insula cortex activity in relation to perceptual mismatch. Activity was decreased within the inferior frontal gyrus, a region implicated in cognitive control. Dysautonomia mediated the link between hypermobility and anxiety. Together, these findings support a neural basis of an autonomic perceptual mismatch model in a clinical sample. This is supported by the engagement of neural systems for emotion-cognition and interoception. This work highlights convergent aspects of neurodiversity, mental health, connective tissue disorders and brain-body interactions relevant to precision healthcare.

Keywords interoception · dysautonomia · brain-body interactions · hypermobility · anxiety · predictive coding

---
[*]Previous: Dept. of Clinical Neuroscience, Brighton and Sussex Medical School, University of Sussex, UK
[†]Sussex Partnership National Health Service Foundation Trust, Brighton, UK
[‡]Sussex Partnership National Health Service Foundation Trust, Brighton, UK

1 Introduction

The coupling of brain and body is evident in cognitive and emotive responses to visceral afferent signals. Interoception, the internal sensing of our milieu intérieur, its representation in brain, and its impact on psychological processes including bodily feelings, is proposed to be central to emotion [1–4]. Afferent interoceptive information is conveyed centrally by viscerosensory nerves that permit the reflexive and allostatic [5] autonomic nervous control of internal physiology [6]. Dysautonomia describes perturbation of adaptive autonomic control and may arise through aberrant interoceptive signalling, representation and regulation. Notably, dysautonomic symptoms are common to many mental and physical conditions.

Anxiety is commonly associated with heightened states of autonomic state of arousal linked to anticipatory fear and worry. The physiological signalling and perception of arousal amplify negative feelings as subjective anxiety symptoms and associated avoidant behaviours. Anxiety is thus linked to uncertainty regarding internal states (generalised anxiety disorder) or external situations (panic and social anxiety disorders) [7]. Arguably, anxious feeling arises from unexplained arousal and the mismatch between anticipated/desired actual interoceptive signalling [8]. Correspondingly, discrepancies in accurately discerning interoceptive signals provide a mechanistic explanation for the precipitation of anxiety by dysregulated bodily states [2, 9, 10].

Interoception provides an empirical framework for the relationship between autonomic dysfunction and anxiety [2, 7, 8, 11]. Recent studies link the expression of anxiety to attenuated interoceptive awareness (interoceptive metacognitive insight), computed as the correspondence (or mismatch) between objective measures of a person's interoceptive sensitivity (from performance accuracy on interoceptive tasks) and their subjective perception of their own interoceptive sensitivity (rated confidence in interoceptive performance accuracy) [12]. Interoceptive mismatch (previously termed interoceptive trait prediction error), has been conceptualised as the discrepancy between self-reported awareness measures of interoception (e.g. assessed through the awareness subscale of the Porges Body Awareness Questionnaire; [13]) relative to behavioural accuracy on interoceptive tests (e.g. heartbeat perception). This interoceptive mismatch is also associated with heightened anxiety symptomatology [14].

The interoceptive predictive coding model (IPC) links aberrant interoceptive processing (uncorrected 'prediction errors') to dysautonomia and anxiety [15, 16]. Heightened attention to physiological responses increases the precision afforded to interoceptive information, increasing the likelihood of errors in autonomic control and interoceptive inference [8, 15, 17]. Unsuccessful top-down suppression of error results in less accurate perception, reflecting a mismatch between expected and actual internal states.

To investigate this mismatch in dysautonomia, we developed an interoceptive measure of autonomic perceptual mismatch (APM) – corresponding to the magnitude of mismatch in objective and subjective sensitivity to autonomic signals.

Joint hypermobility is an outward manifestation of a more general variation in the structural integrity of connective tissues, including collagen and other matrix proteins present throughout the body [18]. Although hypermobility is

common (present in roughly 20% of the general population) some individuals develop symptoms that can affect multiple bodily systems: Hypermobility is associated with chronic pain, fatigue, gastrointestinal disturbance, neurodevelopmental and neuropsychiatric conditions [19–21]. While typically underdiagnosed, the complex comorbidities of symptomatic hypermobility can go unrecognised in individuals, hindering quality-of-life [19, 22–24].

Hypermobility is significantly overrepresented among people with clinical diagnoses of anxiety [25, 26] and present in up to 70% of such patients [18, 27]. Hypermobile-anxious individuals are more vulnerable to side effects from anxiolytics and anti-depressants, and experience autonomic and gastrointestinal-associated symptoms that can amplify affective anxiety [24].

Despite the acknowledgement of this association between hypermobility and anxiety, extending to autonomic and somatic symptoms, a neurobiological account is yet to be elucidated. One such mechanism may exist in interoceptive autonomic perceptual mismatch.

Dysautonomia in hypermobility may manifest as a consequence of reactive autonomic regulation of less elastic vascular tissues [28, 29]. Imprecise feedback control of peripheral blood flow putatively results in physiological symptoms and compensatory autonomic and behavioural responses. These can be interpreted as feelings of distress [30]. Peripheral autonomic dysregulation increases interoceptive surprise, wherein despite a relative reduction in interoceptive accuracy, there is a subjective overestimation of confidence in perceiving interoceptive experiences. Such frequent unanticipated feelings of arousal may fuel negative affect including anxiety [28, 31].

Hypermobile individuals show higher subjective sensitivity to interoceptive sensations [29] and interoceptive accuracy influences the relationship between hypermobility and anxiety level [32]. Based on these observations, we predicted the important interaction between autonomic perceptual mismatch, hypermobility and anxiety will be evident in distinct brain activation patterns.

Orthostatic intolerance (OI) is the onset of autonomic symptoms upon standing, such as palpitations and dizziness, typically associated with increased heart rate and low blood pressure [33]. OI represents a quantifiable measure of dysautonomia [10, 11, 34], has a 78% prevalence in hypermobility [35] and symptoms are higher in anxious individuals [11]. Notably, OI mediates the relationship between hypermobility and diagnosis of anxiety [20], substantiating the premise of altered connective tissue impeding autonomic function.

Functional magnetic resonance imaging (fMRI) has shown engagement of the same brain regions during anxiety states that support interoceptive representations and autonomic control in the limbic system [7]. The insula cortex is a putative mismatch computation site, shown to respond to altered physiological feedback and relay information to anterior cingulate cortex [36, 37].

To date, few brain imaging studies have examined neural mechanisms that may link hypermobility to anxiety through interoception. The presence of hypermobility increases reactivity within insula and related regions associated with anxiety [32]. Hypermobility has also been associated with increased bilateral amygdala volume [29] structural

differences within insula, correlating with increased orthostatic heart rate and interoceptive accuracy in anxious-
hypermobile participants [38].

The present study investigated the brain substrates of autonomic perceptual mismatch using the hypermobility-anxiety
interaction as a clinical model. Using fMRI to identify patterns of regional neural activity that correlate with APM
in the anxiety-hypermobility interaction, we hope to characterise the neural mechanisms that can ultimately inform
precision healthcare.

2 Materials and Methodology

Participants and psychometric measures   Fifty-one participants, matched for age and gender, were recruited to
the study at the Clinical Imaging Sciences Centre at the University of Sussex (Brighton, UK). Of the participants, 26
(51.0%) (mean age 41.4 12.2 SEM, 9 male) had a diagnosis of generalised anxiety disorder (DSM-IV) as confirmed by a
clinician using the Mini-International Neuropsychiatric Interview (MINI; [39]). The remaining 25 (49.0%) participants
(37.8 14.4, 14 male) were healthy controls with no diagnosed psychiatric condition. The Brighton diagnostic criteria
[40] was applied to the classification of hypermobility. Of those with a diagnosis of generalised anxiety disorder, 18
(69.2%) were classified as hypermobile. Six (24.0%) of the healthy controls were classed as hypermobile.

Participant exclusion criteria included MRI incompatibility, neurological condition and any other psychiatric condition
except anxiety and depression in the anxious group. In addition to categorical classification of anxiety and/or hyper-
mobility status, the Beck Anxiety Inventory (BAI; [41]) and the Beighton scale of joint laxity (BS; [42]) were used
to quantify the degree of anxiety and hypermobility respectively. Subjective scores for orthostatic intolerance were
recorded using the orthostatic subscale of the Autonomic Symptoms and Quality of Life scale (AQQoL; [43]).

Clinically anxious participants were recruited from Sussex Partnership NHS trust and via electric bulletin boards.
Controls were recruited via bulletin boards at Sussex and Brighton Universities. The study procedure was ethically
approved by the Brighton and Hove NRES committee (ref 12/LO/1942).

Statistical Analyses   Behavioural results were first assessed using bivariate correlations between continuous variables,
autonomic perceptual mismatch and anxiety score. Subsequently, independent samples t-tests (two-tailed) were
performed to investigate differences in mean autonomic perceptual mismatch scores for hypermobile and anxious
participants versus controls. Psychophysiological interactions were computed using univariate interaction analyses
in the General Linear Model (GLM). Variables were entered as fixed factors (categorical) or dependent variables
(continuous), and perceptual mismatch was entered as the covariate. Sum-of-squares Type III method was used with
intercept included in the model.

Autonomic perceptual mismatch computed as orthostatic intolerance mismatch   Objective autonomic testing
was performed using an active stand test of orthostatic intolerance, which measured heart rate (HR) change from lying
down to one of minute standing. Participant heart rates were recorded using a pulse oximeter (NONIN, Nonin Medical,

Minnesota, USA). Heart rates from lying to standing were recorded at baseline, peak and after one minute of standing.

Changes in heart rate for each participant were calculated as absolute values and proportional to baseline for peak rate

or rate after standing. Subjective scores for orthostatic intolerance were recorded based on the autonomic subsection of

the AQQoLS.

Autonomic perceptual mismatch was computed as the mismatch between signs (OI stand test) and symptoms (AQQoLS)

of orthostatic intolerance in the same framework as interoceptive trait prediction error [12]. This was calculated as

$$abs[Z(proportional HRchange) - Z(AQQoLsscore)] \tag{1}$$

where Z is the standard Z-score. Absolute values were used to investigate the magnitude of error with neural activation.

Final APM scores were assigned to participants as the transformed mismatch between orthostatic intolerance signs and

symptoms, as a type of interoceptive prediction error. The results of all autonomic testing are available in Supplementary

Table 1.

Neuroimaging paradigm: Stimuli and experimental design   The in-scanner task was modified from Umeda et al.

(2009)[44]. Emotional faces were selected and grouped into five classes (angry, afraid, disgusted, neutral, happy) from

the Karolinska Directed Emotional Faces (KDEF; [45]). Faces were presented in an event-related design and were

randomised and counter-balanced across runs. Attendance to faces was ensured by asking participants to determine

whether they could see teeth in the faces or not. In total there were 96 trials per participant; there were 15 trials per

class of emotional face and 21 fixation cross trials used as the implicit baseline which served as the control condition.

Each stochastically ordered trial was 4 seconds long, during which the face remained on screen, and the participant was

expected to respond. The fixation cross duration was also 4 seconds in duration. A total run lasted 384s (6m 24s) and

each participant underwent two runs of the emotional faces task.

Image Acquisition   Neuroimaging took place using a 1.5 Tesla Siemens Avanto Scanner with a 32-channel head

coil (Siemens Medical Solutions, Erlangen, Germany). T1 structural scans were first acquired for each participant

using a magnetisation-prepared rapid gradient-echo acquisition (repetition time TR = 2.73s per volume, echo time TE =

3.57msec, inversion time = 1000ms, flip angle = 7°). T2*-weighted whole-brain functional scans were taken using a

single-shot 2D gradient echo-planar imaging (EPI) sequence. For the functional scans, voxel sizes were 3x3x3mm,

repetition time TR = 2.52s per volume, echo time TE = 43ms. 34 axial slices of 3mm thickness and 0.6mm interslice

gap were taken. Slices were tilted at a 30° flip angle from the intercommissural plane to minimise signal artefacts.

Pre-processing   Imaging data were processed in SPM12 (http://www.fil.ion.ucl.ac.uk/spm) on MATLAB R2021b

(v9.9.0 Mathworks Inc.). Echo planar images were realigned to the mean image for motion correction, scanner drift and

variation. Slice-time correction to slice 6 (which aligned with amygdala) and was performed for all volumes to remove

artefacts. The first two volumes were discarded for scanner equilibration. T2*-functional scans were co-registered with

T1-structural scans for each participant. Images were normalised to the MNI-152 (Montreal Neurologic Institute) brain

space. Spatiotemporal variation was reduced by smoothing the data using an 8mm full-width half-maximum Gaussian

smoothing kernel filter.

Univariate neuroimaging analysis   First- and second-level designs were implemented and analysed under the general

linear model (GLM) in SPM12 on the whole brain. First-level modelling included a regressor for each of the five

emotional face conditions and the implicit baseline (control condition) of each individual. Participant motion from image

realignment (3 translations and 3 rotations) was included as six regressors of no interest for each participant at the first

level. Statistical comparison of voxel-wise parameters was conducted for first-level contrasts within subjects (emotional

face>baseline), and these contrasts for the five emotions were entered into the second level where a full-factorial design

(2x2x5) was used for component interaction analysis.

The first factor had two levels (hypermobile, non-hypermobile), the second factor had two levels (anxious, non-anxious)

and the third factor had five levels (angry, afraid, happy, neutral, sad). Regressors were entered for age and gender. In

this second-level model, autonomic perceptual mismatch was added as a covariate to investigate the correlations and

interactions of autonomic perceptual mismatch on hypermobility and anxiety. All covariates were mean-centred at zero.

Statistical comparison of voxel-wise parameters was conducted for second-level contrasts between subjects

(anxious>non-anxious), (hypermobile>non-hypermobile). A family-wise error cluster correction (FWEc) was per-

formed on the whole brain at $P<0.05$ to correct for multiple comparisons. This reduced the likelihood of type-I errors,

thus minimising false positives. A cluster-forming threshold of $P<0.001$ for cluster-wise false discovery rate (FDR) was

applied to statistical contrast images. Brain activations were interpreted as clusters that produced a significant change in

BOLD response to the emotional faces fMRI task (relative to the baseline of fixation cross).

Clusters were anatomically defined using the SPM Anatomy toolbox v3.0 (Eickhoff et al., 2007). Where labelling

was not available in the anatomy toolbox, the Harvard-Oxford anatomical atlas was used by overlaying the relevant

t-contrasts onto the 'ch2better' template on MRIcron v1.0.20190902 (Rorden and Brett, 2000).

For each participant, the eigenvariates were extracted as the weighted mean of the Blood-Oxygen-Level-Dependency

(BOLD) time series from the peaks of activation by entering '»clustermean=mean(xY.y')' into the command window of

MATLAB. Eigenvariates were averaged over the five conditions for each participant for each t-contrast of interest and

used to generate scatter plots against autonomic perceptual mismatch.

Mediation   The interaction between hypermobility, autonomic perceptual mismatch and anxiety was investigated

further as a secondary analysis using a Baron and Kenny (1986)[46] mediation analysis. Mediation testing conducts

three regression correlations between pairs of variables. If when controlling for one factor (the potential mediator),

the correlation between the remaining factors is reduced, then the controlled factor is said to partially mediate their

relationship. Based on previous findings that autonomic dysfunction mediates the anxiety-laxity link [20], we entered

APM as our model mediator. Consistent with development chronology, hypermobility (status and score) was employed

as the independent variable, appearing in early childhood [27, 47]. Anxiety manifests later, thus being entered as the dependent or outcome variable.

3 Results

3.1 Autonomic testing

Orthostatic intolerance mismatch Both signs (Figure 1A) and symptom scores (Figure 1B) of orthostatic intolerance were higher in anxious-hypermobile participants indicating a possible 'mismatch error' of autonomic dysfunction

Figure 1: Orthostatic Intolerance (OI) in Hypermobility and Anxiety. Objective OI sign measured as the absolute heart rate change after 1 min sitting to standing (A). Subjective OI symptom score from the AQQoLS (B). Error bars represent ±1SE.

Phenotypic correlations of autonomic perceptual mismatch Anxiety score was positively correlated with autonomic perceptual mismatch across all participants (Rpcc=0.425, P=0.002, Figure 2A). When subgrouping by clinical anxiety status, the positive correlation of autonomic perceptual mismatch with anxiety score was significantly higher in those with anxiety (independent samples t-test; t=-2.964, p= 0.005, SED=0.149, Figure 2B). Similarly, the hypermobile group showed a significantly stronger positive correlation between anxiety score and autonomic perceptual mismatch (independent samples t-test; t=-2.13, p=0.038, SED=0.155, Figure 2C).

GLM univariate analysis demonstrated a significant interaction between hypermobility and anxiety on autonomic perceptual mismatch (Figure 3, F=6.04, p=0.002). A significant interaction between hypermobility and autonomic perceptual mismatch was found on anxiety status (F=7.66, p=0.008). This interaction effect was also found on anxiety score (F=4.20, p=0.046) with a main effect of autonomic perceptual mismatch (F=5.70, p=0.021). There was no significant interaction between anxiety status and autonomic perceptual mismatch on hypermobility.

Figure 2: Autonomic perceptual mismatch in the hypermobility-anxiety association. Autonomic perceptual mismatch correlates with anxiety score (A). Anxiety score is higher with increased autonomic perceptual mismatch in clinically anxious (B) and hypermobile (C) individuals.

Figure 3: The hypermobility-anxiety interaction on autonomic perceptual mismatch. Participants classed as anxious-hypermobile have significanlly higher APM scores. Bars represent ±1SE.

The APM score for one participant fell above the 95% percentile (APM = 2.56, Supplementary Table 1). Although still representable of an extreme case of perceptual mismatch in anxiety, we excluded the results of this individual from subsequent statistical neuroimaging analyses. The significant interaction in figure 3 between anxiety and hypermobility on APM remained after exclusion of this participant from the data.

3.2   Univariate functional neuroimaging

Neural correlations with autonomic perceptual mismatch   The first t-contrasts modelled the BOLD response to emotional faces to identify significant clusters of activation that varied with autonomic perceptual mismatch when added as a covariate to the second-level model. Activations were across all individuals, including controls. Autonomic

perceptual mismatch correlated with response to emotional faces in the inferior frontal gyrus (pars triangularis and pars opercularis) and middle insular cortex. Full results of all imaged activations are available in Supplementary Table 2.

Interaction with anxiety   The interaction of anxiety and autonomic perceptual mismatch produced a positive response (regions showing increased activation with APM in anxious participants) within the right anterior cingulate gyrus (Figure 4A) and left middle insular cortex (Figure 4B). Autonomic perceptual mismatch produced a negative response in the right inferior frontal gyrus (Figure 4C) and anterior mid-cingulate cortex (Figure 4D) in anxious participants. Weighted mean BOLD estimates for the anterior cingulate gyrus and mid-insula were positively correlated with APM; inferior frontal gyrus and mid-cingulate estimates were negatively correlated with APM in anxiety (Figure 4E).

Interaction with hypermobility   The interaction of hypermobility and autonomic perceptual mismatch produced a positive response (regions showing increased activation with APM in hypermobile participants) in the left mid-cingulate gyrus (Figure 5A). Autonomic perceptual mismatch produced a negative response in the left inferior frontal gyrus in hypermobile participants (Figure 5B). Weighted mean BOLD estimate for the anterior cingulate gyrus was positively correlated with autonomic perceptual mismatch and inferior frontal gyrus estimates was negatively correlated in hypermobility (Figure 5C).

Mediation   Upon mediation testing, we found autonomic perceptual mismatch to partially mediate the relationship between hypermobility and anxiety status (Figure 6A; categorical variables) when controlled for. Autonomic perceptual mismatch fully mediated the relationship between hypermobility and anxiety scores (Figure 6B; continuous variables) when controlled for.

Figure 4: T-contrast estimates of the interaction between anxiety and autonomic perceptual mismatch. Emotional face stimuli increased activation in the anterior cingulate (A) and mid-insula (B). Activation was decreased in the IFG (C) and mid-cingulate (D). Colour bars represent peak-level t-statistics at FWEc P<0.05. Crosshairs represent MNI co-ordinates of cluster peak activation. Scatter plots show the interaction between APM and anxiety at mean peak-clusters of activation from extracted eigenvariates (E).

Figure 5: T-contrast estimates of the interaction between hypermobility and autonomic perceptual mismatch. Emotional face stimuli induced an increased response in the anterior cingulate (A) and reduced response in the IFG (B). Colour bars represent peak-level t-statistics at FWEc P<0.05. Crosshairs represent MNI co-ordinates of cluster peak activation. Scatter plots show the interaction between APM and hypermobility at mean peak-clusters of activation from extracted eigenvariates (C).

Figure 6: Baron and Kenny mediation. Partial mediation of APM on effect of hypermobility status on anxiety status (A). Full mediation of APM on effect of Beighton score on Beck's anxiety score (B). Reported correlations are standardised beta coefficients.

4 Discussion

Autonomic perceptual mismatch is a novel measure of dysautonomia that builds upon the interoceptive predictive coding framework. Our research has shown that autonomic perceptual mismatch implicates functional brain activity and is positively associated with anxiety. In a clinical sample, we provide evidence that dysautonomia may be a mechanism by which hypermobility evokes anxiety through a discrete set of brain regions that support interoceptive representation and autonomic control.

Our findings are consistent with prior research which showed differences in brain activity and a mediating effect of objective interoception on anxiety, in a non-clinical hypermobile group [32]. However, this previous study did not consider the mismatch between objective and subjective awareness.

Altered emotion-autonomic processing in anterior cingulate cortex Increased anterior cingulate cortex (ACC) activation was observed in both anxious and hypermobile participants with autonomic perceptual mismatch. ACC is implicated in emotion processing, particularly negative emotions like anxiety, and interoceptive accuracy [2, 48–50]. Imaging studies demonstrate high ACC activity in people with high trait anxiety and involvement in the physiological response to stress as a locus of autonomic regulation [51–55]. Our findings support the significant role of ACC in emotion-autonomic regulation, contributing to the development and persistence of anxiety disorders.

A role of mid-insula in interoceptive anxiety Interestingly, we found a positive response to emotional faces in the middle insula cortex in the interaction of APM with anxiety. Mid-insula is often overlooked in interoception research, as the focus is usually on anterior insula as the mismatch comparator [8, 15, 36, 56]. However, some hypotheses suggest mid-insula to implicate polymodal integration of emotion-interoception processing, with somatosensory and bodily afferent information projecting to the anterior insula to inform emotional experience and intensity [57–61]. Functional connections also exist to cingulate and frontal cortices, where autonomic control pertaining to interoceptive information is elicited and feedback integrated [56, 62–64].

Mid-insula is commonly activated in interoceptive neural activations across psychiatric conditions [65]. Its activity correlates with interoceptive accuracy and anxiety scores, indicating a tole in anxiety-related anxiety interoception [37]. The insula is functionally graded, with the mid lobe as an intermediary structure [66]. Interoceptive responses to bodily awareness and organ distension follow this graded response [2, 67].

Paulus and Khalsa (2021[56]) propose re-evaluating the mid-insula's role in integrating autonomic-interoceptive information. Our observed activity patterns suggest potential alterations of this integration, which could result in mismatched models in the brain and symptom anxiety, as per the IPC framework. The mid-insula's connections to cingulate and frontal regions may contribute to this effect.

Reduced frontal control may predispose anxious symptoms Our study revealed reduced response in a distinct cluster in the anterior mid-cingulate cortex (aMCC) in the interaction of APM with anxiety. aMCC is linked to cognitive

function and error detection, co-activating and functionally connected to insula cortex [1, 51, 62, 68–70]. Important for
pain perception [71–74], attenuation of aMCC activity demonstrated reduced awareness of error detection [75]. aMCC
is also a key locus of the Salience Network for internal-external stimulus recognition [36, 76]. In line with this work,
our observation suggests that reduced activity of the mid-cingulate may induce perceptual mismatch through altered
error detection. Connectivity and co-activation to insula may be a path to the onset of subsequent anxious symptoms.

We observed reduced activation of bilateral inferior frontal gyri (IFG) with autonomic perceptual mismatch in both
hypermobile and anxious individuals. IFG is associated with a variety of cognitive functions such as emotion regulation
and response inhibition [77, 78]. IFG is functionally connected to the insula, playing a role in viscero-motor-autonomic
functioning [79]. Thus, perceptual mismatch may implicate these functional networks, resulting in altered autonomic
function and related cognitive processes. The accumulation of interoceptive errors may contribute to decision-making
errors at a higher level, potentially exacerbating anxiety. However, further research is needed to investigate the
mechanisms underlying emotion-autonomic processing in networks involving IFG and insular cortex.

Dysautonomia as a mediator in brain-body interactions   We demonstrate that autonomic perceptual mismatch
mediates the effect of hypermobility on anxiety. Our study adds to existing evidence that interoceptive accuracy [32]
and orthostatic symptoms [20] mediate this association. We have employed the interoceptive mismatch framework
[14], to elucidate the underlying mechanism of this relationship. Our findings support the hypothesis that dysautonomia
in hypermobile individuals, resulting from altered vascular collagen [28], may explain the observed overlap between
hypermobility and anxiety.

Clinical relevance to the anxiety-hypermobility link   The link between connective tissue disorders and psychiatric
conditions requires a holistic approach to understanding brain-body interactions. The 'Neuroconnective phenotype' [26,
80] is a clinical recognition of the hypermobility-anxiety link that combines somatic and sensory symptoms, behaviour,
psychiatry and neurodevelopmental conditions [21, 81]. Autonomic symptoms are higher in neurodevelopmental
conditions [20, 81, 82]) and autism is overrepresented by up to 52% in hypermobility [83].

Our work utilising the Neuroconnective model suggests that somatic symptoms may be linked to differences in sensory
processing, which is common to autism and hypermobility, but needs further empirical research. These findings
highlight the need for personalised screening for various conditions in those displaying autonomic dysfunction in
hypermobility.

We did not consider the influence of medication on our results. Hypermobile individuals are more likely to be
prescribed psychotropic drugs [47], which can increase autonomic symptoms and dysfunction to varying degrees [28].
Nonetheless, Eccles (2016;[28]) demonstrated that robust differences in autonomic function of hypermobile individuals
exist regardless of medication status. Exploring medication effect on brain function in the context of hypermobility and
autonomic dysfunction may implicate personalised treatments and phenotyping that account for individual responses to
psychotropic medication.

Conclusions Our study demonstrates how autonomic dysfunction affects neural activity in emotion-processing and cognitive control regions in hypermobility and anxiety. These findings offer a mechanistic explanation by which imprecise autonomic interoceptive signals from bodily afferents manifest as anxiety through autonomic mediation. The involvement of mid-insula in polymodal error weighting provides novel evidence for interoceptive predictive coding models. These findings have important implications for brain-body interactions in connective tissue disorders contributing to anxiety symptomatology.

References

[1] Craig, A. "How Do You Feel — Now? The Anterior Insula and Human Awareness". In: *Nature Reviews Neuroscience* 10.1 (1 Jan. 2009), pp. 59–70. DOI: 10.1038/nrn2555.

[2] Critchley, H. D. et al. "Neural Systems Supporting Interoceptive Awareness". In: *Nature Neuroscience* 7.2 (2 Feb. 2004), pp. 189–195. DOI: 10.1038/nn1176.

[3] Seth, A. K. and Critchley, H. D. "Extending Predictive Processing to the Body: Emotion as Interoceptive Inference". In: *Behavioral and Brain Sciences* 36.3 (June 2013), pp. 227–228. DOI: 10.1017/S0140525X12002270.

[4] Damasio, A. R. *The feeling of what happens: Body and emotion in the making of consciousness*. Houghton Mifflin Harcourt, 1999.

[5] Sterling, P. "Allostasis: A Model of Predictive Regulation". In: *Physiology & Behavior*. Allostasis and Allostatic Load 106.1 (Apr. 12, 2012), pp. 5–15. DOI: 10.1016/j.physbeh.2011.06.004.

[6] Ueno, D., Ohira, H., and Narumoto, J. "Editorial: Interoception and the Autonomic Nervous System: Investigating Affect, Decision-Making, and Mental Health". In: *Frontiers in Neuroscience* 16 (2023).

[7] Critchley, H. D. "Neural Mechanisms of Autonomic, Affective, and Cognitive Integration". In: *Journal of Comparative Neurology* 493.1 (2005), pp. 154–166. DOI: 10.1002/cne.20749.

[8] Paulus, M. P. and Stein, M. B. "An Insular View of Anxiety". In: *Biological Psychiatry* 60.4 (Aug. 15, 2006), pp. 383–387. DOI: 10.1016/j.biopsych.2006.03.042.

[9] Critchley, H. D. and Garfinkel, S. N. "Interoception and Emotion". In: *Current Opinion in Psychology*. Emotion 17 (Oct. 1, 2017), pp. 7–14. DOI: 10.1016/j.copsyc.2017.04.020.

[10] Owens, A. P. et al. "Emotional Orienting during Interoceptive Threat in Orthostatic Intolerance: Dysautonomic Contributions to Psychological Symptomatology in the Postural Tachycardia Syndrome and Vasovagal Syncope". In: *Autonomic Neuroscience: Basic and Clinical* 212 (July 1, 2018), pp. 42–47. DOI: 10.1016/j.autneu.2018.01.004. pmid: 29519640.

[11] Owens, A. P. et al. "The Genesis and Presentation of Anxiety in Disorders of Autonomic Overexcitation". In: *Autonomic Neuroscience: Basic and Clinical* 203 (Mar. 1, 2017), pp. 81–87. DOI: 10.1016/j.autneu.2016.10.004. pmid: 27865628.

[12]   Garfinkel, S. N. et al. "Knowing Your Own Heart: Distinguishing Interoceptive Accuracy from Interoceptive Awareness". In: *Biological Psychology* 104 (Jan. 1, 2015), pp. 65–74. DOI: 10.1016/j.biopsycho.2014.11.004.

[13]   Porges, S. "Body perception questionnaire". In: *Laboratory of Developmental Assessment, University of Maryland* 10 (1993), s15327752jpa5304_1.

[14]   Garfinkel, S. N. et al. "Interoceptive Dimensions across Cardiac and Respiratory Axes". In: *Philosophical Transactions of the Royal Society B: Biological Sciences* 371.1708 (Nov. 19, 2016), p. 20160014. DOI: 10.1098/rstb.2016.0014.

[15]   Seth, A., Suzuki, K., and Critchley, H. "An Interoceptive Predictive Coding Model of Conscious Presence". In: *Frontiers in Psychology* 2 (2012).

[16]   Critchley, H. and Seth, A. "Will Studies of Macaque Insula Reveal the Neural Mechanisms of Self-Awareness?" In: *Neuron* 74.3 (May 10, 2012), pp. 423–426. DOI: 10.1016/j.neuron.2012.04.012. pmid: 22578492.

[17]   Clark, A. "Whatever next? Predictive Brains, Situated Agents, and the Future of Cognitive Science". In: *Behavioral and Brain Sciences* 36.3 (June 2013), pp. 181–204. DOI: 10.1017/S0140525X12000477.

[18]   Bulbena, A. et al. "Anxiety Disorders in the Joint Hypermobility Syndrome". In: *Psychiatry Research* 46.1 (Jan. 1, 1993), pp. 59–68. DOI: 10.1016/0165-1781(93)90008-5.

[19]   Baeza-Velasco, C. et al. "Association between Psychopathological Factors and Joint Hypermobility Syndrome in a Group of Undergraduates from a French University". In: *International Journal of Psychiatry in Medicine* 41.2 (2011), pp. 187–201. DOI: 10.2190/PM.41.2.g. pmid: 21675349.

[20]   Csecs, J. L. L. et al. "Joint Hypermobility Links Neurodivergence to Dysautonomia and Pain". In: *Frontiers in Psychiatry* 12 (2022).

[21]   Casanova, E. L. et al. "The Relationship between Autism and Ehlers-Danlos Syndromes/Hypermobility Spectrum Disorders". In: *Journal of Personalized Medicine* 10.4 (4 Dec. 2020), p. 260. DOI: 10.3390/jpm10040260.

[22]   Bulbena, A. et al. "Joint Hypermobility, Anxiety and Psychosomatics: Two and a Half Decades of Progress Toward a New Phenotype". In: (Mar. 24, 2015). DOI: 10.1159/000369113.

[23]   Castori, M. "Ehlers-Danlos Syndrome, Hypermobility Type: An Underdiagnosed Hereditary Connective Tissue Disorder with Mucocutaneous, Articular, and Systemic Manifestations". In: *International Scholarly Research Notices* 2012 (Nov. 22, 2012), e751768. DOI: 10.5402/2012/751768.

[24]   Bulbena-Cabre, A. and Bulbena, A. "Anxiety and Joint Hypermobility: An Unexpected Association: Examining This Link Can Improve Diagnosis and Treatment of Both Disorders". In: *Current Psychiatry* 17.4 (Apr. 1, 2018), pp. 15–22.

[25]   Eccles, J. A. et al. "Beyond Bones: The Relevance of Variants of Connective Tissue (Hypermobility) to Fibromyalgia, ME/CFS and Controversies Surrounding Diagnostic Classification: An Observational Study". In: *Clinical Medicine* 21.1 (Jan. 1, 2021), pp. 53–58. DOI: 10.7861/clinmed.2020-0743. pmid: 33479068.

[26] Bulbena-Cabré, A. et al. "Updates on the Psychological and Psychiatric Aspects of the Ehlers–Danlos Syndromes and Hypermobility Spectrum Disorders". In: *American Journal of Medical Genetics Part C: Seminars in Medical Genetics* 187.4 (2021), pp. 482–490. DOI: 10.1002/ajmg.c.31955.

[27] García Campayo, J. et al. "Association Between Joint Hypermobility Syndrome and Panic Disorder: A Case–Control Study". In: *Psychosomatics* 51.1 (Jan. 1, 2010), pp. 55–61. DOI: 10.1016/S0033-3182(10)70659-9.

[28] Eccles, J. A. "Hypermobility and Autonomic Hyperactivity : Relevance for the Expression of Psychiatric Symptoms". PhD thesis. University of Brighton, 2016.

[29] Eccles, J. A. et al. "Brain Structure and Joint Hypermobility: Relevance to the Expression of Psychiatric Symptoms". In: *The British Journal of Psychiatry* 200.6 (June 2012), pp. 508–509. DOI: 10.1192/bjp.bp.111.092460.

[30] Sharp, H. E. C., Critchley, H. D., and Eccles, J. A. "Connecting Brain and Body: Transdiagnostic Relevance of Connective Tissue Variants to Neuropsychiatric Symptom Expression". In: *World Journal of Psychiatry* 11.10 (Oct. 19, 2021), pp. 805–820. DOI: 10.5498/wjp.v11.i10.805.

[31] Pollatos, O., Gramann, K., and Schandry, R. "Neural Systems Connecting Interoceptive Awareness and Feelings". In: *Human Brain Mapping* 28.1 (2007), pp. 9–18. DOI: 10.1002/hbm.20258.

[32] Mallorquí-Bagué, N. et al. "Neuroimaging and Psychophysiological Investigation of the Link between Anxiety, Enhanced Affective Reactivity and Interoception in People with Joint Hypermobility". In: *Frontiers in Psychology* 5 (2014).

[33] Nilsson, D. et al. "Orthostatic Changes in Hemodynamics and Cardiovascular Biomarkers in Dysautonomic Patients". In: *PLOS ONE* 10.6 (June 8, 2015), e0128962. DOI: 10.1371/journal.pone.0128962.

[34] Aslanyan, D. et al. *Dissociative Experiences in Fibromyalgia Are Mediated by Symptoms of Autonomic Dysfunction*. Sept. 11, 2019. DOI: 10.1101/19006320. URL: https://www.medrxiv.org/content/10.1101/19006320v1 (visited on 05/19/2023). preprint.

[35] Gazit, Y. et al. "Dysautonomia in the Joint Hypermobility Syndrome". In: *The American Journal of Medicine* 115.1 (July 1, 2003), pp. 33–40. DOI: 10.1016/S0002-9343(03)00235-3.

[36] Menon, V. and Uddin, L. Q. "Saliency, Switching, Attention and Control: A Network Model of Insula Function". In: *Brain Structure and Function* 214.5 (June 1, 2010), pp. 655–667. DOI: 10.1007/s00429-010-0262-0.

[37] Tan, Y. et al. "The Role of Mid-Insula in the Relationship between Cardiac Interoceptive Attention and Anxiety: Evidence from an fMRI Study". In: *Scientific Reports* 8.1 (1 Nov. 22, 2018), p. 17280. DOI: 10.1038/s41598-018-35635-6.

[38] Eccles, J. et al. "Joint Hypermobility Syndrome and Anxiety Disorder: Structural Brain Correlates". In: *European Psychiatry* 41.S1 (Apr. 2017), S233–S234. DOI: 10.1016/j.eurpsy.2017.01.2246.

[39] Sheehan, D. V. et al. "The Mini-International Neuropsychiatric Interview (MINI): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10". In: *Journal of clinical psychiatry* 59.20 (1998), pp. 22–33.

[40] Grahame, R., Bird, H. A., and Child, A. "The Revised (Brighton 1998) Criteria for the Diagnosis of Benign Joint Hypermobility Syndrome (BJHS)". In: *The Journal of Rheumatology* 27.7 (July 2000), pp. 1777–1779. pmid: 10914867.

[41] Beck, A. T. et al. "An Inventory for Measuring Clinical Anxiety: Psychometric Properties". In: *Journal of Consulting and Clinical Psychology* 56 (1988), pp. 893–897. DOI: 10.1037/0022-006X.56.6.893.

[42] Beighton, P., Solomon, L., and Soskolne, C. L. "Articular Mobility in an African Population." In: *Annals of the Rheumatic Diseases* 32.5 (Sept. 1973), pp. 413–418. pmid: 4751776.

[43] Iodice, V. "The Postural Tachycardia Syndrome: A Multi-System Condition. Clinical Features, Pathophysiology, Genetics and Novel Treatment". In: (Dec. 2013). DOI: 10.25560/38561.

[44] Umeda, S. et al. "Functional MRI investigations of emotional processing and autonomic responses in patients with autonomic hyperactivity". In: *Neuroimage* 47.Supp 1 (2009), S39–S41.

[45] Goeleven, E. et al. "The Karolinska Directed Emotional Faces: A Validation Study". In: *Cognition and Emotion* 22.6 (Sept. 1, 2008), pp. 1094–1118. DOI: 10.1080/02699930701626582.

[46] Baron, R. M. and Kenny, D. A. "The Moderator-Mediator Variable Distinction in Social Psychological Research: Conceptual, Strategic, and Statistical Considerations". In: *Journal of Personality and Social Psychology* 51.6 (Dec. 1986), pp. 1173–1182. DOI: 10.1037//0022-3514.51.6.1173. pmid: 3806354.

[47] Bulbena, A. et al. "Joint Hypermobility Syndrome Is a Risk Factor Trait for Anxiety Disorders: A 15-Year Follow-up Cohort Study". In: *General Hospital Psychiatry* 33.4 (July 1, 2011), pp. 363–370. DOI: 10.1016/j.genhosppsych.2011.03.004.

[48] Drevets, W. C. et al. "Pet Imaging of Serotonin 1A Receptor Binding in Depression". In: *Biological Psychiatry* 46.10 (Nov. 15, 1999), pp. 1375–1387. DOI: 10.1016/S0006-3223(99)00189-4.

[49] Mayberg, H. S. et al. "Cingulate Function in Depression: A Potential Predictor of Treatment Response". In: *NeuroReport* 8.4 (Mar. 3, 1997), p. 1057.

[50] Mayberg, H. S. et al. "Regional Metabolic Effects of Fluoxetine in Major Depression: Serial Changes and Relationship to Clinical Response". In: *Biological Psychiatry* 48.8 (Oct. 15, 2000), pp. 830–843. DOI: 10.1016/S0006-3223(00)01036-2.

[51] Critchley, H. D. et al. "Human Cingulate Cortex and Autonomic Control: Converging Neuroimaging and Clinical Evidence". In: *Brain* 126.10 (Oct. 1, 2003), pp. 2139–2152. DOI: 10.1093/brain/awg216.

[52] Wager, T. D. et al. "Brain Mediators of Cardiovascular Responses to Social Threat: Part I: Reciprocal Dorsal and Ventral Sub-Regions of the Medial Prefrontal Cortex and Heart-Rate Reactivity". In: *NeuroImage*. Brain Body Medicine 47.3 (Sept. 1, 2009), pp. 821–835. DOI: 10.1016/j.neuroimage.2009.05.043.

[53] Wang, L., LaBar, K. S., and McCarthy, G. "Mood Alters Amygdala Activation to Sad Distractors during an Attentional Task". In: *Biological Psychiatry* 60.10 (Nov. 15, 2006), pp. 1139–1146. DOI: 10.1016/j.biopsych.2006.01.021. pmid: 16713587.

[54] Nitschke, J. B. et al. "Anticipatory Activation in the Amygdala and Anterior Cingulate in Generalized Anxiety Disorder and Prediction of Treatment Response". In: *American Journal of Psychiatry* 166.3 (Mar. 2009), pp. 302–310. DOI: 10.1176/appi.ajp.2008.07101682.

[55] Drevets, W. C., Savitz, J., and Trimble, M. "The Subgenual Anterior Cingulate Cortex in Mood Disorders". In: *CNS spectrums* 13.8 (Aug. 2008), pp. 663–681. DOI: 10.1017/s1092852900013754. pmid: 18704022.

[56] Paulus, M. P. and Khalsa, S. S. "When You Don't Feel Right Inside: Homeostatic Dysregulation and the Mid-Insular Cortex in Psychiatric Disorders". In: *American Journal of Psychiatry* 178.8 (Aug. 2021), pp. 683–685. DOI: 10.1176/appi.ajp.2021.21060622.

[57] Kerr, K. L. et al. "Altered Insula Activity during Visceral Interoception in Weight-Restored Patients with Anorexia Nervosa". In: *Neuropsychopharmacology* 41.2 (2 Jan. 2016), pp. 521–528. DOI: 10.1038/npp.2015.174.

[58] Kurth, F. et al. "A Link between the Systems: Functional Differentiation and Integration within the Human Insula Revealed by Meta-Analysis". In: *Brain Structure and Function* 214.5 (June 1, 2010), pp. 519–534. DOI: 10.1007/s00429-010-0255-z.

[59] Simmons, W. K. et al. "Keeping the Body in Mind: Insula Functional Organization and Functional Connectivity Integrate Interoceptive, Exteroceptive, and Emotional Awareness". In: *Human Brain Mapping* 34.11 (June 13, 2012), pp. 2944–2958. DOI: 10.1002/hbm.22113. pmid: 22696421.

[60] Craig, A. D. "How Do You Feel?: An Interoceptive Moment with Your Neurobiological Self". In: *How Do You Feel?* Princeton University Press, Dec. 21, 2014. DOI: 10.1515/9781400852727.

[61] Wiech, K. et al. "Differential Structural and Resting State Connectivity between Insular Subdivisions and Other Pain-Related Brain Regions". In: *PAIN* 155.10 (Oct. 2014), p. 2047. DOI: 10.1016/j.pain.2014.07.009.

[62] Taylor, K. S., Seminowicz, D. A., and Davis, K. D. "Two Systems of Resting State Connectivity between the Insula and Cingulate Cortex". In: *Human brain mapping* 30.9 (Sept. 1, 2009), pp. 2731–2745. DOI: 10.1002/hbm.20705. pmid: 19072897.

[63] Vogt, B. A. "Chapter 3 - Cingulate Cortex in the Three Limbic Subsystems". In: *Handbook of Clinical Neurology*. Ed. by Vogt, B. A. Vol. 166. Cingulate Cortex. Elsevier, Jan. 1, 2019, pp. 39–51. DOI: 10.1016/B978-0-444-64196-0.00003-0.

[64] Reynolds, S. M. and Zahm, D. S. "Specificity in the Projections of Prefrontal and Insular Cortex to Ventral Striatopallidum and the Extended Amygdala". In: *Journal of Neuroscience* 25.50 (Dec. 14, 2005), pp. 11757–11767. DOI: 10.1523/JNEUROSCI.3432-05.2005. pmid: 16354934.

[65] Nord, C. L., Lawson, R. P., and Dalgleish, T. "Disrupted Dorsal Mid-Insula Activation During Interoception Across Psychiatric Disorders". In: *American Journal of Psychiatry* 178.8 (Aug. 2021), pp. 761–770. DOI: 10.1176/appi.ajp.2020.20091340.

[66] Cloutman, L. L. et al. "The Variation of Function across the Human Insula Mirrors Its Patterns of Structural Connectivity: Evidence from in Vivo Probabilistic Tractography". In: *NeuroImage* 59.4 (Feb. 15, 2012), pp. 3514–3521. DOI: 10.1016/j.neuroimage.2011.11.016.

[67]   Stephan, E. et al. "Functional Neuroimaging of Gastric Distention". In: *Journal of Gastrointestinal Surgery* 7.6 (Sept. 1, 2003), pp. 740–749. DOI: 10.1016/S1091-255X(03)00071-4.

[68]   Medford, N. and Critchley, H. D. "Conjoint Activity of Anterior Insular and Anterior Cingulate Cortex: Awareness and Response". In: *Brain Structure and Function* 214.5 (June 1, 2010), pp. 535–549. DOI: 10.1007/s00429-010-0265-x.

[69]   Deen, B., Pitskel, N. B., and Pelphrey, K. A. "Three Systems of Insular Functional Connectivity Identified with Cluster Analysis". In: *Cerebral Cortex* 21.7 (July 1, 2011), pp. 1498–1506. DOI: 10.1093/cercor/bhq186.

[70]   Klein, T., Ullsperger, M., and Danielmeier, C. "Error Awareness and the Insula: Links to Neurological and Psychiatric Diseases". In: *Frontiers in Human Neuroscience* 7 (2013).

[71]   Vogt, B. A., Derbyshire, S., and Jones, A. K. P. "Pain Processing in Four Regions of Human Cingulate Cortex Localized with Co-registered PET and MR Imaging". In: *European Journal of Neuroscience* 8.7 (1996), pp. 1461–1473. DOI: 10.1111/j.1460-9568.1996.tb01608.x.

[72]   Vogt, B. A., Berger, G. R., and Derbyshire, S. W. G. "Structural and Functional Dichotomy of Human Mid-cingulate Cortex". In: *European Journal of Neuroscience* 18.11 (2003), pp. 3134–3144. DOI: 10.1111/j.1460-9568.2003.03034.x.

[73]   Büchel, C. et al. "Brain Systems Mediating Aversive Conditioning: An Event-Related fMRI Study". In: *Neuron* 20.5 (May 1, 1998), pp. 947–957. DOI: 10.1016/S0896-6273(00)80476-6. pmid: 9620699.

[74]   Büchel, C. et al. "Amygdala–Hippocampal Involvement in Human Aversive Trace Conditioning Revealed through Event-Related Functional Magnetic Resonance Imaging". In: *Journal of Neuroscience* 19.24 (Dec. 15, 1999), pp. 10869–10876. DOI: 10.1523/JNEUROSCI.19-24-10869.1999. pmid: 10594068.

[75]   Hester, R., Simões-Franklin, C., and Garavan, H. "Post-Error Behavior in Active Cocaine Users: Poor Awareness of Errors in the Presence of Intact Performance Adjustments". In: *Neuropsychopharmacology* 32.9 (9 Sept. 2007), pp. 1974–1984. DOI: 10.1038/sj.npp.1301326.

[76]   Seeley, W. W. et al. "Dissociable Intrinsic Connectivity Networks for Salience Processing and Executive Control". In: *Journal of Neuroscience* 27.9 (Feb. 28, 2007), pp. 2349–2356. DOI: 10.1523/JNEUROSCI.5587-06.2007. pmid: 17329432.

[77]   Liakakis, G., Nickel, J., and Seitz, R. J. "Diversity of the Inferior Frontal Gyrus—A Meta-Analysis of Neuroimaging Studies". In: *Behavioural Brain Research* 225.1 (Nov. 20, 2011), pp. 341–347. DOI: 10.1016/j.bbr.2011.06.022.

[78]   Swick, D. et al. "Left Inferior Frontal Gyrus Is Critical for Response Inhibition". In: *BMC Neuroscience* 9.1 (Oct. 21, 2008), p. 102. DOI: 10.1186/1471-2202-9-102.

[79]   Du, J. et al. "Functional Connectivity of the Orbitofrontal Cortex, Anterior Cingulate Cortex, and Inferior Frontal Gyrus in Humans". In: *Cortex* 123 (Feb. 1, 2020), pp. 185–199. DOI: 10.1016/j.cortex.2019.10.012.

[80] Bulbena, A. et al. "Psychiatric and Psychological Aspects in the Ehlers–Danlos Syndromes". In: *American Journal of Medical Genetics Part C: Seminars in Medical Genetics* 175.1 (2017), pp. 237–245. DOI: 10.1002/ajmg.c.31544.

[81] Casanova, E. L. et al. *Immune, Autonomic, and Endocrine Dysregulation in Autism and Ehlers-Danlos Syndrome/Hypermobility Spectrum Disorders Versus Unaffected Controls*. June 13, 2019. DOI: 10.1101/670661. URL: https://www.biorxiv.org/content/10.1101/670661v1 (visited on 05/19/2023). preprint.

[82] Eccles, J. A. et al. "Joint Hypermobility and Autonomic Hyperactivity: Relevance to Neurodevelopmental Disorders". In: *Journal of Neurology, Neurosurgery & Psychiatry* 85.8 (Aug. 1, 2014), e3–e3. DOI: 10.1136/jnnp-2014-308883.9.

[83] Savage, G. K. et al. "#3085 Relationship between Variant Connective Tissue (Hypermobility) and Autism Sensory Processing: Externally Oriented Thinking as a Mediator". In: *Journal of Neurology, Neurosurgery & Psychiatry* 92.8 (Aug. 1, 2021), A7–A8. DOI: 10.1136/jnnp-2021-BNPA.19.