1 **A randomized controlled safety and feasibility trial of floatation-REST in anxious**

2 **and depressed individuals**

McKenna M. Garland, M.A.[a,b], Raminta Wilson, M.D., M.P.H.[a], Wesley K. Thompson, Ph.D.[a],
Murray B. Stein, M.D., M.P.H.[c,d,e], Martin P. Paulus, M.D. [a f,], Justin S. Feinstein, Ph.D.[a,f,g],
Sahib S. Khalsa, M.D., Ph.D.[a,f]*

[a]Laureate Institute for Brain Research, University of Tulsa, Tulsa, Oklahoma
[b]Kendall College of Arts and Sciences, University of Tulsa, Tulsa, Oklahoma
[c]Department of Psychiatry, University of California San Diego, La Jolla, California
[d]Herbert Wertheim School of Public Health and Human Longevity Science, University of
California San Diego, La Jolla, California
[e]Psychiatry Service, Veterans Affairs San Diego Healthcare System, San Diego, California
[f]Oxley College of Health Sciences, University of Tulsa, Tulsa, Oklahoma
[g]Float Research Collective, Kihei, Hawaii

21
22
23
24

25 **Corresponding author:** Sahib S. Khalsa, M.D., Ph.D. skhalsa@laureateinstitute.org

Abstract

Background: Reduced Environmental Stimulation Therapy via floatation (floatation-REST) is a

behavioral intervention designed to attenuate exteroceptive sensory input to the nervous system.

Pilot studies in anxious and depressed individuals demonstrated that single sessions of floatation-

REST are safe, well-tolerated, and associated with acute anxiolysis. However, there is not

sufficient evidence of the feasibility of floatation-REST as a repeated intervention.

Methods: We randomized 75 individuals with anxiety and depression to six sessions of

floatation-REST in different formats (pool-REST or pool-REST preferred) or an active

comparator (chair-REST). Feasibility was assessed via adherence rate to the assigned

intervention, tolerability via duration of REST utilization and overall study dropout rate, and

safety via incidence of serious or non-serious adverse events.

Results: Six-session adherence was 85% for pool-REST, 89% for pool-REST preferred, and 74%

for chair-REST. Dropout rates did not differ significantly between the treatment conditions.

Mean session durations were consistently above 50 minutes, and when allowed to choose the

duration and frequency, participants opted to float for an average of 75 minutes. There were no

serious adverse events associated with any intervention. Positive experiences were endorsed

more commonly than negative ones and were also rated at higher levels of intensity.

Conclusions: Taken together, six sessions of floatation-REST appear feasible, well-tolerated, and

safe in anxious and depressed individuals. Floatation-REST induces positively-valenced

experiences with few negative effects. warrant further investigation of clinical efficacy. Larger

randomized controlled trials evaluating markers of clinical efficacy are warranted.

Clinical Trial Registration Identifier: NCT03899090

Introduction

Reduced Environmental Stimulation Therapy via floatation (i.e., floatation-REST) is a

behavioral intervention designed to systematically attenuate exteroceptive sensory input to the

nervous system. Dating back to the 1950's, REST was first introduced to assess how humans

would react to low-level or monotonous stimulation of the nervous system, with subsequent

investigations exploring REST as a therapeutic intervention for habit modification during

smoking cessation or stress reduction [1]. The earliest forms of REST involved strict

confinement to a bed in a dimly lit cubicle for days at a time (i.e., 'chamber-REST') [2] or full-

body submersion in a vertical water tank with the assistance of a breathing helmet [3, 4]. Since

then, several technological developments have considerably expanded the availability of

different REST formats. Currently, the most common form of floatation-REST involves floating

effortlessly in a shallow pool of water heated to skin temperature and saturated with Epsom salt

to increase buoyancy (consequently, it is sometimes also called 'pool-REST'). The environment

is specially engineered to be lightproof, soundproof, and humidity and temperature-controlled, so

that input from visual, auditory, olfactory, gustatory, thermal, tactile, vestibular, and

proprioceptive channels is minimized, as is movement and speech [5]. Chair-REST has also

emerged as an alternative form of REST [6, 7]. In this environment, individuals recline in an

ergonomically engineered chair designed to take pressure off the spinal cord. Typically, these

chairs are placed in dimly lit and quiet rooms, similar but not identical to what would be

experienced during pool-REST. Our previous work has highlighted that although the degree of

sensory attenuation of chair-REST is not as substantial as traditional pool-REST, single sessions

of chair-REST can produce effects on negative affect, anxiety, stress, relaxation, serenity, and

refreshment of a similar nature in clinically anxious individuals [7]. Additionally, chair-REST

induced a similar though weaker pattern of neural changes in resting state functional connectivity

as pool-REST [6]. Such "zero-gravity chairs" have garnered public attention and are now

available in many large department and furniture stores. The availability and utilization of pool-

REST has grown in recent years, with hundreds of recreational "float centers" opening across the

world during the past decade [8]. Despite the increase in public interest and consumption of

various forms of REST, there has been limited research investigating the feasibility of the

technique, especially in clinical populations.

Preliminary insights into the tolerability and safety of floatation-REST in anxious and

depressed individuals are restricted to a few studies. A wait-list control trial reported that

individuals scoring high on self-reported measures of generalized anxiety were largely adherent

to 12 sessions of floatation-REST [9]. Another study found that individuals with stress-related

pain and burnout depression were able to complete up to 33 sessions of floatation-REST [10].

However, neither of these studies assessed adverse events that could have arisen during the

repeated sessions of floatation-REST, nor did they formally evaluate intervention feasibility. In

prior studies, most anxious individuals who were assigned fixed durations of pool-REST or

chair-REST for 45 to 90 minutes in length tolerated this amount [5, 7, 9, 11], but the impact of

flexible assignment on preference has not been investigated. Finally, our initial safety studies

examining the effects of single sessions of pool-REST or chair-REST in clinically anxious

individuals revealed no serious adverse events, suggesting that the brief intervention exposure

was safe and well-tolerated [5, 7]. Regarding safety, it is important to note that there have been

several deaths reported with the use of floatation tanks at recreational float centers associated

with concurrent drug (i.e., ketamine) or alcohol use [12], suggesting the need for inclusion of

substance use screening. There have also been reports of auditory and visual hallucinations

during floatation-REST [13]. However, these are generally described in a positive light, and are

infrequent [1, 5]. Other negative events that have been reported include occasional skin itchiness

or dry mouth, as well as discomfort resulting from accidental introduction of saltwater to the

eyes or open cuts [5, 14]. Despite the common utilization of floatation-REST by the public, and

an emerging clinical trials literature, no studies have systematically evaluated the safety profile

of this intervention across multiple sessions.

This study examined the feasibility, tolerability, and safety of floatation-REST as a

repeatable intervention in clinically anxious and depressed individuals using a randomized

design with an active comparator (chair-REST). Our primary outcome was feasibility. We

hypothesized that feasibility would be reflected by 6-session adherence rates of 80% or above, a

level similar to standard behavioral interventions for anxiety and depression [15, 16]. As an

additional feasibility assessment, we evaluated for potential differences in credibility and

expectancy between the assigned REST interventions. We predicted that all forms of floatation-

REST would be well-tolerated, with low dropout rates (<20%) over the course of the

intervention and 6-month follow-up period. We also predicted that individuals would remain in

their assigned REST environment for the majority of the allotted duration of time. Based on the

prior literature, we predicted that the REST interventions would be safe and associated with

minimal adverse reactions. Finally, as an exploratory aim, we examined the impact of flexible

assignment on floatation-REST preference in terms of the chosen frequency and duration of float

sessions, in order to provide information to help optimize the design of future trials.

Methods

Participant Recruitment

75 treatment-seeking adults with anxiety and depression were recruited through the Laureate

Institute for Brain Research's participant databases ([17] see Supplement for sample size

determination). Inclusion criteria included high levels of current anxiety (as measured by an

OASIS score $\geq$ 6; [18]) and anxiety sensitivity (as measured by an ASI-3 score $\geq$ 24; [19]). Of

note, the initial eligibility criteria of OASIS score $\geq$ 8 and ASI-3 score $\geq$ 29 were lowered in

September 2020 to increase participant recruitment to meet the targeted rate specified in the

funded grant. Depression symptoms were measured via the Patient Health Questionnaire (PHQ-

9, [20]), but an inclusion cutoff score was not specified. Other inclusion criteria included: age

between 18 and 60 years, and either no prior experience of floatation-REST or a minimum of 1

year since the last float session. If taking psychiatric medications or receiving psychotherapy, the

treatment regimen was required to be stable, defined as having taken the medication or been in

therapy for 8 weeks or longer. Daily benzodiazepine use was exclusionary. Participants taking

benzodiazepines or opioids on an as-needed basis had to be willing to abstain from use within 24

hours of each float session. Other exclusion criteria included uncomfortability in water or

receiving blood draws, any skin conditions or open wounds that could cause pain when exposed

to saltwater, diagnosis of a neurological disorder (e.g., epilepsy), a comorbid diagnosis of bipolar

disorder, schizophrenia, or an eating disorder, active suicidality with plan or intent, receiving

current inpatient psychiatric treatment, or moderate to severe substance use disorder (as

determined by the MINI 7.0). Evidence of current drug use was also assessed prior to each

floatation-REST session via urine drug screen testing for amphetamines, methamphetamines,

cocaine, barbiturates, benzodiazepines, methylenedioxymethamphetamine (MDMA), methadone,

opiates, oxycodone, phencyclidine, propoxyphene, and ketamine, and breathalyzer testing for

alcohol; a positive readout on any of these tests was exclusionary. Pregnancy was exclusionary

based on a positive urine pregnancy test. All study procedures were approved by the local ethics

committee (Western Institutional Review Board Protocol #: 20150528) and were performed in

accordance with Declaration of Helsinki. All participants provided their written informed

consent prior to participation and were compensated for their study involvement. The study was

pre-registered on ClinicalTrials.gov (NCT03899090).

Experimental Design and Procedures

Using a randomized parallel study design, participants completed a total of 11 visits. The

study visit timeline is outlined in Figure 1. Participants initially completed a brief telephone

screen to collect demographic information and evaluate potential study eligibility. During visit 1,

official study eligibility was determined using the inclusion/exclusion criterion. After providing

informed consent, participants returned for visit 2, where baseline symptom ratings were

completed via clinician-reported and self-reported scales, and some behavioral testing of cardiac

interoception (results to be reported separately). Visit 2 concluded by unblinding and informing

the participant of their randomized (1:1:1) assignment to one of three intervention conditions:

pool-REST, pool-REST preferred, or chair-REST (i.e., the active comparator, described further

below). The randomization schedule was pre-determined by a statistical consultant uninvolved in

the study before participant recruitment and stored as a blinded electronic list. At the time of

assignment, the study coordinator would unblind and reveal the participant's allocation. At the

start of visit 3, all participants completed the Credibility/Expectancy Questionnaire (CEQ; [21]),

to assess their pre-intervention attitudes towards the assigned intervention arm. Participants then

completed six sessions (visits 3-8) according to their assigned condition. We assessed serious

and non-serious adverse events after each session via a detailed survey of positive and negatively

valenced experiences using a scale drawn from our prior floatation-REST study [5] and via

verbal self-report throughout the study period. Following the sixth and final session (visit 8),

participants returned to the lab for a post-intervention session (visit 9), where clinician-reported

and self-reported symptom ratings were once again collected and behavioral testing of cardiac

interoception was completed. The same clinician and self-reported symptom scales were

completed again at 6-week (visit 10) and 6-month (visit 11) follow-up visits. Follow-up surveys

were initially administered in the laboratory via an electronic tablet, but due to the Covid-19

pandemic these were switched to remote visits starting in May 2020. All surveys were obtained

electronically using REDCap (Research Electronic Data Capture; Version 10.6.19; [22, 23]).

Figure 1. *Study Visit Timeline.*

180
181

Experimental Conditions

Pool-REST. The pool-REST condition, which was delivered at the Laureate Institute for Brain

Research facility, involved prescribed (i.e., fixed) session durations and inter-session intervals

within an open or enclosed circular float pool (Floataway Inc., Norfolk, United Kingdom).

Participants were instructed to complete 1-hour float sessions once per week for six weeks. Each

pool measured 2.44 meters in diameter and 0.28 meters in depth. The water in each pool was

saturated with over 800 kilograms of Epsom salt (magnesium sulphate), creating a specific

gravity of approximately 1.26. This dense salt-water solution allowed for participants to float

effortlessly in a supine position. Both pools were constructed in rooms designed to be lightproof

and soundproof. The air and water temperature of each room was also calibrated to match the

temperature of the skin surface (~35.0 degrees Celsius). More details about how the pool-REST

condition was specially engineered to minimize all manner of external sensory stimulation can

be found in Feinstein et al., 2018a. We allowed participants to choose between the open and

enclosed pool to provide some flexibility for preference and to facilitate the optimal flow of

participants through the study.

Pool-REST Preferred. The pool-REST preferred condition involved increased flexibility of both

the session duration and the frequency. Participants were instructed that they were to complete

six sessions in the open or closed pool, but they were allowed to freely set their float duration for

up to two hours per session. In addition, they were permitted to choose the frequency of their

floating schedule, such that they could float as often as they preferred within a 12-week period,

with the only requirement being that there needed to be a minimum of 24 hours between

sessions.

Chair-REST. During chair-REST, which was delivered at the Laureate Institute for Brain

Research facility, participants reclined in a Zero Gravity Chair (PC510, Classic Power, Series 2,

Human Touch Inc., Long Beach, CA). This active comparator condition was tailored to closely

match the pool-REST condition on many parameters including a supine body position, the same

prescribed session duration and inter-session interval (i.e., 1-hour sessions completed once per

week for six weeks) in a room with reduced light and sound, and a similar instruction set

emphasizing the importance of stillness and wakefulness throughout each session (see

Supplemental Materials for full instruction set). Participants were informed that the chair was

ergonomically designed to take pressure off the spinal cord and used memory foam backing to

help the chair conform to each participant's body shape. To match the instruction set across

REST conditions, the Zero Gravity chair was explicitly referred to as the "float chair" and the act

of lying in the chair was referred to as "floating." The chair was placed in a dark and quiet room,

although the degree of light and sound attenuation was not as strong as that delivered in the pool

conditions (i.e., participants could perceive a low-level of external sound and light). Unlike the

pool conditions, the ambient temperature was maintained at a normal room temperature of ~23.3

degrees Celsius, so participants randomized to the chair-REST condition remained fully clothed

during each session.

Measures

Events Assessment — A 43-item self-report measure ([5]; see Supplemental Material) was used

to assess safety and subjective experiences occurring in each float session. This checklist was

adapted from our previous pilot studies and was designed to assess potential adverse experiences

associated with floatation-REST. 29 of the 43 items assessed the presence of negative physical or

psychological events including itchiness, dry mouth, pain, anxiety/panic, flashbacks, suicidality,

and hallucinations, as well as the presence of positive experiences (e.g., heightened feelings of

joy, refreshment, serenity); these 14 items were included to reduce response biases, as well as to

prevent a sole focus on assessing negative experiences associated with REST. Each item was

rated on four-point scale ("none," "mild," "moderate," or "extreme") using the following

instructions: "*Did you notice or experience an increase in any of these items during or shortly*

after your float today? Please only mark items that showed an increase from your typical day-to-

day experience." For any item that had an endorsement other than "None," participants were

provided a free-response box to describe their experience in more detail. For certain negative

symptoms (strong emotional memories, flashbacks, heightened thoughts related to death, visual

or auditory hallucinations, out-of-body experiences, feelings of detachment, loss of control over

behavior, and flight of ideas/racing thoughts) receiving any endorsement other than "None"

prompted participants with the following question: *"Overall, was this a positive or negative*

experience?" [5]. Items rated as "none" were coded as 0, negatively experienced items ranged

from -3 ("extremely negative") to -1 ("mildly negative"), and positively experienced items

ranged from 1 ("mildly positive) to 3 ("extremely positive").

Credibility/Expectancy Assessment — The Credibility/Expectancy Questionnaire (CEQ; [21])

was used to measure attitudes about treatment feasibility. The CEQ is a 6-item self-report

measure with individual items rated on a 1-9 or a 0%-100% scale. Items assess attitudes about

"how believable, convincing, and logical a treatment is" (i.e., credibility; [24]), and expectations

one holds about how the treatment might influence symptoms (i.e., expectancy).

There were no changes to the primary outcome or the main secondary outcomes reported in the

current study. A number of additional secondary outcomes were specified in May, 2022, prior to

analysis of the data, and will be included in a separate report.

255   **Statistical Analysis**

Feasibility

The primary outcome of feasibility was assessed by calculating adherence to the assigned

sessions via the formula $\hat{p} = \sum_{i=1}^{n} p_i / n$, where p_i was the proportion of completed sessions (out

of 6) for subject i and n was the number of subjects per condition (i.e., 25). To measure

attitudinal aspects of feasibility, pre-treatment credibility beliefs were calculated by taking the

mean of the first three items of the CEQ [25], while expectancy beliefs were captured via a

single item rated on a 0 to 100% scale assessing how much improvement in their symptoms the

participant thought would occur after completing the assigned intervention. Two separate one-

way ANOVAs were performed to assess for between-group differences in pre-intervention

credibility and expectancies, respectively.

Tolerability

Tolerability was assessed in three ways; first, by calculating the overall dropout rate for each

condition (i.e., (number of subjects who withdrew from the study or who were lost to follow-

up)/25). Secondly, a Kaplan-Meier analysis with post-hoc log-rank tests was used to test for

differential dropout rates over time for the three assigned conditions [26]. Third, within-session

tolerability was assessed by fitting a linear mixed models (estimated using REML and the

nloptwrap optimizer) to predict the duration of each session with visit and condition as fixed

effects (formulas: Session Duration (in minutes) ~ Visit * Condition). The model included

subject ID as a random effect (formula: ~1 | Subject ID) and utilized an AR1 covariance

structure. Post-hoc two-sided t-tests with Holm corrections were used to interpret significant

main effects, as well as simple effects for all significant interactions.

Safety

The number of adverse and serious adverse events and their relationship to the intervention were

recorded over the course of the entire study, including during the follow-up period. We then fit a

linear mixed model (estimated using REML and the nloptwrap optimizer) to predict event

magnitude ratings with event, visit, and condition parameters (formulas: Event Intensity Rating ~

Event * Visit * Condition). The models included subject ID and visit as random effects (formula:

~1 | Subject ID/Visit) and utilized an AR1 covariance structure. Post-hoc two-sided t-tests with

Holm corrections (Holm, 1979) were used to interpret significant main effects, as well as simple

effects for all significant interactions. All analyses were performed in RStudio (Version 3.6.0;

[27]).

Pool-REST Preferred

The impact of flexible assignment on preferred frequency of floating for the pool was

measured by calculating the average number of days between float sessions and the

corresponding standard deviation. To assess whether session frequency in this condition was

related to baseline severity of anxiety, depression, and anxiety sensitivity symptoms, we

examined bivariate correlations between float frequency and baseline OASIS, PHQ-9, and ASI-3

scores. Additionally, we examined bivariate correlations between the average session duration

(in minutes) for each intervention and baseline measures of symptom severity (i.e., OASIS,

PHQ-9, and ASI-3).

299   **Results**

Participant Characteristics

Recruitment occurred between February 2019 and October 2021, with a four month

pause starting March 2020 during the Covid-19 pandemic and was terminated upon achievement

of the recruitment target. Sociodemographic and clinical characteristics for the 75 individuals

who were randomized to each intervention arm are summarized in Table 1. A CONSORT

diagram reflecting the flow of participants in each arm is shown in Figure 2. One-way ANOVAs

revealed that the groups did not differ significantly on age, education, medication status (i.e.,

number of psychotropic medications), BMI, or baseline OASIS, ASI-3, or PHQ-9 scores. Chi-

squared tests indicated that there were no significant differences between the conditions

regarding sex, race/ethnicity, or the frequency of clinical diagnosis or rate of psychotherapy

utilization. Most of the sample identified as non-Hispanic White (81.3%). The most common

psychiatric diagnosis in the sample was major depressive disorder (97.3%) followed by

generalized anxiety disorder (50.7%).

Table 1. *Participant Demographics, DSM-5 Diagnostic Comorbidities, and Screening Scores at*

Study Entry

	Chair-REST (n=25)	Pool-REST (n=25)	Pool-REST preferred (n=25)	F	df	p
Demographics						
Sex				$\chi^2 = 0.15$	2	0.93
Male, N (%)	6 (24%)	6 (24%)	5 (20%)			
Female, N (%)	19 (76%)	19 (76%)	20 (80%)			
Age	37.8 (13.4)	33.4 (11.7)	34.3 (9.0)	1.01	72	0.37
Years of Education	15.2 (1.3)	15.1 (1.8)	15.8 (1.7)	1.15	71	0.32

BMI	29.3 (7.3)	28.8 (5.1)	27.9 (6.9)	0.30	70	0.74
Psychiatric Medications[a]	0.7 (1.1)	0.3 (0.6)	1.0 (1.3)	3.00	72	0.06
Receiving Psychotherapy, N(%)	11 (44)	11 (44)	16 (64)	$\chi^2 = 2.67$	2	0.26
Race/Ethnicity, N(%)				$\chi^2 = 8.75$	10	0.56
White/Caucasian	22 (88)	20 (80)	19 (76)			
Black/African American	2 (1)	6 (24)	4 (16)			
Am. Indian/Alaskan Native	5 (20)	5 (20)	4 (16)			
Asian/Pacific Islander	1 (4)	0 (0)	0 (0)			
Hispanic	1 (4)	2 (8)	2 (8)			
Unspecified	0 (0)	0 (0)	2 (8)			
Self-Report Measures						
OASIS Anxiety	10.1 (2.5)	9.4 (2.8)	9.9 (2.5)	0.53	69	0.59
ASI-3 Anxiety Sensitivity	35.6 (13.0)	39.9 (9.4)	40.5 (11.8)	0.04	69	0.97
PHQ-9 Depression	12.5 (4.9)	11.5 (5.5)	12.1 (4.7)	0.28	69	0.76
Psychiatric Diagnosis, N(%)[b]				$\chi^2 = 3.78$	8	0.88
Major depressive disorder	24 (96)	24 (96)	25 (100)			
Generalized anxiety disorder	10 (40)	17 (68)	11 (44)			
Social anxiety disorder	9 (36)	12 (48)	8 (32)			
Posttraumatic stress disorder	9 (36)	8 (32)	7 (28)			
Panic disorder	3 (12)	6 (24)	7 (28)			

Note. Numbers reflect means and standard deviation unless otherwise indicated. BMI = Body

Mass Index; OASIS = Overall Anxiety Severity and Impairment Scale [18] measured at pre-

intervention visit; ASI-3 = Anxiety Sensitivity Index-3 [19], measured at pre-intervention visit;

PHQ-9 = Patient Health Questionairre-9 [20] measured at pre-intervention visit. [a]Numbers

reflect the mean number of prescribed psychiatric medications taken by the participant, measured

at the initial screening. [b]Diagnosis was determined by clinician interview using the MINI version

7.0 [28]. For brevity, only comorbid diagnoses with frequency > 10% across the entire sample (n

= 75) are listed. As this study allowed for comorbid diagnoses, percentage totals will be > 100%.

Figure 2. CONSORT Diagram. Note. C = Completed. W = Withdrew.

Feasibility

Average treatment adherence was 89% (5.3   1.6 sessions) for the pool-REST preferred

condition, 85% (5.1   1.8 sessions) for the pool-REST condition, and 74% (4.4   2.5 sessions)

for the chair-REST condition. One-way ANOVAs revealed that the REST conditions did not

differ significantly on pre-intervention CEQ credibility ($F(2, 67) = 2.42, p = 0.097$) or

expectancy ($F(2, 67) = 0.98, p = 0.379$; see Figures 3a-3b). The average credibility score for the

three groups was 6.67 ± 1.50. On average, the three groups expected the intervention to halve

their symptoms ($50.3 \pm 21.6\%$).

Figure 3. Pre-Treatment Credibility Beliefs and Expectations for Symptom Improvement. *Note.*

Figures reflect pre-intervention ratings for credibility of beliefs (top) and expectations for

symptom improvement (bottom) on the Credibility and Expectancy Questionnaire, according to

the assigned intervention. The black dots and error bars represent the mean and standard error of

the mean, respectively. Grey dashed lines reflect an established average for credibility and

expectancy beliefs of other non-pharmacological treatments (e.g., psychotherapy, see

discussion). There were no significant group differences in levels of pre-treatment credibility or

expectancy beliefs (*ps* > .05).

Tolerability

The Kaplan-Meier survival analyses did not indicate a significant difference in treatment

dropout among the three groups ($p = .40$, Figure 4). Following the randomization at the baseline

visit, the chair-REST condition had the highest dropout rate at 16% (n = 4 individuals), versus

4% (n = 1 individual) for the pool-REST condition and 0% for the pool-REST preferred

condition. Across the six-session intervention, the chair-REST condition demonstrated an overall

dropout rate of 32%, while the pool-REST condition dropout rate was 24%, and the pool-REST

preferred dropout rate was 16%. The dropout rate at the 6-month follow up was 36% for the

chair-REST condition, 28% for the pool-REST condition, and 20% for the pool-REST preferred

condition (1 individual dropped out from each of the groups).

Figure 4. Kaplan-Meier Survival Curve illustrating dropout over time for each REST Condition.

361
362

363    Of note, due to the onset of the Covid-19 pandemic and mandatory shutdowns, two

364 participants were withdrawn from the study by the study investigators. One participant in the

365 Chair-REST condition was withdrawn from the study after session 2 (at day 27), and one

366 participant from the Pool-REST condition was withdrawn following their fifth session (at day

367 73). This dropout information was entered into the analysis, but due to the small amount of

368 dropout and no discernable group pattern, no further corrective action was taken during the

369 analysis.

370    The linear mixed effects analysis of within-session tolerability (i.e., session duration;

371 Figure 5), revealed a main effect of condition ($F(2) = 10.33$, $p < .001$, $\eta_p^2 = .24$), such that

372 session durations were significantly longer for the pool-REST preferred condition ($M = 75.4 \pm$

373 29.4 minutes) than either the chair-REST ($M = 58.4 \pm 4.3$ minutes; $p = .007$, $d = .810$) or pool-

374 REST condition ($M = 53.0 \pm 12.3$ minutes; $p = .0002$, $d = .996$). There were no statistically

375 significant differences in session duration between the chair-REST and pool-REST conditions (p

376 $= .292$). Across the 373 REST sessions administered across the entire study, 317 sessions (85%

377 of total) were 50 minutes or longer in duration.

Figure 5. Session durations. Note. Session durations were significantly longer for the pool-

REST preferred condition than either the chair-REST (p = .007) or pool-REST conditions (p =

.0002). There were no statistically significant differences in session duration between the chair-

REST and pool-REST conditions (p = .292). Duration did not vary as a function of session.

Safety

Four adverse events were recorded during the study. All were determined to be unrelated

to the study intervention. One participant in the chair-REST condition reported two separate

instances of migraines with nausea on the days of their first and second REST session. One

participant in the pool-REST condition who suffered from seasonal allergies reported an ear

infection four days after their third REST session. Finally, one participant in the pool-REST

condition reported a suicide attempt the night before their 6-week follow-up visit. The participant

was subsequently evaluated at a local psychiatric emergency department, where it was

determined that hospital admission was unwarranted.

Regarding the acutely experienced event profile, Figure 6 shows the average magnitude

rating for each event across the three assigned REST conditions. A qualitative inspection of the

data revealed that positive experiences were endorsed more commonly than negative ones and

were also rated at higher levels of intensity, with average intensity ratings in the mild-to-

moderate range for positive experiences and well-below mild for negative experiences. In the

linear mixed effects (LME) quantitative analysis of intensity ratings, we observed a significant

main effect of event ($F(43) = 153.61$, $p < .001$, $\eta_p^2 = .30$) and treatment condition ($F(2) = 3.51$, p

$= .04$, $\eta_p^2 = .09$), which was accounted for by the significant interaction between these variables

($F(86) = 6.43$, $p < .0001$, $\eta_p^2 = .04$). Post-hoc comparisons suggested that feelings of intense

joy/happiness, increased energy, increased ability to focus/concentrate, serenity and

peacefulness, appreciation for life, refreshment, relaxation, silent mind, pain-free existence, and

feelings of flow were experienced more positively in the pool-REST and pool-REST preferred

conditions when compared to the chair-REST condition ($ps < .05$). Stronger positive ratings for

the pool-REST preferred condition were also observed for relaxation and pain-free existence

when compared to the pool-REST condition ($ps < .05$). The opposite trend was found for

feelings of flow, increased empathy and compassion for others, and joy/happiness, such that

participants in the pool-REST condition tended to rate these items more positively than their

pool-REST preferred counterparts ($ps < .05$). The pool-REST group also rated feelings of

empathy and compassion for others more positively than their chair-REST counterparts ($p < .01$).

Figure 6. *Events Checklist. Note.* Event anchors ranged from "Extremely Negative" to

"Extremely Positive" (anchors not shown). Asterisk (*) indicates significant group difference in

rating magnitude.

Regarding events that typically carry negative connotations, post-hoc comparisons

suggested that itchiness was experienced more negatively in the pool-REST and pool-REST

preferred conditions than the chair-REST condition ($ps < .05$). The pool-REST preferred

condition also rated feelings of intense fear and panic more negatively than the chair-REST

condition ($p = .045$). Interestingly, the pool-REST and pool-REST preferred conditions endorsed

experiencing feelings of detachment more positively than the chair-REST condition ($ps < .001$).

Stronger positive ratings were also observed in the pool-REST preferred condition than the

chair-REST condition for hallucinations ($p = .03$). Of note, the reports of hallucinations and

feelings of detachment occurred infrequently and tended to be described in neutral or positive

terms. For example, descriptions of hallucinations commonly mentioned visually experienced

colors and shapes, such as "*Colorful visions; shapes, people, and characters,*" "*Saw green,*

crystal-like geometric shapes off and on (eyes were closed)," and "*I experienced those colorful*

flashes and dots similar to when you close your eyes and press lightly on your eyes." Feelings of

detachment were typically described using feelings of relaxation, weightlessness, and

disconnection from space and time, including: "*I felt weightless. I felt like nothing else mattered.*

It was just me," "*Felt very relaxed like everything was far away,*" "*A similar dozing detachment*

as when you're half asleep for a really nice nap," and "*I felt like I was in my own world and the*

real world was frozen in time."

436
437   The linear mixed model analysis also revealed a significant interaction between event and

438   visit ($F(214) = 1.40$, $p = .0001$, $\eta_p^2 = .02$). Specifically, post-hoc analyses showed a weakening

439   in the magnitude of positive events for all three interventions over the course of the study for the

440   following events: serenity and peacefulness, appreciation for life, refreshment, relaxation, silent

441   mind, pain-free existence, and feelings of flow ($ps < .05$).

Pool-REST Preferred

444   When allowed to select the frequency of sessions, the pool-REST preferred participants

445   chose to float at a frequency of once every 12 days, on average ($SD = 9.33$), and for an average

446   float duration of 75 minutes ($M = 75.4 \pm 29.4$ minutes). Exploratory analyses indicated that

447   session frequency and session duration were not significantly associated with baseline measures

448   of anxiety, depression, or anxiety sensitivity (ps > .05).

450   **Discussion**

This study represents the first randomized controlled trial examining the feasibility,

tolerability, and safety of repeated sessions of floatation-REST in individuals with anxiety and

depression. We observed evidence of a favorable feasibility, tolerability, and safety profile for

the application of floatation-REST and chair-REST in this outpatient sample. Positive

experiences were endorsed more commonly than negative ones and were also rated at higher

levels of intensity, with no serious adverse events associated with any intervention. Taken

together, six sessions of floatation-REST appear feasible, well-tolerated, and safe in anxious and

depressed outpatient individuals.

Our primary outcome of adherence suggested that the feasibility of each form of REST,

indexed via adherence to the allocated six-sessions, was adequate, ranging from 85-89% for the

pool-REST conditions to 74% for the chair-REST condition. These levels of adherence are

comparable to traditional behavioral interventions for anxiety and depression, whose treatment

completion rates hover around 80% [16]. Each REST intervention was perceived to be

moderately credible, with participants having positive expectations for symptom improvement.

The observed levels of credibility and expectancy beliefs are also consistent with those typically

seen for other behavioral interventions such as exposure therapy (range of credibility beliefs =

5.95-7.47, range of expectancy beliefs = 50.50-67.90; [21, 25, 29]).

The low dropout rates in this behavioral intervention study provides additional evidence

that repeated sessions of REST are well-tolerated. Dropout rates in the post randomization

setting prior to intervention onset were lowest for the pool interventions (4% dropout for pool-

REST, and 0% dropout for pool-REST preferred), while the chair-REST pretreatment dropout

rate was the highest (16%) and similar to those observed to cognitive behavioral therapy

(dropout rates ranging from 11-21%, [15]). Dropout rates during the administration of the REST

intervention (32% for chair-REST, 24% for pool-REST, and 16% for pool-REST preferred) were

also comparable to rates seen in non-pharmacological and pharmacological interventions such as

cognitive behavioral therapy (19-36%, [15]), meditation (33-44%, [30]), acceptance and

commitment therapy (15%, [31]), and antidepressant medications (37-49%, [32]).

Using session duration as an additional proxy of intervention tolerability, we saw very

few individuals in the pool-REST and chair-REST conditions who had a prescribed session

duration (i.e., 60 minutes) terminate early, with the majority of individuals utilizing their full

time allotment. For the pool-REST preferred group, who was given greater flexibility in session

duration (i.e., allowed to float for up to two hours), we observed longer average session

durations. Although the preferred condition was given an additional 60 minutes, they tended to

float about 20 minutes longer than the prescribed session conditions on average. Moreover,

session duration was unrelated to symptom severity at baseline, indicating that individuals with

higher levels of anxiety, depression, or anxiety sensitivity were no more or less likely to utilize

longer durations (i.e., in a self-medicating or symptom-dependent fashion). Taken together, these

findings may suggest that a 60-minute float is a suitable duration for future clinical trials in a

clinically anxious and depressed sample. However, it remains to be determined whether there is

additional utility of more time (e.g., 75 minutes), or alternatively, whether similar effects could

be achieved with a shorter durations (e.g., 30 or 45 minutes).

The safety findings from this trial are consistent with previous single-session studies of

pool-REST and chair-REST which found few negative events and no serious adverse events [5,

7, 14]. Our study expands upon this work by showing a favorable safety profile across repeated

sessions of floatation-REST. We also demonstrated that positive experiences were endorsed

more commonly than negative ones, and at higher levels of intensity, in the mild-to-moderate

range. Importantly, while all three intervention groups consistently rated positive effects of the

intervention, the pool-REST groups tended to rate positive events more strongly, especially

regarding feelings of peace and tranquility. These findings provide support for the overall safety

of floatation-REST delivered in a pool or in a Zero Gravity chair setting. Moreover, the

comparable intensity ratings of many negative events but weaker ratings of positive events

within the chair-REST condition suggests that the chair is a suitable active comparator.

There were differential event profiles between the pool-REST conditions and the chair-

REST condition, such that participants in the pool conditions reported stronger feelings of

tranquility (e.g., refreshment, peacefulness, serenity) than those in the chair-REST condition and

tended to rate itchiness and fear/panic more negatively than their chair-REST counterparts. The

most common negative events associated with pool-REST were itchiness and dry mouth, which

is consistent with our previous pilot study [5], and may be related to the high salt concentration

in the pools. In addition, several experiences that traditionally carry negative connotations,

including audiovisual hallucinations and feelings of detachment, tended to be described in a

positive light in the pool conditions. We do not believe the hallucinations reported by

participants in this study reflect hallucinations in the true sense of the term and do not consider

them indicative of psychosis-spectrum symptoms. Rather, these visual and auditory sensations

may be the result of the brain 'filling in the gaps,' i.e., generating sensory representations in the

face of absent exteroceptive input to the nervous system. Finally, the repeated nature of this

intervention allowed for a thorough examination of event profiles over time. An attenuation in

the magnitude of several positive events was observed at later sessions for all three interventions,

raising the possibility that the increased familiarity may reduce the novelty of the intervention

and the magnitude of positive events.

This investigation had several limitations that must be acknowledged. First, four

individuals in the chair-REST condition and one participant in the pool-REST condition

withdrew from the study after randomized allocation to their respective intervention and did not

complete the CEQ. These dropouts may suggest the presence of unmet expectations, particularly

for the individuals allocated to the chair-REST condition given that the largest dropout occurred

in this group. Alternatively, pre-intervention anticipatory anxiety could have played a role in

withdrawal, although we did not receive such reports from participants. To avoid the limitation

of missing data in the future, subsequent studies should aim to collect the CEQ measure

immediately following randomization. Another limitation of the current study is the modest

sample size. While our sample size was consistent with those used in previous floatation-REST

trials, larger groups may be needed for an adequately powered survival analysis for the purposes

of tolerability assessment. As such, the current study may be underpowered to detect meaningful

group differences in dropout rate. We examined six-session feasibility of REST whereas prior

clinical trials have typically used more sessions (e.g., 12). Thus, the measurement of feasibility

and tolerability during the employment of a larger number of sessions may be advisable in future

trials. Additionally, most of our sample was comprised of Caucasian, highly educated females,

thereby limiting generalizability claims to the broader population. Finally, due to the fact that

participants were financially compensated for their participation, we did not examine structural

aspects of feasibility relevant to the real-world delivery of behavioral therapies, including

financial and time commitments that might otherwise be important for treatment engagement.

540   **Conclusion**

541   The findings from this clinical trial suggests that six sessions of REST in a chair or pool

542   environment are feasible, well-tolerated, and safe in clinically anxious and depressed outpatient

543   individuals. Larger randomized controlled trials evaluating markers of clinical efficacy are

544   warranted.

Funding

This research was supported by an award from the National Center for Complimentary and Integrative Health (R34AT009889). The authors were also supported by the National Institute of Mental Health (R01MH127225, K23MH112949 to SSK), National Institute of General Medical Sciences Center Grant Award Number (1P20GM121312 to MPP, JSF, SSK), and The William K. Warren Foundation. The funding sources had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; or decision to submit the manuscript for publication.

Author Contributions

McKenna Garland: Writing—Original draft preparation, investigation, formal analysis, visualization. Raminta Wilson: Investigation. Wesley Thompson: Writing—Review and editing. Murray Stein: Writing—Review and editing. Martin Paulus: Writing—Review and editing, supervision. Justin Feinstein: Conceptualization, methodology, resources, writing—review and editing, supervision, project administration, funding acquisition. Sahib Khalsa: Conceptualization, methodology, resources, writing—review and editing, supervision, project administration, funding acquisition.

Declaration of Conflicting Interests

The authors declare that there are no conflicts of interests.

Acknowledgments

We would like to thank Henry Yeh, Jamie Rhudy, Emily Adamic, and Joel Greenshields for consultation on coding and statistical matters. We would also like to acknowledge Laura Garrison for her contributions to data collection. Finally, we would like to thank the custodial

References

1.   Suedfeld P, Coren S. Perceptual isolation, sensory deprivation, and rest: Moving

introductory psychology texts out of the 1950s. Canadian Psychology/Psychologie canadienne.

1989;30(1):17.

2.   Rossi AM. General methodological considerations. Sensory deprivation: Fifteen years of

research. 1969:16-46.

3.   Lilly JC, Shurley JT. Experiments in solitude, in maximum achievable physical isolation

with water suspension, of intact healthy persons. Psychophysiological aspects of space flight.

1961:238-47.

4.   Shurley JT. Profound experimental sensory isolation. American Journal of Psychiatry.

1960;117(6):539-45.

5.   Feinstein JS, Khalsa SS, Yeh HW, Wohlrab C, Simmons WK, Stein MB, et al.

Examining the short-term anxiolytic and antidepressant effect of Floatation-REST. PLoS One.

2018;13(2):e0190292. Epub 20180202. doi: 10.1371/journal.pone.0190292. PubMed PMID:

29394251; PubMed Central PMCID: PMCPMC5796691.

6.   Al Zoubi O, Misaki M, Bodurka J, Kuplicki R, Wohlrab C, Schoenhals WA, et al. Taking

the body off the mind: Decreased functional connectivity between somatomotor and default-

mode networks following Floatation-REST. Hum Brain Mapp. 2021;42(10):3216-27. Epub

20210409. doi: 10.1002/hbm.25429. PubMed PMID: 33835628; PubMed Central PMCID:

PMCPMC8193533.

7.   Khalsa SS, Moseman SE, Yeh H-W, Upshaw V, Persac B, Breese E, et al. Reduced

environmental stimulation in anorexia nervosa: an early-phase clinical trial. Frontiers in

psychology. 2020:2534.

8.   Solutions FT, LLC FO. State of the Float Industry Report. Float Tank Solutions; 2020.

9.   Jonsson K, Kjellgren A. Promising effects of treatment with flotation-REST (restricted

environmental stimulation technique) as an intervention for generalized anxiety disorder (GAD):

a randomized controlled pilot trial. BMC complementary and alternative medicine. 2016;16(1):1-

12.

10.   Bood SÅ, Sundequist U, Kjellgren A, Nordström G, Norlander T. Effects of flotation rest

(restricted environmental stimulation technique) on stress related muscle pain: are 33 flotation

sessions more effective than 12 sessions? Social Behavior and Personality: an international

journal. 2007;35(2):143-56.

11.   Feinstein JS, Khalsa SS, Yeh H, Al Zoubi O, Arevian AC, Wohlrab C, et al. The

elicitation of relaxation and interoceptive awareness using floatation therapy in individuals with

high anxiety sensitivity. Biological psychiatry: cognitive neuroscience and neuroimaging.

2018;3(6):555-62.

12.   Lann MA, Martin A. An unusual death involving a sensory deprivation tank. Journal of

forensic sciences. 2010;55(6):1638-40.

13.   Norlander T, Kjellgren A, Archer T. The experience of flotation-REST as a function of

setting and previous experience of altered state of consciousness. Imagination, Cognition and

Personality. 2000;20(2):161-78.

14.   Forgays DG, Belinson MJ. Is flotation isolation a relaxing environment? Journal of

Environmental Psychology. 1986;6(1):19-34.

15.   Fernandez E, Salem D, Swift JK, Ramtahal N. Meta-analysis of dropout from cognitive

behavioral therapy: Magnitude, timing, and moderators. Journal of Consulting and Clinical

Psychology. 2015;83(6):1108.

16.   Swift JK, Greenberg RP, Tompkins KA, Parkin SR. Treatment refusal and premature

termination in psychotherapy, pharmacotherapy, and their combination: A meta-analysis of head-

to-head comparisons. Psychotherapy. 2017;54(1):47.

17.   Victor TA, Khalsa SS, Simmons WK, Feinstein JS, Savitz J, Aupperle RL, et al. Tulsa

1000: a naturalistic study protocol for multilevel assessment and outcome prediction in a large

psychiatric sample. BMJ open. 2018;8(1):e016620.

18.   Campbell-Sills L, Norman SB, Craske MG, Sullivan G, Lang AJ, Chavira DA, et al.

Validation of a brief measure of anxiety-related severity and impairment: the Overall Anxiety

Severity and Impairment Scale (OASIS). Journal of affective disorders. 2009;112(1-3):92-101.

19.   Taylor S, Zvolensky MJ, Cox BJ, Deacon B, Heimberg RG, Ledley DR, et al. Robust

dimensions of anxiety sensitivity: development and initial validation of the Anxiety Sensitivity

Index-3. Psychological assessment. 2007;19(2):176.

20.   Kroenke K, Spitzer RL, Williams JB. The PHQ 9: validity of a brief depression severity

measure. Journal of general internal medicine. 2001;16(9):606-13.

21.   Devilly GJ, Borkovec TD. Psychometric properties of the credibility/expectancy

questionnaire. Journal of behavior therapy and experimental psychiatry. 2000;31(2):73-86.

22.   Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic

data capture (REDCap)—a metadata-driven methodology and workflow process for providing

translational research informatics support. Journal of biomedical informatics. 2009;42(2):377-81.

23.   Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, O'Neal L, et al. The REDCap

consortium: Building an international community of software platform partners. Journal of

biomedical informatics. 2019;95:103208.

24.   Kazdin AE. Therapy outcome questions requiring control of credibility and treatment-

generated expectancies. Behavior Therapy. 1979;10(1):81-93.

25.   Thompson-Hollands J, Bentley KH, Gallagher MW, Boswell JF, Barlow DH. Credibility

and outcome expectancy in the unified protocol: Relationship to outcomes. Journal of

Experimental Psychopathology. 2014;5(1):72-82.

26.   Kaplan EL, Meier P. Nonparametric estimation from incomplete observations. Journal of

the American statistical association. 1958;53(282):457-81.

27.   Team R. RStudio: Integrated Devleopment for R. . Boston, MA: RStudio, PBC; 2020.

28.   Sheehan D, Janavs J, Baker R, Harnett-Sheehan K, Knapp E, Sheehan M. Mini

international neuropsychiatric interview. Tampa: University of South Florida. 1994.

29.   Arch JJ, Twohig MP, Deacon BJ, Landy LN, Bluett EJ. The credibility of exposure

therapy: Does the theoretical rationale matter? Behaviour research and therapy. 2015;72:81-92.

30.   Krisanaprakornkit T, Sriraj W, Piyavhatkul N, Laopaiboon M. Meditation therapy for

anxiety disorders. Cochrane Database of Systematic Reviews. 2006;(1).

31.   Ong CW, Lee EB, Twohig MP. A meta-analysis of dropout rates in acceptance and

commitment therapy. Behaviour research and therapy. 2018;104:14-33.

32.   Machado M, Iskedjian M, Ruiz I, Einarson TR. Remission, dropouts, and adverse drug

reaction rates in major depressive disorder: a meta-analysis of head-to-head trials. Current

medical research and opinion. 2006;22(9):1825-37.

657