Child Anemia in Cambodia: A Descriptive Analysis of Temporal and Geospatial Trends and Logistic Regression-Based Examination of Factors Associated with Anemia in Children Aged 6-59 Months

Samnang Um[1,2*], Michael R. Cope[3], and Jonathan A. Muir[4]

1 The National Institute of Public Health, Tuol Kork District, Phnom Penh, Cambodia, **2** The Center for Population Studies, Faculty of Social Science and Humanities, Royal University of Phnom Penh, Cambodia, **3** Department of Sociology, Brigham Young University, Provo, Utah, United States, **4** The Global Health Institute, Emory University, Atlanta, Georgia, United States

*umsamnang56@gmail.com

Abstract

Anemia in children remains a public health concern in developing countries. In Cambodia, the prevalence of anemia among children decreased slightly from 55.2% in 2014 to 49% in 2019. We examined descriptively temporal and geospatial trends of anemia and used logistic regression to analyze potential associations between individual and household characteristics and anemia among children aged 6-59 months using data from the Cambodia Demographic and Health Survey (CDHS) for 2005, 2010, and 2014. Survey weights were applied to account for the complex survey design of the CDHS. We used logistic regression to assess factors associated with anemia among children aged 6-59 months. The prevalence of anemia among children decreased from 62.2% in 2005 to 56.6% in 2014. The proportion of the Cambodian child population with anemia was highest in Pursat (84.3%) for 2005, Kampong Thom (67%) for 2010, and Preah Vihear and Steung Treng (68.6%) for 2014, respectively. Factors independently associated with increased odds of having anemia included children with mothers who were anemic (adjusted odds ratio (AOR) =1.77, 95% CI: 1.58 - 1.97); children who were male (AOR=1.20, 95% CI: 1.07 - 1.33), underweight (AOR=1.24, 95% CI: 1.14 - 1.57), stunted (AOR=1.24, 95% CI: 1.09 - 1.41), or had a recent episode of fever (AOR=1.16, 95% CI: 1.03 - 1.31). Children had decreased odds of having anemia if they were aged 12-23 months (AOR=0.68; 95% CI: 0.55 -0.86), 24-35 months (AOR=0.21; 95% CI: 0.17 - 0.26), 36-47 months (AOR=0.17; 95% CI: 0.13 - 0.21) or aged 48-59 months (AOR=0.15; 95% CI: 0.12 - 0.19) compared to infants; they were also less likely to have anemia if they were from a wealthier household (AOR=0.64; 95% CI: 0.50 - 0.84), and children who took drugs for the intestinal parasite were decreased by 14% (AOR = 0.86; 95% CI: 0.89-0.93) than those who did not take drugs. Anemia remains highly prevalent among children aged 6-59 months in Cambodia. Public health interventions and policies to alleviate anemia should be prioritized to address these factors.

Keywords: anemia prevalence, children, trends and determinants, Cambodia

Introduction

Anemia is a serious public health problem worldwide [1]. According to the World Health Organization (WHO) criteria, children aged 6-59 months are considered anemic if their hemoglobin level is below 11.0 grams per liter (g/dL) [2]. Anemia in children is a major cause of adverse health consequences such as stunted growth, impaired cognitive development, compromised immunity, disability and increased risk of morbidity and mortality [3]. Globally, anemia affected 40% (269 million) of children aged 6-59 months in 2019 [1]. About 43% of all children in low-income and middle-income countries were diagnosed with anemia in 2019 [1]. Moreover, a higher prevalence of anemia has occurred in the South Asian region, with 52% (87 million) of children presenting with anemia in 2019 [1]. An estimated 145,073 children die annually worldwide due to iron deficiency from anemia [4]. Similar to many low-income and middle-income nations, anemia is a serious public health problem; among children aged 6-59 months in Cambodia, the proportion of those with anemia slightly decreased from 55.2% to 49% between 2014, and 2019, respectively [5].

Anemia has significant consequences for human health, as well as for social and economic development. Previous studies found than children with very severe anemia were 4.3 times more likely to die compared to children without anemia [6]. The causes of anemia in developing countries are typically grouped into three broad categories: nutritional deficiencies, infectious diseases, and genetic hemoglobin disorders [7]. In many low- and middle-income economies, the prevalence of anemia varies by socioeconomic factors such as maternal education, household wealth status, maternal occupation, and residence [8-12]. Some studies have demonstrated evidence of anemia associated with child age and sex [10,11,13]. Moreover, children who had experienced fever, recent diarrhea had a higher likelihood of having anemia [14,15]. Serval

studies found maternal anemia increased the odds of having anemia among children [14-16].

Evidence from an analysis of nationally representative DHS data of 47 countries showed that in

over 60% of countries, children with off☐premises water access had significantly increased odds

of anemia; children exposed to surface water had higher odds of anemia in over a quarter of

countries [17].

The prevalence of anemia in Cambodia has slowly decreased in children aged under five years.

To further reduce rates of anemia in Cambodia, an improved understanding of its geographical

distribution and associated risk factors is required. An enhanced understanding will help identify

populations at greater risk for anemia and prioritize geographic areas for targeted interventions.

To our knowledge, there are no published peer-reviewed studies that assess social and

demographic factors associated with anemia among children aged 6-59 months in Cambodia

over time. One prior study on anemia patterns and risk factors utilized data from CDHS 2014

[15]. This study included all children aged 6-59 months and pooled DHS data in South and

Southeast Asian countries. An additional study aimed to describe trends in the nutritional status

of children under age five in Cambodia based on four CDHS surveys taken from 2000 to 2014 to

assess the effect of inequality (child's age, child's sex, mother's education, place of residence

and wealth index) on nutrition [18]. Given the paucity of scholarship addressing this health

concern among Cambodian children aged 6-59 months, in this study, we aimed to assess

temporal and geospatial trends of anemia among children aged 6-59 months in Cambodia and

examine factors associated with anemia among children aged 6-59 months in Cambodia.

Understanding these factors may further support policy development and programs with more

effective strategies and interventions in Cambodia to reduce the prevalence of anemia and

associated health risks such as infant and child mortality among children aged 6-59 months.

Materials and methods

Ethics Statement

The original CDHS protocol was approved by the National Ethical Committee for Health Research, Ministry of Health of the Cambodia (Ref: 056 NECHR) and the Institutional Review Board (IRB) of ICF in Rockville, Maryland, USA. Moreover, for the protection of using the secondary data in this study, the research proposal was reviewed and provided ethical approval (Ref: 225 NECHR) by the National Ethics Committee for Health Research, Ministry of Health of the Cambodia [19]. Also, the CDHS data are publicly accessible and were made available to us upon request through the DHS website at (https://dhsprogram.com/data/available-datasets.cfm) [20]. Interviewers explained the objective of the survey to parents/guardians of each participant under 18 years of age, and written informed consent was obtained from parents/guardians before data collection. Also, the study contains no individual identifiers that could affect the confidentiality of the participants, and the data were used for analysis purposes only. Details of the questionnaires, procedures, and methodology used in this survey can be found elsewhere [21-23].

Data sources and procedures

We used existing children's data from the 2005, 2010, and 2014 CDHS. The CDHS is a nationally representative population-based household survey that is regularly conducted roughly every 5 years. The survey typically uses two-stage stratified cluster sampling to collect samples from all of the Cambodian provinces, which have been divided into sampling domains. In the first stage, clusters, or enumeration areas (EAs), that represent the entire country are randomly selected from the sampling frame using estimates of probability proportional to cluster size

(PPS). The second stage then involves the systematic sampling of households listed in each

cluster or EA. Details of CDHS survey reported have been described elsewhere [21-23].

Data about the maternal and children demographic, their behaviors, their access to health care,

and their household characteristics, as well as the health status of children, were collected when

the enumerators identified children in the selected households. Anemia testing was performed on

women and men aged 15-49, as well as children aged 6-11 months when parents or caretakers

consented to the test. Blood samples were drawn using a finger prick or a heel prick in the case

of women and men aged 15-49 and children aged 6-59 months [24]. Hemoglobin analysis was

done on site using a battery-operated portable HemoCue 201+analyzer [24]. From the total of

23,687 children who were under five years old, 21,390 were of 6-59 months of age. Data on the

children's hemoglobin level were available; ultimately, data for 10,434 children 6-59 months

were included in the final analysis.

Measures

Anemia status among children aged 6-59 months was the outcome variable. The original variable

for anemia level in the CHDS was recorded as a categorical variable representing "not anemic,"

"mild anemia," "moderate anemia," and "severe anemia." Then, the original variable was

recoded into the binary variable "Anemia," with mild, moderate, and severe anemia coded as

Anemic = 1 and non-Anemic = 0.

Right-hand sided variables included individual-level (maternal and child) and household-level

characteristics. **Maternal characteristics** included age (coded as 1 = 15-24 (reference), 2 = 25-

34, and 3= 35-49 years); education (coded as 0 = no education (reference), 1 = primary, and 2=

secondary or above); marital status (coded as 0 = not married and 1 = married, which included

divorced, single, widowed, and separated); occupation (coded as 0 = not working (reference), 1 =

agricultural, which included self-employed, 2 = professional, which combined clerical, technical,

managerial, sales, and services; and 3 = manual, which included skilled and unskilled manual

work; maternal anemia (0 = non anemic and 1 = anemic); and smoking (coded as 0 = non-

smoker and 1 = smoker).

Child's characteristics included sex (coded as 1= male and 2 = female); age (coded as 1= 6-11

(reference), 2 = 12-23, 3 = 24-35, 4 = 36-47, and 5 = 48-59 months); birth order (coded as 1 = 1-

3 (reference), 2 = 4-5 and 3 = 6 or more); place of birth (coded as 1 = public (reference), 2 =

private, 3 = home); stunting (coded as 0 = not stunted and 1 = stunted); underweight (coded as 0

= not underweight and 1 = underweight); and wasting (coded as 0 = not wasted and 1 = wasted);

recent fever (coded as 0 = no and 1 = yes), recent diarrhea (coded as 0 = no and 1 = yes);

Vitamin A supplementation (coded as 0 = no and 1 = yes); intestinal parasite medication (coded

as 0 = no and 1 = yes); and Iron supplementation within the last 6 months (coded as 0 = no and 1

= yes).

Household characteristics included the DHS wealth index, coded as an ordinal level variable

with richest =1 (reference), richer = 2, middle = 3, poorer = 4, and poorest = 5 (we opted to use

the wealth index that was provided by each respective survey year of the CDHS as opposed to

calculating a wealth index across the pooled data. This decision was based on prior research that

found wealth status designation was comparable across the CDHS provided indices and an index

based on data pooled across years) [23]; source of drinking water (coded as 1 = unimproved and

0 = improved), type of toilet (coded as 1 = unimproved and 0 = improved; and residence (coded

as 1 = urban and 2 = rural). Cambodia's domains/provinces were regrouped for analytic purposes

into a categorical variable with 4 geographical regions that were coded as plains = 1 (reference),

Tonle Sap = 2, costal/sea = 3, and mountains = 4 (Plains included Phnom Penh, Kampong

Cham/Tbong Khmum, Kandal, Prey Veng, Svay Rieng, and Takeo; Tonle Sap included Banteay

Meanchey, Kampong Chhnang, Kampong Thom, Pursat, Siem Reap, Battambang, Pailin, and

Otdar Meanchey; Coastal/sea included Kampot, Kep, Preah Sihanouk, and Koh Kong; and

Mountains incnluded Kampong Speu, Kratie, Preah Vihear, Stung Treng, Mondul Kiri, and

Ratanak Kiri). Survey year was coded as 1 = 2005 (reference), 2 = 2010, and 3 = 2014.

Analytic strategy

Statistical analyses were performed using STATA version 17 (Stata Crop 2019, College Station,

TX) [25]. We formally incorporated the DHS's complex sample design using the "survey"

package; all estimations were carried out using the *svy* command in our descriptive and logistic

regression analyses. Key maternal, child and household characteristics were described using

weighted frequency distributions for specific survey years and for all years pooled together. A

temporal series of maps illustrating provincial variations in the prevalence of anemia over time

were created using ArcGIS software version 10.8 [26]. Shapefiles for administrative boundaries

in Cambodia are publicly accessible through the DHS website [27]. Data quality assurance and

minor data cleaning followed standard procedures in preparation for statistical analysis [28].

Cross-tabular weighted frequency distributions with Chi-square tests were used to assess the

association between the variables of interest (including maternal, child household characteristics)

and anemia status. Variables associated with anemia status with a p-value ≤ 0.05 were used in

the multivariate logistic regression analysis [29]. Women's age, residence, and survey years were

included in the multivariate regression based on the literature and prior knowledge. Unadjusted

logistic regression was used to analyze associations between anemia and maternal bio-

demographic and household characteristics, geographical regions, and health-related factors.

Results are reported as odds ratios (OR) with 95% confidence intervals (CI). Adjusted logistic

regression was then used to assess factors associated with anemia after adjusting for other

variables in the model. Results from the final adjusted model are reported as adjusted odds ratios

(AOR) with 95% confidence intervals and corresponding p-values. Results from the final

adjusted logistic regression model were considered statistically significant based on a p-value

less than 0.05 and 95% confidence intervals that did not cross an AOR of 1.0. Potential multi-

collinearity between right-hand sided variables was checked using a continuous version of the

outcome variable and evaluating VIF scores after fitting an Ordinary Least Squares regression

model (**S1-table**).

Results

A total of 10,294 children aged 6-59 months were included in the pooled data, there were 2,933

in 2005, 3,407 in 2010, and 3,955 in 2014, respectively. Overall, 49.2% of children were female,

23% had a mother who had at least completed secondary education, 46.5% had a mother who

was anemic, 3.7% had a mother who smoked, 86% resided in rural areas, and 26% belonged to

the poorest households (**Table 1**). The overall prevalence of anemia among children decreased

slightly from 62.2% in 2005 to 56.6% in 2014 (**Fig 1**). The extent of this downward trend varied

across the four geographical regions; the highest prevalence of anemia in the Tonle Sap region in

2005 (69.5%). The coastal region had the lowest prevalence of anemia across all survey years,

with prevalence levels in 2005 that were lower than the prevalence levels of other regions in

2014 (**Fig 1**). Anemia was highest among children in Pursat (84.3%), in Kampong Thom

province (67%), and in Preah Vihear and Steung Treng (68.6%) for 2005, 2010 and 2014,

respectively (**Fig 2, S2-Table**). While the lowest prevalence of anemia among children was

observed in Kampot and Kep provinces (48.3%), in Pursat (39.7%), and in Phnom Penh (41.3%)

for 2005, 2010 and 2014.

Table 1. Descriptive Statistics of Individual and Household Socio-demographic Characteristics

Variables	2005 (n=2,933)		2010 (n=3,407)		2014 (n=3,955)		2005-2014 (n=10,294)	
	Freq.	%	Freq.	%	Freq.	%	Freq.	%
Maternal Characteristics								
Age (years)								
15-20	127	4.3	170	5.0	210	5.3	507	4.9
20-34	1,962	66.9	2,529	74.2	3,097	78.3	7,587	73.7
35-49	844	28.8	708	20.8	648	16.4	2,200	21.4
Education								
No education	719	24.5	649	19.0	537	13.6	1,905	18.5
Primary	1,730	59.0	1,944	57.1	2,191	55.4	5,865	57.0
Secondary plus	484	16.5	814	23.9	1,226	31.0	2,365	23.0
Marital status								
Not married	132	4.5	155	4.5	179	4.5	466	4.5
Married	2,801	95.5	3,252	95.5	3,775	95.4	9,828	95.5
Occupation								
Not working	19	0.6	523	15.4	988	25.0	1,530	14.9
Agricultural	2,146	73.2	1,845	54.2	1,447	36.6	5,439	52.8
Professional	563	19.2	658	19.3	861	21.8	2,082	20.2
Manual	205	7.0	381	11.2	658	16.6	1,243	12.1
Anemia status								
Non anemic	1,438	49.0	1,898	55.7	2,151	54.4	5,488	53.3
Anemic	1,495	51.0	1,509	44.3	1,783	45.1	4,787	46.5
Smoking								
No-smoker	2,767	94.3	3,300	96.9	3,842	97.1	9,908	96.3
Smoker	166	5.7	107	3.1	113	2.9	386	3.7
Child Characteristics								
Place of birth								
Public	453	15.4	1,411	41.4	2,759	69.8	4,623	44.9
Private	113	3.9	325	9.5	541	13.7	979	9.5
Home	2,368	80.7	1,671	49.0	655	16.6	4,693	45.6
Age								
6-11	324	11.0	409	12.0	455	11.5	1,188	11.5
12-23	696	23.7	786	23.1	959	24.2	2,440	23.7
24-35	620	21.1	763	22.4	874	22.1	2,257	21.9
36-47	673	22.9	710	20.8	852	21.5	2,236	21.7
48-59	621	21.2	739	21.7	815	20.6	2,174	21.1
Sex								
Female	1,485	50.6	1,632	47.9	1,943	49.1	5,060	49.2
Male	1,448	49.4	1,775	52.1	2,011	50.8	5,234	50.8
Birth order								
1-2	1,903	64.9	2,638	77.4	3,345	84.6	7,887	76.6
3-5	610	20.8	502	14.7	439	11.1	1,551	15.1
6+	420	14.3	267	7.8	171	4.3	857	8.3
Stunted								
Not stunted	1,555	53.0	1,945	57.1	2,575	65.1	6,075	59.0

Stunted	1,305	44.5	1,387	40.7	1,311	33.1	4,003	38.9
Wasted								
Not wasted	2,629	89.6	2,970	87.2	3,518	89.0	9,117	88.6
Wasted	232	7.9	362	10.6	368	9.3	961	9.3
Underweight								
Not Underweight	1,997	68.1	2,366	69.4	2,892	73.1	7,256	70.5
Underweight	863	29.4	965	28.3	994	25.1	2,822	27.4
Recent fever								
No	1,860	63.4	2,383	69.9	2,774	70.1	7,017	68.2
Yes	1,073	36.6	1,024	30.1	1,180	29.8	3,277	31.8
Recent diarrhea								
No	2,357	80.4	2,856	83.8	3,413	86.3	8,626	83.8
Yes	576	19.6	551	16.2	541	13.7	1,668	16.2
Vitamin A supplement								
No	1,925	65.6	1,041	30.6	1,226	31.0	4,193	40.7
Yes	1,008	34.4	2,366	69.4	2,729	69.0	6,102	59.3
Iron supplement								
No	2,901	98.9	3,352	98.4	3,711	93.8	9,964	96.8
Yes	32	1.1	55	1.6	244	6.2	331	3.2
Parasite medication								
No	2,155	73.5	1,438	42.2	1,586	40.1	5,180	50.3
Yes	778	26.5	1,969	57.8	2,363	59.7	5,109	49.6
Household Characteristics								
Place of residence								
Urban	380	13.0	523	15.4	545	13.8	1,448	14.1
Rural	2,553	87.0	2,884	84.6	3,409	86.2	8,846	85.9
Wealth index								
Poorest	763	26.0	895	26.3	1,011	25.6	2,669	25.9
Poorer	668	22.8	708	20.8	792	20.0	2,168	21.1
Middle	551	18.8	639	18.8	764	19.3	1,955	19.0
Richer	482	16.4	605	17.8	670	16.9	1,757	17.1
Richest	469	16.0	560	16.4	717	18.1	1,746	17.0
Source of drinking water								
Improved	1,537	52.4	1,839	54.0	2,006	50.7	5,381	52.3
Unimproved	1,396	47.6	1,568	46.0	1,949	49.3	4,913	47.7
Type of toilet facility								
Improved	573	19.5	1,177	34.5	1,934	48.9	3,684	35.8
Non-Improved	2,360	80.5	2,230	65.5	2,021	51.1	6,610	64.2
Region								
Plain	1,326	45.2	1,655	48.6	1,770	44.8	4,751	46.2
Tonle Sap	998	34.0	1,066	31.3	1,264	32.0	3,328	32.3
Coastal	215	7.3	210	6.2	236	6.0	661	6.4
Plateau/Mountain	395	13.5	476	14.0	685	17.3	1556	15.1

Notes: Data Source: CDHS 2005, 2020, and 2014. Survey weights applied to obtain weighted percentages across survey years and pooled data.

Fig 1. Overall and regional trends of anemia among children age 6-59 months by survey year. Notes: Data Source: CDHS 2005, 2020, and 2014. Survey weights applied to obtain weighted percentages across survey years and pooled data.

Fig 2. Geographical distribution of anemia among children 6-59 months. Map created using ArcGIS software version 10.8 [26]. Shapefiles for administrative boundaries in Cambodia are publicly accessible through DHS website [20]. Notes: Survey weights applied to obtain weighted percentages across survey years and pooled data.

Preliminary analyses included evaluation of cross-tabular frequency distributions combined with Chi-square tests (see **Table 2**). Maternal age was significantly associated with the risk of anemia in children in survey years 2010, and 2014; the largest percentage of children with anemia were those with younger mothers aged less than 19 years (71.3% in 2005, 71.4% in 2010, and 71.6% in 2014). Maternal educational attainment was significantly associated with anemia in children in all survey years; the largest percentage of children with anemia were generally those whose mothers had no formal education (68.5% in 2005, 58.3% in 2010, and 57.2% in 2014). Maternal anemia was also significantly associated with anemia in children; the prevalence of anemia among children who had anemic mothers was 69% in 2005, 63.1% in 2010, and 64% in 2010. Anemia was related to children's ages, with a higher prevalence in the age group of 6 to 11 months (6-11 months: 84.3% in 2005, 83.3% in 2010, and 80.6% in 2014; 12-23 months: 79.1%, 77.1%, and 73.7%, respectively). In addition, anemia among male children was higher than female children: 64.4% in 2005, 58% in 2010, and 58.1% in 2014. Stunted child had higher prevalence of anemia (65.3% in 2005, 57.2% in 2010 and 61.5% in 2014) as did children who were underweight (65% in 2005, 60% in 2010, and 61.3% in 2014). Having had fever or diarrhea in the last 2 weeks was associated with anemia among children in all of the survey years; among children with a history of fever, 67.2% in 2005, 61.1% in 2010, and 62.4% in 2014 were anemic; among children with diarrhea, 70.4% in 2005, 68.8% in 2010, and 66.4% in 2014 were anemic, respectively. Deworming for intestinal parasites was also associated with anemia in children. Results show that among children who had taken drugs for intestinal parasites in the last 6 months, 55.3% in 2005, 52.1% in 2010, and 51.7% in 2014 were anemic. The proportion of anemic children in rural areas was higher than in urban areas in all three years, at 62.5% in 2005,

58% in 2010, and 59% in 2014. Anemia was also associated with household toilet facility; across

the survey years, 64.5% in 2005, 58.5% in 2010 and 62.5% in 2014 of children in households

that reported having no toilet facility were anemic. The proportion of children with anemia was

highest among children in the poorest households across all the survey years: 69.4% in 2005,

60.4% in 2010 and 65% in 2014 were anemic, respectively.

After adjusting for other individual or household-level characteristics (see Table 3), children

aged 6-59 months in Cambodia were more likely to have anemia if their mothers were likewise

anemic (AOR=1.77, 95% CI: 1.58 - 1.97) compared with children of non-anemic mothers; if

they were male comprared to female (AOR=1.20, 95% CI: 1.07 - 1.33) ; underweight (95% CI:

1.14 - 1.57) or stunted (95% CI: 1.09 - 1.41) compared to children who were not underweight or

stunted; if they had recently experienced a fever (AOR=1.16, 95% CI: 1.03 - 1.31) or a bout of

diarrhea (AOR=1.16, 95% CI: 0.97 - 1.38) compared to children who had not recently

experienced a fever or diarrhea; or if they were residing in rural areas (AOR=1.10, 95% CI: 0.92

- 1.31) compared to children residing in urban areas. Children were less likely to have anemia if

they were older, came from a wealthier household, or had medications for intestinal paraites in

the last 6 months. With respect to age, children aged 12-23 months (AOR=0.68; 95% CI: 0.55 -

0.86), 24-35 months (AOR=0.21; 95% CI: 0.17 - 0.26), 36-47 months (AOR=0.17; 95% CI: 0.13

- 0.21) and 48-59 months (AOR=0.15; 95% CI: 0.12 - 0.19) were less likely to be anemic than

those aged 6-11 months. Children from the wealthiest households had lower odds of being

anemic compared with those in the poorest households (AOR=0.66; 95% CI: 0.50 - 0.86). The

odds of being anemic were lower by 14% (AOR = 0.86; 95% CI: 0.89-0.93) among children who

took medications for intestinal parasites compared to those who did not take similar medications.

Table 2. Chi-square test results for associations between child anemia and child, maternal, and household characteristics

Variables	2005 (n=2,933)			2010 (n=3,407)			2014 (n=3,955)			2005-2014 (n=10,294)		
	Not Anemic	Anemic	p-value	Not Anemic	Anemic	p-value	Not Anemic	Anemic	p-value	Not Anemic	Anemic	p-value
Maternal Characteristics												
Age (years)												
15-20	28.7	71.3	0.241	28.6	71.4	<0.001	28.2	71.8	0.002	28.4	71.6	<0.001
20-34	38.3	61.7		44.0	56.0		43.9	56.1		42.5	57.5	
35-49	38.0	62.0		48.7	51.3		46.1	53.9		43.8	56.2	
Education												
No education	31.5	68.5	<0.001	41.7	58.3	<0.001	42.8	57.2	<0.001	38.2	61.8	<0.001
Primary	37.7	62.3		43.4	56.6		41.1	58.9		40.8	59.2	
Secondary plus	47.6	52.4		48.0	52.0		47.9	52.1		47.9	52.5	
Marital status												
Not married	42.7	57.3	0.294	38.2	61.8	0.201	48.6	51.4	0.239	43.5	56.5	0.592
Married	37.6	62.4		44.5	55.5		43.2	56.8		42.0	58.0	
Occupation												
Not working	31.9	68.1	0.310	44.4	55.6	0.348	40.3	59.7	<0.001	41.6	58.4	<0.001
Agricultural	36.6	63.4		42.6	57.4		38.4	61.6		39.1	60.9	
Professional	41.4	58.6		47.4	52.6		52.0	48.0		47.7	52.3	
Manual	41.0	59.0		45.9	54.1		48.1	51.9		46.3	53.7	
Anemia status												
Non anemic	44.8	55.2	<0.001	36.9	50.0	<0.001	49.6	50.4	<0.001	48.4	51.6	<0.001
Anemic	31.1	68.9		50.0	63.1		36.0	64.0		34.8	65.2	
Smoking												
No-smoker	38.1	61.9	0.266	44.5	55.5	0.102	43.8	56.2	0.011	42.5	57.5	<0.001
Smoker	33.2	66.8		33.9	66.1		29.9	70.1		32.4	67.6	
Child Characteristics												
Place of birth												
Public	38.8	61.2	0.081	42.1	57.9	0.058	43.2	56.8	0.041	42.4	57.6	<0.001
Private	49.5	50.5		51.6	48.4		48.6	51.4		49.7	50.3	
Home	37.1	62.9		44.5	55.5		40.1	59.9		40.1	59.9	
Age												
6-11	15.7	84.3	<0.001	16.7	83.3	<0.001	19.4	80.6	<0.001	17.5	82.5	<0.001
12-23	20.9	79.1		22.9	77.1		26.3	73.7		23.7	76.3	
24-35	42.3	57.7		54.3	45.7		49.4	50.6		49.1	50.9	
36-47	48.7	51.3		58.1	41.9		54.1	45.9		53.8	46.2	
48-59	51.9	48.1		58.2	41.8		59.5	40.5		56.9	43.1	
Sex												
Female	39.9	60.1	0.063	42.1	53.5	0.039	45.1	54.9	0.099	44.0	56.0	0.002
Male	35.6	64.4		46.5	57.9		41.9	58.1		40.2	59.8	

Variables	2005 (n=2,933)			2010 (n=3,407)			2014 (n=3,955)			2005-2014 (n=10,294)		
	Not Anemic	Anemic	p-value	Not Anemic	Anemic	p-value	Not Anemic	Anemic	p-value	Not Anemic	Anemic	p-value
Birth order												
1-2	39.0	61.0	0.240	44.5	55.5	0.717	43.9	56.1	0.435	42.9	57.1	0.040
3-5	36.8	63.2		42.0	58.0		41.5	58.5		39.8	60.2	
6+	33.8	66.2		45.2	54.8		38.9	61.1		38.3	61.7	
Stunted												
Not stunted	40.6	59.4	0.008	44.8	55.2	0.368	46.0	54.0	<0.001	44.3	55.7	<0.001
Stunted	34.7	65.3		42.8	57.2		38.5	61.5		38.8	61.2	
Wasted												
Not wasted	37.4	62.6	0.132	44.8	55.2	0.032	44.5	55.5	0.002	42.5	57.5	0.028
Wasted	43.9	56.1		37.8	62.2		34.2	65.8		37.9	62.1	
Underweight												
Not Underweight	39.1	60.9	0.087	45.5	54.5	0.041	45.2	54.8	0.003	43.6	56.4	<0.001
Underweight	35.2	64.8		40.2	59.8		38.7	61.3		38.1	61.9	
Recent fever												
No	40.7	59.3	<0.001	46.4	53.6	<0.001	45.9	54.1	<0.001	44.7	55.3	<0.001
Yes	32.8	67.2		38.9	61.1		37.6	62.4		36.5	63.6	
Recent diarrhea												
No	39.8	60.2	<0.001	46.7	53.3	<0.001	45.0	55.0	<0.001	44.1	55.9	<0.001
Yes	29.6	70.4		31.2	68.8		33.6	66.4		31.4	68.6	
Vitamin A supplement												
No	38.7	61.3	0.315	43.4	56.6	0.635	39.0	61.0	0.007	39.9	60.1	0.007
Yes	36.2	63.8		44.5	55.5		45.4	54.6		43.5	56.5	
Iron supplement												
No	37.9	67.6	0.577	44.1	49.8	0.464	43.7	60.8	0.280	42.1	59.6	0.614
Yes	32.4	62.1		50.2	55.9		39.2	56.3		40.4	57.9	
Parasite medication												
No	35.3	64.7	<0.001	39.2	60.8	<0.001	36.3	63.7	<0.001	36.7	63.3	<0.001
Yes	44.7	**55.3**		47.9	**52.1**		48.3	**51.7**		47.6	**52.5**	
Household Characteristics												
Place of residence												
Urban	39.8	60.2	0.608	54.8	45.2	<0.001	56.6	43.4	<0.001	51.5	48.5	<0.001
Rural	37.5	62.5		42.3	57.7		41.3	58.7		40.5	59.5	
Wealth index												
Poorest	30.6	69.4	<0.001	39.6	60.4	<0.001	35.0	65.0	<0.001	35.3	64.7	<0.001
Poorer	34.9	65.1		41.4	58.6		38.5	61.5		38.3	61.7	
Middle	38.9	61.1		41.8	58.2		45.4	54.6		42.4	57.6	
Richer	42.0	58.0		45.4	54.6		45.9	54.1		44.7	55.3	
Richest	48.1	51.9		56.4	43.6		56.4	43.6		54.2	45.8	
Source of drinking water												
Improved	38.1	61.9	0.763	45.4	54.6	0.265	44.0	56.0	0.556	42.8	57.2	0.235
Unimproved	37.4	62.6		42.8	57.2		42.8	57.2		41.3	58.7	

Variables	2005 (n=2,933)			2010 (n=3,407)			2014 (n=3,955)			2005-2014 (n=10,294)		
	Not Anemic	Anemic	p-value	Not Anemic	Anemic	p-value	Not Anemic	Anemic	p-value	Not Anemic	Anemic	p-value
Type of toilet facility												
Improved	47.4	52.6	<0.001	49.2	50.8	<0.001	49.8	50.2	<0.001	49.2	50.8	<0.001
Non-Improved	35.5	64.5		41.5	58.5		37.4	62.6		38.1	61.9	
Region												
Plain	42.7	57.3	<0.001	45.3	54.7	0.572	44.4	55.6	0.223	44.3	55.7	0.004
Tonle Sap	30.5	69.5		42.3	57.7		44.7	55.3		39.7	60.3	
Coastal	43.4	56.6		45.8	54.2		41.3	58.7		43.4	56.6	
Mountain	36.9	63.1		43.7	56.3		39.2	60.8		40.0	60.0	

Notes: Data Source: CDHS 2005, 2020, and 2014. Survey weights applied to obtain weighted percentages across survey years and pooled data.

To evaluate potential effect modification of statistically signficiant associations by time and

space, interaction models were estimated for each of the statistically significant factors from the

adjusted analysis presented in Table 3. These factors were separately interacted with region and

year. Almost all of these analyses resulted in interaction terms that were not statistically

significant, indicating that the corresponding factors were associated with child anemia

regardless of the region of Cambodia where a child lived or the year in which the data were

collected. The one exception was the interaction between parasite treament and region, which

resulted in statistically significant interaction that was visualized with predicted probabilities (see

Fig. 4). The predicted probabilities indicate that treatment for parasites was protective in the

mountainous and tonle sap regions of Cambodia as children who had recently received a

treament in these regions had significantly lower probability of anemia compared to other

children.

Table 2. Unadjusted and adjusted analyses of child anaemia in Cambodia

Variables	Unadjusted (N=10,294)		Adjusted (N=10,051)	
	OR	95%CI	AOR	95%CI
Mothers age (years)				
15-20	Ref.	Ref.	Ref.	Ref.
20-34	0.54***	(0.41 - 0.70)	0.92	(0.68-1.25)
35-49	0.51***	(0.38 - 0.68)	0.87	(0.62-1.21)
Maternal education				
No education	Ref.	Ref.	Ref.	Ref.
Primary	0.89	(0.79 - 1.02)	1.00	(0.87-1.15)
Secondary and Higher	0.67***	(0.58 - 0.78)	0.96	(0.80-1.16)
Maternal marital status				
Not married	Ref.	Ref.		
Married	1.06	(0.85 - 1.33)		
Maternal occupation				
Not working	Ref.	Ref.	Ref.	Ref.
Agricultural	1.11	(0.94 - 1.30)	1.09	(0.91-1.30)
Professional	0.78**	(0.65 - 0.93)	1.11	(0.91-1.35)
Manual and unskilled	0.83	(0.68 - 1.00)	1.07	(0.86-1.33)
Maternal anemia				
Non anemic	Ref.	Ref.	1.00	(1.00-1.00)
Anemic	1.76***	(1.59 - 1.95)	**1.77*****	**(1.58-1.97)**
Maternal smoking				
No-smoker	Ref.	Ref.	Ref.	Ref.
Smoker	1.54**	(1.19 - 1.99)	1.22	(0.91-1.64)
Place of birth				
Public	Ref.	Ref.	Ref.	Ref.

Variables	Unadjusted (N=10,294)		Adjusted (N=10,051)	
	OR	95%CI	AOR	95%CI
Private	0.74**	(0.63 - 0.89)	0.93	(0.76-1.14)
Home	1.10	(0.99 - 1.22)	0.95	(0.83-1.10)
Child age				
6-11	Ref.	Ref.		
12-23	0.68**	(0.55 - 0.85)	**0.68**[***]	**(0.54-0.86)**
24-35	0.22***	(0.18 - 0.27)	**0.21**[***]	**(0.17-0.26)**
36-47	0.18***	(0.15 - 0.22)	**0.17**[***]	**(0.13-0.21)**
48-59	0.16***	(0.13 - 0.20)	**0.15**[***]	**(0.12-0.19)**
Child sex				
Female	Ref.	Ref.	Ref.	Ref.
Male	1.17**	(1.06 - 1.29)	**1.20**[**]	**(1.07-1.33)**
Birth order				
1-2	Ref.	Ref.		
3-5	1.14*	(0.99 - 1.31)	1.07	(0.91-1.28)
6+	1.21*	(1.00 - 1.47)	1.09	(0.85-1.40)
Child stunted				
Not stunted	Ref.	Ref.	Ref.	Ref.
Stunted	1.25***	(1.13 - 1.39)	**1.24**[**]	**(1.09-1.41)**
Child wasted				
Not wasted	Ref.	Ref.	Ref.	Ref.
Wasted	1.21*	(1.02 - 1.44)	0.92	(0.75-1.12)
Child underweight				
Not underweight	Ref.	Ref.	Ref.	Ref.
Underweight	1.25***	(1.12 - 1.40)	**1.27**[**]	**(1.10-1.47)**
Recent fever				
No	Ref.	Ref.	Ref.	Ref.
Yes	1.41***	(1.27 - 1.57)	**1.16**[*]	**(1.03-1.31)**
Recent diarrhea				
No	Ref.	Ref.	Ref.	Ref.
Yes	1.72***	(1.48 - 2.00)	1.16	(0.97-1.38)
Vitamin A supplement				
No	Ref.	Ref.	Ref.	Ref.
Yes	0.86**	(0.77 - 0.96)	1.04	(0.91-1.20)
Taken iron				
No	Ref.	Ref.	Ref.	Ref.
Yes	1.08	(0.81 - 1.43)	1.30	(0.95-1.79)
Parasite medication				
No	Ref.	Ref.	Ref.	Ref.
Yes	0.64***	(0.58 - 0.71)	**0.86**[*]	**(0.76-0.98)**
Place of residence				
Urban	Ref.	Ref.	Ref.	Ref.
Rural	1.56***	(1.35 - 1.81)	1.10	(0.92-1.31)
Wealth index				
Poorest	Ref.	Ref.	Ref.	Ref.
Poorer	0.88*	(0.76 - 1.01)	0.95	(0.81-1.11)
Middle	0.74***	(0.63 - 0.87)	0.84	(0.70-1.01)
Richer	0.68***	(0.58 - 0.79)	0.85	(0.68-1.06)
Richest	0.46***	(0.39 - 0.54)	**0.66**[**]	**(0.50-0.86)**
Source of drinking water				
Improved	Ref.	Ref.		
Unimproved	1.06	(0.96 - 1.18)		
Type of toilet facility				
Improved	Ref.	Ref.	Ref.	Ref.
Non-improved	1.57***	(1.42 - 1.74)	1.17	(0.99-1.38)
Region				
Plains	Ref.	Ref.	Ref.	Ref.

Variables	Unadjusted (N=10,294)		Adjusted (N=10,051)	
	OR	95%CI	AOR	95%CI
Tonle Sap	1.21**	(1.06 - 1.37)	1.10	(0.96-1.26)
Coastal	1.03	(0.86 - 1.24)	1.05	(0.86-1.27)
Plateau/Mountain	1.19*	(1.03 - 1.37)	1.01	(0.86-1.19)
Survey years				
2005	Ref.	Ref.		
2010	0.77**	(0.67 - 0.88)	**0.85***	**(0.72-0.99)**
2014	0.79**	(0.69 - 0.91)	0.90	(0.76-1.07)

Notes: Data Source: CDHS 2005, 2020, and 2014. Survey weights applied to obtain weighted percentages across year and pooled data. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Fig 3. Adjusted analyses of the odds of a child aged 6-59 months being anaemic in Cambodia (**N=10,051**)

Fig 4. Predicted probability of child anemia in Cambodia by region and parasitic treatment (**N=10,051**)

Discussion

This study shows that the prevalence of anemia among children in Cambodia declined only slightly over between 2005 and 2014; it increased between the surveys in 2010 and 2014 and still affects a majority of the children aged 6-59 months. The overall prevalence of anemia among children was 62.2% in 2005, 55.8% in 2010, and 56.6% in 2014, which is similar to the prevalence of anemia among children under five in South Asia (61.5% in 2005, 57.4% in 2010, 54.4% in 2014, and 52% in 2019) [1,30]. According to WHO (2008), the cut-off values for public health significance are such that when the prevalence of anemia is 40% and above, the country is considered to have a serious public health problem [2]. This study suggests that Cambodia has not achieved national nutrition targets, which were aimed at reducing the prevalence of anemia among children by 42% and 36% in 2015 and 2020, respectively [31]. Although Cambodia has had a comprehensive national nutrition policy, strategy, and plan of action since 2008 [31], the implementation and enforcement of these policies is lacking in the country's remote settings [31]. In particular, the lack of nationwide interventions to address the problem of anemia among Cambodian children is a major public health challenge.

Our study found several factors associated with increased odds of having anemia among children

aged 6-59 months in Cambodia, including maternal anemia, being male, being stunted, being

underweight, and having a history of fever diarrhea in the last 2 weeks. However, children living

in the richest wealth quintiles and children aged 12-59 months were less likely to have anemia.

Maternal anemia increased the odds of having anemia among children. This finding is consistent

with previous studies in Laos [14], South and Southeast Asia [15], Southern Africa [16], and

Uganda [32]. There are several explanations for this. For instance, maternal anemia during

pregnancy may contribute to low birth weight and preterm birth, both of which increase the risk

of childhood anemia [33]. Maternal anemia may also reduce iron content in breast milk [34].

Children have lower iron intake from breast milk if their mothers are anemic [35]. Moreover,

mother and child share the same socioeconomic environment, and when the child is on family

food, the child's diet quality may be similar to that of the mother [14,15,18,36]. For example,

women who live in low-income families have problems purchasing and providing an iron-rich

diet and other micro-nutrient-rich food for themselves and their children, which can increase the

risk of developing childhood anemia [8,15,32].

Children aged 12-59 months were less likely to have anemia as compared to those of 6-11

months. This is consistent with other studies showing that anemia was more prevalent among

children aged 6-11 months. This reduction in the number of children becoming anemic continues

with age, possibly because the concentration of iron and other micronutrients in human milk is

relatively low [37]. Inadequate iron intake contributes to the development of anemia among

infants aged 12 months after exclusive breastfeeding for 6 months [38]. Male children have

increased odds of having anemia. Similar results were found in studies in Laos [14] and sub-

Saharan Africa [39]. This may be explained by physiological variations in gender or due to

longitudinal and faster growth rates in boys than in girls [14,39], which requires a higher demand

for iron for their bodies [40], particularly during the early years of life [41].

In addition, stunted and underweight children had increased odds of having anemia. Malnutrition

is a major public health challenge in Cambodia. A prevalence of stunted youth decreased from

32% to 22%, and those were underweight from 25% to 16.3%, among children under five years

of age in 2014 and 2021-22 [42]. Being stunted and underweight is associated with acute and

chronic malnutrition, which share common causes with anemia [43], and these factors are

aggravated by food insecurity and poverty. Food insecurity and inadequate consumption of

micronutrients such as iron, vitamin B12, and folate affect the nutritional status of children,

which contributes to the development of anemia [43]. Therefore, public health interventions

should be targeted at improving the nutritional status of children by introducing diverse

complementary foods enriched with micronutrients such as iron, vitamin B12, and folate from

infancy.

Children with a history of fever had increased odds of having anemia, which is consistent with

the results in previous studies conducted in Laos [14], South and Southeast Asia [15], and

Southern Africa [16]. Fever among under-five children in Cambodia was a common symptom of

acute and chronic inflammatory disease [31]. Cambodia is one of the malaria-endemic areas in

Southeast Asia [44]. An association between childhood anemia and inflammatory infections,

which includes fever, cough, and worm infestation, has been well established [45]. The possible

pathway of this association is due to iron losses during the time of infections [46]. One previous

study reported that iron-deficiency anemia may depend on inadequate iron-rich food intake,

increased iron use in the body, and iron depletion due to parasitic infection [45]. Also, previous

studies have reported that malaria increases erythrocyte destruction with a simultaneous failure

of the bone marrow thus jeopardizing the ability to compensate for the losses [46].

Children of rich and middle-class households had lower odds of having anemia compared to poor

households. This finding is also consistent with previous studies [14,15,18,36]. This may

plausibly reflect improved household nutritional status linked to SES [32]. Better household

income allows for living in good social and economic conditions, and hence it is easy for higher

income families to provide a diverse diet, adequate care, and health services to their children [16].

Children who took drugs for intestinal parasites in previous 6 months had lower odds of having

anemia compared to children who did not take drugs for intestinal parasites. It is consistent with

studies in Thailand [47] and sub-Saharan Africa [48].This may be due to the fact that intestinal

parasites can induce anemia, and treating intestinal parasites with medication can reduce the

incidence of anemia in children [49].

This study was conducted using pooled Cambodia Demographic and Health surveys from 2005,

2010, and 2014, which enabled us to evaluate the temporal and geographical trends in anemia

among children aged 6-59 months using nationally representative surveys, which enables us to

generalize our findings to children in Cambodia. Therefore, these findings could be used to

determine factors associated with anemia among children aged 6-59 months throughout

Cambodia. However, our study has some limitations. First, the cross-sectional study design does

not make it possible to establish a cause-effect relationship. Second, this study relied on

hemoglobin as the measure of anemia; further studies should consider other red blood cell

indices [50]. Third, this study included self-reported data for the occurrence of diarrhea and fever;

therefore, we could not completely rule out recall bias or social desirability bias in the case of

those findings. Fourth, since this study is based on secondary data analysis, several known risk

factors for anemia, including parasitic infections such as malaria, and dietary information were

not included in this study. Lastly, the HemoCue device has been widely used in field settings

where resources are limited since it is portable, simple to use, and reasonably priced, However,

air bubbles, too much blood on the cuvettes' back sides, overfilling the cuvettes, and inadequate

mixing of the samples can all provide inaccurate results [51]. Additionally, there is variability in

Hemoglobin (Hb) and measurements dependent on Hemocue models, which may have affected

the prevalence of anemia presented in this study [51]. Finally, the most recent round of CDHS

data with information on anemia status was gathered in 2014 (Hemoglobin measurements were

not collected in CDHS 2022). This may no longer reflect present prevalence of anemia among

children in Cambodia. Future research should further evaluate these relationships and the general

prevalence of anemia among young Cambodian children aged 6-59 months once additional data

become available.

Conclusion

Our findings suggest that the high prevalence of anemia among children aged 6-59 months in

Cambodia declined only slightly over the 10-year period between 2005 and 2014. Anemia

among children continues to be a major public health concern in Cambodia. This study indicated

that childhood anemia was associated with various factors, including being male, being younger,

being stunted, being underweight, having a history of fever or diarrhea in the last 2 weeks,

maternal anemia, being a member of the poorest families, did not take drugs for intestinal

parasites and having a place of residence in a rural setting. Public health practitioners and policy

makers should consider demographic groups represented by these characteristics to produce

better targeted interventions to address anemia among children in Cambodia.

Funding Statement

The authors received no specific funding for this work.

Competing Interests

The authors have declared that no competing interests exist.

Data Availability

Our study used 2005, 2010, 2014 and 2021-22 Cambodia Demographic and Health Survey

(CDHS) datasets. The DHS data are publicly available from the website at (URL:

https://www.dhsprogram.com/data/available-datasets.cfm).

The Shapefiles for administrative boundaries in Cambodia are publicly accessible through DHS

website at (URL: https://spatialdata.dhsprogram.com/covariates).

Acknowledgments

The authors are grateful to DHS-ICF for the gathering, cleaning, and provision of the data used

for this paper. We are indebted to Dr. Chhea Chhorvann and Dr. Heng Sopheab from the

National Institute of Public Health, Phnom Penh, Cambodia for their encouragement, feedback,

and   support.

Reference

1.   Stevens GA, Paciorek CJ, Flores-Urrutia MC, Borghi E, Namaste S, Wirth JP, et al. National, regional, and global estimates of anaemia by severity in women and children for 2000–19: a pooled analysis of population-representative data. The Lancet Global Health. 2022;10(5):e627-e39.

2.   WHO. World Health Organization, Anaemia. Available online: [https://www.who.int/data/nutrition/nlis/info/anaemia] 2023.

3.   WHO. Global nutrition targets 2025: anaemia policy brief (WHO/NMH/NHD/14.4). Geneva: World Health Organization; 2014.

4.   Roser M, Ritchie H, Dadonaite B. Child and infant mortality. Our world in data https://ourworldindataorg/child-mortality (accessed January 2023). 2013.

5.   World Health Organization-Global Health Observatory Data Repository, Appears in: Anaemia in children < 5 years. Available online: [https://www.who.int/data/gho/data/indicators/indicator-details/GHO/anaemia-in-children-under-5-years-number]. Accessed March 2023.

6.   Chami N, Hau DK, Masoza TS, Smart LR, Kayange NM, Hokororo A, et al. Very severe anemia and one year mortality outcome after hospitalization in Tanzanian children: A prospective cohort study. PloS one. 2019;14(6):e0214563. Epub 2019/06/21. doi: 10.1371/journal.pone.0214563. PubMed PMID: 31220109; PubMed Central PMCID: PMCPmc6586275.

7.   Chaparro CM, Suchdev PS. Anemia epidemiology, pathophysiology, and etiology in low- and middle-income countries. Annals of the New York Academy of Sciences. 2019;1450(1):15-31. Epub 2019/04/23. doi: 10.1111/nyas.14092. PubMed PMID: 31008520; PubMed Central PMCID: PMCPmc6697587.

8.   Shimanda PP, Amukugo HJ, Norström F. Socioeconomic factors associated with anemia among children aged 6-59 months in Namibia. Journal of public health in Africa. 2020;11(1):1131. Epub 2020/11/20. doi: 10.4081/jphia.2020.1131. PubMed PMID: 33209233; PubMed Central PMCID: PMCPmc7649727.

9.   Balarajan Y, Ramakrishnan U, Özaltin E, Shankar AH, Subramanian S. Anaemia in low-income and middle-income countries. The lancet. 2011;378(9809):2123-35.

10.   Cornet M, Le Hesran J-Y, Fievet N, Cot M, Personne P, Gounoue R, et al. Prevalence of and risk factors for anemia in young children in southern Cameroon. The American journal of tropical medicine and hygiene. 1998;58(5):606-11.

11.   Foote EM, Sullivan KM, Ruth LJ, Oremo J, Sadumah I, Williams TN, et al. Determinants of anemia among preschool children in rural, western Kenya. The American journal of tropical medicine and hygiene. 2013;88(4):757.

12.   Khan JR, Awan N, Misu F. Determinants of anemia among 6-59 months aged children in Bangladesh: evidence from nationally representative data. BMC pediatrics. 2016;16:3. Epub 2016/01/13. doi: 10.1186/s12887-015-0536-z. PubMed PMID: 26754288; PubMed Central PMCID: PMCPmc4707771.

13.   Stevens GA, Bennett JE, Hennocq Q, Lu Y, De-Regil LM, Rogers L, et al. Trends and mortality effects of vitamin A deficiency in children in 138 low-income and middle-income countries between 1991 and 2013: a pooled analysis of population-based surveys. The Lancet Global Health. 2015;3(9):e528-e36.

14.   Keokenchanh S, Kounnavong S, Midorikawa K, Ikeda W, Morita A, Kitajima T, et al. Prevalence of anemia and its associated factors among children aged 6–59 months in the Lao People's Democratic Republic: A multilevel analysis. PloS one. 2021;16(3):e0248969.

15.   Sunuwar DR, Singh DR, Pradhan PMS, Shrestha V, Rai P, Shah SK, et al. Factors associated with anemia among children in South and Southeast Asia: a multilevel analysis. BMC Public Health. 2023;23(1):343. doi: 10.1186/s12889-023-15265-y.

16.   Ntenda PAM, Nkoka O, Bass P, Senghore T. Maternal anemia is a potential risk factor for anemia in children aged 6-59 months in Southern Africa: a multilevel analysis. BMC Public Health. 2018;18(1):650. Epub 2018/05/24. doi: 10.1186/s12889-018-5568-5. PubMed PMID: 29788935; PubMed Central PMCID: PMCPmc5964691.

17.   Kothari MT, Coile A, Huestis A, Pullum T, Garrett D, Engmann C. Exploring associations between water, sanitation, and anemia through 47 nationally representative demographic and health

surveys. Annals of the New York Academy of Sciences. 2019;1450(1):249-67. Epub 2019/06/25. doi:
10.1111/nyas.14109. PubMed PMID: 31232465; PubMed Central PMCID: PMCPmc6771505.
18. Greffeuille V, Sophonneary P, Laillou A, Gauthier L, Hong R, Hong R, et al. Persistent
inequalities in child undernutrition in Cambodia from 2000 until today. Nutrients. 2016;8(5):297.
19. Um S, Sopheab H, Yom A, Muir JA. Trends and factors associated with anemia among pregnant
women aged 15-49 years old in Cambodia: Data analysis of the Cambodia Demographic and Health
Surveys. medRxiv. 2022:2022.09. 08.22279710.
20. DHS Program. The Demographic and Health Surveys (DHS) Program; ICF: Rockville, MD, USA,
2020; Available online: [https://dhsprogram.com/data/available-datasets.cfm].
21. NIS, N., National Institute of Public Health, National Institute of Statistics [Cambodia] and ORC
Macro. 2006. Cambodia Demographic and Health Survey 2005. Phnom Penh, Cambodia and Calverton,
Maryland, USA: National Institute of Public Health, National Institute of Statistics and ORC Macro. 2005.
22. NIS, National Institute of Statistics, Directorate General for Health, and ICF Macro, 2011.
Cambodia Demographic and Health Survey 2010. Phnom Penh, Cambodia and Calverton, Maryland,
USA: National Institute of Statistics, Directorate General for Health, and ICF Macro. 2010.
23. Statistics NIo. Cambodia Demographic and Health Survey 2014. 2015.
24. Demographic and health surveys methodology, Biomarker field manual, ICF International:
[Internet]. Calverton; 2012. Available from:
https://dhsprogram.com/pubs/pdf/DHSM7/DHS6_Biomarker_Manual_9Jan2012.pdf.
25. StataCorp. 2021. Stata Statistical Software: Release 17. College Station, TX: StataCorp LLC.
26. Desktop A. Release 10.8. 1.(2020). ESRI (Environmental Systems Research Institute), Redlands,
CA.
27. DHS program, Spatial Data Repository Boundaries. Avaiable online:
[https://spatialdata.dhsprogram.com/covariates/].
28. Cunningham, S. A. and Muir, J. A. (In Press). The Cambridge Handbook of Research Methods
and Statistics for the Social and Behavioral Sciences: Volume 1: Building a Program of Research, volume
1, book section Data Cleaning. Cambridge University Press.
29. Chhea C, Ir P, Sopheab H. Low birth weight of institutional births in Cambodia: Analysis of the
Demographic and Health Surveys 2010-2014. PloS one. 2018;13(11):e0207021. Epub 2018/11/09. doi:
10.1371/journal.pone.0207021. PubMed PMID: 30408102; PubMed Central PMCID: PMCPmc6224106.
30. WHO. The global prevalence of anaemia in 2011. Geneva: World Health Organization. 2015.
31. National Maternal and Child Health Center; National Nutrition Program; Fast Track Road Map
for Improving Nutrition 2014 - 2020. Available online [https://nmchc.gov.kh/wp-
content/uploads/2021/07/FTRM-for-Nutrition_Eng_Final-May-2014with-signature.pdf]. .
32. Nankinga O, Aguta D, Kabahuma C. Trends and determinants of anemia in Uganda: further
analysis of the Demographic and Health Surveys. DHS Working Papers. 2019;(149).
33. Abioye AI, McDonald EA, Park S, Ripp K, Bennett B, Wu HW, et al. Maternal anemia type
during pregnancy is associated with anemia risk among offspring during infancy. Pediatric research.
2019;86(3):396-402.
34. Kumar A, Rai AK, Basu S, Dash D, Singh JS. Cord blood and breast milk iron status in maternal
anemia. Pediatrics. 2008;121(3):e673-e7.
35. Skau JK, Grenov B, Chamnan C, Chea M, Wieringa FT, Dijkhuizen MA, et al. Stunting, wasting
and breast-feeding as correlates of body composition in Cambodian children at 6 and 15 months of age.
British Journal of Nutrition. 2019;121(6):688-98.
36. Pasricha S-R, Black J, Muthayya S, Shet A, Bhat V, Nagaraj S, et al. Determinants of anemia
among young children in rural India. Pediatrics. 2010;126(1):e140-e9.
37. Friel J, Qasem W, Cai C. Iron and the breastfed infant. Antioxidants. 2018;7(4):54.
38. Wang F, Liu H, Wan Y, Li J, Chen Y, Zheng J, et al. Prolonged exclusive breastfeeding duration
is positively associated with risk of anemia in infants aged 12 months. The Journal of nutrition.
2016;146(9):1707-13.
39. Bjertness E, Wu T, Stigum H, Nafstad P. Acute mountain sickness, arterial oxygen saturation and
heart rate among Tibetan students who reascend to Lhasa after 7 years at low altitude: a prospective
cohort study. BMJ open. 2017;7(7):e016460.

40.   Zuffo CRK, Osório MM, Taconeli CA, Schmidt ST, Silva BHCd, Almeida CCB. Prevalence and
risk factors of anemia in children. Jornal de pediatria. 2016;92:353-60.
41.   Konstantyner T, Roma Oliveira TC, de Aguiar Carrazedo Taddei JA. Risk factors for anemia
among Brazilian infants from the 2006 National Demographic Health Survey. Anemia. 2012;2012.
42.   National Institute of Statistics (NIS) [Cambodia], Ministry of Health (MoH) [Cambodia], and ICF.
2022. Cambodia Demographic and Health Survey 2021–22 Key Indicators Report. Phnom Penh,
Cambodia, and Rockville, Maryland, USA: NIS, MoH, and ICF.
43.   Engidaye G, Melku M, Yalew A, Getaneh Z, Asrie F, Enawgaw B. Under nutrition, maternal
anemia and household food insecurity are risk factors of anemia among preschool aged children in Menz
Gera Midir district, Eastern Amhara, Ethiopia: a community based cross-sectional study. BMC public
health. 2019;19(1):1-11.
44.   National Centre for Parasitology Entomology and Malaria Control (CNM) National Malaria
Porgram Review. CNM; Phnom Penh, Cambodia: 2019.
45.   Alamneh TS, Melesse AW, Gelaye KA. Determinants of anemia severity levels among children
aged 6–59 months in Ethiopia: Multilevel Bayesian statistical approach. Scientific Reports.
2023;13(1):4147. doi: 10.1038/s41598-022-20381-7.
46.   Menendez C, Fleming AF, Alonso PL. Malaria-related anaemia. Parasitol Today. 2000;16:469–
76.
47.   Watthanakulpanich D, Maipanich W, Pubampen S, Sa-Nguankiat S, Pooudouang S,
Chantaranipapong Y, et al. Impact of hookworm deworming on anemia and nutritional status among
children in Thailand. The Southeast Asian journal of tropical medicine and public health. 2011;42(4):782-
92. Epub 2012/02/04. PubMed PMID: 22299460.
48.   Tesema GA, Worku MG, Tessema ZT, Teshale AB, Alem AZ, Yeshaw Y, et al. Prevalence and
determinants of severity levels of anemia among children aged 6–59 months in sub-Saharan Africa: A
multilevel ordinal logistic regression analysis. PloS one. 2021;16(4):e0249978.
49.   Girum T, Wasie A. The Effect of Deworming School Children on Anemia Prevalence: A
Systematic Review and Meta-Analysis. The open nursing journal. 2018;12:155-61. Epub 2018/09/11. doi:
10.2174/1874434601812010155.   PubMed   PMID: 30197721;   PubMed   Central   PMCID:
PMCPmc6110060.
50.   Ntenda PAM, Chuang K-Y, Tiruneh FN, Chuang Y-C. Multilevel analysis of the effects of
individual-and community-level factors on childhood Anemia, severe Anemia, and hemoglobin
concentration in Malawi. Journal of tropical pediatrics. 2018;64(4):267-78.
51.   Rappaport AI, Karakochuk CD, Whitfield KC, Kheang KM, Green TJ. A method comparison
study between two hemoglobinometer models (Hemocue Hb 301 and Hb 201+) to measure hemoglobin
concentrations and estimate anemia prevalence among women in Preah Vihear, Cambodia. International
journal of laboratory hematology. 2017;39(1):95-100.

Supplementary Material 1

S1 Table. Results of checking multicollinearity using Variance Inflation Factor

Variable	VIF
Wealth quintile	2.13
Type of toilet facility	1.96
Children underweight	1.63
Birth order	1.56
Mother age	1.50
Children stunted	1.45
Mother education	1.36
Region	1.24
Children wasted	1.2
Place of residence	1.18
Recent diarrhea	1.11
Child age	1.10
Recent fever	1.09
Source of drinking water	1.09
Maternal anemia	1.02
Sex of child	1.00

587 S2 Table. **Prevalence of Anemia in the 19 domains in CDHS 2005 2010 and 2014**

Provinces	CDHS 2005 (n=2,933)		CDHS 2010 (n=3,407)		2014 (n=3,955)	
	%Anemia	95% CI	%Anemia	95% CI	%Anemia	95% CI
Banteay Meanchey	70.6	(61.0-78.7]	52.7	(42.9-62.2]	41.5	(35.6-47.6]
Kampong Cham	57.8	(50.1-65.2]	56.8	(47.7-65.5]	63.9	(57.7-69.7]
Kampong Chhnang	58.6	(49.1-67.5]	62.5	(56.9-67.7]	57.5	(50.6-64.1]
Kampong Speu	63.0	(54.6-70.7]	55.1	(46.2-63.6]	64.1	(55.3-72.0]
Kampong Thom	74.1	(68.7-78.9]	67.0	(57.6-75.2]	67.1	(58.5-74.7]
Kandal	57.2	(48.4-65.5]	56.5	(46.6-65.9]	59.8	(50.1-68.8]
Kratie	59.4	(51.8-66.5]	62.2	(54.8-69.0]	50.7	(43.8-57.6]
Phnom Penh	54.3	(38.9-68.9]	48.2	(41.7-54.7]	41.3	(34.6-48.4]
Prey Veng	57.0	(48.1-65.5]	51.6	(43.3-59.9]	54.5	(44.7-64.0]
Pursat	**84.3**	(77.3-89.4]	39.7	(33.1-46.8]	64.2	(56.0-71.6]
Siem Reap	78.0	(70.5-84.0]	61.4	(54.3-68.0]	50.5	(44.0-56.9]
Svay Rieng	68.0	(57.8-76.8]	66.1	(56.2-74.7]	50.9	(40.8-60.9]
Takeo	55.5	(47.8-62.9]	53.2	(41.5-64.5]	54.6	(45.4-63.4]
Odar Meanchey	73.4	(57.5-84.9]	60.0	(51.9-67.7]	63.1	(55.4-70.1]
Battambong/Pailin	54.6	(47.2-61.8]	55.1	(46.7-63.3]	50.1	(38.6-61.6]
Kampot/Kep	48.3	(41.3-55.3]	51.7	(41.5-61.8]	59.7	(50.5-68.3]
Sihanoukville/Koh Kong	74.7	(66.7-81.3]	58.8	(49.7-67.4]	57	(49.7-63.9]
Preah Vihear/Stung Treng	67.5	(60.3-73.9]	54.7	(48.2-61.1]	68.6	(61.1-75.2]
Mondulkiri/Rattanakiri	62.6	(56.3-68.5]	53.1	(47.1-59.1]	57.1	(48.4-65.5]
Overall	62.2	(59.8-64.6]	55.8	(53.5-58.1]	56.6	(54.4-58.7]

2005

2010

2014

% Anemia
- 40 - 44
- 45 - 49
- 50 - 59
- 60 - 69
- 70 - 79
- 80 - 89

Adjusted Odds Ratios