# Evaluating the Risk of Conflict on Recent Ebola Outbreaks in Guinea and the Democratic Republic of the Congo

Gina E C Charnley[1,2,3*], Nathan Green[4], Ilan Kelman[3,5,6], Espoir B Malembaka[1,7], Katy A M Gaythorpe[2,8]

1. Bloomberg School of Public Health, Johns Hopkins University, Baltimore, United States

2. School of Public Health, Imperial College London, London, United Kingdom

3. Institute for Global Health, University College London, London, United Kingdom

4. Department of Statistical Science, University College London, London, United Kingdom

5. Institute for Disaster Risk Reduction, University College London, London, United Kingdom

6. University of Agder, Kristiansand, Norway

7. Centre for Tropical Diseases and Global Health (CTDGH), Université Catholique de Bukavu, Bukavu, The Democratic Republic of the Congo

8. MRC Centre for Global Infectious Disease Analysis, Imperial College London, London, United Kingdom

* Corresponding author

**Abstract**

*Background*

Reducing Ebola virus transmission relies on the ability to identify cases and limit contact with

infected bodily fluids through biosecurity, safe sex practices, safe burial and vaccination. Armed

conflicts can complicate outbreak detection and interventions due to widespread disruption to

governments and the populations. Guinea and the Democratic Republic of the Congo (DRC) have

historically reported the largest and the most recent Ebola virus outbreaks. Understanding if

conflict played a role in these outbreaks may help in identifying key risks factors to improve

disease control.

*Methods*

We used data from a range of publicly available data sources for both Ebola virus cases and

conflict events from 2018 to 2021 in Guinea and the DRC. We fitted these data to conditional

logistic regression models using the Self-Controlled Case Series methodology to evaluate the

magnitude in which conflict increased the risk of reported Ebola virus cases in terms of incidence

rate ratio. We re-ran the analysis sub-nationally, by conflict sub-event type and tested any lagged

effects.

*Results*

Conflict was significantly associated with an increased risk of reported Ebola virus cases in both

the DRC and Guinea in recent outbreaks. The effect was of a similar magnitude at 1.88- and 1.98-

times increased risk for the DRC and Guinea, respectively. The greatest effects (often higher than

the national values) were found in many conflict prone areas and during protest/riot-related conflict

events. Conflict was influential in terms of Ebola virus risk from 1 week following the event and

remained important by 10 weeks.

*Conclusion*

Extra vigilance is needed following protests and riot-related conflict events in terms of Ebola virus

transmission. These events are highly disruptive, in terms of access to transportation and

healthcare and are often in urban areas with high population densities. Additional public health

messaging around these types of conflict events, relating to the risks and clinical symptoms may

be helpful in reducing transmission. Future work should aim to further understand and quantify

conflict severity and intensity, to evaluate dose-response relationships in terms of disease risk.

**Keywords**

Ebola virus, Conflict, Outbreaks, Guinea, the Democratic Republic of the Congo

**Background**

Ebola is a virus belonging to the *Filoviridae* family and was first identified in 1976 in what are now

the Democratic Republic of the Congo (DRC) and South Sudan [1,2]. Ebola virus disease (EVD)

causes acute haemorrhagic fever, leading to mortality rates of ~50%, in the absence of any

specific treatment. The virus is spread through human-to-human transmission via contact with

infected body fluids and fomites, with an average incubation period of 8 to 10 days [3,4].

Since its discovery, there have been several EVD outbreaks (with sporadic cases in Europe and

North America) [5]. The worst recorded outbreaks, in terms of total cases and deaths, were the

2013-2016 West Africa outbreak, particularly in Sierra Leone, Liberia and Guinea and the 2018-

2020 outbreak in the DRC. The most recent EVD outbreaks have been reported in Uganda (*Sudan*

*Orthoebolavirus)*, Guinea (*Zaire Orthoebolavirus)* and the DRC (*Zaire Orthoebolavirus)*, all of

which occurred in 2021 and 2022 [6].

EVD control is achieved through identifying cases and interventions which reduce the risk of

contact with infected people, via biosecurity in EVD treatment centres, safe burial practices,

reducing sexual transmission [7], timely detection (via strategically positioned decentralized

laboratories) [8], contact tracing and post-mortem testing to identify EVD-related deaths, genomic

sequencing (to infer spatiotemporal transmission dynamics) [9,10], and more recently with the

rVSVΔG-ZEBOV-GP Ebola virus envelope glycoprotein vaccine and monoclonal antibody

therapeutics [11,12]. Conflict can impact these control measures, reducing the capacity to identify

and respond to outbreaks [13]. Furthermore, conflict can cause disruption to a range of basic

services including transport, healthcare and water, sanitation and hygiene (WASH) [14].

Targeted attacks on healthcare centres and workers are not limited to specific diseases or

geographic regions [15] and there have been several reported attacks on Ebola treatment centres,

creating fear and avoidance in accessing them [16]. To protect staff and patients, security forces

and military personnel are often required to carry out healthcare activities [10,15,17]. Additionally,

fear and mistrust can lead to further attacks, sabotage and disruption of healthcare facilities and

services. Distrust and dissatisfaction in the government, the military and international involvement

via non-governmental organisations (NGOs) has also reduced the willingness to follow public

health guidance and has increased vaccine hesitancy and misinformation [7,18].

In previous EVD outbreaks in the DRC and Guinea, conflict has been an ongoing issue to greater

or lesser extents [19,20]. The DRC is currently experiencing armed conflict, particularly but not

limited to the Kivu provinces. The conflict has been ongoing since the 1990s due to instability

within the region, with many armed groups currently active along with governments' involvement

[21,22]. Guinea has experienced conflict, albeit to a lesser extent than in the DRC, mainly due to

politically driven violence and demonstrations during post-election periods and excessive force by

security forces [23].

Due to Guinea and the DRC having both the largest recorded outbreaks and the most recent EVD

outbreaks [6], along with reported conflict, there is a need to further understand the mechanisms

and evaluate the up-to-date conflict-related risks for recent EVD resurgences. Additionally, it is

essential to understand the timing of conflict-related risks for EVD transmission and any lagged

effects. Increasing our understanding of the magnitude and timing of risks can help EVD response

planning and identify when extra surveillance may be needed.

Despite evidence showing possible links and effects of conflict on EVD, research to date is limited

on magnitude, timings and mechanisms through which conflict impacts the risk of EVD, particularly

qualitatively. Here, we use a methodology that has proved robust in evaluating the relationship

between disease and conflict in previous studies [24] to address the following research aims:

1.  Evaluate the risk of conflict on reported EVD cases during the 2018 to 2020 outbreak in the

88            DRC and the 2021 outbreak in Guinea, both nationally and sub-nationally.

89        2.  Evaluate if certain conflict types were more important than others in impacting EVD cases

90            and suggest mechanisms for any differences found.

3.  Examine the timing of conflict-related risk on EVD, including lagged effects.

**Methods**

*Spatial Administrative Units*

Subnational administrative boundaries differ between Guinea and the DRC, Table 1 presents each

country's definitions of subnational regions.

**Table 1** Subnational administrative unit names for Guinea and the Democratic Republic of the

Congo (DRC).

| Administrative Unit | Country | Name |
|---|---|---|
| 1 | Guinea | Region |
| 1 | DRC | Province |
| 2 | Guinea | Prefecture |
| 2 | DRC | Territory |
| 3 | Guinea | Sub-prefecture |
| 3 | DRC | Sector |
| 4 | Guinea | Settlement |
| 4 | DRC | Settlement |

*Datasets*

For the DRC, EVD data were collated from the DRC Ministry of Health (MoH) mailing lists [25,26]

and from the Humanitarian Emergency Response Africa (HEMA) dataset [27]. The temporal scale

of the data was daily and ranged from 04 April 2018 to 11 July 2020 for the MoH data and 06

February 2021 to 14 April 2021 for the HEMA data. The spatial scale for the DRC data was to

administrative level 2 for the DRC MoH data and to administrative level 1 for the HEMA dataset

(Table 1), including new reported case and death counts. For Guinea, the data ranged from 14

February 2021 to 19 June 2021 and included new and cumulative daily case counts by

administrative level 1 (Table 1), extracted from the Government of Guinea situation reports [28].

For both the DRC and Guinea, temporal and spatial scales and ranges were chosen based on data

availability and all data which were available were utilized for the analysis.

Conflict data were exported from the Armed Conflict Location & Event Data Project's (ACLED,

Wisconsin, United States) data export tool [29]. ACLED was chosen over other data sources (e.g.,

Uppsala Data Conflict Program [30]), as it contained more detailed categorisation of conflict event

type, which were fundamental to the aims of this work. For each country, data were available from

1997 to the present, with each data entry point equating to a reported conflict. Conflicts were

reported on a daily scale to longitude and latitude (in degrees), and by administrative level 1, 2, 3

and 4  (Table 1). The data were categorised by conflict event type (riots, protest, battles,

explosions/remote violence, strategic developments and violence against civilians) and sub-event

type and provided information on number of conflict fatalities (definitions of the event types,

according to ACLED, are provided in Additional file 1).

Due to the granularity of the different datasets and for compatibility with the model (a single

number was needed to define the time of each EVD case/conflict, see Model Structure and Fitting

and Additional file 2), the temporal scale was set to continuous weeks from 01 January 2018 to 30

June 2021 (183 continuous weeks). Week number was not restarted at the beginning of the

calendar year, to account for outbreaks/conflict which endured over a December to January period.

Administrative level 1 for Guinea and administrative level 2 for the DRC were selected as the

spatial granularity for model fitting (Table 1), due to data availability and discrepancies in finer

scale location names among datasets. Weeks rather than continuous days were chosen, due to

the potential for reporting lags in both the EVD and conflict datasets and due to the incubation

period of Ebola virus (>1 week).

*Model Structure and Fitting*

We used the Self-Controlled Case Series (SCCS) methodology which investigates the association

between an exposure and an outcome event. The method aims to estimate the effect by

comparing the relative incidence of the adverse events (EVD case) within an exposure period of

hypothesised excess risk (conflict), compared to all other times (peace, according to the dataset

used). The method is a case only method, meaning cases are the only subjects used to test the

hypothesised effect. The method has the advantage of not needing separate controls (e.g., those

not infected), by automatically controlling for fixed confounders that remain constant over the

observational period. Cases are treated as individuals and then stratified in the modelling approach

to control for the confounders [31]. The method uses conditional logistic regression (R function

clogit(), package "survival" [32]) to estimate the effect of an exposure (conflict event) on an

outcome event (EVD case, either suspected or confirmed), compared to all other times (peace) in

an observation period (0-183 weeks). Maximum likelihood estimations were used, based on an

assumed Poisson probability distribution [33,34]. All models were fit using R Studio with R version

4.3.0.

Both the exposure and event were transformed to binary outcomes, either being reported in the

data in a specific week and administrative unit (1), or not (0), assuming that if a conflict or EVD

case occurred, then it would have been correctly identified and reported. For conflict, the definition

of a conflict event was the same for both countries. As the method is case-only, both a conflict

event and EVD case had to be reported in the same administrative level (administrative unit 1 for

Guinea and administrative unit 2 for the DRC, see Table 1) to be included.

To control for the fixed confounders, each exposure and event (in the same administrative unit)

was allocated a unique number and used as a stratified term in the model. The event was then

assigned a pre-exposure (start of observation period to exposure), exposure (exposure week) and

post-exposure (exposure to end of observation period) period (see Additional file 2 for fitted data

setup). The intervals between the three periods were logarithmically transformed and used as an

offset term in the model. Offsetting the interval accounts for the possibility that a longer interval

would result in a greater opportunity for the event (EVD case) to occur by chance, not due to its

association with conflict (or peace).

The outcome event data consist of values $n_{ijk}$, which take the values 1 or 0 corresponding

respectively to whether an EVD case was reported or not, for individual $i = 1, \dots N$, in the $j$th week,

$j = 1, \dots, J$, with conflict exposure $k = 1,2$, corresponding to peace or conflict. The length of time

spent in a particular interval $ijk$ is given by $e_{ijk}$. The likelihood, $L$, is a function of $\beta_k$ which are the

coefficients for conflict state $k$, and $\alpha_j$ the coefficients for week $j$. The total likelihood is given by the

following,

$$L = \prod_{i=1}^{N} \prod_{j=1,\dots,J;k=1,2}^{\square} \left( \frac{\exp(\alpha_j + \beta_k)e_{ijk}}{\sum_{j,k}^{\square} \exp(\alpha_j + \beta_k)e_{ijk}} \right)^{n_{ijk}}.$$

In standard survival analysis, $e_{ijk}$ can be thought of as an offset term. Notice that non-time varying

coefficients would drop-out of this self-controlled model. This is equivalent to a product multinomial

likelihood. Hence, $\exp(\beta_2 - \beta_1)$ is the relative incidence for conflict, controlling for time ($\beta_0$ = peace,

based on the assumption that peace has no effect on EVD cases).

The model outcome variable was a log rate ratio, which was then exponentially transformed to

incidence rate ratio (IRR), with a 95% confidence interval. IRR is the ratio of the incidence rates for

individuals in a population during times of conflict exposure compared to times of peace (according

to the dataset). The incidence rate is the rate at which incidence increases or decreases in a

specific person-time (administrative level-week). IRR values of 1 indicate similar incidence rates in

the conflict exposure period compared to the non-conflict exposure period, whereas an IRR > 1

suggests an increased risk and IRR < 1 indicates a decreased risk. Therefore, IRR quantifies the

magnitude of the association in which conflict increased/decreased the risk of EVD transmission.

IRR values were considered statistically significant at $p < 0.05$.

*By Sub-National Unit, Event Type and Lag*

The model was fit several times to evaluate the different research questions above, either by sub-

setting the data, or altering the way in which the exposure period was defined. The datasets were

subset by the sub-national administrative level of the fitted data (see *Datasets* and Table 1) and

the analysis repeated, to understand if certain geographic regions saw a greater magnitude of

effect between Ebola and conflict. Similarly, conflict sub-event type (and conflict event type, results

in Additional file 3) was also subset and the analysis repeated, to evaluate if certain types of

conflict had a greater effect on EVD reported cases. Conflict event sub-type was chosen for the

main analysis, rather than event type, as the event type names are generally more ambiguous and

less descriptive, making the interpretation of the results more challenging.

The impact of lag was explored, based on the assumption that the effect of a conflict event on EVD

cases may not be immediate (EVD's incubation period is 8-10 days [3]) and may be long-lasting.

Different exposure periods were tested, extending the period from 1 week following the conflict to

2, 4, 6, 8 and 10 weeks following the exposure, moving beyond previous work on EVD that tested

the effect of conflict to 4 weeks [18]. However, a longer exposure period may result in a diluting

effect due to a greater chance of an event (EVD case) being reported in an exposure period

(conflict). To account for this, we tested the effect of a single week, 2, 4, 6, 8 and 10 weeks

following the week the exposure was reported, rather than extending the exposure period.

*Quantifying conflict severity and proximity*

To explore mechanisms through which the most impactful types of conflict (battles, protests and

riots) may impact EVD transmission, we quantified conflict severity using two metrics. First, we

presented conflict fatalities as a metric of severity (higher mortality = more severe conflict), both in

terms of raw numbers for each country and conflict event type and as a proportion of total conflict

fatalities. Second, we measured the proximity of all battles, protests and riots to major cities by the

shortest distance between the coordinates of the conflict and the city (measured in longitude and

latitude as degrees) in both the DRC [35] and Guinea [36]. We then compared the difference

between the average distance for each conflict event type, to the total average distance for all

event types, over the observation period of 2018 to 2022. The distance was measured using the

Haversine method, which assumes a spherical earth, ignoring ellipsoidal effects (R functions

distHaversine() and distm(), package "geosphere" [37]). We hypothesised that conflict types which

more often occurred near more people (e.g., closer to major cities), would be more severe, and

therefore have a bigger effect on disease transmission.

**Results**

*Datasets*

During the observation period, there were 12,237 EVD cases reported in 94 weeks (out of 183

weeks in the observation period) across 14 administrative 2 units for the DRC (all in Nord-Kivu,

Sud-Kivu, Ituri and Équateur). For Guinea, there were 18 weeks (out of 183) which reported 1,797

EVD cases in two administrative 1 units (N'Zérékoré and Conakry). These cases corresponded to

168 (DRC) and 86 (Guinea) weeks of reported conflict exposure in the dataset during the

observation period for the same corresponding administrative levels as the reported EVD cases. In

the DRC, most EVD cases were reported in late 2018 to late 2019, while for Guinea, most cases

were reported in 2021 (see Fig. 1).

**Fig. 1** Weeks which reported **A,** Ebola virus disease (EVD) cases and **B,** conflict exposure for the

Democratic Republic of the Congo (DRC) and Guinea (GUI).

In the DRC, EVD cases for the 2018-2020 outbreak were commonly reported in areas of high

conflict frequency, namely the Kivu provinces. Similar results were found for Guinea, with all EVD

cases being reported in the two regions that reported the highest conflict event frequency (Conakry

and N'Zérékoré), but with overall lower levels of reported conflict compared to the DRC. The most

frequent conflict sub-event types (event types) in the fitted data varied by country. For the DRC,

these included armed clashes (battles), attacks (violence against civilians) and peaceful protests

(protests). In Guinea, the most reported conflict sub-event types (event types) were violent

demonstrations (riots), peaceful protests (protests) and mob violence (riots) (Table 2).

**Table 2** Frequency of conflict event types & sub-event types in the data fitted to the model for the

Democratic Republic of the Congo (DRC) and Guinea.

| Country | Event Type | Sub-Event Type | Frequency |
|---------|-----------|----------------|-----------|
| Guinea | Riots | Violent demonstration | 108 |
| DRC | Battles | Armed clash | 106 |
| DRC | Violence against civilians | Attack | 74 |
| Guinea | Protests | Peaceful protest | 72 |
| Guinea | Riots | Mob violence | 27 |
| Guinea | Violence against civilians | Attack | 27 |
| DRC | Protests | Peaceful protest | 22 |
| Guinea | Protests | Protest with intervention | 20 |
| DRC | Violence against civilians | Abduction/forced disappearance | 15 |
| Guinea | Battles | Armed clash | 11 |

| Guinea | Strategic developments | Looting/property destruction | 11 |
|---|---|---|---|
| DRC | Strategic developments | Looting/property destruction | 8 |
| Guinea | Strategic developments | Arrests | 6 |
| DRC | Battles | Government regains territory | 6 |
| DRC | Protests | Protest with intervention | 6 |
| DRC | Riots | Mob violence | 6 |
| Guinea | Protests | Excessive force against protesters | 5 |
| DRC | Riots | Violent demonstration | 5 |
| DRC | Strategic developments | Change to group/activity | 4 |
| DRC | Strategic developments | Agreement | 3 |
| DRC | Strategic developments | Arrests | 3 |
| DRC | Violence against civilians | Sexual violence | 3 |
| Guinea | Strategic developments | Other | 2 |
| DRC | Battles | Non-state actor overtakes territory | 2 |
| DRC | Strategic developments | Headquarters or base established | 2 |
| Guinea | Strategic developments | Disrupted weapons use | 1 |
| Guinea | Violence against civilians | Abduction/forced disappearance | 1 |
| Guinea | Violence against civilians | Sexual violence | 1 |
| DRC | Explosions/Remote violence | Remote explosive/landmine/IED | 1 |
| DRC | Protests | Excessive force against protesters | 1 |
| DRC | Strategic developments | Disrupted weapons use | 1 |
| DRC | Strategic developments | Non-violent transfer of territory | 1 |
| DRC | Strategic developments | Other | 1 |

*Model Output*

Conflict was significantly associated with an increased risk of reported EVD cases in both the DRC

and Guinea between 2018 and 2021. The magnitude of effect was similar for both countries, with

conflict increasing the risk of EVD by 1.88 (1.76-2.02 95%Confidence Interval (CI)) times in the

DRC, and 1.98 (1.11-3.51 95%CI) times in Guinea.

*By Sub-National Unit*

The effect of conflict on EVD varied sub-nationally in terms of magnitude and uncertainty (Fig. 2).

In the first week of the reported conflict exposure, the administrative units with the highest reported

effect were Mbandaka (Équateur, DRC), Mwenga (Sud-Kivu, DRC), Goma (Nord-Kivu, DRC),

Butembo (Nord-Kivu, DRC) and N'Zérékoré (Guinea) with IRR values of 4.26 (0.58-31.2 95%CI),

2.49 (1.87-3.31 95%CI), 2.19 (1.69-2.83 95%CI), 2.12 (1.76-2.57 95%CI) and 2.06 (1.13-3.77

95%CI), respectively. Some areas did not present significant results, often having higher

uncertainty, including Aru, Bolomba, Ingende, Mbandaka in the DRC and Conakry in Guinea.

**Fig. 2** Incidence rate ratio (IRR) for the effect of conflict event exposure on EVD cases for the

Democratic Republic of the Congo (DRC) and Guinea (GUI) during the week the conflict was

reported at sub-national administrative unit.

*By Event Type*

The greatest impact for conflict event sub-type was armed clashes in Guinea (IRR = 3.45, 0.80-

14.8 95%CI) (Fig. 3); however, this was not statistically significant, along with 12 other conflict sub-

types in Guinea (abduction/forced disappearances, arrests, attacks, change to group/activity,

disrupted weapons use, excessive force against protesters, looting/property destruction, mob

violence, other, peaceful protests, protests with intervention and sexual violence). Violent

demonstrations had the greatest significant impact on EVD cases in Guinea causing a 2.38 (1.04-

5.41 95%CI) times increased risk of EVD.

Only three conflict event sub-types were not significantly associated with EVD incidence in the

DRC: air/drone strikes, disrupted weapons use and headquarters or base establishments (mainly

due to very small sample sizes). For the conflict sub-types that did have a significant effect, shelling/artillery/missile attack had the biggest effect on EVD cases at 3.44 (1.27-9.3195%CI) times increased risk, followed by peaceful protests (IRR = 3.11, 2.66-3.64 95%CI), excessive force against protesters (2.83, 1.60-5.03 95%CI), looting/property destruction (2.69, 2.18-3.31 95%CI), protests with interventions (2.68, 2.05-3.52 95%CI) and violent demonstrations (2.58, 2.00-3.34 95%CI) (Fig. 3). For the results regarding conflict event type (rather than sub-event types), see Additional file 3.

**Fig. 3** Incidence rate ratio (IRR) for the effect of conflict event exposure on EVD cases for the Democratic Republic of the Congo (DRC) and Guinea (GUI) during the week the conflict was reported by conflict sub-event type.

*By Lag*

All lag periods were found to be statistically significant for both countries. Extending the lag periods caused the effect of conflict on EVD cases to be almost eliminated (IRR = 1) by 10 weeks in the DRC, with an IRR of 1.05 (1.02-1.09 95%CI). In Guinea, the risk of EVD cases decreased to 1.41 (0.99-2.01 95%CI) times at 4 weeks and then increased to 1.72 (1.38-2.14, 95%CI) by 10 weeks. The uncertainty was far greater in Guinea with slightly less significant results (according to p values) compared to the DRC (see Fig. 4).

**Fig. 4** Incidence rate ratio (IRR) for the effect of conflict event exposure on EVD cases for the Democratic Republic of the Congo (DRC) and Guinea (GUI) from the week the conflict was reported to 10 weeks following the exposure, by extending the exposure periods.

The lagged exposure periods for a single week 2, 4, 6, 8 and 10 weeks following the exposure resulted in conflict having a greater and more statistically significant (according to p values) impact on EVD cases (Fig. 5), compared to extending the periods (from 1 week to 10 weeks). All lag periods saw an increased risk from week 1 to week 10, with Guinea having a greater effect

(although with wider ranging uncertainty), potentially due to the increase already found in Guinea

at week 4 with the extended exposure periods (Fig. 4). The risk of EVD cases increased from 1.88

(1.76-2.02 95%CI) to 1.96 (1.83-2.10 95%CI) times in the DRC and from 1.96 (1.11-3.51 95%CI) to

3.30 (2.12-5.13 95%CI) times in Guinea by the 10th week following the event.

**Fig. 5** Incidence rate ratio (IRR) for the effect of conflict event exposure on EVD cases for the

Democratic Republic of the Congo (DRC) and Guinea (GUI) from the week the conflict was

reported to the 10th week following the conflict report, as a single week.

*Quantifying conflict severity and proximity*

Table 3 shows the full results for the conflict fatalities and proximity analysis for each of the three

most significant event types (battles, riots and protests). In the DRC, battles caused 56% of all

conflict fatalities in the fitted data here (n = 7,900), compared to 0.19% (n = 27) and 3.06% (n =

430) for protests and riots, respectively. In Guinea, 64% (n = 189) of conflict fatalities were caused

by riots, followed by 17.2% (n = 51) in battles and 7.12% (n = 21) in protests.

On average, over the four-year period, battles were further away from major cities in both

countries, followed by protests and riots in the DRC and riots and then protests in Guinea (for

maps, see Additional file 4: Fig. S2). Compared to the average distance between all conflict events

and major settlements, battles were 6.33km further away in the DRC and 17.2km further away in

Guinea. In the DRC, protests were 2.99km closer and riots 3.33km closer, compared to the

average distance. For Guinea, protests and riots were 11.4km and 5.73km closer, respectively.

**Table 3** Battles, riots and protests in terms of EVD risks by number of conflict fatalities, the

percentage of total fatalities and the difference in average proximity from conflict events to major

settlements (event type average distance minus all conflicts average distance) for the full four-year

observation period for each country.

| Country | Event Type | Fatalities Frequency | Fatality Percentage | Difference in proximity |
|---------|-----------|---------------------|---------------------|------------------------|
| DRC | Battles | 7,900 | 56.2% | 6.33km |
| DRC | Riots | 430 | 3.05% | -3.33km |
| DRC | Protests | 27 | 0.19% | -2.99km |
| Guinea | Battles | 51 | 17.3% | 17.2km |
| Guinea | Riots | 189 | 64.1% | -5.73km |
| Guinea | Protests | 21 | 7.11% | -11.4km |

**Discussion**

During recent EVD outbreaks in 2018 to 2021, we quantified the risk of conflict event exposure and

found an increased risk at a national level by a similar magnitude in the DRC and Guinea, at 1.88

and 1.98 times, respectively. The uncertainty in the IRR values for Guinea throughout all the

analyses was much wider than the DRC. In the distribution of the exposures and events data fit to

the model, there was overlap with areas of high reported conflict and EVD case frequency. Several

of the regions/territories analysed had higher IRR values than the national risk: Mbandaka,

Mwenga, Goma, Butembo (all in the DRC) and N'Zérékoré (in Guinea), which ranged from 2.06 to

4.26.

We evaluated if certain types of conflict may be more impactful than others, to contribute to our

understanding of how conflict can change disease risk. The conflict sub-event types with the

greatest impacts in terms of EVD risk were armed clashes in Guinea and shelling/artillery/missile

attack in the DRC. However, both armed clashes and shelling/artillery/missile attacks had the

largest uncertainty. Violent demonstrations had the strongest and most significant impact in

Guinea, while in the DRC, peaceful protests, excessive force against protesters, looting/property

destruction, protests with interventions and violent demonstrations had the largest, most significant

impact, with little uncertainty.

The cornerstone of EVD control is through identifying cases and reducing contact with infected

bodily fluids (and vaccination) [8,10]. Conflict can impact these control measures in multiple ways.

In previous EVD outbreaks, weak health systems (due to access, supplies, safety of staff and

patients and fear), complex socio-cultural and political environments (creating mistrust in

governments and NGOs) and ineffective messaging and community engagement have all created

problems in EVD response [20,38]. Additionally, there have been difficulties in administering EVD

vaccines due to damaged transportation and infrastructure, although continued efforts aim to reach

those most at risk [11,16,39]. Conflict and EVD can create a positive feedback loop, and it has

been previously observed that conflict increases more during EVD outbreaks compared to controls

[18,19]. A suggested explanation is that EVD outbreaks may further destabilise fragile regions and

fuel conflicts. For example, mistrust increased in the DRC during the 2018-2019 EVD outbreak due

to the government's decision to ban populations in the affected provinces from participating in the

presidential election [18,19].

A potential explanation for the large number of protest-related impacts on EVD cases found here,

may be due to these events being more disruptive to a greater number of people. Protests rely on

large congregations of people, increasing contact among individuals and therefore fueling EVD

transmission. Furthermore, protests and demonstrations may be more likely to impact

transportation, education and healthcare, due to the widespread disruption they can cause in

dense urban areas [40,41]. Disruption of healthcare and transportation may be particularly

damaging in controlling EVD, as it could complicate contact tracing and delay testing, diagnosis

and treatment. Further evidence for this explanation may be that within these very impactful sub-

events, looting/property destruction was also found highly impactful and important, which is often

the product of protests and demonstrations and results in damage to healthcare infrastructure and

supplies.

During 2018 to 2021, battles were the furthest away from major settlements in both countries,

compared to protests and riots. Proximity may impact the effect of the conflict on the population,

both in terms of the number of people affected and the magnitude of the effect. Proximity may also

add further evidence to the above hypothesis that protests and demonstrations, although

potentially less violent, may be more impactful than battles and armed clashes. Battles (in terms of

the ACLED's definition [29]) are more likely to be fought on pre-defined battle "grounds", which are

less likely to be close to settlements. Alternatively, these settlements, including their healthcare

and testing facilities are quickly displaced and abandoned following the onset of conflict due to the

immediate threat to life [42]. The absence of testing facilities may result in cases not being

identified and reported, therefore making battles appear less impactful to EVD transmission.

A potential explanation for why remote violence/explosions and battles were very impactful but

often with higher uncertainty is that these types of conflict occurred less frequently, but when they

did, they had a very large impact on EVD cases. The fewer events therefore resulted in the large

uncertainty and more data (if available) may help to understand if the larger impact of these conflict

types is consistent through time. Additionally, peoples' behaviour and risks must be considered

during more "extreme" forms of violence such as battles and explosions, compared to protests and

demonstrations. Violent conflict may have a very large initial impact (accounting for the high IRR

values), but then the risk stabilises in the area as civilians take precautions (e.g., displacement) to

protect their life [43].

Alternatively, there is the potential for more violent forms of conflict (such as battles and

explosions) to decrease the risk of EVD. People may be more likely to stay at home during these

events or be subject to curfews and not mix in public spaces, decreasing their exposure to the

pathogen [44]. Furthermore, severe conflict may be more fatal (of both civilians and combatants),

reducing the susceptible pool for EVD cases. More work is needed to understand how to quantity

conflict severity for scientific research purposes, which could be used to identify dose-response

relationships with disease, while still considering that conflicts impact individuals differently in terms

of disruption and severity.

When extending the exposure periods, the effect of conflict on EVD cases was almost eliminated

by 10 weeks in the DRC, while in Guinea, the risk of EVD cases decreased to 4 weeks and then

increased by 10 weeks. However, it appears that extending the lag periods may have been diluting

the effect of conflict on EVD cases, as changing the lag periods to a single week made all the lag

periods for both countries more significant and impactful. These results track with previous

research which found synchronicity and correlation between the reporting of EVD cases and

conflict in the DRC and suggest that the effect of conflict may be long-lasting [11,19].

The results show that conflict is still highly influential multiple months following the event, and in

some instances, conflict had a greater and more significant effect on EVD with time. A potential

explanation may be due to the disruption caused by the conflict event having a cumulative effect.

Previous findings suggest that conflict lengthens the EVD treatment process, e.g., longer times to

detection, isolation, treatment and vaccination, creating a lasting effect and prolonging outbreaks

[18]. Therefore, time would be needed for EVD cases to increase and be reported, after elevated

transmission.

There are several limitations to the datasets used here, the first being the assumption that if EVD

or conflict occurs, it is correctly identified (e.g., confirmed and suspected EVD cases were

included) and reported on the correct day with no delay, where the transmission event/symptom

onset occurred. These assumptions are unlikely, as EVD cases may be missed if cases never

seek formal medical assistance or are in hard-to-reach areas. Furthermore, the categorisation of

conflict events is subjective (category definitions were included in Additional file 1) and several

categories are ambiguous. Data availability was limited and sub-setting the data (by sub-national

unit or conflict sub-event type) led to small sample sizes in some instances, resulting in wide

uncertainty, statistical insignificance or the models not being able to fit, particularly in Guinea.

However, generally the findings were replicated in both countries, in terms of sub-event type, the

effects of lag and geographic distribution of risk.

The methodology accounts for confounders by fixing them across the administrative unit. It is not

as effective across multiple administrative units. Sub-national units are different geographical

sizes, with different population sizes and risk factors. Therefore, comparing different

regions/territories can create misleading results. Additionally, the use of vaccination during these

more recent outbreaks [45,46], makes comparison to previous work challenging. While data on

Ebola vaccination are available, the use of ring vaccination makes it difficult to incorporate

vaccination data into the model used here (as a binary outcome) [38,46]. For example, the sub-

national region with the largest IRR here was Équateur, part of the DRC which has received

relatively little vaccination, compared to the eastern provinces [38].

**Conclusions**

The research presented here highlights the increased risk that conflict has had on EVD cases in

the DRC and Guinea in recent resurgences. The risks of conflict on EVD was spatially

heterogenous and appears to last for several months and may be longer-lasting. Protests and riot-

related conflict event types were highly influential, potentially being more disruptive to more people,

compared to more violent forms of conflict e.g., battles. More research to understand how best to

quantify conflict severity and intensity could provide further evidence for these mechanisms.

Extra vigilance is needed following protest- and riot-related conflict events for EVD transmission.

Continued efforts are needed to try and account for the long-term effect of conflict on EVD and

other infectious diseases. Effective public health messaging and scaling up testing in urban areas

may help to mitigate these risks, including information on how to reduce transmission in dense

urban areas and early clinical symptoms to be aware of. Ebola treatment centres should be

prepared for an increase in cases following conflict events for multiple months, with sufficient

vaccination stocks made available to areas most at risk.

**List of abbreviations**

ACLED - Armed Conflict Location & Event Data Project

CI – Confidence Interval

DRC – Democratic Republic of the Congo

EVD – Ebola Virus Disease

GUI – Guinea

HEMA - Humanitarian Emergency Response Africa

IRR – Incidence Rate Ratio

MoH – Ministry of Health

NGOs – Non-Governmental Organisations

SCCS – Self-Controlled Case Series

WASH – Water, Sanitation and Hygiene

**Declarations**

*Ethics approval and consent to participate*

Although the work presented here uses human data, all the data used are from public available

data sources, which are none identifiable and therefore ethical approval was waived for this

analysis.

*Consent for publication*

Not applicable

*Availability of data and materials*

The datasets analysed during this study are all publicly available and references throughout the

manuscript. Further code related to the methods and data visualization here are available at:

https://github.com/GinaCharnley/ebola_conflict

*Competing interests*

The authors declare that they have no competing interests.

*Funding*

The work presented here received no specific funding, however we acknowledge joint Centre funding from the UK Medical Research Council and Department for International Development [MR/R0156600/1]. The funder played no role in the design of the study and collection of data, the analysis, and interpretation of data and in writing the manuscript.

*Authors' contributions*

GECC conceptualised and designed the study, collated the data sources, analysed the data, wrote the manuscript and incorporated feedback. IK contributed to interpreting the findings, provided expertise on conflicts and global health and revised several drafts of the manuscript. NG provided statistical expertise on the methodology, assisted in writing the manuscript, analysing the data and revised several drafts. EBM contributed to interpreting the findings, provided expertise on conflicts and Ebola in the DRC and revised several drafts of the manuscript. KAMG co-conceptualised and designed the study, provided data sources, provided expertise on epidemiological methods, contributed to interpreting the findings and revised several drafts of the manuscript. All authors read and approved the final manuscript.

*Acknowledgements*

We would like to acknowledge and thank those working in the field who collected and collated the data sources used here, namely the Ministère de la Santé Publique de la République Démocratique du Congo and Ministère de la Santé et de l'Hygiène Publique de Guinée. Additionally, we would like to thank Sophie Meakin for collating the situation reports used in some of the DRC data used here.

**References**

1. Pattyn SW, Jacob W, Van der Groen G, Piot P, Courteille G. Isolation of Marburg-like virus from a case of haemorrhagic fever in Zaire. Lancet. 1977:573-4.

2.  Bowen ET, Platt GS, Lloyd G, Baskerville A, Harris WJ, Vella EE. Viral haemorrhagic fever

in southern Sudan and northern Zaire. Preliminary studies on the aetiological agent. Lancet.

1977:571-3.

3.  Jacob ST, Crozier I, Fischer WA, Hewlett A, Kraft CS, Vega MA, et al. Ebola virus disease.

Nat. Rev. Dis. Primers. 2020 Feb 20;6(1):13.

4.  World Health Organization. Ebola virus disease. https://www.who.int/news-room/fact-

sheets/detail/ebola-virus-disease (2021).

5.  Pourrut X, Kumulungui B, Wittmann T, Moussavou G, Délicat A, Yaba P, et al. The natural

history of Ebola virus in Africa. Microbes Infect. 2005;7(7-8):1005-14.

6.  Centres for Disease Control and Prevention (CDC). History of Ebola Disease Outbreaks.

https://www.cdc.gov/vhf/ebola/history/chronology.html (2022)

7.  Adepoju P. Lessons from Ebola as DRC grapples with conflict, measles, and covid-19. BMJ.

2020 Jul 21;370.

8.  Mukadi-Bamuleka D, Mambu-Mbika F, De Weggheleire A, Edidi-Atani F, Bulabula-Penge J,

Mfumu MMK, et al. Efficiency of Field Laboratories for Ebola Virus Disease Outbreak during

Chronic Insecurity, Eastern Democratic Republic of the Congo, 2018-2020. Emerg. Infect.

Dis. 2023;29(1):1-9.

9.  Kinganda-Lusamaki E, Black A, Mukadi DB, Hadfield J, Mbala-Kingebeni P, Pratt CB, et al.

Integration of genomic sequencing into the response to the Ebola virus outbreak in Nord

Kivu, Democratic Republic of the Congo. Nat. Med. 2021;27(4):710-716.

10. Mukadi-Bamuleka D, Sanogo YO, Bulabula-Penge J, Morales-Betoulle ME, Fillon P,

Woodruff P, et al. RDT Working Group. Postmortem Surveillance for Ebola Virus Using

OraQuick Ebola Rapid Diagnostic Tests, Eastern Democratic Republic of the Congo, 2019-

2020. Emerg. Infect. Dis. 2022;28(2):420-424.

11. Wells CR, Pandey A, Ndeffo Mbah ML, Gaüzère BA, Malvy D, Singer BH, et al. The

exacerbation of Ebola outbreaks by conflict in the Democratic Republic of the Congo. PNAS.

2019;116(48):24366-72.

12. Mbala-Kingebeni P, Aziza A, Di Paola N, Wiley MR, Makiala-Mandanda S, Caviness K, et

al. Medical countermeasures during the 2018 Ebola virus disease outbreak in the North Kivu

and Ituri Provinces of the Democratic Republic of the Congo: a rapid genomic assessment.

Lancet Infect, Dis. 2019;19(6):648-657.

13. Wise PH, Barry M. Civil war & the global threat of pandemics. Daedalus. 2017;146(4):71-84.

14. Gonzalez-Torres A, Esposito E. Epidemics and conflict: Evidence from the Ebola outbreak

in Western Africa. Available at SSRN 3544606. 2016.

15. van Stekelenburg BC, De Cauwer H, Barten DG, Mortelmans LJ. Attacks on health care

workers in historical pandemics and COVID-19. Disaster Med. Public Health Prep.

2023;17:e309.

16. Oppenheim B, Lidow N, Ayscue P, Saylors K, Mbala P, Kumakamba C, et al. Knowledge

and beliefs about Ebola virus in a conflict-affected area: Early evidence from the North Kivu

outbreak. J. Glob, Health. 2019 Dec;9(2).

17. Mukadi-Bamuleka D, Nkuba-Ndaye A, Mbala-Kingebeni P, Ahuka-Mundeke S, Muyembe-

Tamfum JJ. Impact of ebola epidemics on the daily operation of existing systems in eastern

Democratic Republic of the Congo: A brief review. J Med Econ. 2024:1-5.

18. Vinck P, Pham PN, Bindu KK, Bedford J, Nilles EJ. Institutional trust and misinformation in

the response to the 2018–19 Ebola outbreak in North Kivu, DR Congo: a population-based

survey. Lancet Infect. Dis. 2019;19(5):529-36.

19. Kraemer MU, Pigott DM, Hill SC, Vanderslott S, Reiner RC, Stasse S, et al. Dynamics of

conflict during the Ebola outbreak in the Democratic Republic of the Congo 2018–2019.

BMC Med. 2020;18(1):1-0.

20. McPake B, Witter S, Ssali S, Wurie H, Namakula J, Ssengooba F. Ebola in the context of

conflict affected states and health systems: case studies of Northern Uganda and Sierra

Leone. Confl. Health. 2015;9(1):1-9.

21. Global Conflict Tracker. Instability in the Democratic Republic of Congo.

https://www.cfr.org/global-conflict-tracker/conflict/violence-democratic-republic-congo

(2023).

22. Human Rights Watch. World Report 2023.

https://www.hrw.org/sites/default/files/media_2023/01/World_Report_2023_WEBSPREADS

_0.pdf (2023).

23. Human Rights Watch. Guinea Events of 2020. https://www.hrw.org/world-

report/2021/country-chapters/guinea (2020).

24. Charnley GEC, Jean K, Kelman I, Gaythorpe KAM, Murray KA. Association between

Conflict and Cholera in Nigeria and the Democratic Republic of the Congo. Emerg. Infect.

Dis. 2022;28(12):2472-81.

25. Meakin S. Data for the 2018 outbreak of Ebola in the Democratic Republic of the Congo in

North-Kivu and Ituri Provinces. https://github.com/sophiemeakin/ebola_drc (2019).

26. Ministère de la Santé de la République Démocratique du Congo. Mailing list.

https://us13.campaign-

archive.com/home/?u=89e5755d2cca4840b1af93176&id=aedd23c530 (2019).

27. Human Emergency Response Africa (HEMA). Democratic Republic of the Congo: Ebola

(2021) Subnational cases. https://data.humdata.org/dataset/democratic-republic-of-congo-

ebola-2021-subnational-cases (2021).

28. ACAPS. GUINEA EBOLA EVD 2021 SUBNATIONAL CASES, DEATHS,

HOSPITALISATIONS, CONTACT TRACING, AND VACCINATIONS.

https://www.acaps.org/guinea-ebola-evd-2021-subnational-cases-deaths-hospitalisations-

contact-tracing-and-vaccinations (2021).

29. Armed Conflict Location & Event Data Project (ACLED, Wisconsin, United States). Data

Export Tool. https://acleddata.com/data-export-tool/ (2023).

30. Eck K. In data we trust? A comparison of UCDP GED and ACLED conflict events datasets.

Coop. Confl. 2012;47(1):124-41.

31. Whitaker HJ, Paddy Farrington C, Spiessens B, Musonda P. Tutorial in biostatistics: the

self-controlled case series method. Stat. Med. 2006;25(10):1768-97.

32. Therneau T. A Package for Survival Analysis in R. R package version 3.5-5.

https://CRAN.R-project.org/package=survival (2023).

33. Mostofsky E, Coull BA, Mittleman MA. Analysis of observational self-matched data to

examine acute triggers of outcome events with abrupt onset. Epidemiol. (Cambridge,

Mass.). 2018;29(6):804.

34. Xu S, Zeng C, Newcomer S, Nelson J, Glanz J. Use of fixed effects models to analyze self-

controlled case series data in vaccine safety studies. J. Biom. Biostat. 2012:006.

35. UN Office for the Coordination of Humanitarian Affairs - Field Information Services Section.

Democratic Republic of the Congo - major cities.

https://data.humdata.org/dataset/democratic-republic-of-the-congo-major-cities (2019).

36. Simplemaps. Guinea Cities Database. https://simplemaps.com/data/gn-cities (2023).

37. Hijmans RJ. geosphere: Spherical Trigonometry. R package version 1.5-18.

https://CRAN.R-project.org/package=geosphere (2022).

38. Bausch DG. The need for a new strategy for Ebola vaccination. Nat. Med. 2021;27(4):580-1.

39. Moran B. Fighting Ebola in conflict in the DR Congo. Lancet. 2018;392(10155):1295-6.

40. AL JAZEERA AND NEWS AGENCIES. Several killed as protests rock Goma in eastern

DRC. https://www.aljazeera.com/news/2021/12/21/several-killed-as-protests-rock-goma-in-

eastern-drc (2021).

41. AFP, AL JAZEERA. Protests against military administration paralyse Guinea capital.

https://www.aljazeera.com/news/2022/7/29/protests-against-military-administration-

paralyse-guinea-capital (2022).

42. Relief Web. Democratic Republic of the Congo faces acute health crisis amid rising

violence. https://reliefweb.int/report/democratic-republic-congo/democratic-republic-congo-

faces-acute-health-crisis-amid-rising-violence (2023).

43. Relief Web. After evacuating Kayna, DRC, MSF is trying to return to assist displaced in

Kanyabayonga. https://reliefweb.int/report/democratic-republic-congo/after-evacuating-

kayna-drc-msf-trying-return-assist-displaced (2004).

44. Radio France Internationale. Clashes erupt in Guinea Conakry after opposition call for anti-

junta protests. https://www.rfi.fr/en/africa/20220818-clashes-erupt-in-guinea-conakry-after-

opposition-call-for-anti-junta-protests (2022).

45. Maxmen A. Ebola vaccine approved for use in ongoing outbreak. Nature. 2017 May 30;10.

46. World Health Organization. Ebola virus disease: Vaccines. https://www.who.int/news-

room/questions-and-answers/item/ebola-vaccines (2020).