Quantifying the Risk of Conflict on Recent Ebola Outbreaks in Guinea and the Democratic Republic of the Congo

Gina E C Charnley[1,2,3*], Nathan Green[4], Ilan Kelman[3,5,6], Espoir B Malembaka[1,7], Katy A M Gaythorpe[2,8]

1. Bloomberg School of Public Health, Johns Hopkins University, Baltimore, United States

2. School of Public Health, Imperial College London, London, United Kingdom

3. Institute for Global Health, University College London, London, United Kingdom

4. Department of Statistical Science, University College London, London, United Kingdom

5. Institute for Disaster Risk Reduction, University College London, London, United Kingdom

6. University of Agder, Kristiansand, Norway

7. Centre for Tropical Diseases and Global Health (CTDGH), Université Catholique de Bukavu, Kukavu, Democratic Republic of the Congo

8. MRC Centre for Global Infectious Disease Analysis, Imperial College London, London, United Kingdom

* Corresponding author

Abstract

Background

Reducing Ebola virus transmission relies on the ability to identify cases and limit contact with

infected bodily fluids through biosecurity, safe sex practices, safe burial and vaccination. Armed

conflicts can complicate outbreak detection and interventions due to the widespread disruption

they cause to the government and the population. Guinea and the Democratic Republic of the

Congo (DRC) have historically reported the largest and the most recent Ebola virus outbreaks.

Understanding if conflict played a role in these outbreaks may help in identifying key risks factors

to improve disease control.

Methods

We used data from a range of publicly available data sources for both Ebola virus cases and

conflict events from 2018 to 2021 in Guinea and the DRC. We fitted these data to conditional

logistic regression models using the Self-Controlled Case Series methodology, to quantify the

magnitude in which conflict increased the risk of reported Ebola virus cases in terms of incidence

rate ratio. We re-ran our analysis sub-nationally, by conflict sub-event type and tested the effect of

lag.

Results

Conflict was significantly associated with an increased risk of reported Ebola virus cases in both

the DRC and Guinea in recent outbreaks. The effect was of a similar magnitude at 1.88 and 1.98

times increased risk for the DRC and Guinea, respectively. The greatest effects (often higher than

the national values) were found in the most conflict prone areas and during protest/riot-related

conflict events. Conflict was influential in terms of Ebola virus risk from 1 week following the event

and remained important and, in some cases, more so by 10 weeks.

Conclusion

Extra vigilance is needed following protests and riot-related conflict events in terms of Ebola virus

transmission. These events are highly disruptive, in terms of access to transportation and

healthcare and are often in urban areas with high population densities. Additional public health

messaging around these types of conflict events, relating to the risks and clinical symptoms may

be helpful in reducing transmission. Future work should aim to further understand and quantify

conflict severity and intensity, to evaluate dose-response relationships in terms of disease risk.

Keywords

Ebola virus, Conflict, Outbreaks, Guinea, the Democratic Republic of the Congo

Background

Ebola is a virus belonging to the *Filoviridae* family and was first identified in 1976 in what are now

the Democratic Republic of the Congo (DRC) and South Sudan [1,2]. Ebola virus disease (EVD)

causes acute haemorrhagic fever, leading to mortality rates of ~50%, in the absence of any

specific treatment. The virus is spread through human-to-human transmission via contact with

infected body fluids and fomites, with an average incubation period of 8 to 10 days [3,4].

Since its discovery, there have been several EVD outbreaks (with sporadic cases in Europe and

North America) [5]. The worst recorded outbreaks, in terms of total cases and deaths have been

the 2014-2016 West Africa outbreak, particularly in Sierra Leone, Liberia and Guinea and the

2018-2020 outbreak in the DRC. The most recent EVD outbreaks have been reported in Uganda

(*Sudan ebolavirus)*, Guinea (*Zaire ebolavirus)* and the DRC (*Zaire ebolavirus)*, all of which

occurred in 2021 and 2022 [6].

The control of EVD is through identifying cases and interventions which reduce the risk of contact

with infected people's body fluids, via biosecurity in EVD treatment centres, safe burial practices,

reducing sexual transmission [7], timely detection (via strategically positioned decentralized

laboratories) [8], contact tracing and post-mortem testing to identify EVD-related deaths, genomic

sequencing (to infer spatiotemporal transmission dynamics) [9,10] and more recently with the

rVSVΔG-ZEBOV-GP Ebola virus envelope glycoprotein vaccine and monoclonal antibody

therapeutics [11,12]. Conflict can impact these control measures, reducing the capacity to identify

and respond to outbreaks [13]. Furthermore, conflict can cause disruption to a range of basic

services including transport, healthcare and water, sanitation and hygiene (WASH) [14].

Targeted attacks on healthcare centres and workers are not limited to specific diseases or

geographic regions [15] and there have been several reported attacks on Ebola treatment centres,

creating fear and avoidance in accessing them [16]. To protect staff and patients, security forces

and military personnel are often required to carry out healthcare activities [15]. Additionally, fear

and mistrust can lead to further attacks, sabotage and disruption of healthcare facilities and

services. Distrust and dissatisfaction in the government, the military and international involvement

via non-governmental organisations (NGOs) has also reduced the willingness to follow public

health guidance and has increased vaccine hesitancy and misinformation [7,17].

In previous EVD outbreaks in the DRC and Guinea, conflict has been an ongoing issue to greater

or lesser extents [18,19]. The DRC is currently experiencing armed conflict in the eastern region,

namely the Kivu provinces, which has been ongoing since the 1990s due to instability within the

region, with many armed groups currently active along with governments' involvement [20,21].

Guinea has experienced conflict, albeit to a lesser extent than in the DRC, mainly due to politically

driven violence and demonstrations during post-election periods and excessive force by security

forces [22].

Due to Guinea and the DRC having both the largest recorded outbreaks and the most recent EVD

outbreaks [6], along with reported conflict, there is a need to further understand the mechanisms

and quantify the up-to-date conflict-related risks for recent EVD resurgences. Additionally, it is

essential to understand the timing of conflict-related risks for EVD transmission and any lagged

effects. Increasing our understanding of the magnitude and timing of risks can help EVD response

planning and identify when extra surveillance may be needed.

Despite evidence showing possible links and effects of conflict on EVD, research to date is limited

on magnitude, timings and mechanisms through which conflict impacts the risk of EVD, particularly

qualitatively. Here, we use a methodology that has proved robust in quantifying the relationship

between disease and conflict in previous studies [23] to address the following research aims:

1. Quantify the effect of conflict in terms of increasing the risk of reported EVD cases during

the 2018 to 2020 outbreak in the DRC and the 2021 outbreak in Guinea, both nationally and

sub-nationally.

2. Evaluate if certain conflict types were more important than others in impacting EVD cases.

3. Examine the timing of conflict-related risk on EVD, including lagged effects, from the onset

of the conflict report to the EVD case report.

4. Suggest mechanisms for any changes in risk relating to conflict type and EVD.

Methods

Datasets

EVD data were taken from multiple sources. For the DRC, data were collated from the DRC

Ministry of Health (MoH) mailing lists [24,25] and from the Humanitarian Emergency Response

Africa (HEMA) dataset [26]. The temporal scale of the data was daily and ranged from 04/04/2018

to 11/07/2020 for the MoH data and 06/02/2021 to 14/04/2021 for the HEMA data. The spatial

scale for the DRC data was to administrative level 2 (territory) for the DRC MoH data and to

administrative level 1 (province) for the HEMA dataset, including new reported case and death

counts. For Guinea, the data ranged from 14/02/2021 to 19/06/2021 and included new and

cumulative daily case counts to administrative level 1 (region), extracted from the Government of

Guinea situation reports [27]. For both the DRC and Guinea, temporal and spatial scales and

ranges were chosen based on data availability and all data which were available were utilized for

the analysis.

For conflict, data were exported from the Armed Conflict Location & Event Data Project's (ACLED)

data export tool [28]. ACLED was chosen over other data sources (e.g., Uppsala Data Conflict

Program [29]), as it contained more detailed categorisation of conflict event type, which were

fundamental to the aims of this work. For each country, data were available from 1997 to the

present, with each data entry point equating to a reported conflict. Conflicts were reported on a

daily temporal scale to longitude and latitude (in degrees), plus administrative level 1 (province in

DRC, region in Guinea), 2 (territory in DRC, prefecture in Guinea), 3 (sector in DRC, sub-

prefecture in Guinea) and 4 (settlement in both countries). The data were categorised by conflict

event type (riots, protest, battles, explosions/remote violence, strategic developments and violence

against civilians) and sub-event type and provide information on number of conflict fatalities

(definitions of the event types, according to ACLED, are provided in Additional file 1).

Due to the granularity of the different datasets and for compatibility with the model (a single

number was needed to define the time of each EVD case/conflict, see Model Structure and Fitting

and Additional file 2), the temporal scale was set to continuous weeks from 01/01/2018 to

30/06/2021 (183 continuous weeks). Week number was not restarted at the beginning of the

calendar year, to account for outbreaks/conflict which endured over a December to January period.

Administrative level 1 (region) for Guinea and administrative level 2 (territory) for the DRC were

selected as the spatial granularity for model fitting, due to data availability and discrepancies in

finer scale location names among datasets. Weeks rather than continuous days were chosen, due

to the potential for reporting lags in both the EVD and conflict datasets and due to the incubation

period of Ebola virus (>1 week).

Model Structure and Fitting

We used the Self-Controlled Case Series (SCCS) methodology which investigates the association

between an exposure and an outcome event. The method aims to estimate the effect by

comparing the relative incidence of the adverse events (EVD case) within an exposure period of

hypothesised excess risk (conflicts), compared to all other times (peace, according to the dataset

used). The method is a case only method and has the advantage of not needing separate controls,

by automatically controlling for fixed confounders that remain constant over the observational

period using stratification [30]. The method uses conditional logistic regression (R function clogit(),

package "survival" [31]) to estimate the effect of an exposure (conflict event) on an outcome event

(EVD case, either suspected or confirmed), compared to all other times (peace) in an observation

period (0-183 weeks). Maximum likelihood estimations were used, based on an assumed Poisson

probability distribution [32,33]. All models were fit using R Studio with R version 4.3.0.

Both the exposure and event were transformed to binary outcomes, either being reported in the

data in a specific week and administrative unit (1), or not (0), assuming that if a conflict or EVD

case occurred, then it would have been correctly identified and reported. For conflict, the definition

of a conflict event was the same for both countries. As the method is case-only, both a conflict

event and EVD case had to be reported in the same administrative level (region for Guinea and

territory for the DRC) to be included.

To control for the fixed confounders, each exposure and event (in the same administrative unit)

was allocated a unique number (indiv) and used as a stratified term in the model. The event was

then assigned a pre-exposure (start of observation period to exposure), exposure (exposure week)

and post-exposure (exposure to end of observation period) period (see Additional file 2 for fitted

data setup). The intervals between the three periods were logarithmically transformed and used as

an offset term in the model. Offsetting the interval accounts for the possibility that a longer interval

would result in a greater opportunity for the event (EVD case) to occur by chance, not due to its

association with conflict (or peace).

The outcome event data consist of values n_{ijk}, which take the values 1 or 0 corresponding

respectively to whether an EVD case was reported or not, for individual $i = 1, ... N,$ in the jth week,

$j = 1, ..., J,$ with conflict exposure $k = 1,2,$ corresponding to peace or conflict. The length of time

spent in a particular interval ijk is given by e_{ijk}. The likelihood, L, is a function of β_k which are the

coefficients for conflict state k, and α_j the coefficients for week j. The total likelihood is given by,

$$L = \prod_{i=1}^{N} \prod_{j=1,\dots,J; k=1,2} \left(\frac{\exp(\alpha_j + \beta_k)e_{ijk}}{\sum_{j,k}^{\square} \exp(\alpha_j + \beta_k)e_{ijk}} \right)^{n_{ijk}}.$$

In standard survival analysis, e_{ijk} can be thought of as an offset term. Notice that non-time varying

coefficients would drop-out of this self-controlled model. This is equivalent to a product multinomial

likelihood. Hence, $\exp(\beta_2 - \beta_1)$ is the relative incidence for conflict, controlling for time (β_0 = peace,

based on the assumption that peace has no effect on EVD cases).

The model outcome variable was a log rate ratio, which was then exponentially transformed to

incidence rate ratio (IRR), with a 95% confidence interval. IRR is the ratio of the incidence rates for

individuals in a population during times of conflict exposure compared to times of peace (according

to the dataset). The incidence rate is the rate at which incidence increases or decreases in a

specific person-time (administrative level-week). IRR values of 1 indicate similar incidence rates in

the exposure compared to the un-exposed, whereas an IRR > 1 suggests an increased risk and

IRR < 1 indicates a decreased risk. Therefore, IRR quantifies the magnitude of the association in

which conflict increased/decreased the risk of EVD cases. IRR values were considered statistically

significant at $p < 0.05$.

By Sub-National Unit, Event Type and Lag

The model was fit several times to evaluate the different research questions above, either by sub-

setting the data, or altering the way in which the exposure period was defined. The datasets were

subset to sub-national administrative level (regions for Guinea and territories for the DRC) and the

analysis repeated, to understand if certain geographic regions saw a greater magnitude of effect

between Ebola and conflict. Similarly, conflict sub-event type (and conflict event type, results in

Additional file 3) was also subset and the analysis repeated, to evaluate if certain types of conflict

had a greater effect on EVD reported cases. Conflict event sub-type was chosen for the main

analysis, rather than event type, as the event type names are generally more ambiguous and less

descriptive, making the interpretation of the results more challenging.

The impact of lag was explored, based on the assumption that the effect of a conflict event on EVD

cases may not be immediate (EVD's incubation period is 8-10 days) and may be long-lasting.

Different exposure periods were tested, extending the period from 1 week following the conflict to

2, 4, 6, 8 and 10 weeks following the exposure, moving beyond previous work on EVD that tested

the effect of conflict to 4 weeks [18]. However, a longer exposure period may result in a diluting

effect due to a greater chance of an event (EVD case) being reported in an exposure period

(conflict). To account for this, we tested the effect of a single week, 2, 4, 6, 8 and 10 weeks

following the week the exposure was reported, rather than extending the exposure period.

Quantifying conflict severity and proximity

To suggest mechanisms through which the most impactful types of conflict (battles, protests and

riots) may be impacting EVD transmission, we quantified conflict severity using two metrics. First,

we presented conflict fatalities, both in terms of raw numbers for each country and conflict event

type and as a proportion of total conflict fatalities. Second, we measured the proximity of all battles,

protests and riots to major cities by the shortest distance between the coordinates of the conflict

and the city (measured in longitude and latitude as degrees) in both the DRC [34] and Guinea [35].

We then compared the difference between the average distance for each conflict event type, to the

total average distance for all event types, over the observation period of 2018 to 2022. The

distance was measured using the Haversine method, which assumes a spherical earth, ignoring

ellipsoidal effects (R functions distHaversine() and distm(), package "geosphere" [36]).

Results

Datasets

During the observation period, there were 12,237 EVD cases reported in 94 weeks (out of 183

weeks in the observation period) across 14 administrative 2 units for the DRC (all in Nord-Kivu,

Sud-Kivu, Ituri and Équateur). For Guinea, there were 18 weeks (out of 183) which reported 1,797

EVD cases in two administrative 1 units (N'Zérékoré and Conakry). These cases corresponded to

168 (DRC) and 86 (Guinea) weeks of reported conflict exposure in the dataset during the

observation period for the same corresponding administrative levels as the reported EVD cases. In

the DRC, most EVD cases were reported in late 2018 to late 2019, while for Guinea, most cases

were reported in 2021 (see Fig. 1).

Fig. 1 Weeks which reported **A,** Ebola virus disease (EVD) cases and **B,** conflict exposure for the

Democratic Republic of the Congo (DRC) and Guinea (GUI).

In the DRC, EVD cases for the 2018-2020 outbreak were commonly reported in areas of high

reported conflict event frequency, namely the Kivu provinces. Similar results were found for

Guinea, with all EVD cases being reported in the two regions that reported the highest conflict

levels (Conakry and N'Zérékoré), but with overall lower levels of reported conflict. The most

frequent conflict sub-event types (event types) in the fitted data varied by country. For the DRC,

these included armed clashes (battles), attacks (violence against civilians) and peaceful protests

(protests). In Guinea, the most reported conflict sub-event types (event types) were violent

demonstrations (riots), peaceful protests (protests) and mob violence (riots) (Table 1).

Table 1 Frequency of conflict event types & sub-event types in the data fitted to the model for the

Democratic Republic of the Congo (DRC) and Guinea.

Country	Event Type	Sub-Event Type	Frequency
Guinea	Riots	Violent demonstration	108
DRC	Battles	Armed clash	106
DRC	Violence against civilians	Attack	74
Guinea	Protests	Peaceful protest	72
Guinea	Riots	Mob violence	27
Guinea	Violence against civilians	Attack	27
DRC	Protests	Peaceful protest	22
Guinea	Protests	Protest with intervention	20
DRC	Violence against civilians	Abduction/forced disappearance	15
Guinea	Battles	Armed clash	11
Guinea	Strategic developments	Looting/property destruction	11
DRC	Strategic developments	Looting/property destruction	8
Guinea	Strategic developments	Arrests	6
DRC	Battles	Government regains territory	6
DRC	Protests	Protest with intervention	6
DRC	Riots	Mob violence	6

Guinea	Protests	Excessive force against protesters	5
DRC	Riots	Violent demonstration	5
DRC	Strategic developments	Change to group/activity	4
DRC	Strategic developments	Agreement	3
DRC	Strategic developments	Arrests	3
DRC	Violence against civilians	Sexual violence	3
Guinea	Strategic developments	Other	2
DRC	Battles	Non-state actor overtakes territory	2
DRC	Strategic developments	Headquarters or base established	2
Guinea	Strategic developments	Disrupted weapons use	1
Guinea	Violence against civilians	Abduction/forced disappearance	1
Guinea	Violence against civilians	Sexual violence	1
DRC	Explosions/Remote violence	Remote explosive/landmine/IED	1
DRC	Protests	Excessive force against protesters	1
DRC	Strategic developments	Disrupted weapons use	1
DRC	Strategic developments	Non-violent transfer of territory	1
DRC	Strategic developments	Other	1

Model Output

Conflict was significantly associated with an increased risk of reported EVD cases in both the DRC

and Guinea between 2018 and 2021. The magnitude of effect was similar for both countries, with

conflict increasing the risk of EVD by 1.88 (1.76-2.02 95%Confidence Interval (CI)) times in the

DRC, and 1.98 (1.11-3.51 95%CI) times in Guinea.

By Sub-National Unit, Event Type and Lag

The effect of conflict on EVD varied sub-nationally in terms of magnitude and uncertainty (Fig. 2).

In the first week of the reported conflict exposure, the administrative units with the highest reported

effect were Mbandaka (Équateur, DRC), Mwenga (Sud-Kivu, DRC), Goma (Nord-Kivu, DRC),

Butembo (Nord-Kivu, DRC) and N'Zérékoré (Guinea) with IRR values of 4.26 (0.58-31.2 95%CI),

2.49 (1.87-3.31 95%CI), 2.19 (1.69-2.83 95%CI), 2.12 (1.76-2.57 95%CI) and 2.06 (1.13-3.77

95%CI), respectively. Some areas did not find significant results and often had high uncertainty,

including Aru, Bolomba, Ingende, Mbandaka in the DRC and Conakry in Guinea.

Fig. 2 Incidence rate ratio (IRR) for the effect of conflict event exposure on EVD cases for the

Democratic Republic of the Congo (DRC) and Guinea (GUI) during the week the conflict was

reported at sub-national administrative unit.

By Event Type

The greatest impact for conflict event sub-type was armed clashes in Guinea (IRR = 3.45, 0.80-

14.8 95%CI) (Fig. 3); however, this was not statistically significant, along with 12 other conflict sub-

types in Guinea (abduction/forced disappearances, arrests, attacks, change to group/activity,

disrupted weapons use, excessive force against protesters, looting/property destruction, mob

violence, other, peaceful protests, protests with intervention and sexual violence). Violent

demonstrations had the greatest significant impact on EVD cases in Guinea causing a 2.38 (1.04-

5.41 95%CI) times increased risk of EVD.

Only three conflict event sub-types were not significantly associated with EVD incidence in the

DRC: air/drone strikes, disrupted weapons use and headquarters or base establishments. For the

conflict sub-types that did have a significant effect, shelling/artillery/missile attack had the biggest

effect on EVD cases at 3.44 (1.27-9.3195%CI) times increased risk, followed by peaceful protests

(IRR = 3.11, 2.66-3.64 95%CI), excessive force against protesters (2.83, 1.60-5.03 95%CI),

looting/property destruction (2.69, 2.18-3.31 95%CI), protests with interventions (2.68, 2.05-3.52

95%CI) and violent demonstrations (2.58, 2.00-3.34 95%CI) (Fig. 3). For the results regarding

conflict event type (rather than sub-event types), see Additional file 3.

Fig. 3 Incidence rate ratio (IRR) for the effect of conflict event exposure on EVD cases for the

Democratic Republic of the Congo (DRC) and Guinea (GUI) during the week the conflict was

reported by conflict sub-event type.

By Lag

All lag periods were found to be statistically significant for both countries. Extending the lag periods

caused the effect of conflict on EVD cases to be almost eliminated (IRR = 1) by 10 weeks in the

DRC, with an IRR of 1.05 (1.02-1.09 95%CI). In Guinea, the risk of EVD cases decreased to 1.41

(0.99-2.01 95%CI) times at 4 weeks and then increased to 1.72 (1.38-2.14, 95%CI) by 10 weeks.

The uncertainty was far greater in Guinea with slightly less significant results (according to p

values) compared to the DRC (see Fig. 4).

Fig. 4 Incidence rate ratio (IRR) for the effect of conflict event exposure on EVD cases for the

Democratic Republic of the Congo (DRC) and Guinea (GUI) from the week the conflict was

reported to 10 weeks following the exposure, by extending the exposure periods.

The lagged exposure periods for a single week 2, 4, 6, 8 and 10 weeks following the exposure

resulted in conflict having a greater and more statistically significant (according to p values) impact

on EVD cases (Fig. 5), compared to extending the periods (from 1 week to 10 weeks). All lag

periods saw an increased risk from week 1 to week 10, with Guinea having a greater effect

(although with wider ranging uncertainty), potentially due to the increase already found in Guinea

at week 4 with the extended exposure periods (Fig. 4). The risk of EVD cases increased from 1.88

(1.76-2.02 95%CI) to 1.96 (1.83-2.10 95%CI) times in the DRC and from 1.96 (1.11-3.51 95%CI) to

3.30 (2.12-5.13 95%CI) times in Guinea by the 10th week following the event.

Fig. 5 Incidence rate ratio (IRR) for the effect of conflict event exposure on EVD cases for the

Democratic Republic of the Congo (DRC) and Guinea (GUI) from the week the conflict was

reported to the 10th week following the conflict report, as a single week.

Quantifying conflict severity and proximity

Table 2 shows the full results for the conflict fatalities and proximity analysis for each of the three

most significant event types (battles, riots and protests). In the DRC, battles caused 56% of all

conflict fatalities in the fitted data here (n = 7,900), compared to 0.19% (n = 27) and 3.06% (n =

430) for protests and riots, respectively. In Guinea, 64% (n = 189) of conflict fatalities were caused

by riots, followed by 17.2% (n = 51) in battles and 7.12% (n = 21) in protests.

317 On average, over the four-year period, battles were further away from major cities in both

318 countries, followed by protests and riots in the DRC and riots and then protests in Guinea (for

319 maps, see Additional file 4: Fig. S2). Compared to the average distance between all conflict events

320 and major settlements, battles were 6.33km further away in the DRC and 17.2km further away in

321 Guinea. In the DRC, protests were 2.99km closer and riots 3.33km closer, compared to the

322 average distance. For Guinea, protests and riots were 11.4km and 5.73km closer, respectively.

324 **Table 2** Battles, riots and protests in terms of EVD risks by number of conflict fatalities, the

325 percentage of total fatalities and the difference in average proximity from conflict events to major

326 settlements (event type average distance minus all conflicts average distance) for the full four-year

327 observation period for each country.

Country	Event Type	Fatalities Frequency	Fatality Percentage	Difference in proximity
DRC	Battles	7,900	56.2%	6.33km
DRC	Riots	430	3.05%	-3.33km
DRC	Protests	27	0.19%	-2.99km
Guinea	Battles	51	17.3%	17.2km
Guinea	Riots	189	64.1%	-5.73km
Guinea	Protests	21	7.11%	-11.4km

329 **Discussion**

330 During recent EVD outbreaks in 2018 to 2021, conflict event exposure increased the risk of

331 reported cases at a national level by a similar magnitude in the DRC and Guinea, at 1.88 and 1.98

332 times, respectively. The uncertainty in the IRR values for Guinea throughout all the analyses was

333 much wider than the DRC. In the distribution of exposure and events fit to the model, there was

334 overlap with areas of highest reported conflict and EVD case frequency. Several of the

335 regions/territories analysed had higher IRR values than the national risk: Mbandaka, Mwenga,

336 Goma, Butembo (all in the DRC) and N'Zérékoré (in Guinea), which ranged from 2.06 to 4.26.

Evaluating if certain types of conflict may be more impactful than others contributes to our

understanding of how conflict can impact disease. The conflict sub-event types with the greatest

impacts in terms of EVD risk were armed clashes in Guinea and shelling/artillery/missile attack in

the DRC. However, both armed clashes and shelling/artillery/missile attacks had the largest

uncertainty. Violent demonstrations had the strongest and most significant impact in Guinea, while

in the DRC, peaceful protests, excessive force against protesters, looting/property destruction,

protests with interventions and violent demonstrations had the largest, most significant impact, with

little uncertainty.

Conflict-related disruption

The cornerstone of EVD control is through identifying cases and reducing contact with infected

bodily fluids (and vaccination) [8,10]. Conflict can impact these control measures in multiple ways.

In previous EVD outbreaks, weak health systems (due to access, supplies, safety of staff and

patients and fear), complex socio-cultural and political environments (creating mistrust in

governments and NGOs), ineffective messaging and ineffective community engagement have all

created problems in EVD response [19,37]. Additionally, there have been difficulties in

administering EVD vaccines due to damaged transportation and infrastructure, although continued

efforts aim to reach those most at risk [11,16,38]. Conflict and EVD can create a positive feedback

loop, where EVD exacerbates the conflict and conflict exacerbates EVD. It has previously been

observed that conflict increases much more during EVD outbreaks compared to controls [17,18]. A

suggested explanation is that EVD outbreaks may further destabilise fragile regions and fuel

conflicts. For example, mistrust increased in the DRC during the 2018-2019 EVD outbreak due to

the government's decision to ban populations in the affected provinces from participating in the

presidential election [17,18].

A potential explanation for the large number of protest-related impacts on EVD cases found here,

may be the bigger impact these types of events would have on a greater number of people.

Protests rely on large congregations of people and crowds, increasing contact among individuals

and therefore fueling EVD transmission. Furthermore, protests and demonstrations may be more

likely to impact transportation, education and healthcare, due to the widespread disruption they can

cause in dense urban areas [39,40]. Disruption of healthcare and transportation may be particularly

damaging in controlling EVD, as it could complicate contact tracing and delay testing, diagnosis

and treatment. Further evidence for this explanation may be that within these very impactful sub-

events, looting/property destruction was also found highly impactful and important, which is often

the product of protests and demonstrations and results in damage to healthcare infrastructure and

supplies.

Proximity to the conflict

During 2018 to 2021, battles were the furthest away from major settlements in both countries,

compared to protests and riots. Proximity may impact the effect of the conflict on the population,

both in terms of the number of people affected and the magnitude of the effect. Proximity may also

add further evidence to the above hypothesis that protests and demonstrations, although

potentially less violent, may be more impactful than battles and armed clashes. Battles (in terms of

the ACLED's definition [28]) are more likely to be fought on pre-defined battle "grounds", which are

less likely to be close to settlements, or these settlements, including their healthcare and testing

facilities are quickly displaced and abandoned following the onset of conflict due to the immediate

threat to life [41]. The absence of testing facilities may result in cases not being identified and

reported, therefore making battles appear less impactful to EVD transmission. Meanwhile, protests

and demonstrations aim to cause as much disruption to as many people as possible.

Severity of the conflict

A potential explanation for why remote violence/explosions and battles were very impactful but

often with higher uncertainty is that these types of conflict occurred less frequently, but when they

did, they had a very large impact on EVD cases. The fewer events therefore resulted in the large

uncertainty and a longer dataset (if available) may help to understand if the larger impact of these

conflict types is consistent through time. Additionally, peoples' behaviour and risks must be

considered during more "extreme" forms of violence such as battles and explosions, compared to

protests and demonstrations. Violent conflict may have a very large initial impact (accounting for

the high IRR values), but then the risk stabilises in the area as civilians take precautions (e.g.,

displacement) to protect their life [42].

Alternatively, there is the potential for more violent forms of conflict (such as battles and

explosions) to decrease the risk of EVD. People may be more likely to stay at home during these

events or be subject to curfews and not mix in public spaces, decreasing their exposure to the

pathogen [43]. Furthermore, severe conflict may be more likely to cause fatalities (of both civilians

and combatants), reducing the susceptible pool for EVD cases. More work is needed to

understand how to quantity conflict severity for scientific research purposes, which could be used

to identify dose-response relationships with disease, while still considering that all conflict should

be considered highly disruptive and severe to those living in the affected area.

Duration of conflict related EVD risks

When extending the exposure periods, the effect of conflict on EVD cases was almost eliminated

by 10 weeks in the DRC, while in Guinea, the risk of EVD cases decreased to 4 weeks and then

increased by 10 weeks. However, it appears that extending the lag periods may have been diluting

the effect of conflict on EVD cases, as changing the lag periods to a single week made all the lag

periods for both countries more significant and impactful. These results track with previous

research which found synchronicity and correlation between the reporting of EVD cases and

conflict in the DRC and suggest that the effect of conflict may be long-lasting [11,18].

The results show that conflict is still highly influential multiple months following the event, and in

some instances, conflict had a greater and more significant effect on EVD with time. A potential

explanation may be due to the disruption caused by the conflict event having a cumulative effect.

Previous findings suggest that conflict lengthens the EVD treatment process; e.g., longer time to

detection, isolation, treatment and vaccination, creating a lasting effect and prolonging outbreaks

[18]. Therefore, time would be needed for EVD cases to increase and be reported, after elevated

transmission.

Limitations

There are several limitations to the datasets used here, the first being the assumption that if EVD

or conflict occurs, it is correctly identified (e.g., confirmed and suspected EVD cases were

included) and reported on the correct day with no delay. These assumptions are unlikely, as EVD

cases may be missed if cases never seek formal medical assistance or are in hard-to-reach areas.

Furthermore, the categorisation of conflict events is subjective (category definitions were included

in Additional file 1) and several categories are ambiguous. Data availability was limited and

therefore sub-setting the data (by sub-national region or conflict sub-event type) resulted in limited

data to fit to the model. Small sample sizes meant that some results had wide uncertainty, were

statistically insignificant or could not be calculated, particularly in Guinea. However, generally the

findings were replicated in both countries, in terms of sub-event type, effects of lag and geographic

distribution of risk.

The methodology accounts for confounders by fixing them across the administrative unit. It is not

as effective across multiple administrative units. Sub-national units are different geographical

sizes, with different population sizes and risk factors. Therefore, comparing different

regions/territories can create misleading results. Additionally, the use of vaccination during these

more recent outbreaks [44,45], makes comparison to previous work challenging. While data on

Ebola vaccination are available, the use of ring vaccination means it is difficult to incorporate into

the model used here (as a binary outcome) [37,45]. For example, the sub-national region with the

largest IRR here was in Équateur, part of the DRC which has received relatively little vaccination,

compared to the eastern provinces [37].

Conclusions

The research presented here shows the statistically significant impact that conflict, a known risk factor for infectious disease transmission, has had on EVD cases in the DRC and Guinea in recent resurgences. The effect appears long-lasting (<10 weeks and perhaps longer), with certain geographic regions having >4 times the risk of EVD cases (compared to the national risk of 1.88 and 1.98 for the DRC and Guinea, respectively). Protests and riot-related conflict event types were highly influential and statistically significant here, potentially being far more disruptive to more people, compared to battles. More research to understand how best to quantify conflict severity and intensity could provide further evidence for these mechanisms.

Extra vigilance is needed following protest- and riot-related conflict events for EVD transmission. Continued efforts are needed to try and account for the long-term effect of conflict on EVD and other infectious diseases. Effective public health messaging and scaling up testing in urban areas may help to mitigate these risks, including information on how to reduce transmission in dense urban areas and early clinical symptoms to be aware of. Ebola treatment centres should be prepared for an increase in cases following conflict events. The risk may be elevated for multiple months, with sufficient vaccination stocks made available to areas most at risk.

List of abbreviations

ACLED - Armed Conflict Location & Event Data Project

CI – Confidence Interval

DRC – Democratic Republic of the Congo

EVD – Ebola Virus Disease

GUI – Guinea

HEMA - Humanitarian Emergency Response Africa

IRR – Incidence Rate Ratio

MoH – Ministry of Health

NGOs – Non-Governmental Organisations

SCCS – Self-Controlled Case Series

WASH – Water, Sanitation and Hygiene

Declarations

Ethics approval and consent to participate

Although the work presented here uses human data, all the data used are from public available

data sources, which are none identifiable and therefore ethical approval was waived for this

analysis.

Consent for publication

Not applicable

Availability of data and materials

The datasets analysed during this study are all publicly available and references throughout the

manuscript. Further code related to the methods and data visualization here are available at:

https://github.com/GinaCharnley/ebola_conflict

Competing interests

The authors declare that they have no competing interests.

Funding

The work presented here received no specific funding, however we acknowledge joint Centre

funding from the UK Medical Research Council and Department for International Development

[MR/R0156600/1]. The funder played no role in the design of the study and collection of data, the

analysis, and interpretation of data and in writing the manuscript.

Authors' contributions

GECC conceptualised and designed the study, collated the data sources, analysed the data, wrote

the manuscript and incorporated feedback. IK contributed to interpreting the findings, provided

expertise on conflicts and global health and revised several drafts of the manuscript. NG provided

statistical expertise on the methodology, assisted in writing the manuscript, analysing the data and

revised several drafts. EBM contributed to interpreting the findings, provided expertise on conflicts

and Ebola in the DRC and revised several drafts of the manuscript. KAMG co-conceptualised and

designed the study, provided data sources, provided expertise on epidemiological methods,

contributed to interpreting the findings and revised several drafts of the manuscript. All authors

read and approved the final manuscript.

Acknowledgements

We would like to acknowledge and thank those working in the field who collected and collated the

data sources used here, namely the Ministère de la Santé Publique de la République

Démocratique du Congo and Ministère de la Santé et de l'Hygiène Publique de Guinée.

Additionally, we would like to thank Sophie Meakin for collating the situation reports used in some

of the DRC data used here.

References

1. Pattyn SW, Jacob W, Van der Groen G, Piot P, Courteille G. Isolation of Marburg-like virus from a case of haemorrhagic fever in Zaire. Lancet. 1977:573-4.

2. Bowen ET, Platt GS, Lloyd G, Baskerville A, Harris WJ, Vella EE. Viral haemorrhagic fever in southern Sudan and northern Zaire. Preliminary studies on the aetiological agent. Lancet. 1977:571-3.

3. Jacob ST, Crozier I, Fischer WA, Hewlett A, Kraft CS, Vega MA, et al. Ebola virus disease. Nat. Rev. Dis. Primers. 2020 Feb 20;6(1):13.

4. World Health Organization. Ebola virus disease. https://www.who.int/news-room/fact-sheets/detail/ebola-virus-disease (2021).

5. Pourrut X, Kumulungui B, Wittmann T, Moussavou G, Délicat A, Yaba P, et al. The natural history of Ebola virus in Africa. Microbes Infect. 2005;7(7-8):1005-14.

6. Centres.for Disease Control and Prevention (CDC). History of Ebola Disease Outbreaks.

https://www.cdc.gov/vhf/ebola/history/chronology.html (2022)

7. Adepoju P. Lessons from Ebola as DRC grapples with conflict, measles, and covid-19. BMJ.

2020 Jul 21;370.

8. Mukadi-Bamuleka D, Mambu-Mbika F, De Weggheleire A, Edidi-Atani F, Bulabula-Penge J,

Mfumu MMK, et al. Efficiency of Field Laboratories for Ebola Virus Disease Outbreak during

Chronic Insecurity, Eastern Democratic Republic of the Congo, 2018-2020. Emerg. Infect.

Dis. 2023;29(1):1-9.

9. Kinganda-Lusamaki E, Black A, Mukadi DB, Hadfield J, Mbala-Kingebeni P, Pratt CB, et al.

Integration of genomic sequencing into the response to the Ebola virus outbreak in Nord

Kivu, Democratic Republic of the Congo. Nat. Med. 2021;27(4):710-716.

10. Mukadi-Bamuleka D, Sanogo YO, Bulabula-Penge J, Morales-Betoulle ME, Fillon P,

Woodruff P, et al. RDT Working Group. Postmortem Surveillance for Ebola Virus Using

OraQuick Ebola Rapid Diagnostic Tests, Eastern Democratic Republic of the Congo, 2019-

2020. Emerg. Infect. Dis. 2022;28(2):420-424.

11. Wells CR, Pandey A, Ndeffo Mbah ML, Gaüzère BA, Malvy D, Singer BH, et al. The

exacerbation of Ebola outbreaks by conflict in the Democratic Republic of the Congo. PNAS.

2019;116(48):24366-72.

12. Mbala-Kingebeni P, Aziza A, Di Paola N, Wiley MR, Makiala-Mandanda S, Caviness K, et

al. Medical countermeasures during the 2018 Ebola virus disease outbreak in the North Kivu

and Ituri Provinces of the Democratic Republic of the Congo: a rapid genomic assessment.

Lancet Infect, Dis. 2019;19(6):648-657.

13. Wise PH, Barry M. Civil war & the global threat of pandemics. Daedalus. 2017;146(4):71-84.

14. Gonzalez-Torres A, Esposito E. Epidemics and conflict: Evidence from the Ebola outbreak

in Western Africa. Available at SSRN 3544606. 2016.

15. van Stekelenburg BC, De Cauwer H, Barten DG, Mortelmans LJ. Attacks on health care

workers in historical pandemics and COVID-19. Disaster Med. Public Health Prep.

2023;17:e309.

16. Oppenheim B, Lidow N, Ayscue P, Saylors K, Mbala P, Kumakamba C, et al. Knowledge

and beliefs about Ebola virus in a conflict-affected area: Early evidence from the North Kivu

outbreak. J. Glob, Health. 2019 Dec;9(2).

17. Vinck P, Pham PN, Bindu KK, Bedford J, Nilles EJ. Institutional trust and misinformation in

the response to the 2018–19 Ebola outbreak in North Kivu, DR Congo: a population-based

survey. Lancet Infect. Dis. 2019;19(5):529-36.

18. Kraemer MU, Pigott DM, Hill SC, Vanderslott S, Reiner RC, Stasse S, et al. Dynamics of

conflict during the Ebola outbreak in the Democratic Republic of the Congo 2018–2019.

BMC Med. 2020;18(1):1-0.

19. McPake B, Witter S, Ssali S, Wurie H, Namakula J, Ssengooba F. Ebola in the context of

conflict affected states and health systems: case studies of Northern Uganda and Sierra

Leone. Confl. Health. 2015;9(1):1-9.

20. Global Conflict Tracker. Instability in the Democratic Republic of Congo.

https://www.cfr.org/global-conflict-tracker/conflict/violence-democratic-republic-congo

(2023).

21. Human Rights Watch. World Report 2023.

https://www.hrw.org/sites/default/files/media_2023/01/World_Report_2023_WEBSPREADS

_0.pdf (2023).

22. Human Rights Watch. Guinea Events of 2020. https://www.hrw.org/world-

report/2021/country-chapters/guinea (2020).

23. Charnley GEC, Jean K, Kelman I, Gaythorpe KAM, Murray KA. Association between

Conflict and Cholera in Nigeria and the Democratic Republic of the Congo. Emerg. Infect.

Dis. 2022;28(12):2472-81.

24. Meakin S. Data for the 2018 outbreak of Ebola in the Democratic Republic of the Congo in

North-Kivu and Ituri Provinces. https://github.com/sophiemeakin/ebola_drc (2019).

25. Ministère de la Santé de la République Démocratique du Congo. Mailing list.

https://us13.campaign-

archive.com/home/?u=89e5755d2cca4840b1af93176&id=aedd23c530 (2019).

26. Human Emergency Response Africa (HEMA). Democratic Republic of the Congo: Ebola

(2021) Subnational cases. https://data.humdata.org/dataset/democratic-republic-of-congo-

ebola-2021-subnational-cases (2021).

27. ACAPS. GUINEA EBOLA EVD 2021 SUBNATIONAL CASES, DEATHS,

HOSPITALISATIONS, CONTACT TRACING, AND VACCINATIONS.

https://www.acaps.org/guinea-ebola-evd-2021-subnational-cases-deaths-hospitalisations-

contact-tracing-and-vaccinations (2021).

28. Armed Conflict Location & Event Data Project (ACLED). Data Export Tool.

https://acleddata.com/data-export-tool/ (2023).

29. Eck K. In data we trust? A comparison of UCDP GED and ACLED conflict events datasets.

Coop. Confl. 2012;47(1):124-41.

30. Whitaker HJ, Paddy Farrington C, Spiessens B, Musonda P. Tutorial in biostatistics: the

self-controlled case series method. Stat. Med. 2006;25(10):1768-97.

31. Therneau T. A Package for Survival Analysis in R. R package version 3.5-5.

https://CRAN.R-project.org/package=survival (2023).

32. Mostofsky E, Coull BA, Mittleman MA. Analysis of observational self-matched data to

examine acute triggers of outcome events with abrupt onset. Epidemiol. (Cambridge,

Mass.). 2018;29(6):804.

33. Xu S, Zeng C, Newcomer S, Nelson J, Glanz J. Use of fixed effects models to analyze self-

controlled case series data in vaccine safety studies. J. Biom. Biostat. 2012:006.

34. UN Office for the Coordination of Humanitarian Affairs - Field Information Services Section.

Democratic Republic of the Congo - major cities.

https://data.humdata.org/dataset/democratic-republic-of-the-congo-major-cities (2019).

35. Simplemaps. Guinea Cities Database. https://simplemaps.com/data/gn-cities (2023).

36. Hijmans RJ. geosphere: Spherical Trigonometry. R package version 1.5-18.

https://CRAN.R-project.org/package=geosphere (2022).

37. Bausch DG. The need for a new strategy for Ebola vaccination. Nat. Med. 2021;27(4):580-1.

38. Moran B. Fighting Ebola in conflict in the DR Congo. Lancet. 2018;392(10155):1295-6.

39. AL JAZEERA AND NEWS AGENCIES. Several killed as protests rock Goma in eastern

DRC. https://www.aljazeera.com/news/2021/12/21/several-killed-as-protests-rock-goma-in-

eastern-drc (2021).

40. AFP, AL JAZEERA. Protests against military administration paralyse Guinea capital.

https://www.aljazeera.com/news/2022/7/29/protests-against-military-administration-

paralyse-guinea-capital (2022).

41. Relief Web. Democratic Republic of the Congo faces acute health crisis amid rising

violence. https://reliefweb.int/report/democratic-republic-congo/democratic-republic-congo-

faces-acute-health-crisis-amid-rising-violence (2023).

42. Relief Web. After evacuating Kayna, DRC, MSF is trying to return to assist displaced in

Kanyabayonga. https://reliefweb.int/report/democratic-republic-congo/after-evacuating-

kayna-drc-msf-trying-return-assist-displaced (2004).

43. Radio France Internationale. Clashes erupt in Guinea Conakry after opposition call for anti-

junta protests. https://www.rfi.fr/en/africa/20220818-clashes-erupt-in-guinea-conakry-after-

opposition-call-for-anti-junta-protests (2022).

44. Maxmen A. Ebola vaccine approved for use in ongoing outbreak. Nature. 2017 May 30;10.

45. World Health Organization. Ebola virus disease: Vaccines. https://www.who.int/news-

room/questions-and-answers/item/ebola-vaccines (2020).