Title

scRNA-Seq reveals elevated interferon responses and TNF-α signaling via NFkB in
monocytes in children with uncomplicated malaria

Authors: Collins M. Morang'a[1], Riley S. Drake[2,3,4,5], Vincent N. Miao[2,3,4,5], Nancy K. Nyakoe[1],
Dominic S.Y. Amuzu[1], Vincent Appiah[1], Yaw Aniweh[1], Yaw Bediako[1], Saikou Y. Bah[6], Alex
K. Shalek[2,3,4,5], Gordon A. Awandare[1], Thomas D. Otto[6, *] & Lucas Amenga–Etego[1 ,*]

Affiliations
[1]West African Centre for Cell Biology of Infectious Pathogens (WACCBIP), Department of
Biochemistry, Cell and Molecular Biology, University of Ghana, Legon, Accra, Ghana
[2]Program in Health Sciences & Technology, Harvard Medical School & Massachusetts
Institute of Technology, Boston, MA 02115, USA
[3]Institute for Medical Engineering & Science (IMES), Department of Chemistry, and Koch
Institute for Integrative Cancer Research, MIT, Cambridge, Massachusetts, 02139, USA
[4]Ragon Institute of MGH, MIT and Harvard, Cambridge, Massachusetts, USA
[5]Broad Institute of MIT and Harvard, Cambridge, Massachusetts, USA
[6]School of Infection & Immunity, MVLS, University of Glasgow, Glasgow, UK

*Correspondence thomasdan.otto@glasgow.ac.uk & lamengaetego@ug.edu.gh

Abstract

Malaria causes significant morbidity and mortality worldwide, disproportionately impacting
sub-Saharan Africa. Disease phenotypes associated with malarial infection can vary widely,
from asymptomatic to life-threatening. To date, prevention efforts, particularly those related
to vaccine development, have been hindered by an incomplete understanding of which
factors impact host immune responses resulting in these divergent outcomes. Here, we
conducted a field study in 224 malaria positive individuals (rapid diagnostic test - RDT) from
a high transmission area in Ghana, to determine immunological factors associated with
uncomplicated malaria "patients" compared to healthy individuals in the community
"controls". Generally, the patients had higher parasite density levels although it had a
negative correlation with age, suggesting that, is a key indicator of disease pathogenesis.
We applied single-cell RNA-sequencing to compare the immunological phenotypes of 18,176
peripheral blood mononuclear cells (PBMCs) isolated from a subset of the patients and
controls (n=11/224), matched on location, age, sex, and parasite density. On average,
patients were characterized by a higher fractional abundance of monocytes and an
upregulation of innate immune responses, including those to type I and type II interferons
and tumor necrosis factor-alpha (TNF-α) signaling via NFκB. Further, in the patients, we
identified more putative interactions between antigen-presenting cells and proliferating CD4
T cells and naïve CD8 T cells driven by MHC-I and MHC-II signaling pathways, respectively.
Together, these findings highlight transcriptional differences between immune cell subsets
associated with malaria that may help guide the development of improved vaccines and new
therapeutic interventions for individuals residing in endemic areas.

Background

In 2022, global estimates of malaria cases and deaths have increased to 249 million cases
and 608 000 deaths [1]. However, the development of an effective vaccine to address this
global health threat remains challenging due to an incomplete understanding of the parasite's
biology and limited knowledge of which host factors influence clinical responses to infection.

In malaria endemic communities, individuals may harbour malaria infections with mild to no
symptoms warranting treatment, here referred to as healthy community controls. Such
infections may be cleared naturally or progress to a uncomplicated malaria, where symptoms
become profound enough to necessitate medical intervention. Instructive factors include
environmental exposures, transmission intensity, host and parasite genetics, host-pathogen
interactions, and host immune responses [2–5]. Illustratively, upregulation of interferon
responses and p53 gene expression can attenuate inflammation and protect children from
fever [6]; and, when comparing children with asymptomatic and severe malaria, the genes
most upregulated in severe cases are related to immunoglobin production and interferon
signaling [7]. As reviewed previously, studies have postulated that interferons can orchestrate
immune regulatory networks to dampen inflammatory responses and restrict humoral
immunity, thus playing a critical role as a wedge that determines protection versus
permissiveness to malaria infection [8,9].

Similarly, it has been shown that the number and phenotype of cells responding to infection
can vary with exposure to *Plasmodium* [10]. For example, Africans, who tend to have higher
levels of exposure, have been shown to exhibit metabolic and platelet activation during
malaria infection as compared to typically infection-naïve Europeans [10]. Similarly, children
who experience high cumulative malaria episodes show upregulation of interferon-inducible
genes and immunoregulatory cytokines, suggesting an immune modification to prevent
immunopathology and severe outcomes during new infections [11]. Beyond differences in
exposure and infection history, the strain responsible for each infection can also alter
immune response dynamics and disease pathogenesis [12,13].

Since so many factors can influence host response dynamics to infection (e.g., exposure, the
timing of infection), some studies have implemented tightly regulated models of malarial
infection, such as controlled human malaria infections (CHMI). CHMI studies have identified
several pathways, including toll-like receptor signaling [14], platelet activation [10], interferon
signaling [10,15,16], and B-cell receptor signaling, that are involved in immunological modulation
of *Plasmodium falciparum* infections [6]. Although CHMI enable more controlled examinations
of host-pathogen dynamics post-malaria infection, clinically relevant differences can arise
between responses seen in CHMI and natural exposure due to unresolved
immunopathological mechanisms elicited during *P. falciparum* infection [10]. Thus, studies
involving natural cohorts provide a better avenue to understand vaiability in immune
responses developed through repeated exposure and how they influence disease
phenotypes. Besides, immunity to malaria develops very slowly through repeated infections,
and can wane quickly if individuals leave a malaria endemic areas, suggesting that
continuous natural exposure to malaria antigens is important for development of long term
immunity [11]. Collectively, these studies demonstrate the importance of obtaining a more

comprehensive understanding of the host and pathogen factors that influence immune
responses to inform the development of new therapeutic approaches and improved vaccines.

To date, most genomic analyses of immunological responses to malarial infection have been
performed in heterogeneous cell populations of blood, brain, liver, or spleen tissues [17]. The
majority of these studies have been conducted in children and the studies show that
symptomatic infections, as mentioned above, are characterized by upregulated expression of
genes involved in interferon signaling, antigen presentation, neutrophil-associated
signatures, and B cell modules relative to healthy controls [17]. Adults, meanwhile, present
slightly varying responses: symptomatic Malian adults, compared to naïve individuals, had
upregulated B cell receptor signaling but more modest upregulation of interferon responses,
while symptomatic Cameroonian adults showed marked induction of genes related to
interleukins and apoptosis compared to presymptomatic individuals [18,19]. These
inconsistencies may be related to patient history/exposure or differences in cellular
composition influencing clinical course through a combination of direct and indirect
responses. The emergence of single-cell transcriptomics provides a unique opportunity to
examine the sources of this variability [20] by profiling abundance and transcriptomic variation
across immune cell populations in individuals with high malaria exposure but divergent
clinical phenotypes. Moreover, by examining the expression of ligands, receptors, and genes
involved in intercellular signaling, we can identify the critical mediators of immune responses
and the pathogenesis of malaria for subsequent validation [21].

Here, we present a comparative analysis of peripheral blood mononuclear cells (PBMCs)
phenotypes in children from two related surveys conducted in 2019. An active case detection
of *P. falciparum* infections at the community level (controls) and passive case detection at
the health facility level for patients with uncomplicated malaria (patients) in an endemic area
in northern Ghana. Our data describe in unprecedented detail, cell subsets and signaling
pathways associated with disease severity to provide new insights into the immune response
mechanisms that influence the course of *P. falciparum* infections in young children.

Results

Clinical characteristics of study participants

In this study, we defined "controls" as healthy individuals who tested *Plasmodium* positive by RDT in the community. We defined "patients" as individuals with uncomplicated malaria who visited the hospital/health center from the same community and tested *Plasmodium* positive by RDT and were treated on outpatient basis. All samples were collected from the same region, Upper East region of Ghana which is a high transmission area. Overall, 224 individuals were surveyed, including 40% (90/224) of the participants who were community healthy controls and 60% (134/224) of the participants who were patients with uncomplicated malaria **(Figure 1a, Supplementary Table 1).** Although most participants were children between 1-15 years, there was no significant difference between the median age of patients compared to the controls (Wilcoxon rank-sum test, P=0.74) *(Supplementary Table 1).* But, there was a significant difference in the median parasite density of patients compared to the controls (Wilcoxon rank-sum test, P<0.001) *(Supplementary Table 1).* Further, the study sought to determine if the patients had higher parasite densities than the controls regardless of age. In general, there was negative correlation between parasite density and age regardless of phenotype up to age 25 years *(Figure 1b).* Under 3 years, the patients tended to have lower parasite densities, but these were still higher than their controls counterparts. After about age 10 parasite densities fell gradually and plateaued around age 25 but with high variability between the groups *(Figure 1b).* The correlation between parasite density and malaria patients was statistically significant (R2 = -0.38, P<0.001), but the correlation between parasite density and age was not statistically significant in the control group **(Figure 1b)**.

Profiling pediatric malaria immune-cell populations using single-cell analysis

In order to examine global differences in cellular composition, gene expression and intercellular communication between the two groups, we matched individuals based on age (aged 4-8 years), sex, and parasite density for both patients and controls and performed single-cell RNA-seq (scRNA-seq) (***Figure 1a, Supplementary Table 2)***. There was no significant difference in median parasite density between patients and controls in the matched individuals (Wilcoxon rank-sum test, P>0.71) (***Supplementary Table 2)***. In total, we generated 18,176 high-quality single-cell profiles across eleven children with *P. falciparum* infections, allowing us to ascertain differences in expression patterns of immune response genes that might influence disease pathogenesis. Each sample was profiled using Seq-Well S[3], a portable, simple massively parallel scRNA-Seq method [22]. The resulting data were filtered to remove cells based on the fractional abundance of mitochondrial genes (<30%) and transcripts expressing in <20 cells. After variable gene selection, dimensionality reduction, clustering, cluster removal, and reclustering (***Methods***), we retained 18,303 transcripts and identified 10 distinct cell subsets in the 18,176 cells, across the two groups of children (***Figure 1c; Supplementary Figure 1a***).

We manually annotated these 10 clusters using known RNA marker genes to identify B cells, CD4 T cells, CD8 T cells, natural killer (NK) cells, monocytes (Mono), and dendritic cells (DC) (***Supplementary Figure 1b and 1c***). To identify and enumerate cell subsets in our dataset at higher resolution, we opted to map our query dataset to an annotated multimodal

reference dataset of PBMCs. First, we confirmed that all the cell subsets identified using
manual annotation were present in the resultant UMAP (**Supplementary Figure 1d**)**.** As
expected, our reference mapped dataset recapitulated all PBMC subsets, including B, CD4 T
cells, CD8 T cells, NK cells, Mono, and DC (these subsets are used throughout the work;
Figures 1c and 1d)**.** We identified several subclusters, such as intermediate, memory, and
naïve B cells; naïve, proliferating, effector memory and central memory CD8 and CD4 T
cells; proliferating CD56+ NK cells; CD14+ and CD16+ monocytes (Mono); plasmacytoid
(pDC) and conventional (cDC) dendritic cells, and other cell subsets (**Figures 1e and 1f**)**.**
Since the reference dataset has only annotated two Mono clusters (CD14+ and CD16+), we
hypothesized that there might be additional transcriptional heterogeneity describing actively
responding Mono subpopulations. Therefore, further sub-clustering was done which resolved
the Mono into 3 large subpopulations (Mono 1, Mono 2, Mono 3) and 1 small cluster (Mono
4) (**Figures 1g, 1h and 1i**) based on previously reported markers [23]. Taken together, these
data distinguish nearly all distinct cell subsets that were present in PBMCs of children in both
the patients and controls.

Differences in relative cellular composition between the groups

Next, we asked whether there were significant differences in the relative proportions of cell
types between the patients and control group. We found that relative cell proportions of the
major cell subsets (B, CD4 T, CD8 T, NK, Mono, and DC) varied between individuals in each
group (**Figures 2a and 2b, Supplementary Table 3**). The patients exhibited elevated levels
of circulating Mono while the controls had higher proportions of circulating B cells (Dirichlet-
multinomial regression, P<0.01; **Figure 2a, Supplementary Table 3**). Further analysis of the
B cell subsets showed that the abundance of naïve and intermediate B cells was significantly
reduced in the patient group compared to the control group (Dirichlet-multinomial regression,
P<0.05; **Figure 2b, Supplementary Table 3**). We also found a significant expansion of both
CD14+ and CD16+ Mono subsets in patients compared to the control group (Dirichlet-
multinomial regression, P<0.01; **Figure 2b, Supplementary Table 3**). Although there is
evident variation in cellular proportions of T lymphocytes among all the individuals (**Figures
2a and 2b**)**,** we did not observe any significant difference in proportions of either CD4 or CD8
T cells between the groups (Dirichlet-multinomial regression, P>0.05; **Figure 2b,
Supplementary Table 3**). However, the proportions of naïve and central memory CD4 T
cells were significantly higher in the patients compared to the control group (Dirichlet-
multinomial regression, P<0.01; **Figure 2b, Supplementary Table 3**). NK cell frequency was
also higher in patients suggesting that they may play a role in disease progression (Dirichlet-
multinomial regression, P>0.05; **Figure 2b, Supplementary Table 3**). Among NK cells, the
proliferating and CD56+ subsets were higher in patients compared to controls, but these
differences were not statistically significant (Dirichlet-multinomial regression, P>0.05; **Figure
2b, Supplementary Table 3**). Overall, the minor subsets of T cells and other cell types with
low frequencies did not show differences in proportions between the groups but the main cell
subsets had significant differences in proportions between patients and controls.

Comparative analysis of inflammatory responses in children with malaria

Having identified shifts in the composition of circulating immune cells between the patients
and controls, we next asked whether gene expression differed within each immune cell
subset between the two groups. Comparing patients to controls, we observed the largest

transcriptional changes (measured by pairwise DE across cell types with adjusted P value <
0.05 and log fold change > 2) within B cells and Mono (**Figures 2c, Supplementary Table
4**)**.** Apart from B cell function genes, there was a general trend towards upregulation of
inflammatory genes in B and T cells in patients relative to control group, including *S100A8,*
CXCL8, and *S100A9* (**Figure 2c**). Significant transcriptional changes were also observed in
Mono, with genes such as *IFITM3, FCER1G,* and *CCL4* being upregulated in patients
compared to the control group (**Figure 2c**). Patients were also associated with the
upregulation of Major Histocompatibility Complex I (MHC-I) genes such as *HLA-A* and *HLA-*
C which are involved in antigen presentation in Mono (**Figure 2c**). In CD4 and CD8 T cells,
there was increased expression of some inflammatory factor signalling genes such as
CXCL8 and *NFKBIA* in patients relative to the control group*,* suggesting direct sensing of
parasite products during clinical presentation (**Figure 2c**). Using gene set enrichment
analyses (GSEA), we found that the patients had robust induction of several innate immune
response pathways such as TNF-α signaling via NF-κB, TGF-β signaling, IL6-JAK-STAT
pathway, complement, IL2-STAT5 signaling, inflammatory response, interferon-α response
(IFN-α), and interferon-γ response (IFN-γ) (**Figure 3a, Supplementary Table 5**). We
observed that although each cell type was enriched in one or more of these pathways, there
was a unique molecular signature of the genes involved in each. Upregulation of IFN-γ and
IFN-α response pathways in Mono were characterized by increased expression of genes
such as *IFITM2, IFITM3, IL10RA,* and *TNFAIP3,* while in NK cells they were typified by
genes such as *NFKBIA, CD69*, and *ISG20* (**Figure 3b & 3c, Supplementary Table 5**). Mono
and natural killer cells upregulated TNF-α signaling via the NF-κB pathway with the induction
of genes related to this pathway such as *IL1B* and *TNFAIP3* for Mono, and *IL7R, CD44,* and
NFKBIA for NK cells (**Figure 3b & 3c, Supplementary Table 6**). Inflammatory responses in
Mono were characterized by *IL10RA, IL1B*, and *CXCL8* while in NK cells they were driven by
CD69, IL7R, CXCL8, and *NFKBIA* among others (**Figure 3b & 3c, Supplementary Table 5**).
Thus, the enrichment of unique genes for each cell subset for similar pathways suggests a
specific but concerted contribution of each cell subset toward the innate immune response in
patients.

Relative enrichment of ISGs gene modules in monocytes of patients relative to
controls

Since IFN genes were significantly upregulated in Mono patients relative to the control group,
we next sought to determine if entire gene modules were enriched. Interferon stimulated
genes (ISGs) modules scores were significantly higher in B cells, DC, CD4 T cells, and Mono
in patients compared to the control group (**Wilcoxon, adjusted P < 0.01 for all
comparisons, Figure 4a, 4b, 4d, and 4e**)*;* however, there were no significant differences in
ISG module scores in CD8 T cells and NK cells. Further examination of intra- and inter-
individual variation in these module scores revealed substantial intra-individual variation in
cells from the same participant and between cells of the same type from different participants
(**Supplementary Figure 1e**)**.** Overall, our data show that Mono play a significant role in
defining malaria patients compared to control participants from the same community through
induction of the ISGs gene modules.

Role of MHC-I and MHC-II signaling pathways in cell-to-cell interactions

Next, we used our single-cell data to infer putative axes of cell-to-cell communication using signaling ligands, cofactors, and receptors. First, we discerned cell-to-cell interactions in the patients and found that the number of interactions (ligand-receptor) originating from primary innate immune cells such as DC and Mono were greater than those originating from non-antigen presenting cells (**Supplementary Table 6**). However, our data show very few inferred cellular communication networks in the control group (**Supplementary Table 6**). This analysis suggests a role for Mono as antigen-presenting cells in orchestrating pro-inflammatory responses by interacting with proliferating CD4 T cells, intermediate B cells, effector memory T cells, and naïve CD8 T cells in the patient group (**Figure 5a**). Conventional DC also produced factors that interact with proliferating and effector memory CD4 T cells respectively, suggesting a concerted effort by antigen-presenting cells to activate the immune response in patients (**Figure 5a**). Communication probabilities indicated that MHC-I and MHC-II play a role in these interactions among other pathways. The most significant receptor-ligand pairs for HLA-A, HLA-B, HLA-C, HLA-E, HLA-G, and HLA-F ligands for MHC class I include CD8A, CD8B, LILRB2, and LILRB1 (**Figure 5b**). The leading intercellular ligand-receptor pairs with CD4 T cells as signal receivers were distinct HLA genes, with the highest relative contribution being driven by HLA-DRA and HLA-DRB1 (**Figure 5c**). The other minor signaling pathways that were important in patients include MIF, RESISTIN, ANNEXIN, GALECTIN, ADGRE5, APP, CD22, CD45, SELPLG, CD99, CLEC, and TNF signaling networks. For the TNF signaling pathway, the CD56$^+$ NK cells showed to be interacting with Mono and also with proliferating CD4 T cells, effector memory CD4 and CD8 T cells, and cDC (**Figure 5d**). This cell communication network was mediated by *TNF* in the sender cells and *TNFRSF1B* in the receiver subsets (**Figure 5e**), and this corroborates the DE results (**Figure 5g**). We examined the expression levels of *TNFRSF1B* across all the cell subsets and found that indeed it was expressed in all the receiver cells (**Figure 5f**). Only the pDC and CD16 Mono showed cell-to-cell interactions with naïve and intermediate B cells and might be playing a role in B cell activation and development in the control group through MHC class II molecules (**Figure 5g and 5h**). Therefore, exposure of innate immune cells to parasite ligands may potentially activate intracellular signaling cascades through cell-to-cell interactions to induce rapid expression of a variety of innate immune genes.

Discussion

Here, we recruited 224 participants malaria from Navrongo, a high malaria transmission area with seasonal fluctuations [24]. Interestingly, most of participants indicated that they use long lasting insecticide treated mosquito nets (LLINs), which helps to explain the low frequency of infections; after screening 1,000 individuals in the community, <10% of them were positive for *P. falciparum* as community healthy controls, suggesting a reduction in the malaria infection reservoir. The National malaria elimination programme (NMEP) distributes LLINs as part of strategy interventions, including community-based seasonal malaria chemoprevention initiatives for children under 5 years in order to reduce the malaria burden in this area [24]. The ability of insecticide treated nets (ITNs) to interrupt malaria transmission has been shown in large scale studies, which demonstrated that modern housing and ITNs could reduce malaria infections by 1% and 16%, respectively [25]. Further, we investigated the relationship between age and parasite density, and found that parasite densities tended to decrease with age, but the levels were generally higher in patients compared to controls in this high transmission intensity area [5].

To better understand cellular responses driving these divergent clinical phenotypes, we performed scRNA-seq on PBMCs samples from eleven of the 224 individuals among the two groups, controlling for group variability driven by age, fever and parasite density. This enabled us to identify a potential role for interferon responses and TNF-α signaling via NFκB in Mono during the clinical manifestation of pediatric malaria infection. We also found differences in the fractional abundances of PBMC cell subsets, with patients characterized by a proportional increase in Mono while controls had a higher proportion of circulating B cells. We showed cellular level variations in the expression of innate immune modules within and between individuals as well as between clinical phenotypes. Further, we identified a role for Mono and other innate immune cells through MHC-I and MHC-II molecules in driving cell-to-cell interactions with CD8 and CD4 T cells respectively. Together, our work recontextualizes the function of the innate immune cells in malaria, demonstrates how variable their responses can be, and links specific acute phase response signaling pathways to clinical presentation.

Differential gene expression comparing patients and controls across cell types revealed a significant upregulation of genes associated with innate immunity in different cell types. We show that CCL3 and CCL4 (also known as macrophage inflammatory protein MIP-1α and MIP-1β respectively) were upregulated in Mono of patients, suggesting their possible role in modulating clinical disease [26]. CXCL8, the most potent human neutrophil attracting/activation chemokine [27] was also highly upregulated in B cells, CD4, and CD8 T cells. Other studies have shown that circulating levels of CXCL8 and CCL4 correlated with parasite density, and when found in the cerebrospinal fluid they can predict cerebral malaria mortality [13,28–30]. Furthermore, the adaptive immune cell subsets (B cells and T cells) in the patient group expressed two alarmins (S100A8 and S100A9) that are known to form calprotein heterodimer, an endogenous TLR4 ligand; this could suggest a possible role to silence hyperinflammation [31]. We also show significant expression of FCER1G in B cells, Mono, and DC in patients, which is induced by IFN-γ and encodes for a gamma chain of the FC receptor and it is suggested to play an important role in controlling parasitemia [6]. Collectively, our data imply that both adaptive and innate immune cells cooperatively play a role during the pathogenesis of malaria in patients when compared to healthy controls.

322
We showed that several immune-related pathways are activated by *Plasmodium* infection
and disease including the TNF-α signaling via NFκB pathway, IFN-α/γ responses, IL2-
STAT5 signaling, and inflammatory response pathway in patients. Since the parasite life
cycle involves repeated red cell invasion and rupture, the release of pyrogenic cytokines that
drive these pathways such as interleukins, interferons, and TNF in Mono and NK cells, can
signify pathophysiological events occurring in malaria patients [13,32]. These observations could
also mean that children who patients were sampled quite early during the onset of the
disease progression trajectory [12]. Our data are consistent with those previously described by
integrating whole blood transcriptomics, flow cytometry, and plasma cytokine analysis [6], and
our results further identify the cell subsets in which these pathways were more enriched. We
show that each of the cell subsets has a unique signature of genes enriched in these
immunogenic pathways with minimal sharing. Several studies have shown similar innate
immune response pathways in individuals with malaria such as whole blood transcriptomics
of the Fulani of West Africa [33], children repeatedly exposed to malaria [6,11], controlled human
malaria infection (CHMI) studies [16], and even mice models [31]. We have now confirmed some
of these observations and demonstrated that in the patient state, robust upregulation of
certain genes in specific cell subsets is associated with systemic inflammatory responses.
Innate immune cells, such as Mono, DC, and NK cells, appear to be most reactive in
patients, probably due to continuous exposure in a high transmission area as suggested by
other studies [7,31,34].
343
By collating gene modules of interferon-stimulated genes (ISGs), we show that there is a
differencial expression between patients and controls across different cell subsets. ISGs are
normally produced as a function of interferon responses (IFNs) [8], which we observe to be
enriched in patients. IFNs are produced primarily by DC to activate ISGs in other cells [35], and
we observed that B cells, T cells, Mono, and DC have higher ISG module scores in patients
compared to controls. Notably, our data show that each cell or cell subset responds
differently upon IFN activation with varying transcriptional responses of an ISG module
between individuals. This variability was also observed for cytokine modules, NF-κB target
modules, and HLA modules. Similarly, a previous CHMI study observed striking inter-
individual variation in immune cell composition and immune responses, demonstrating that
an individual can have a unique immune fingerprint [10]. Thus, the variations in immune
responses that we observed could be attributed to the complexity of the *P. falciparum* life
cycle with several developmental erythrocytic stages, duration of infections, intensity of
infection in each individual, genetic factors, genetic variation in immune response genes
among other factors [12,36]. These findings on inter-individual variability in immune responses
could provide insights when considering the design and evaluation of interventions that target
host immunity in the control of malaria.
361
Our scRNA-Seq data enabled us to quantitatively infer and analyze cell-to-cell
communication networks across all the innate and adaptive immune cells (Jin et al.,
2021). This analysis enabled us to uncover coordinated interactions between innate and
adaptive immune cells through various ligands. The cell-to-cell interactions in patients were
driven by MHC class I and II signaling pathways, whereby antigen-presenting cells were
shown to have more interactions with proliferating CD4 and naive CD8 T cells. The
importance of HLA genes has long been demonstrated by Hill and colleagues who
associated HLA-Bw53 antigen and DRB1*1302–DQB1*0501 haplotype to independently
protect against severe malaria in West Africa [37]. Thus, our observations on cell-cell

interaction involving HLA molecules and T cells support the importance of these molecules
during *P. falciparum* infection and disease progression, consistent with the observed varying
degrees of interactions in patients compared to control groups. We also showed that within
patient group, there are contrasting interactions between various HLA I and HLA II molecules
with CD8 or CD4 T cell receptors respectively, which could be related to their tight regulation
and antigen-presenting ability [38,39]. Activation of CD4 and CD8 T cells has been correlated
with protective immunity to malaria, and they can differentiate into several functionally distinct
subsets in the presence of various cytokines [40]. It was not surprising that we identified
different fractional abundances of CD4 and CD8 T cell subsets in patients compared to
control group of children, but we demonstrate that ultimately this results in varying degrees of
interactions with Mono or DC. Future work should seek to identify the mechanisms that result
in these variations and their impact in orchestrating phagocytic and humoral responses as
this critical knowledge gap will be important in developing T cell-based malaria vaccines.

Overall, by using scRNA-seq on PBMCs obtained from patients and controls in a high
transmission area, this work sheds light on the interplay between peripheral immune cells
during uncomplicated malaria, uncovering the genes and immune pathways in specific cell
types that might play a significant role in defining the outcomes of infection. Data presented
here demonstrate that the patients with uncomplicated malaria were characterized by the
presence of inflammatory response signatures in specific cell types compared to the control
group. The results could also suggest that in the control group, a muted innate immune
response or disease tolerance mechanism plays a role in enabling children to harbour
malaria parasites in high malaria transmission areas without developing uncomplicated
malaria [41]. The findings are relevant for guiding the development of malaria vaccines, as well
as immunotherapeutics for alleviating uncomplicated malaria disease and preventing
progression to severe disease.

Methods

Study design and sample collection

In 2019, we conducted a cross-sectional active case detection survey in the Kassena-Nankana Municipality of the Upper East Region of Ghana, to recruit children with uncomplicated malaria and controls with *P. falciparum* infections. About 1000 community members were screened for malaria infection using a CareStart™ PfHRP2-based malaria Rapid Diagnostic Test (RDT, Access Bio, NY, USA). Positive cases in the community were defined as "controls", since these individuals hadn't sought treatment within the past two weeks . Similarly, a passive case detection of uncomplicated malaria cases was carried out using the same mRDTs to screen individuals presenting at the Navrongo War Memorial Hospital outpatient department. Individuals who tested positive for malaria and who provided written informed consent were recruited into the study as defined as "patients". Five milliliters of whole blood was collected for PBMC isolation and for thick and thin blood smears for parasite identification and quantification using microscopy. Linear regression models was used to determine the relatisohsip between parasite density and age in patients and controls in R (version 4.2.1).

Of the 224 individuals recruited in both arms of the study, five control participants and six patients were selected for single-cell transcriptomic analysis. Due to the observed clear differences in clinical presentation between the groups driven by fever, headache and parasite density, the 11 children selected for single-cell analysis had close similarity in these factors. PBMCs were isolated in ACD tubes and spun at 2,000 revolutions per minute (rpm) for 10 minutes, and the leukocytes layer was transferred to 15 mL. The leukocytes were mixed with phosphate-buffered saline (PBS) and layered on 3 mL of lymphoprep in a 15 mL falcon tube. The layered cells were spun for 30 minutes at 800 g without breaks and harvested carefully by taking the buffy layer into another falcon tube. The PBMCs were washed twice with PBS and stored in freezing media. During thawing, complete media (RP10) with 20% Fetal Bovine Serum (FBS) was prepared by diluting 20 mLs of FBS in 80 mL of Roswellpark Memorial Institute (RPMI) media [42]. PBMCs were removed from Liquid Nitrogen to the -80 °C freezer and then thawed during each experiment. Thawing was done by placing the vial in a clean water bath at 37°C until a small crystal of frozen cells was visible. The tubes were cleaned with 70% ethanol, and the contents were transferred to 10 mL of RP10 gently to minimise stressing the cells. The cells were centrifuged at 500g for 10 minutes and resuspended in RP10. Cell viability was estimated using Haemocytometer and PBMCs were used after 1 hr of resting in the incubator at 37 °C.

Seq-Well scRNA-Seq Workflow

Seq-Well scRNA-Seq S^3 workflow was performed according to the published methods [22,43]. In brief, 5×10^5 PBMCs from each patient were dispensed into a single array containing barcoded mRNA capture beads (Supplementary Figure 1). The arrays were sealed with a

Polycarbonate Track Etch (PCTE) membrane (pore size of $0.01\mu M$), allowing cells to remain
separated through the lysis and hybridization steps. mRNA transcripts were hybridized and
recovered for reverse transcription using the Maxima H Minus Reverse Transcriptase in the
first strand synthesis step. Exonuclease (I) was used to remove excess primers and mRNA
was captured via poly-T priming of the poly-A mRNA. The captured mRNA underwent first-
strand synthesis to generate single-stranded cDNA while bound to the beads. Enzymes with
terminal transferase were used to create 3' overhangs and three cytosines. The overhangs
are used in template switching, whereby a SMART sequence is appended to the overhang
on both ends of the cDNA molecule during the first strand synthesis. Some templates fail to
switch, resulting in loss of the mRNA; hence they are chemically denatured using 0.1M
NaOH with random octamer with the SMART sequence in 5' orientation, and a second strand
is synthesized. Whole transcriptome amplification of the cDNA was performed using the
KAPA HiFi PCR master mix (Kapa Biosystems). Libraries were pooled and purified using
AgenCourt AMPure XP Beads. The quality of the library was assessed using Agilent Tape
Station with D5000 High Sensitivity tapes and reagents. Samples were barcoded as
described in the Nextera XT DNA (Illumina, USA) segmentation method. Tagmentation was
important because, after cDNA amplification and clean-up, there are usually very long cDNA
molecules that need to be fragmented to be sequenced by Illumina. The Nextera XT DNA
tagmentation method is effective and allows for the addition of adaptors and multiplex
indexes at both ends of each fragment [22]. Finally, the amplified library was purified using
SPRI beads, pooled, and sequenced using the NextSeq500 kit (Illumina, USA). Paired-end
sequencing was performed with a read structure of 20 bp read one, 50 bp read two, and 8 bp
index one as recommended for Seq-Well. The targeted sequencing depth was 100 million
reads for all samples.

Processing sequencing reads

The raw data were converted to demultiplexed FastQ files using bcl2fastq (Version 5, Terra
Workspace) using the Nextera XT indices and then aligned to the hg19 human genome
using STAR aligner (Version 2.7.9) within the Broad Institute DropSeq workflow (Version 11,
Terra Workspace). The data was cleaned using Cell Bender (V 0.2.0) with default settings, to
remove ambient RNA [44]. The raw expression matrices and sample information were loaded
into the open-source statistical software R (R version 4.2.1). An array with 45,691 gene
features for 22,819 cells described data collected across 11 samples. The data were filtered
to include only features expressed in more than 20 cells, and the resultant matrix described
18,303 gene features across 22,819 cells. A Seurat (Version 4.0) object was created, and
the metadata was added to it to identify the participants [45]. Cell cycle scoring was performed
and computation of the percentage of mitochondria genes before integration. The object from
each participant was transformed individually within the object using SCTransform followed
by the selection of integration features, finding the anchors, and finally combined integration.
Principal component analysis was performed to reduce the dimensionality of the data in
order to identify clusters of cells with similar transcriptomic profiles. Clusters and cluster
resolution were determined using FindNeighbors and a customized FindClusters function
that showed that the best resolution was 0.523, with an average silhouette score of: 0.2 and
11 clusters. One cluster showed no cluster-specific genes and was removed as multiplets,
leaving 18,176 cells. The remaining clusters were reclustered and re-embedded, resulting in
10 clusters with a resolution of 0.292, and an average silhouette score of: 0.301. The
average number of transcripts and expressed genes were evaluated per cluster using half

violin and boxplots. The clusters were projected to a two-dimensional space using the
Uniform Manifold Approximation and Projection (UMAP) [46] algorithm in Seurat.

Reference-based mapping

Immune cell subsets were identified using common cell markers to identify the Mono, T cells,
B cells, NK cells, DC, and other immune cell populations. Uniform Manifold Approximation
and Projection for Dimensional Reduction (UMAP) was used to embed the cell populations
and color code based on the expression of surface markers. The clustered PBMC dataset in
this study (query) was mapped to a reference CITE-Seq dataset of 162,000 PBMCS
measured with 228 antibodies [45]. The query data were projected into the same dimensional
space as the reference dataset, thus separating the cells into the cell types present in the
reference dataset. The method first projected the reference data transformation onto the
query data, followed by the application of KNN-based identification of mutual nearest
neighbors (anchors) between the reference and query. On an L2-normalized dimensional
space, the reference data transferred continuous data onto the query data to annotate the
scRNA data based on a weighted vote classifier. For visualization, reference-based UMAP
embedding was used, considering that all the immune cell populations are well represented.

Analyzing differences in samples

Cluster/sample composition was calculated to determine the proportion of cells per cluster
and per cell type. Cell subsets that were significantly different between patient and control
groups were identified by computing Dirichlet Regression using the DirichReg function in
DirichletReg Package in R [47]. Differentially expressed (DE) genes were computed using the
FindMarkers function on Seurat (Version 4.0), which we used to determine differentially
expressed genes in the patient and control groups using MAST with significance at $(P<0.05)$
and log fold change of > 0.2. Control 4 was not included in the DE analysis due to different
levels of cytokine module scores compared to the other controls participants
(***Supplementary Figure 1e)***. DE genes were visualized using volcano-like plots and
heatmaps to compare all the cell types between patients and controls. The fgsea (R-
package) was used to analyze the pre-ranked gene set enrichment analysis (GSEA). Module
scores for HLA genes, ISG, NFκB target genes, and cytokines were analyzed using the
AddModuleScore function in the Seurat R package. Statistical differences in module scores
between the patients and control groups for each cell subset was computed using Wilcoxon
sign-rank test with Bonferroni correction. Boxplots were used to visualize the module scores
for each cell, denoting the median and interquartile range.

Cell-to-cell Interaction using CellChat

CellChat (Version 1.1.1) was used to quantitatively infer and analyze cell-to-cell
communication networks [21]. Statistically significant intercellular communication between cell
groups was identified using permutation tests, and interactions with a significance level of
less than 0.05 were considered significant [21]. Heatmaps were used to visualize each
signaling pathway and their cell-cell communications, highlighting the number of interactions,
the sources (ligands) of the interactions, and the receivers (receptors) of the interactions.
The relative contribution of each ligand-receptor pair to the overall signaling was shown in
bar plots. The relative contribution provides a measure of a particular ligand-receptor

interaction in a particular cell-cell signaling network. This measure demonstrates the
importance or significance of the interaction in mediating cell communication between the
cell types and potential functional relationships. It is calculated by comparing the expression
levels of different cell receptor and ligand genes between the cell types while accounting for
all the possible interaction pairs within a signaling network.

Acknowledgments

531 We acknowledge the study participants for contributing to the study, and the staff of
532 Navrongo Health Research Centre, who provided support for this study. We are grateful to
533 Felix Ansah, Jersley Chirawurah, and Jonas Kengne, for their contributions to the data
534 collection and critical review of the work. All data were stored and analyzed on the University
535 of Ghana's high-performance computing system (Zuputo).

Author's contributions

537 G.A.A, TDO, LA, YB, and AKS contributed to design and conceptualization; CMM, LA, DA,
538 YA, and NKN contributed to sample collection and processing. VM, RD, and CMM
539 contributed to performing Seq-Well experiments; CMM performed the data analysis and
540 drafted the manuscript. VM, VA, SB, NKN, TDO, LA, and AKS contributed to data analysis
541 and drafting of the manuscript. All authors read and approved the manuscript.

Funding

The study was funded by a DELTAS Africa grant (DEL-15-007: Awandare). The DELTAS
Africa Initiative is an independent funding scheme of the African Academy of Sciences
(AAS)'s Alliance for Accelerating Excellence in Science in Africa (AESA) and supported by
the New Partnership for Africa's Development Planning and Coordinating Agency (NEPAD
Agency) with funding from the Wellcome Trust (107755/Z/15/Z: Awandare) and the UK
government. TDO was supported by the Wellcome Trust (104111/Z/14/Z & A). The study
was also supported by the Ragon Institute of MGH, MIT, and Harvard (AKS). The funders
had no role in the study design and interpretation of the results. The views expressed in this
publication are those of the author(s) and not necessarily those of AAS, NEPAD Agency,
Wellcome Trust, or the UK government.

Availability of the data

554 All data generated or analyzed during this study are included in this published article and its
555 supplementary information files. The scRNA-Seq data are available at https://cellatlas-
556 cxg.mvls.gla.ac.uk/PBMC.Pediatric.Malaria.Ghana.

Ethics approval and consent to participate

558 Ethical clearance was obtained from the Noguchi Memorial Institute of Medical Research,
559 University of Ghana **(IRB 0000908),** and the Ghana Health Service **(GHS-ERC 008/02/19).**
560 All participants provided written informed consent before inclusion in the study.

Consent for publication

562 Not applicable.

Competing interests

References

1. WHO. *World malaria report 2022. World Health Organisation.* (2022).

2. Babiker, H. A., Gadalla, A. A. H. & Ranford-Cartwright, L. C. The role of asymptomatic P. falciparum parasitaemia in the evolution of antimalarial drug resistance in areas of seasonal transmission. *Drug Resist. Updat.* **16**, 1–9 (2013).

3. Bousema, T., Okell, L., Felger, I. & Drakeley, C. Asymptomatic malaria infections: Detectability, transmissibility and public health relevance. *Nat. Rev. Microbiol.* **12**, 833–840 (2014).

4. Nyarko, P. B. & Claessens, A. Understanding Host–Pathogen–Vector Interactions with Chronic Asymptomatic Malaria Infections. *Trends Parasitol.* **37**, 195–204 (2021).

5. Ademolue, T. W., Aniweh, Y., Kusi, K. A. & Awandare, G. A. Patterns of inflammatory responses and parasite tolerance vary with malaria transmission intensity. *Malar. J.* **16**, (2017).

6. Tran, T. M. *et al.* A Molecular Signature in Blood Reveals a Role for p53 in Regulating Malaria-Induced Inflammation. *Immunity* **51**, 750-765.e10 (2019).

7. Boldt, A. B. W. *et al.* The blood transcriptome of childhood malaria. *EBioMedicine* **40**, 614–625 (2019).

8. Mooney, J. P., Wassmer, S. C. & Hafalla, J. C. Type I Interferon in Malaria: A Balancing Act. *Trends in Parasitology* vol. 33 257–260 (2017).

9. Sebina, I. & Haque, A. Effects of type I interferons in malaria. *Immunology* vol. 155 176–185 (2018).

10. de Jong, S. E. *et al.* Systems analysis and controlled malaria infection in Europeans and Africans elucidate naturally acquired immunity. *Nat. Immunol.* **22**, 654–665 (2021).

11. Bediako, Y. *et al.* Repeated clinical malaria episodes are associated with modification of the immune system in children. *BMC Med.* **17**, (2019).

12. Crompton, P. D. *et al.* Malaria immunity in man and mosquito: Insights into unsolved mysteries of a deadly infectious disease. *Annu. Rev. Immunol.* **32**, 157–187 (2014).

13. Ioannidis, L. J., Nie, C. Q. & Hansen, D. S. The role of chemokines in severe malaria: More than meets the eye. *Parasitology* **141**, 602–613 (2014).

14. Kazmin, D. *et al.* Systems analysis of protective immune responses to RTS,S malaria vaccination in humans. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 2425–2430 (2017).

15. Milne, K. *et al.* Mapping immune variation and var gene switching in naive hosts infected with plasmodium falciparum. *Elife* **10**, 1–31 (2021).

16. Loughland, J. R. *et al.* Transcriptional profiling and immunophenotyping show sustained activation of blood monocytes in subpatent Plasmodium falciparum infection. *Clin. Transl. Immunol.* **9**, e1144 (2020).

17. Lee, H. J. *et al.* Transcriptomic Studies of Malaria: a Paradigm for Investigation of Systemic Host-Pathogen Interactions. *Microbiol. Mol. Biol. Rev.* **82**, (2018).

18. Tran, T. M. *et al.* Transcriptomic evidence for modulation of host inflammatory responses during febrile Plasmodium falciparum malaria. *Sci. Rep.* **6**, 1–12 (2016).

19. Ockenhouse, C. F. *et al.* Common and divergent immune response signaling pathways discovered in peripheral blood mononuclear cell gene expression patterns in presymptomatic and clinically apparent malaria. *Infect. Immun.* **74**, 5561–5573 (2006).

20. Hie, B. *et al.* Computational Methods for Single-Cell RNA Sequencing. *Annu. Rev. Biomed. Data Sci.* **3**, 339–364 (2020).

21. Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nat. Commun.* **12**, 1–20 (2021).

22. Hughes, T. K. *et al.* Second-Strand Synthesis-Based Massively Parallel scRNA-Seq Reveals Cellular States and Molecular Features of Human Inflammatory Skin Pathologies. *Immunity* **53**, 878-894.e7 (2020).

23. Villani, A. C. *et al.* Single-cell RNA-seq reveals new types of human blood dendritic
cells, monocytes, and progenitors. *Science (80-.).* **356**, (2017).
24. Amenga-Etego, L. N. *et al.* Temporal evolution of sulfadoxine-pyrimethamine
resistance genotypes and genetic diversity in response to a decade of increased
interventions against Plasmodium falciparum in northern Ghana. *Malar. J.* **20**, 152
(2021).
25. Tusting, L. S. *et al.* Housing Improvements and Malaria Risk in Sub-Saharan Africa: A
Multi-Country Analysis of Survey Data. *PLoS Med.* **14**, (2017).
26. Awandare, G. A. *et al.* Increased levels of inflammatory mediators in children with
severe Plasmodium falciparum malaria with respiratory distress. *J. Infect. Dis.* **194**,
1438–1446 (2006).
27. Cambier, S., Gouwy, M. & Proost, P. The chemokines CXCL8 and CXCL12: molecular
and functional properties, role in disease and efforts towards pharmacological
intervention. *Cell. Mol. Immunol.* **20**, 217–251 (2023).
28. Abrams, E. T. *et al.* Host Response to Malaria During Pregnancy: Placental Monocyte
Recruitment Is Associated with Elevated β Chemokine Expression. *J. Immunol.* **170**,
2759–2764 (2003).
29. Armah, H. B. *et al.* Cerebrospinal fluid and serum biomarkers of cerebral malaria
mortality in Ghanaian children. *Malar. J.* **6**, 1–17 (2007).
30. Ayimba, E. *et al.* Proinflammatory and regulatory cytokines and chemokines in infants
with uncomplicated and severe Plasmodium falciparum malaria. *Clin. Exp. Immunol.*
166, 218–226 (2011).
31. Nahrendorf, W., Ivens, A. & Spence, P. J. Inducible mechanisms of disease tolerance
provide an alternative strategy of acquired immunity to malaria. *Elife* **10**, (2021).
32. Gazzinelli, R. T., Kalantari, P., Fitzgerald, K. A. & Golenbock, D. T. Innate sensing of
malaria parasites. *Nat. Rev. Immunol.* **14**, 744–757 (2014).
33. Quin, J. E. *et al.* Major transcriptional changes observed in the Fulani, an ethnic group
less susceptible to malaria. *Elife* **6**, (2017).
34. Gray, J. C. *et al.* Profiling the antibody immune response against blood stage malaria
vaccine candidates. *Clin. Chem.* **53**, 1244–1253 (2007).
35. Spaulding, E. *et al.* STING-Licensed Macrophages Prime Type I IFN Production by
Plasmacytoid Dendritic Cells in the Bone Marrow during Severe Plasmodium yoelii
Malaria. *PLoS Pathog.* **12**, e1005975 (2016).
36. Awandare, G. A. *et al.* MIF (Macrophage Migration Inhibitory Factor) promoter
polymorphisms and susceptibility to severe malarial anemia. *J. Infect. Dis.* **200**, 629–
637 (2009).
37. Hill, A. V. S. *et al.* Common West African HLA antigens are associated with protection
from severe malaria. *Nature* **352**, 595–600 (1991).
38. Guermonprez, P., Valladeau, J., Zitvogel, L., Théry, C. & Amigorena, S. Antigen
presentation and T cell stimulation by dendritic cells. *Annu. Rev. Immunol.* **20**, 621–
667 (2002).
39. Rock, K. L., Reits, E. & Neefjes, J. Present Yourself! By MHC Class I and MHC Class
II Molecules. *Trends Immunol.* **37**, 724–737 (2016).
40. Kurup, S. P., Butler, N. S. & Harty, J. T. T cell-mediated immunity to malaria. *Nat. Rev.*
Immunol. **19**, 457–471 (2019).
41. Nideffer, J. *et al.* Disease Tolerance Acquired Through Repeated Plasmodium
Infection Involves Epigenetic Reprogramming of Innate Immune Cells. *bioRxiv*
2023.04.19.537546 (2023) doi:10.1101/2023.04.19.537546.
42. Hønge, B. L., Petersen, M. S., Olesen, R., Møller, B. K. & Erikstrup, C. Optimizing
recovery of frozen human peripheral blood mononuclear cells for flow cytometry. *PLoS*
One **12**, 1–17 (2017).
43. Gierahn, T. M. *et al.* Seq-Well: Portable, low-cost rna sequencing of single cells at
high throughput. *Nat. Methods* **14**, 395–398 (2017).
44. Fleming, S. J., Marioni, J. C. & Babadi, M. CellBender remove-background: A deep
generative model for unsupervised removal of background noise from scRNA-seq
datasets. *bioRxiv* 791699 (2019) doi:10.1101/791699.

45.   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-
3587.e29 (2021).
46.   McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and
Projection for Dimension Reduction. (2018).
47.   Simmons, S. Cell Type Composition Analysis: Comparison of statistical methods.
bioRxiv 2022.02.04.479123 (2022).

680
Figures

Figure 1 | Analysis of scRNA-Seq data from uncomplicated malaria patients and
community healthy controls. a) Experimental flow showing that PBMCs were collected
from eleven individuals out of 224 based on the modelling and matching the patients and
controls by age, sex, and parasite density. **b)** Regression analysis between parasite density
and age for patients (grey) and controls (blue). **c)** Uniform manifold approximation and
projection (UMAP) plot of 22,819 cells from eleven participants colored by identities of 10 cell
clusters; mainly B cells, T cells, and Mono. **d)** Expression levels of cluster-defining marker
genes organized by color intensity to show the average expression of the marker in that
particular cell type and the proportion of cells with non-zero expression shown by the size of
the dot. **e)** Markers used to annotate the subclusters to various cell subsets showing
average expression and fraction of cells expressing the marker. **f)** Reference mapped
dataset showing the predicted subclusters of B, CD4 T, CD8 T, NK, Mono, and DC cell
subsets. Reference-defined cell subsets were generated from CITE-seq reference of
162,000 PBMCS measured using 228 antibodies [45]. **g)** UMAP of re-clustered and re-
embedded Mono showing four subclusters of the CD14 and CD16 Mono. **h)** Markers used to
identify monocyte subclusters. **i)** Mono top 10 highly expressed genes in each subcluster.

Figure 2 | Profiling of immune cells from patients compared to the controls. a).
Relative cell proportions of the major cell subsets within patients and control groups.
Statistical tests were conducted using the Dirichlet Multinomial Regression in the
DirichletReg package in R [47]. The dots represent individual proportions while the color
scheme represents the patients and control groups. **b)** Relative proportions of minor cell
subsets compared between patients and controls. Cell proportions per group and P-value are
shown in **Supplementary Table 1. c)** Violin-like plots showing genes that are differentially
expressed between patients and controls. The x-axis shows the Log_2 fold change against the
cell subsets (y-axis) – i.e., B cells, Mono, CD4 T cells, CD8 T cells, other T cells, dendritic
cells (DC) and natural killer **(**NK) cells. The color scheme is based on the upregulated (up
patients) and downregulated (down patients) genes in patients and the size of the point
represents the adjusted P value. The frequency shows the number of comparisons in which
the gene is significantly expressed in the cell subset.

Figure 3 | Pathway analysis using geneset enrichment method a) Pathway analysis
using an immunologic signature geneset enrichment analysis (GSEA) and the color scheme
is based on the normalized enrichment score of genes DEG in patients. **b)** Dot plots showing
some of the leading edge genes in IFN-γ and IFN-α response, TNF-α signaling via NFκB &
inflammatory response pathways in Mono and, **c**) NK cells. Dot size represents the fraction
of cell subsets expressing a given gene. The dot color indicates scaled average expression
by gene column.

Figure 4 | Module score analysis of innate immune gene modules. a) Boxplot showing
interferon-stimulated gene (ISG) module scores per cell subset compared between patients
and controls in B cells, **b)** CD4 T cells, **c)** CD8 T cells, **d)** DC e) Mono, **f)** NK cells. Module
scores are computed using the AddModuleScore function in the Seurat R package.
Statistical significance between the patients and controls of each cell subset was computed
using Wilcoxon sign-rank test with Bonferroni correction. Non significant differences are
indicated by ns.

Figure 5 | Primary innate immune cells dominate the cell-to-cell interactions with other
cell subsets. a) Heatmap showing the number of interactions between the PBMCs cell
subsets. The y-axis shows the signal senders and the x-axis shows the signal receivers. **b)**
Relative contribution of ligand receptor pairs in patients within the MHC class I signaling
pathway and **(c)** MHC class II signaling networks, respectively. A higher relative contribution
indicates the magnitidue of contribution of the ligand-receptor and its significant role in the
MHC I or II signalling networks. **d)** Cell communications through the TNF signaling pathway
and the arrows indicate signal sender to receiver. **e)** Relative contribution of the TNF-
TNFRSF1B ligand-receptor pair towards the TNF signaling pathway. **f)** Violin plots showing
the expression levels of the TNFRSF1B in the Seurat object for the cell subclusters. **g)**
Heatmap comparison showing the overall signaling between all cell subclusters and the
number of interactions. **h)** Relative contribution of MHC class II signaling pathway in the
control group.

Supplementary Figure 1 | Cluster identification and annotation. a) UMAP of 22,819 cells
from all participants, showing 10 clusters in the dataset (following iterative Louvian
clustering). **b)** Dot plots showing genes used to manually annotate the clusters and show the
fraction of cells expressing it and the non-zero expression. Dot size represents the fraction of
cell types (rows) expressing a given gene (columns). The dot color indicates scaled average
expression by gene column. **c**) UMAP colored by various manually annotated clusters based
on the cell markers **d)** UMAP showing cell clusters identified from a reference-mapped
dataset but labeled with the manually annotated cluster identities. **e)** Heatmap showing
overall module score for each cell, and grouped based on each participant and all the cell
types, and overall study groups. The color scheme represents a scale for module scores.

748
749 **Supplementary Tables**
750
Supplementary Table 1: Descriptive statistics of all study participants
Supplementary Table 2: Descriptive statistics of matched study participants for scRNA-Seq
Supplementay Table 3: Dirichlet regression analysis of proportions of various cell types
compared between the two groups
Supplementary Table 4: Differentially expressed genes between patients and controls
Supplementary Table 5: Pathway analysis of differentially expressed genes between
patients and control group
Supplementary Table 6: Leading pathways in cell-cell interaction analysis for the patient
group
Supplementary Table 7: Leading pathways in cell-cell interaction analysis for the control
group
762