Research Article

Acute ischemic Stroke in Tsutsugamushi: Understanding the Underlying Mechanisms and Risk Factors

Dain Kim[1*], Yeon Hee Cho[1*], Ji Yeon Chung, MD[2], Young Seo Kim, MD[3], Byoung-Soo Shin MD, PhD[4,5], Hyun Goo Kang MD, PhD[4,5†]

[1]Jeonbuk National University Medical School, Jeonju, Republic of Korea

[2]Department of Neurology, Chosun University School of Medicine, Gwangju, South Korea

[3]Department of Neurology, Wonkwang University School of Medicine, Iksan, South Korea

4Department of Neurology, Jeonbuk National University Medical School and Hospital, Jeonju, Republic of Korea

[5]Research Institute of Clinical Medicine of Jeonbuk National University Biomedical Research Institute of Jeonbuk National University Hospital, Jeonju, Republic of Korea

* These authors have contributed equally to this work.

Short title: Acute ischemic stroke in tsutsugamushi patients

Word count: 5427 words

[†]**Corresponding author:**

Hyun Goo Kang MD, PhD

Department of Neurology and Research Institute of Clinical Medicine of Jeonbuk National University Biomedical Research Institute of Jeonbuk National University Hospital, 20 Geonji-ro, Deokjin-gu, Jeonju-si, Jeonbuk-do 54907, South Korea

Tel: +82-63-250-1590; Fax: +82-63-251-9363; E-mail: hgkang@jbnu.ac.kr

ORCID: 0000-0001-5443-3635

ABSTRACT

Background: Tsutsugamushi, also known as scrub typhus, is an acute infectious febrile disease common in the Asia-Pacific region. Common symptoms of tsutsugamushi include lymphadenopathy, fever, and myalgia. Although tsutsugamushi rarely causes acute ischemic stroke (AIS), it has been reported that acute infection increased the likelihood of AIS development. Our study hypothesized that tsutsugamushi infection could be a trigger for AIS.

Method: We retrospectively examined patients diagnosed with AIS within 2 weeks of tsutsugamushi diagnosis at three hospitals over a 15-year period. We categorized patients who developed AIS while being treated for tsutsugamushi as the case group; patients who did not develop AIS while being treated for tsutsugamushi at the same hospital and were of similar age and sex to the case group were categorized as the control group. The case and control groups consisted of 22 and 66 participants, respectively. When a scattered pattern was observed or lesions were found in two or more vascular territories on diffusion-weighted imaging, the pattern was defined as embolic. Other patterns were defined as nonembolic.

Results: Among the 19 patients, excluding 3 with TIA, 15 (78.9%) showed an embolic pattern. Although fever was common at the time of the visit in the control group, it was less common in the case group (93.8% vs. 27.3%, p<0.001). Multivariate analysis revealed that higher D-dimer at the time of hospitalization with tsutsugamushi was associated with AIS development in tsutsugamushi patients (adjusted OR, 1.58; 95% CI, 1.06─2.34; p=0.023).

Discussion: AIS in patients with tsutsugamushi showed an embolic rather than a non-embolic pattern on brain MRI. It was more likely to occur in patients with stroke risk factors. These patients were unlikely to have fever or high D-dimer levels. We speculated that D-dimers

played an important role in the pathophysiology, as tsutsugamushi infection increased the likelihood of AIS.

Introduction

Tsutsugamushi, also known as scrub typhus, is an acute infectious febrile disease caused by the bacterium *Orientia tsutsugamushi*; it is prevalent in Asia and the islands of the Pacific and Indian Oceans.[1] Particularly, in 2022, 6,232 patients with tsutsugamushi were reported in South Korea, with approximately 5,000 cases per year in the last five years, thus showing a high prevalence.[2]

Common symptoms of tsutsugamushi include eschar, lymphadenopathy, fever, headache, myalgia, and gastrointestinal symptoms; severe symptoms of the condition include interstitial pneumonia, acute respiratory distress syndrome, meningoencephalitis, acute kidney injury, and disseminated intravascular coagulation.[1] Although acute ischemic stroke (AIS) is a rare symptom of tsutsugamushi, some patients present with both tsutsugamushi and AIS in clinical practice. Similar cases have been reported infrequently. We hypothesized that tsutsugamushi infection could be a causal factor of AIS because previous studies have reported that acute infections, of which tsutsugamushi infection is a type, could increase the occurrence of AIS.[3]. This study aimed to investigate the specific mechanisms underlying the development of AIS in patients with tsutsugamushi.

The main pathophysiology of tsutsugamushi affecting blood vessels involves endothelial damage and vasculitis. However, given that AIS rarely develops in patients with tsutsugamushi, we suspected that there may be other mechanisms besides vasculitis. This study hypothesized that the coagulation disorder induced by tsutsugamushi leads to AIS, based on the results of previous studies, which suggested that tsutsugamushi could cause a disorder of the coagulation pathway[4] and that thrombogenesis is an infection's main mechanism of inducing stroke.[5] To test this hypothesis, we analyzed the data of

tsutsugamushi patients with AIS.

Methods

Patient selection

This study retrospectively evaluated patients who were hospitalized for tsutsugamushi at three regional hub hospitals in different urban areas over a 15-year period (2005-2019) and were diagnosed with AIS within 2 weeks of tsutsugamushi diagnosis. We collated and analyzed the symptoms, medical history, and laboratory findings of the patients at the time of admission using the electronic medical record (EMR). We categorized patients who developed AIS while being hospitalized and treated for tsutsugamushi as the case group and those who did not develop AIS while being hospitalized and treated for tsutsugamushi at the same hospital and were similar in age and sex to the control group. As a result, this study categorized 22 and 66 patients into the case and control groups, respectively. This study was conducted in accordance with the Declaration of Helsinki, and approved by the Institutional Review Board in Jeonbuk National University Hospital (CUH 2023-05-001). The requirement for informed consent was waived owing to the retrospective nature of the study.

Definition

Patients were diagnosed with tsutsugamushi when the *O. tsutsugamushi* gene was detected using polymerase chain reaction (PCR), *O. tsutsugamushi* was identified in the serum, or IgM antibodies against *O. tsutsugamushi* were detected. The AIS group included patients with acute ischemic stroke confirmed by brain magnetic resonance imaging (MRI) and patients with transient ischemic stroke (TIA). However, this study excluded patients with subacute or chronic stroke. TIA was defined as the presence of stroke-related neurological symptoms, such as dysarthria or hemiplegia, which improved within 24 h. Patient symptoms were identified based on the presenting symptoms at the time of hospitalization. This study

examined basic demographic information (age and sex), presence of eschar owing to tsutsugamushi, and various tsutsugamushi-related symptoms (fever, headache, fatigue, anorexia, dyspepsia, nausea, vomiting, abdominal pain, diarrhea, arthralgia, myalgia, sore throat, back pain, dyspnea, cough, and thirst). In addition, we recorded the antibiotics used for tsutsugamushi treatment, measured the time from admission to the development of AIS symptoms, and investigated the risk factors for stroke (hypertension, diabetes mellitus, atrial fibrillation, dyslipidemia, coronary arterial disease, drinking history, and smoking history) at the time of admission. Regarding stroke risk factors, a patient was defined to have a risk factor as an underlying disease if the patient had been diagnosed to have a risk factor and was receiving medication for it or the patient met the relevant diagnostic criteria after hospitalization even though the patient had no diagnosis of a risk factor. Hypertension was defined as a state when the blood pressure measured at rest during hospitalization was □130/80 mmHg.[6] Diabetes mellitus was diagnosed if the 8-h fasting blood glucose level was 126 mg/dL or higher, HbA1c was 6.5% or higher, or blood glucose was 200 mg/dL or higher after 2 h of oral glucose tolerance test.[7] Atrial fibrillation included all patients who had atrial fibrillation confirmed by electrocardiogram performed after admission or those previously diagnosed with atrial fibrillation. Dyslipidemia was diagnosed if low density lipoprotein (LDL) was 160 mg/dL or higher, triglyceride was 200 mg/dL or higher, high density lipoprotein (HDL) was 40 mg/dL or lower, or total cholesterol was 240 mg/dL or higher.[8-10] Coronary artery disease was diagnosed if abnormal findings were confirmed by coronary angiography. If a patient's response showed that the patient had consumed alcohol or cigarettes within 1 year of hospitalization, the patient was defined as having a drinking or smoking risk factor. aspartate aminotransferase (AST), alanine aminotransferase (ALT), white blood cell (WBC), hemoglobin (Hg), and platelet (PLT), which are laboratory findings

generally known to show abnormal values in tsutsugamushi patients,[11] and prothrombin time (PT), activated partial thromboplastin time (aPTT), D-dimer, fibrinogen, and fibrin degradation product (FDP), which are laboratory findings related to blood coagulation that may affect ischemic stroke, were investigated based on blood tests at the time of hospitalization.

Imaging analysis

This study investigated the pattern of lesions caused by stroke by collecting and analyzing brain MRIs of patients admitted for tsutsugamushi and diagnosed with acute ischemic stroke or TIA during treatment. Subsequently, we examined the presence of atrial fibrillation and D-dimer levels, which are factors believed to be associated with embolic patterns. When a scattered pattern (indicating that a high signal intensity is scattered over multiple territories) was observed on diffusion weighted imaging (DWI) or when there were lesions in two or more different vascular territories, it was defined as an embolic pattern. Other types were defined as non-embolic.

Statistical analysis

First, a control group was selected based on age and sex matching of patients with a disposition similar to that of the case group. Age and sex matching was performed using propensity score matching and the nearest neighbor matching algorithm, and 66 patients were selected as the control group using a 1:3 matching ratio. Afterward, we compared the demographics, clinical symptoms, and stroke risk factors of patients with tsutsugamushi infection with or without AIS. We used Pearson's chi-squared or Fisher's exact tests to analyze categorical variables and the Student's t-test was used for continuous variables. Continuous variables were summarized as means with standard deviations or medians with

interquartile ranges, whereas categorical data were expressed as counts and percentages. Multivariate analysis was performed to identify the independent factors associated with tsutsugamushi presenting with AIS. To avoid variable selection caused by spurious correlations, only variables showing a potential association (p<0.1) in univariate analysis were included as potential factors associated with tsutsugamushi presenting with AIS in the multivariate logistic regression model. Statistical significance was set at p<0.05 (two-tailed). All statistical analyses were performed using SPSS 21.0 (IBM Corporation, Armonk, NY, USA).

Results

Among the 22 tsutsugamushi patients who developed AIS, 14 were females (63.6%) and 8 were males (36.4%); mean age was 74.4 ± 13.4 years, with the youngest patient being 26 years old and the oldest being 93 years old (Table 1). Antibiotics used for the treatment of tsutsugamushi included doxycycline in 16 patients (72.7%), azithromycin in four patients (18.2%), and rifampicin in two patients (9.1%). The average time from the onset of tsutsugamushi symptoms to that of AIS was 3.91 ± 4.54 days; meanwhile, the time of onset in four patients was □10 days. In addition, factors associated with vascular risk were current alcohol consumption in two (9.1%) participants, current smoking status in two (9.1%), hypertension in 12 (54.5%), dyslipidemia in seven (31.8%), diabetes mellitus in five (22.7%), atrial fibrillation in five (22.7%), and coronary artery disease in two (9.1%).

This study described the brain MRI characteristics of 19 patients with tsutsugamushi and AIS, excluding three patients with TIA in whom no lesions were observed on DWI (Table 2). Among the 19 patients, 15 (78.9%) showed an embolic pattern, whereas four (21.1%) patients had a non-embolic pattern, indicating that an embolic pattern was more common. Sixteen (84.2%) patients had elevated D-dimer levels above the normal level (0.5 mg/L) and six (31.6%) patients had atrial fibrillation as an underlying condition. Among the 16 patients with elevated D-dimer levels, 13 (81.3%) had an embolic pattern and five (31.3%) had atrial fibrillation. Among the six patients with atrial fibrillation, five had elevated D-dimer levels and one did not; all six patients showed an embolic pattern. None of the 19 patients had a history of cancer.

This study compared patients with tsutsugamushi infection, with and without AIS (Table 3). The mean age of the target group was 74.41 ± 13.36 years, compared with 74.18 ± 12.81

years of the control group. At the time of admission, fever was commonly observed in tsutsugamushi patients without AIS but was significantly less common in tsutsugamushi patients with AIS (93.8 vs. 27.3%, respectively; p<0.001). In addition, headache was significantly less common in tsutsugamushi patients with AIS than in those without AIS (69.8 vs. 31.8%; p=0.002). The incidence of sore throat (31.8 vs. 9.1%; p=0.036), dyspnea (21.2 vs. 0%; p=0.018), and thirst (82.9 vs. 25.0%; p<0.001) was lower in tsutsugamushi patients with AIS than in those without AIS. Tsutsugamushi patients with AIS were more likely to have stroke risk factors than those without AIS. Particularly, tsutsugamushi patients with AIS were significantly more likely to have hypertension (16.1 vs. 54.5%; p=0.001), diabetes mellitus (5.3 vs. 22.7%; p=0.021), atrial fibrillation (4.5 vs. 22.7%; p=0.01), and dyslipidemia (7.6% vs. 31.8%; p=0.004) than did tsutsugamushi patients without AIS. Regarding laboratory findings, tsutsugamushi patients with AIS showed significantly higher aPTT (32.31 ± 12.27 vs. 40.54 ± 13.47; p=0.009), D-dimer (0.87 ± 1.27 vs. 8.14 ± 10.47; p=0.005), and FDP (2.27 ± 0.69 vs. 42.01 ± 59.55; p=0.009) than did tsutsugamushi patients without AIS. The results in Table 3 are presented as a bar graph (Figure 1).

The present study performed multivariate analysis to identify factors associated with the development of AIS in patients with tsutsugamushi (Table 4). Hypertension, diabetes mellitus, atrial fibrillation, and dyslipidemia, which are known risk factors for stroke, did not show significant results. However, higher D-dimer level at the time of hospitalization owing to tsutsugamushi was found to affect the development of AIS in tsutsugamushi patients (adjusted OR, 1.58; 95% CI, 1.06–2.34; p=0.023).

Discussion

This retrospective study aimed to determine the effects of tsutsugamushi infection on AIS onset. The co-occurrence of tsutsugamushi and AIS is rare and only a few case reports have reported their co-occurrence.[12,13] This study collected the data of 22 tsutsugamushi patients with AIS from three hub university hospitals over a 15-year period, compared them with the data of 66 tsutsugamushi patients without AIS (control), and identified the relationship between tsutsugamushi infection and the development of AIS.

Among the 22 patients who developed AIS after tsutsugamushi infection, four patients developed AIS more than 10 days after the onset of tsutsugamushi symptoms (Table 1). The diagnosis of AIS was delayed for Patient 1 because although the patient had syncope, a symptom of ischemic stroke, two days before the diagnosis of tsutsugamushi, it was before hospitalization. Diagnosis was delayed in Patients 10 and 11 because they were not hospitalized in the neurology department, which delayed their neurological evaluation. Patient 14 developed AIS shortly after tsutsugamushi treatment and was discharged from hospital, resulting in a late diagnosis. AIS was diagnosed in 18 patients after tsutsugamushi infection for < 10 days (Table 1).

According to a recent multi-institute registry study in South Korea, in patients with stroke due to "other determined etiology" such as hypercoagulable state, the prevalence of stroke risk factors was 47.5, 23.8, 20.6, 3.2, and 21.2% for hypertension, dyslipidemia, diabetes, atrial fibrillation, and current smoker, respectively.[14] The results of our study showed that the prevalence of risk factors was 54.5, 31.8, 22.7, 22.7, 9.1, and 9.1% for hypertension, dyslipidemia, diabetes mellitus, atrial fibrillation, coronary artery disease, and current

smoking, respectively, which was generally higher than those of stroke patients due to "other determined etiology" (Table 3). This difference was likely due to the tsutsugamushi patients with AIS in this study having an infectious disease (tsutsugamushi). Because people with multiple concomitant diseases or poor immunity due to an underlying disease are more likely to have an infectious disease, the patients in the present study might have had more underlying diseases than the general population.

The incidence of fever was significantly lower in tsutsugamushi patients with AIS than in those without AIS at the time of admission (93.8 vs. 27.3%, p<0.001; Table 3), which was believed to be related to the sequence of symptoms resulting from tsutsugamushi infection. Fever, one of the most common symptoms of tsutsugamushi, appears relatively early after infection,[15] whereas coagulopathy is a late manifestation of the illness.[12] Considering the sequence of these symptoms and the fact that fever is a mechanism for protecting the body against infection,[16] the development of fever early in the course of tsutsugamushi infection is thought to trigger the body's defense mechanism against *O. tsutsugamushi* in the early stages of infection. However, we suspect that patients who did not develop fever early in the course of their infection did not adequately deal with the bacteria; instead, the disease progressed to a coagulation disorder, which is a late manifestation. This hypothesis agreed with previous reports that disseminated intravascular coagulation was more common in tsutsugamushi patients with severe illness than those with less severe illness.[4] In addition to fever, the incidence rate of headache, sore throat, dyspnea, and thirst was significantly lower in tsutsugamushi patients with AIS than in those without AIS (Table 3, Figure 1). This finding was expected because headache, sore throat, dyspnea, and thirst are secondary symptoms of fever.

Regarding stroke risk factors, tsutsugamushi patients with AIS had significantly higher rates of hypertension, diabetes mellitus, atrial fibrillation, and dyslipidemia than those without AIS (Table 3 and Figure 1). These results suggest that tsutsugamushi infection is more likely to cause AIS in patients with stroke risk factors than in those without the risk factors. Several previous studies have reported that common acute infections trigger ischemic stroke.[3,17] However, we believe that the triggering action of these infections targets stroke risk factors. For example, if a patient with underlying atrial fibrillation is infected with tsutsugamushi, a thrombus is more likely to develop due to an increase in heart rate in response to the infection, thus increasing the likelihood of AIS development. This can be explained by the results of previous studies that showed that thrombus formation was the pathogenesis of atrial fibrillation to induce AIS,[18] heart rate increased during infection, and when the resting heart rate increased, the risk of incident venous thromboembolism and inflammatory and coagulation factors increased.[19] This means that tsutsugamushi infection acts as a trigger, further aiding the patient's pre-existing risk factors to cause AIS. Our hypothesis needs to be confirmed in a larger, systematic study in the future.

The aPTT, D-dimer, and FDP levels in the target patient group were significantly higher than those in the control group (p=0.009, 0.005, and 0.009, respectively). Elevated aPTT, D-dimer, and FDP levels were associated with coagulation disorders. Therefore, the results suggested that the mechanism of tsutsugamushi infection to induce AIS was related to coagulation diseases, which agreed with the results of previous studies.[4] Furthermore, from multivariate logistic analysis, we identified D-dimer elevation as a factor influencing the development of AIS after tsutsugamushi infection (adjusted OR, 1.58; 95% CI, 1.06 - 2.34; p=0.023). Dardiotis et al.[20] reported that the secretion of inflammatory cytokines by cancer

triggered an inflammatory response, which created a hypercoagulable state, resulting in systemic and cerebral arterial or venous thrombosis that could induce ischemic stroke. In addition, it was reported that in patients with such cancer-related strokes, D-dimer elevation was observed, and brain images showed infarction of multifocal lesions.[20] Moreover, Jackson et al.[21] reported that inflammation and hemostatic response were both activated in response to infection. Based on the results of these previous studies and the present study, we believed that tsutsugamushi patients with a higher D-dimer level were more likely to develop AIS than those with a lower D-dimer level at the time of admission and progress to an embolic stroke due to their hypercoagulable state. This study examined D-dimer levels in 19 of 22 AIS patients with tsutsugamushi infection, after excluding three patients without imaging information, and found that the D-dimer level of 16 patients exceeded the threshold of 0.5 mg/L (Table 2). Among the 16 patients, 13 showed an embolic pattern on brain MRI, which supported our hypothesis (Table 2). However, we could not conclude whether higher D-dimer levels induced embolic stroke. Because atrial fibrillation, another risk factor, can cause a cardioembolic stroke by abnormally promoting thromboembolism,[22] the five patients with atrial fibrillation, elevated D-dimer level ($\geq$ 0.5 mg/dL), and an embolic pattern lesions on DWI could have developed cardioembolic stroke due to atrial fibrillation. However, the eight patients who showed embolic-pattern lesions with D-dimer elevation but without atrial fibrillation could serve as evidence for our proposed hypothesis. Although previous studies have reported that tsutsugamushi infection can elevate coagulation factors such as D-dimer and cause DIC to induce bleeding,[4] its association with ischemic stroke has not been reported. Additionally, some previous studies have hypothesized that tsutsugamushi infection-induced coagulopathy affects the onset of stroke,[12,13] but all of these studies were case reports. We believe that the novelty of this study lies in its evaluation of the association between

tsutsugamushi infection and AIS by analyzing brain images and data.

This study had several limitations. First, we could not control for all confounding variables because it was a retrospective study. Therefore, this study could not regulate variables other than tsutsugamushi infection that could cause AIS. Future prospective cohort studies are needed to improve the reliability of our results. Second, because the data were collected at a tertiary hospital, many patients had severe diseases, which could have introduced selection bias. However, because most people in South Korea visit tertiary hospitals when they suspect stroke, it is difficult to find cases of co-occurrence of tsutsugamushi and AIS in small primary care clinics. Therefore, we believe that the results will be similar even after including patients from primary and secondary hospitals. Third, it is necessary to exclude the possible influence of atrial fibrillation history on the stroke pattern to explain the correlation between D-dimer and embolic patterns in the stroke pattern of tsutsugamushi patients. However, this study could not compare patients with atrial fibrillation history and those without atrial fibrillation history. It is necessary to conduct a histological study to demonstrate that excessive thrombin production due to DIC, rather than atherosclerosis, causes embolic stroke. However, there was a limitation in that histological samples could not be obtained from stroke patients. Finally, the sample size was small because the co-occurrence of tsutsugamushi and AIS is a rare condition. There could have been a bias in the statistics because of the small sample size. For the same reason, we could not perform a statistical analysis to determine whether the brain MRI embolic pattern of the target group was associated with D-dimer levels. However, it was meaningful to collect data from 22 patients because no other study has collected data from as many tsutsugamushi patients with AIS.

Conclusion

AIS in tsutsugamushi patients tended to appear in an embolic pattern, rather than in a non-embolic pattern, on brain MRI and occurred more easily in patients with many stroke risk factors. Moreover, these patients were less likely to have a fever and had higher D-dimer levels. We believe that D-dimer could play an important role in the pathophysiology of tsutsugamushi infection and increase the occurrence of AIS. In this study, we believe that Tsutsugamushi infection increased the likelihood of developing AIS in individuals with stroke risk factors. We suspect that if a patient does not develop fever early in the course of tsutsugamushi infection, the patient may develop AIS after a few days. In conclusion, we believe that if a tsutsugamushi patient has an elevated D-dimer level or does not have fever on admission, the possibility of AIS should be considered.

Acknowledgments

Funding

This work was supported by the Technology Development Program (S3174187), funded by the Ministry of SMEs and Startups (MSS, Korea).

Disclosures

None

References

1. Walker DH. Scrub typhus - scientific neglect, ever-widening impact. *N Engl J Med*. 2016;375:913-915. doi:10.1056/NEJMp1608499.

2. Agency KDCaP. *Surveillance Statistics of Infectious Diseases for Week 10, 2023*. Public Health Weekly Report. 16; 2023.

3. Sebastian S, Stein LK, Dhamoon MS. Infection as a stroke trigger: associations between different organ system infection admissions and stroke subtypes. *Stroke*. 2019;50:2216-2218. doi:10.1161/STROKEAHA.119.025872.

4. Lee HJ, Park CY, Park SG, Yoon NR, Kim DM, Chung CH. Activation of the coagulation cascade in patients with scrub typhus. *Diagn Microbiol Infect Dis*. 2017;89:1-6. doi:10.1016/j.diagmicrobio.2017.06.011.

5. Grau AJ, Boddy AW, Dukovic DA, Buggle F, Lichy C, Brandt T, Hacke W; CAPRIE Investigators. Leukocyte count as an independent predictor of recurrent ischemic events. *Stroke*. 2004;35:1147-1152. doi:10.1161/01.STR.0000124122.71702.64.

6. Brook RD, Rajagopalan S. *ACC/AHA/AAPA/ABC/ACPM/AGS. APhA/ASH*; 2017*/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. A report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines*. Journal of the American Society of Hypertension: JASH, 2018. 12(3): p. 238-238.

7. ElSayed NA, Aleppo G, Aroda VR, Bannuru RR, Brown FM, Bruemmer D, Collins BS, Hilliard ME, Isaacs D, Johnson EL, et al. 2. Classification and Diagnosis of Diabetes: Standards of Care in Diabetes-2023. *Diabetes Care*. 2023;46 Supplement 1: S19-S40. doi:10.2337/dc23-S002.

8. Grundy SM, Stone NJ, Bailey AL, Beam C, Birtcher KK, Blumenthal RS, Braun LT, de

Ferranti S, Faiella-Tommasino J, Forman DE, et al. AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA guideline on the management of blood cholesterol: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation*. 2019. 2018;139:e1082-e1143.

9. Ishibashi S, Yamashita S, Arai H, Araki E, Yokote K, Suganami H, Fruchart JC, Kodama T; K-877-04 Study Group. Effects of K-877, a novel selective PPARα modulator (SPPARMα), in dyslipidaemic patients: a randomized, double blind, active- and placebo-controlled, phase 2 trial. *Atherosclerosis*. 2016;249:36-43. doi:10.1016/j.atherosclerosis.2016.02.029.

10. Wiviott SD, Raz I, Bonaca MP, Mosenzon O, Kato ET, Cahn A, Silverman MG, Zelniker TA, Kuder JF, Murphy SA, et al. Dapagliflozin and cardiovascular outcomes in type 2 diabetes. *N Engl J Med*. 2019;380:347-357. doi:10.1056/NEJMoa1812389.

11. Kim D-M. Clinical features and diagnosis of scrub typhus. *Infect Chemother*. 2009;41:315-322. doi:10.3947/ic.2009.41.6.315.

12. Chung JH, Yun NR, Kim DM, Lee JW, Yoon SH, Kim SW. Reportscrub typhus and cerebrovascular injury: A phenomenon of delayed treatment? *Am J Trop Med Hyg*. 2013;89:119-122. doi:10.4269/ajtmh.13-0094.

13. Park JH, Lee DK, Kwon O-H. Cerebral Infarction associated with scrub typhus. *Korean J Stroke*. 2007;9:168-170.

14. Kim H, Kim JT, Lee JS, Kim BJ, Kang J, Lee KJ, Park JM, Kang K, Lee SJ, Kim JG, Cha JK, Kim DH, et al. Stroke of other determined etiology: results from the nationwide multicenter stroke registry. *Stroke*. 2022;53:2597-2606. doi:10.1161/STROKEAHA.121.037582.

15. Peter JV, Sudarsan TI, Prakash JA, Varghese GM. Severe scrub typhus infection: clinical features, diagnostic challenges and management. *World J Crit Care Med*. 2015;4:244-250. doi:10.5492/wjccm.v4.i3.244.

16. Tan CL, Knight ZA. Regulation of body temperature by the nervous system. *Neuron*. 2018;98:31-48. doi:10.1016/j.neuron.2018.02.022.

17. Das S, Ghosh KC, Pulai S, Pulai D, Bhar D, Gangopadhyay PK. Systemic infection and inflammation as trigger factors of ischemic stroke. *Ann Neurosci*. 2011;18:17-20. doi:10.5214/ans.0972.7531.1118106.

18. Violi F, Pastori D, Pignatelli P. Mechanisms and management of thrombo-embolism in atrial fibrillation. *J Atr Fibrillation*. 2014;7:1112. doi:10.4022/jafib.1112.

19. Awotoye J, Fashanu OE, Lutsey PL, Zhao D, O'Neal WT, Michos ED. Resting heart rate and incident venous thromboembolism: the Multi-Ethnic Study of Atherosclerosis. *Open Heart*. 2020;7:e001080. doi:10.1136/openhrt-2019-001080.

20. Dardiotis E, Aloizou AM, Markoula S, Siokas V, Tsarouhas K, Tzanakakis G, Libra M, Kyritsis AP, Brotis AG, Aschner M, et al. Cancer-associated stroke: pathophysiology, detection and management (Review). *Int J Oncol*. 2019;54:779-796. doi:10.3892/ijo.2019.4669.

21. Jackson SP, Darbousset R, Schoenwaelder SM. Thromboinflammation: challenges of therapeutically targeting coagulation and other host defense mechanisms, The Journal of the American Society of Hematology. *Blood*. 2019;133:906-918. doi:10.1182/blood-2018-11-882993.

22. Lip GY. Does atrial fibrillation confer a hypercoagulable state? *Lancet*. 1995;346:1313-1314. doi:10.1016/s0140-6736(95)92339-x.

Figure legends

Figure 1. Comparison of symptoms and stroke risk factors between controls (tsutsugamushi without stroke) and patients (tsutsugamushi with stroke). Conventional symptoms of tsutsugamushi with and without acute ischemic stroke (A). Gastrointestinal symptoms of tsutsugamushi with and without acute ischemic stroke (B). Other symptoms of tsutsugamushi with and without acute ischemic stroke (C). Stroke risk factors in controls and patients (D). A statistically significant variant is expressed as *, ** or ***. * means p-value <0.05, and ** means p-value <0.01, and *** means p-value <0.001.

Table 1. Baseline characteristics of patients with tsutsugamushi and acute ischemic stroke

	Sex	Antibiotics	Symptom onset to stroke (day)	Alc	Smk	HTN	DM	A.fib	DL	CAD
1	F	Dox	14	0	0	0	0	0	0	0
2	F	Dox	1	0	0	1	1	1	1	0
3	M	Rip	1	0	1	0	0	0	0	0
4	M	Rip	9	0	0	1	1	0	1	1
5	M	Dox	1	0	1	0	0	0	0	0
6	F	Dox	1	0	0	0	0	1	0	0
7	F	Dox	1	0	0	1	0	0	1	0
8	F	Dox	1	0	0	1	1	0	1	0
9	M	Azi	1	0	0	1	0	1	0	1
10	F	Dox	11	0	0	0	0	0	0	0
11	F	Dox	10	0	0	1	0	0	1	0
12	M	Dox	1	0	0	1	0	0	1	0
13	F	Dox	5	0	0	1	0	0	0	0
14	F	Dox	13	0	0	0	0	0	0	0
15	M	Dox	1	0	0	0	0	0	0	0
16	F	Dox	3	0	0	1	1	0	1	0
17	F	Dox	1	0	0	1	0	1	0	0
18	M	Dox	0	1	0	0	1	0	0	0
19	M	Azi	7	1	0	0	0	0	0	0
20	F	Azi	1	0	0	1	0	1	0	0
21	F	Dox	0	0	0	1	0	0	0	0
22	F	Azi	3	0	0	0	0	0	0	0

F, female; M, male; Dox, doxycycline; Rip, rifampicin; Azi, azithromycin; Alc, alcohol; Smk, smoking; HTN, hypertension; DM, diabetes mellitus; A.fib, atrial fibrillation; DL, dyslipidemia; CAD, coronary artery disease

Table 2. Diffusion Weighted Imaging of a patient with acute ischemic stroke, with tsutsugamushi and stroke patterns (embolic, non-embolic) and risk factors of the patients

Patient	MR image	lesion pattern	Atrial fibrillation	D-dimer (mg/L)
1		non-embolic	-	0.279
2		embolic	+	5.003

3

 | embolic | - | 257.136 |
| 4 | | embolic | + | 1.050 |

5		embolic	-	1.570
6		embolic	+	1.152

| 7 |

 | embolic | - | 0.438 |

8

  embolic - 2.057

9		embolic	+	0.210

| 10 |

 | non-embolic | - | 34.32 |
| 11 | | embolic | - | 1.4 |

| 12 |

 | embolic | - | 18.96 |
| 14 |

 | embolic | - | 6.27 |

15   embolic   -   5

16

 non-embolic - 9.48

| 17 | | embolic | + | 5.83 |
| 19 | | non-embolic | - | 5.74 |

21

 embolic   +   2.67

| 22 |

 | embolic | - | 17.47 |

Table 3. Comparison of tsutsugamushi with and without acute ischemic stroke

	Tsutsugamushi without stroke (n = 66)	Tsutsugamushi with stroke (n = 22)	P value
Female	39 (59.1)	14 (63.6)	0.706
Age	74.18 ± 12.81	74.41 ± 13.36	0.943
Eschar	52 (83.9)	17 (77.3)	0.524
Symptoms			
Fever	61 (93.8)	6 (27.3)	<0.001
Headache	44 (69.8)	7 (31.8)	0.002
Fatigue	39 (59.1)	17 (77.3)	0.125
Anorexia	46 (69.7)	13 (59.1)	0.359
Dyspepsia	16 (41.0)	7 (33.3)	0.559
Nausea	15 (29.4)	4 (18.2)	0.316
Vomiting	6 (11.8)	3 (13.6)	1
Abdominal pain	5 (7.6)	1 (4.5)	1
Diarrhea	2 (3.0)	2 (9.1)	0.259
Arthralgia	14 (22.2)	5 (23.8)	1
Myalgia	38 (60.3)	12 (54.5)	0.636
Sore throat	21 (31.8)	2 (9.1)	0.036
Back pain	7 (10.6)	2 (9.1)	1
Dyspnea	14 (21.2)	0 (0.0)	0.018
Cough	15 (23.8)	2 (10.5)	0.335
Thirst	29 (82.9)	4 (25.0)	<0.001
Stroke risk factors			
Hypertension	9 (16.1)	12 (54.5)	0.001
Diabetes mellitus	3 (5.3)	5 (22.7)	0.021
Atrial fibrillation	3 (4.5)	5 (22.7)	0.01
Dyslipidemia	5 (7.6)	7 (31.8)	0.004
Coronary artery disease	3 (5.5)	2 (9.1)	0.612
Alcohol	2 (5.1)	2 (9.1)	0.615
Smoking	3 (7.9)	2 (9.1)	1
Laboratory findings			
PT	12.47 ± 1.06	14.18 ± 7.53	0.302
aPTT	32.31 ± 12.27	40.54 ± 13.47	0.009
D-dimer	0.87 ± 1.27	8.14 ± 10.47	0.005
Fibrinogen	292.96 ± 98.15	252.62 ± 106.69	0.132
FDP	2.27 ± 0.69	42.01 ± 59.55	0.009

AST	194.18 ± 860.00	99.82 ± 113.20	*0.611*
ALT	140.67 ± 629.12	58.95 ± 55.11	*0.546*
WBC	8.37 ± 3.48	9.54 ± 4.62	*0.213*
Hg	12.23 ± 1.85	11.49 ± 2.02	*0.117*
PLT	170.40 ± 94.30	176.55 ± 165.69	*0.831*
Creatinine	1.28 ± 0.71	1.44 ± 0.67	*0.373*

PT, prothrombin time; aPTT, activated partial thromboplastin time; FDP, fibrinogen degradation product; AST, aspartate aminotransferase; ALT, alanine aminotransferase; WBC, white blood cell; Hg, hemoglobin; PLT, platelet;

Table 4. Multivariate analysis of factors associated with tsutsugamushi with acute ischemic stroke

	Crude OR (95% CI)	P value	Adjusted OR (95% CI)	P value
Hypertension	6.27 (2.08 - 18.85)	0.001	2.64 (0.24 - 29.65)	0.432
DM	5.29 (1.15 - 24.49)	0.033	5.86 (0.53 - 64.50)	0.149
A.fib	5.29 (1.14 - 24.48)	0.033	4.55 (0.61 - 33.75)	0.139
Dyslipidemia	4.48 (1.24 - 16.21)	0.022	0.95 (0.05 - 17.94)	0.971
D-dimer	1.74 (1.22 - 2.47)	0.002	1.58 (1.06 - 2.34)	0.023

DM, diabetes mellitus; A.fib, atrial fibrillation.

(A)

Conventional symptom

(B)

GI symptom

(C)

Other symptom

(D)

Stroke risk factor