HIV self-testing positivity rate and linkage to confirmatory testing and care: a telephone survey in Côte d'Ivoire, Mali and Senegal

Arsène Kouassi Kra[1*], Arlette Simo Fotso[1,2], Nicolas Rouveau[1], Mathieu Maheu-Giroux[3], Marie-Claude Boily[4], Romain Silhol[4], Marc d'Elbée[1,5], Anthony Vautier[6] and Joseph Larmarange[1,2] on behalf of the ATLAS team

[1] Centre Population et Développement (Ceped), Université Paris Cité, Institut de Recherche pour le Développement (IRD), Inserm, France
[2] Institut National d'études Démographiques (Ined), France
[3] Department of Epidemiology and Biostatistics, School of Population and Global Health, McGill University, Montréal, QC, H3A 1A2, Canada
[4] MRC Centre for Global Infectious Disease Analysis, School of Public Health, Imperial College London, London, United Kingdom
[5] National Institute for Health and Medical Research UMR 1219, Research Institute for Sustainable Development EMR 271, Bordeaux Population Health Centre, University of Bordeaux, France
[6] Solidarité Thérapeutique et Initiatives pour la Santé (Solthis), Sénégal

*Corresponding author
Correspondence: arsene.krakouassi@ceped.org

ABSTRACT

HIV self-testing (HIVST) empowers individuals by allowing them to decide when and where to test and with whom to share their results. From 2019 to 2022, the ATLAS program distributed ~ 400 000 HIVST kits in Côte d'Ivoire, Mali, and Senegal. It prioritised key populations, including female sex workers and men who have sex with men, and encouraged secondary distribution of HIVST to their partners, peers and clients.

To preserve the confidential nature of HIVST, use of kits and HIVST results were not systematically tracked. Therefore, an anonymous phone survey was conducted to estimate HIVST positivity rates and linkage to confirmatory testing and care. This two-step survey involved an initial recruitment phase from March to June 2021 where participants were encouraged via leaflets to call a free phone number and complete a sociobehavioural questionnaire.

This was followed by a second phase in September and October 2021, where participants who reported a reactive HIVST result were re-contacted to complete a further questionnaire. Of the 2 615 participants recruited during the first phase, 89.7% reported consistent results (2 visible lines and result interpreted as reactive; one line and interpreted as non-reactive; or no/one line and interpreted as invalid). HIVST positivity rates varied between 2.4% to 9.1% based on calculation methods (i.e. self-interpreted result or reported number of lines, inclusion or exclusion of don't knows and refusals).

The second phase saw 78 out of 126 eligible participants complete the questionnaire. Of the 27 who reported a consistent reactive result in the first phase, 15 (56%, 95%CI: 36 to 74%)

underwent confirmatory HIV testing, with 12 (80%) confirmed as HIV-positive, all of whom began antiretroviral treatment.

The confirmation rate of HIVST results was fast, with 53% doing so within a week and 91% within three months of self-testing. Two-thirds (65%) went to a general public facility, and one-third to a facility dedicated to key populations.

The ATLAS HIVST distribution strategy reached people living with HIV in West Africa. Linkage to confirmatory testing remained sub-optimal in these first years of HIVST implementation. However, if confirmed HIV-positive, almost all initiated treatment. HIVST constitutes a relevant complementary tool to existing screening services.

Keywords: AIDS, HIV self-testing, HIV testing services, diagnosis, knowledge of status, awareness fo status, gay, bisexual, sex work, linkage to confirmatory testing and care, phone-based survey, key populations, West Africa.

Introduction

Early testing followed by successful linkage to antiretroviral treatment for those diagnosed with HIV can drastically reduce the risk of onward HIV transmission and mortality [1–6]. In 2021, according to the United Nations Program for HIV/AIDS (UNAIDS), 81% of the adult population living with HIV in West and Central Africa knew their status. Only 77% of them were on antiretroviral treatment [7], below the 95-95-95 UNAIDS targets for 2030 [8].

HIV self-testing (HIVST) is the process by which a person who wants to know their HIV status collects their own sample (oral fluid or blood), performs the test, and then interprets the results themself, often in a private setting [9]. It is an innovative tool that empowers individuals and guarantees the confidentiality of the test [10]. Individuals may decide when and where to test and with whom they want to share their result. It has been shown to be effective in screening populations vulnerable to HIV acquisition and transmission that are often hardly reached through conventional approaches [11–13]. Since 2016, the World Health Organization (WHO) has recommended HIVST as a complementary testing approach [14].

Following the experience gained in Eastern and Southern Africa under the STAR project [15–21], the UNAIDS funding agency sought to stimulate HIVST in West Africa. The ATLAS programme (*AutoTest de dépistage du VIH : Libre d'Accéder à la connaissance de son Statut*) aimed to promote, implement, and expand HIV self-testing in Côte d'Ivoire, Mali, and Senegal. Country national prevalence was comparatively low in West Africa in 2021: 1.9% (1.7%-2.2%) in Côte d'Ivoire, 0.8% (0.6%-1%) in Mali, and 0.3% (0.3%-0.4%) in Senegal according to Unaids[22].

In the ATLAS project's catchment areas, HIVST was integrated into existing testing policies, programs, and activities. A total of 397 367 HIVST kits were distributed free of charge between July 2019 and February 2022 as part of the national AIDS strategy in these three countries. At the time of ATLAS' implementation, in 2019, only small-scale pilot studies on HIVST had previously been conducted in Senegal and Côte d'Ivoire, and none existed in Mali.

The design of the different delivery channels and the priority populations were developed with country stakeholders: national AIDS programs/councils, international institutions including the WHO, international and national non-governmental organisations involved in local HIV programs, and civil society and community leaders. ATLAS HIVST distribution was organised through eight different operational delivery channels (figure 1): five were facility-based (delivery of HIVST kits through public or community-based health facilities), and three used community-based approaches involving outreach activities engaging female sex workers (FSW), men who have sex with men (MSM), and people who use drugs (PWUD) [23]. Peer educators conducted these outreach activities through group activities (e.g. talks, discussion groups, night visits, social events, or parties) and face-to-face activities (e.g. home visits). Outreach activities represented the majority (~85%) of ATLAS's distribution volume.

ATLAS activities relied on both primary and secondary distribution. HIVST kits were distributed by peer educators and healthcare professionals to primary contacts for their personal use (primary distribution). With secondary distribution, primary contacts were provided HIVST kits and invited to redistribute them to their peers, partners, and clients. These secondary contacts were often members of key populations that can be more difficult to engage in HIV prevention, along with other peripheral vulnerable populations. This chain-referral distribution of HIVST implies that end-users were not limited to primary contacts.

Only oral self-testing (OraQuick HIV Self-Test®) has been distributed through ATLAS. OraSure Technologies, the manufacturer of the OraQuick test, accompanies each HIVST kit with a user manual for result interpretation. OraQuick HIVST should be interpreted as follow: reactive if two lines (C & T) are visible, even barely; non-reactive if only the C (control) line is visible; invalid if no line is visible or if only the T (test) line is visible.

In addition to the manufacturer's instructions (figure 2), locally adapted brochures and explanatory videos in French and local languages have been developed to help users perform the test, interpret the result and know what actions should be taken following a non-reactive, a reactive or indeterminate result. They also encouraged people with a reactive HIVST to seek confirmatory HIV testing and care. Free phone lines have been set up in each country, and operators of these lines were trained about HIVST.

Figure 2. Guidelines for interpreting HIVST result, extracted from the English version of the manufacturer instructions for use (OraQuick HIV Self-Test®)

To preserve the anonymity and confidentiality of HIVST and not impede their use, ATLAS has decided, in line with WHO recommendations, not to track the use and outcomes of distributed HIVST kits directly. Such tracking can be logistically challenging and costly and could limit the distribution, redistribution and use of HIVST [24]. Without systematic tracking, it was challenging to obtain information on who was using the HIVST, the results of the tests and the linkage to confirmatory testing and treatment. These are crucial indicators to assess program effectiveness and impact. For instance, the positivity rate can be related to the yield of new individuals diagnosed with HIV and would suggest that the testing modality is indeed reaching those in need. Diagnosed individuals must seek confirmatory testing and be linked to care to maximise health benefits and decrease onward transmission.

We conducted an innovative survey by setting up an anonymous and free telephone platform in Côte d'Ivoire, Mali and Senegal while preserving anonymity and encouraging voluntary participation. This survey among ATLAS HIVST users showed that HIVST secondary distribution was feasible and acceptable. Participants reported that they appreciated the ease of use of HIVST, its discretion and the fact that they are autonomous in carrying out the test. Finally, HIVST appeared as a relevant additional approach for those

134 usually distant from community activities and HIV testing services, and has the potential to reach, beyond
135 key populations, partners, clients, and other groups vulnerable to HIV [25].
136   A complementary survey was conducted among those with an HIVST reactive result. Here we report on
137 HIVST's positivity rates and linkage to confirmatory testing and care.
138

Materials and Methods

Sources of data

The ATLAS program embedded multiple research activities, from qualitative studies to economic
analyses, which have been described in detail elsewhere [26–30].
It included a voluntary phone survey. Between mid-March and mid-June 2021, dedicated survey flyers
were distributed with the HIVST kits inviting self-test users in each country to call a toll-free number to
complete a questionnaire (phase 1). All calls from the three countries were rerouted to a telephone platform
located in Abidjan and operated by Ipsos Côte d'Ivoire, which was selected following an international call for
tenders. The questionnaire, which lasted 20 to 30 minutes, collected information on sociodemographic
characteristics of HIVST users, testing history, sexual and preventive behaviours, HIVST use and difficulties
encountered. Participation in the survey was rewarded with 2 000 XOF (≈3.40 USD) of phone communication
credit. In order to participate in the survey, participants had to be of legal age to use an HIVST on their own
without parental permission (16 years in Côte d'Ivoire, 18 years in Mali, and 15 years in Senegal) and had to
have used an HIVST provided to them through the ATLAS project.
As the survey was anonymous, there was a risk that some HIVST users may participate more than once or
that individuals who have never used HIVST tried to participate to receive the financial incentive. To limit
these risks, several measures were taken: (i) the leaflet distributed with the HIVST kits had a unique 9-digit
number generated by the research team that was requested prior to participation in the survey; (ii) the same
unique number could not be used twice; (iii) the financial incentive was only paid out once the questionnaire
was fully completed (however individuals could refuse to answer any particular question); (iv) the same
telephone number could not be used twice to receive the telephone credit. These unique 9-digit numbers
were generated non-sequentially and were grouped by country, delivery channel and implementing partner.
Thus, any unique number could indirectly identify the delivery channel where the HIVST kit was initially
dispensed.
In total, 2 615 participants were recruited for phase 1 [25]. During the interviews, each participant was
asked about the number of lines that appeared when reading the HIVST result and their self-interpretation of
it (reactive or non-reactive). Those who reported two lines or a reactive result were asked for their consent
to be called back a few months later to participate in a complementary survey and, if consented, to provide a
phone contact.
Between September 27th and October 22th 2021, eligible participants who agreed to be re-contacted
were phoned to complete a 5-minute questionnaire (phase 2) on linkage to confirmatory testing and care.
The interviews were conducted in either French, English, Bambara, or Wolof. On-the-fly translation into
other local languages was also available. Compensation of XOF 2 000 (≈3.40 USD) in the form of telephone
credit was given to participants who completed the phase 2 questionnaire. Unlike in phase 1, it was not a
financial incentive: participants were informed about it only at the end of the interview. Interviews were not
audio-recorded. Questionnaires' responses were captured on a computer and stored in a database managed
by PAC-CI, an Ivorian research institute with expertise in clinical research.
At the end of the survey, collected telephone numbers (for appointments and rewards) were deleted
from the database. All procedures have been described in a publicly available data management plan
(https://dmp.opidor.fr/plans/3354/export.pdf).

Data analysis

Based on phase 1 participants' self-reports, we distinguished those having reported HIVST results
consistent with both the reported number of visible lines and the reported self-interpretation (2 visible lines
and result interpreted as reactive; one line and interpreted as non-reactive; or no/one line and interpreted
as invalid), an inconsistent result, or a partial result (they refused to answer or answered they didn't know to
one or both questions).

To estimate HIVST positivity rates, we separately considered the self-interpreted results and the reported
number of lines on the HIVST. For each source, we made three hypotheses (low, central and high) about
"don't know" and refusals (DK-R). Using self-reported results, the low hypothesis considered DK-R as non-
reactive, and the high hypothesis as reactive, while DK-R were excluded from both the numerator and the
denominator in the central hypothesis. Using the reported number of visible lines, the low hypothesis
considered DK-R as one line, and the high hypothesis as two lines, while DK-R were excluded from both the
numerator and the denominator in the central hypothesis. Positivity rates were stratified by respondents'
gender, country, and distribution channel.

We described the selection of eligible participants for phase 2 questionnaires and corresponding
participation rates. To assess any participation bias, characteristics of phase 2 participants were compared
with individuals eligible for phase 2 but who did not participate and with phase 1 participants not eligible for
phase 2.

Among phase 2 eligible participants who completed their questionnaire, linkage to confirmatory testing,
the proportion being confirmed HIV positive and treatment initiation were described, stratified by the
reported number of lines and self-interpreted HIVST result in phase 1 questionnaire. Confidence intervals
(95% confidence interval, 95%CI) were computed using Wilson's method with Yate's continuity correction.

We also explored (i) for those who did not link to confirmatory testing, the main reported reason; and (ii)
for those who did link to confirmatory testing, the type of facility was confirmatory testing was performed
and the time between HIVST and confirmatory testing.

All analyses have been performed using R version 4.2.2 [31]. A dedicated anonymised dataset and the
corresponding R script are available on Zenodo (https://doi.org/ 10.5281/zenodo.7986077) to allow
replication of the analysis.

. Ethics
ATLAS research protocol (version 3.0, October 8 2020) has been approved by the WHO Ethical Research
Committee (January 12, 2021, reference: ERC 0003181), the National Ethics Committee for Life Sciences and
Health of Côte d'Ivoire (November 27, 2020, reference: 191-20/MSHP/CNESVS-km, IRB:000111917), the
Ethics Committee of the Faculty of Medicine and Pharmacy of the University of Bamako, Mali (November 16,
2020, reference: 2020/254/CE/FMPOS/FAPH), and the National Ethics Committee for Health Research of
Senegal (January 26, 2021, protocol SEN19/32, n°8 MSAS/CNERS/Sec).

215   **Results**

HIVST results
Of the 2 615 participants recruited in phase 1, 2 346 (89.7%) reported a self-interpreted HIVST result
consistent with their reported number of visibles lines on the HIVST: 2 292 (88.0%) reported one line and
self-interpreted it as non-reactive, 50 (1.9%) two lines and reactive and 4 (0.2%) none or one line self-
interpreted as invalid (table 1). In contrast, 48 (1.8%) reported inconsistent answers: 10 (0.4%) one line and
self-interpreted it as reactive, 35 (1.3%) two lines and non-reactive and 3 (0.1%) no line and non-reactive.

Finally, 221 (8.5%) reported a partial result: 147 (5.6%) reported 0, 1 or 2 lines, but did not know how to
interpret the result or refused to answer; 46 (1.7%) self-interpreted their result, but did not know or refused
to report the number of lines; and 28 (1.1%) did not know or refused to answer to both questions

Table 1. Self-reported HIV self-test (HIVST) result, reported number of lines on the HIVST, and positivity rates according to different hypotheses among participants of the first phase of the survey in Côte d'Ivoire, Mali, and Senegal (2021).

Phase 1 participants	2615 (100%)
Consistent answer (C)	*2346 (89.7%)*
2 lines / reactive[†] (C_1)	50 (1.9%)
1 line / non-reactive (C_2)	2 292 (88%)
0-1 line/ invalid (C_3)	4 (0.2%)
Inconsistent answer (I)	*48 (1.8%)*
1 line / reactive[†] (I_1)	10 (0.4%)
0 line / reactive[†] (I_2)	0 (0%)
2 lines/ non-reactive[†] (I_3)	35 (1.3%)
0 line / non-reactive (I_4)	3 (0.1%)
2 lines/ invalid[†] (I_5)	0 (0%)
Partial answer (P)	*221 (8.5%)*
0 line / DK-R (P_1)	1 (<0.1%)
1 line / DK-R (P_2)	117 (4.5%)
2 lines/ DK-R[†] (P_3)	29 (1.1%)
DK-R / reactive[†] (P_4)	2 (<0.1%)
DK-R / non-reactive (P_5)	44 (1.7%)
DK-R / invalid (P_6)	0 (0%)
DK-R / DK-R (P_7)	28 (1.1%)
Positivity Rate	
Based on self-interpreted test results	
Low hypothesis (DK-R as not reactive) $(C_1 + I_1 + I_2 + P_4) / n$	62 / 2615 (2.4 %)
Central hypothesis (DK-R excluded) $(C_1 + I_1 + I_2 + P_4) / (C + I + P_4 + P_5 + P_6)$	62 / 2440 (2.5 %)
High hypothesis (DK-R as reactive) $(C_1 + I_1 + I_2 + P_1 + P_2 + P_3 + P_4 + P_7) / n$	237 / 2615 (9.1 %)
Based on the reported number of lines	
Low hypothesis (DK-R as 1 line) $(C_1 + I_3 + I_5 + P_3) / n$	114 / 2615 (4.4 %)
Central hypothesis mid (DK-R excluded) $(C_1 + I_3 + I_5 + P_3) / (C + I + P_1 + P_2 + P_3)$	114 / 2541 (4.5 %)
High hypothesis (DK-R as 2 lines) $(C_1 + I_3 + I_5 + P_3 + P_4 + P_5 + P_6 + P_7) / (C + I + P_1 + P_2 + P_3)$	188 / 2615 (7.2 %)

†: Eligible for phase 2 Survey
DK: don't know. R: refused to answer

HIVST positivity rates

Based on self-interpreted HIVST results, the overall positivity rate was 2.4% when DK-R were considered non-reactive (Table 2, low hypothesis). By excluding DK-R from the numerator and the denominator (central hypothesis), the positivity rate increased to 2.5%. Considering DK-R as reactive (high hypothesis) increased the positivity rate to 9.1%. Estimates based on the reported number of visible lines on the HIVST were 4.4%, 4.5% and 7.2%, respectively, for low, central and high hypotheses. Positivity rates ranged from 1.8% to 9.8% in Côte d'Ivoire, 3.5% to 7.8% in Mali, and 1.2% to 15.0% in Senegal depending on the hypothesis (e.g., low or high; Table 2).

Positivity rates were higher among participants recruited through community-based distribution channels: 4.8% for men and 4.9% for women in the MSM-based channels, and 4.6% for men and 4.2% for women in the FSW-based channels (central hypothesis based on the number of lines). In the other distribution channels (PWUD-based and facility-based), positivity rates were slightly lower: 3.1% for men and 2.9% for women.

Participation in phase 2
During phase 1, 126 individuals reported two lines or self-interpreted their result as reactive, thus were
identified as eligible for phase 2 (table 1). Among them, 6 had refused to be re-contacted after phase 1
(figure 3). Among the 120 (95%) who agreed to be re-contacted, 24 (20%) were unreachable at the time of
the phase 2 survey, and 96 (80%) were successfully re-contacted. Among the latest, 89 (93%) accepted to
participate in the phase 2 survey. Ten dropped out before the end of the interview, and 1 disconnected and
was unreachable afterwards. As a result, 78 participants completed phase 2 questionnaire.
The participants who completed the phase 2 questionnaire had similar sociodemographic characteristics
(e.g. country, sex, distribution channel, age group, marital status) compared to those eligible for phase 2, but
that did not complete it, and to phase 1 participants not eligible for phase 2 (table S1). For most participants
(86%), phase 2 questionnaire was completed between 4 and 6 months after phase 1 questionnaire (table S5).
253
254

256   **Figure 3.** Flow chart of the participant selection process for the 2[nd] phase of the survey in Côte d'Ivoire,
257   Mali, and Senegal (2021).

Table 2. Positivity rates based on self-interpreted HIVST results or the reported number of visible lines, by distribution channel, gender and country, among participants of the first survey phase in Côte d'Ivoire, Mali, and Senegal (2021).

			MSM-based channels		FSW-based channels		Others delivery channels		Total
			Man	Woman	Man	Woman	Man	Woman	
Positivity rate based on self-reported HIVST results	Low	Côte d'Ivoire	2.5% (16/650)	1.4% (1/73)	1.5% (5/339)	1.2% (3/245)	0% (0/60)	0% (0/23)	1.8% (25/1 390)
		Mali	4.6% (14/306)	0% (0/29)	1.9% (5/269)	3.9% (14/360)	9.1% (1/11)	0% (0/9)	3.5% (34/984)
		Senegal	4.9% (2/41)	0% (0/1)	0% (0/12)	0% (0/80)	0% (0/66)	2.4% (1/41)	1.2% (3/241)
		Overall	3.2% (32/997)	1.0% (1/103)	1.6% (10/620)	2.5% (17/685)	0.7% (1/137)	1.4% (1/73)	2.4% (62/2 615)
	Central	Côte d'Ivoire	2.7% (16/597)	1.4% (1/71)	1.6% (5/311)	1.4% (3/221)	0% (0/58)	0% (0/21)	2.0% (25/1 279)
		Mali	4.7% (14/301)	0% (0/29)	1.9% (5/257)	4.1% (14/345)	9.1% (1/11)	0% (0/9)	3.6% (34/952)
		Senegal	6.1% (2/33)	0% (0/1)	0% (0/11)	0% (0/65)	0% (0/61)	2.6% (1/38)	1.4% (3/209)
		Overall	3.4% (32/931)	1.0% (1/101)	1.7% (10/579)	2.7% (17/631)	0.8% (1/130)	1.5% (1/68)	2.5% (62/2 440)
	High	Côte d'Ivoire	10.6% (69/650)	4.1% (3/73)	9.7% (33/339)	11% (27/245)	3.3% (2/60)	8.7% (2/23)	9.8% (136/1 390)
		Mali	6.2% (19/306)	0% (0/29)	6.3% (17/269)	8.1% (29/360)	9.1% (1/11)	0% (0/9)	6.7% (66/984)
		Senegal	24.0% (10/41)	0.0% (0/1)	8.3% (1/12)	19.0% (15/80)	7.6% (5/66)	9.8% (4/41)	15.0% (35/241)
		Overall	9.8% (98/997)	2.9% (3/103)	8.2% (51/620)	10.0% (71/685)	5.8% (8/137)	8.2% (6/73)	9.1% (237/2 615)
Positivity rate based on the reported number of visible lines	Low	Côte d'Ivoire	4.2% (27/650)	5.5% (4/73)	4.7% (16/339)	2.0% (5/245)	0% (0/60)	4.3% (1/23)	3.8% (53/1 390)
		Mali	4.9% (15/306)	3.4% (1/29)	4.5% (12/269)	5.3% (19/360)	9.1% (1/11)	0% (0/9)	4.9% (48/984)
		Senegal	12.2% (5/41)	0% (0/1)	0% (0/12)	5.0% (4/80)	4.5% (3/66)	2.4% (1/41)	5.4% (13/241)
		Overall	4.7% (47/997)	4.9% (5/103)	4.5% (28/620)	4.1% (28/685)	2.9% (4/137)	2.7% (2/73)	4.4% (114/2 615)
	Central	Côte d'Ivoire	4.2% (27/641)	5.5% (4/73)	4.8% (16/331)	2.1% (5/241)	0% (0/60)	4.5% (1/22)	3.9% (53/1 368)
		Mali	5.0% (15/298)	3.4% (1/29)	4.5% (12/264)	5.5% (19/344)	9.1% (1/11)	0% (0/9)	5.0% (48/955)
		Senegal	13.2% (5/38)	0% (0/1)	0% (0/10)	5.3% (4/75)	5.3% (3/57)	2.7% (1/37)	6.0% (13/218)
		Overall	4.8% (47/977)	4.9% (5/103)	4.6% (28/605)	4.2% (28/660)	3.1% (4/128)	2.9% (2/68)	4.5% (114/2 541)
	High	Côte d'Ivoire	5.5% (36/650)	5.5% (4/73)	7.1% (24/339)	3.7% (9/245)	0% (0/60)	8.7% (2/23)	5.4% (75/1 390)
		Mali	7.5% (23/306)	3.4% (1/29)	6.3% (17/269)	9.7% (35/360)	9.1% (1/11)	0% (0/9)	7.8% (77/984)
		Senegal	19.5% (8/41)	0% (0/1)	16.7% (2/12)	11.2% (9/80)	18.2% (12/66)	12.2% (5/41)	14.9% (36/241)
		Overall	6.7% (67/997)	4.9% (5/103)	6.9% (43/620)	7.7% (53/685)	9.5% (13/137)	9.6% (7/73)	7.2% (188/2 615)

DK: don't know. R: refusals. FSW: female sex workers, MSM: men having sex with men, PR: positivity rate. Shaded cells indicated cells with less than 50 participants.
Low hypothesis: DK-R as non-reactive or 1 line. Central hypothesis: DK-R excluded from the numerator and the denominator. High hypothesis: DK-R as reactive or 2 lines.

Linkage to confirmatory testing and care

Overall, 34 of the 78 participants of phase 2 survey (44%) reported having performed confirmatory testing. Linkage was higher for those who reported 2 lines and correctly self-interpreted their result as reactive: 56% (95%CI: 36-74%), followed by those who reported two lines but did not know or refused to report their test interpretation with a linkage rate of 44% (95%CI: 22-69%). Not surprisingly, linkage to confirmatory testing was lower (36%; 95%CI: 19-57%) among those who reported 2 lines but incorrectly self-interpreted the result as non-reactive (table 3). Finally, among the 8 participants who reported none or one line or did not know how many lines and incorrectly self-interpreted the result as reactive, only 2 linked to confirmatory testing.

The main reason given for not linking to confirmatory testing was that "their HIVST was non-reactive" (18/44, 41%, and although 8 of these 18 reported a reactive result in phase 1 questionnaire); followed by "not knowing that a confirmation test was required" (10/44, 23%); and "not having time" (8/44, 18%) (table S2).

When participants were linked to confirmatory testing, it was a short time after performing their HIVST: 53% linked in less than one week and 91% in less than 3 months (table S3).

Most participants (65%) performed their confirmatory testing in a general public facility (health centre, hospital, clinic or maternity), and 35% chose a community-based clinic or health centre dedicated to key populations (table S4).

Among the 34 linked to confirmatory testing, 19 (56%, 95%CI: 38-72%) were confirmed HIV-positive, and 18 (95%, 95%CI; 72-100%) initiated antiretroviral treatment. Among the 27 who reported a consistent reactive result in the phase 1 questionnaire, 15 (56%, 95%CI: 36-74%) linked to confirmatory test, 12 (80%) were confirmed HIV-positive and all started treatment (100%).

289　**Table 3**. Linkage to confirmatory testing, proportion being confirmed HIV positive and treatment initiation, by reported number of lines and self-interpreted HIVST
290　result among eligible participants of the second phase of the survey who completed their questionnaire in Côte d'Ivoire, Mali, and Senegal (2021).

Reported number of lines/ self-interpreted HIVST result	Completed phase 2	Linked to confirmatory testing		Confirmed HIV positive		Initiated ART	
	n	n (%)	95%CI	n (%)	95%CI	n (%)	95%CI
Overall	78	34 (44%)	33% to 55%	19 (56%)	38% to 72%	18 (95%)	72% to 100%
2 lines / reactive	27	15 (56%)	36% to 74%	12 (80%)	51% to 95%	12 (100%)	70% to 100%
1 line / reactive	7	1 (14%)	1% to 58%	0 (0%)	0% to 80%		
2 lines / non-reactive	25	9 (36%)	19% to 57%	3 (33%)	9% to 69%	3 (100%)	31% to 100%
2 lines / DK-R	18	8 (44%)	22% to 69%	4 (50%)	22% to 78%	3 (75%)	22% to 99%
DK-R / reactive	1	1(100%)	5% to 100%	0 (0%)	0% to 95%		

291　DK: don't know. R: refuse to answer. CI: confidence interval.
292

Discussion

Our study reveals that the strategy deployed by the ATLAS program, through secondary distribution
of HIVST and targeted channels, successfully reached people living with HIV in West Africa. However,
linkage to confirmatory HIV testing and access to care remained sub-optimal during these initial years of
HIVST implementation. Moreover, among participants who confirmed their reactive self-test result with a
traditional facility-based HIV test, a significant proportion quickly proceeded with this confirmation (more
than half in less than a week and the vast majority in less than three months). Furthermore, if individuals
were confirmed HIV-positive, almost all began antiretroviral treatment.
The implementation of a telephone survey, aimed at gathering information from HIVST users while
preserving anonymity and without interfering with secondary distribution, has proven to be very useful
to evaluate the ATLAS project. However, its high cost makes it difficult to integrate it into national
strategies for assessing the impact of HIVST. Nevertheless, other impact evaluation methods, such as
data triangulation[27] and modelling, may prove more suitable for routine monitoring of HIVST's impacts.
In our study, most participants (90%) demonstrated a consistent interpretation between the number
of lines reported and the declared result of their HIVST. However, 2% of them reported an inconsistent
interpretation of the results. Among them, a small number reported the presence of two visible lines,
suggesting potential issues in interpreting the number of visible lines on HIVST kits. In the context of the
ATLAS program, the distribution strategy combining primary and secondary approaches has led many
HIVST users to perform their HIVST without receiving advice from a healthcare professional or a trained
peer educator. Although the HIVST is not designed to require supervision, it is essential to have received
information on its use before proceeding with the test. However, the lack of supervision is likely
insufficient to explain the inconsistencies observed [32]. Some inconsistencies may result from a
misunderstanding of the terms "*reactive*" and "*non-reactive*", particularly considering that HIVST was a
new tool in our context and that traditional terms used to describe conventional HIV testing are
"*positive*" and "*negative*". This possible misunderstanding of the terms is also highlighted by the fact that
8 participants reported a "*reactive*" result in phase 1 questionnaire and then in phase 2 that their test
was "*non-reactive*" as the main reason for not linking to confirmatory testing. Specific qualitative
interviews or focus groups discussion with HIVST users could help better understand how they perceive
different terms.
According to our estimates, HIVST positivity rates range from 1.8% to 9.8% in Côte d'Ivoire, 3.5% to
7.8% in Mali, and 1.2% to 15.0% in Senegal depending on how missing results (e.g., "*don't know*" and
refusals) are classified. It is important to interpret these HIV positivity rates while considering the
treatment-adjusted prevalence (i.e., removing those on treatment from the numerator and denominator
of HIV prevalence), a more reliable indicator for evaluating the effectiveness of targeted screening
programs [33]. In West Africa, the treatment-adjusted prevalence remained relatively low in 2021: 0.6%
in Côte d'Ivoire, 0.7% in Mali, and 0.06% in Senegal, according to UNAIDS data
(https://aidsinfo.unaids.org/). Our results suggest that the ATLAS HIVST distribution strategy successfully
reached people living with HIV. In 2021, the observed HIV positivity rates for conventional HIV testing
were 1.4% in Côte d'Ivoire, 2.2% in Mali, and 1.0% in Senegal [34]. These rates were slightly lower to
those we identified with HIVST based on our lower (conservative) assumption. Between 2020 and 2021,
ATLAS implementing partners collected spontaneous feedback from HIVST users. This unpublished data
collection was non-systematic and varied from one partner to another. Among 4 463 documented
feedbacks, HIVST was reactive for 188 cases (4.2%), consistent with our estimates based on the reported
number of visible lines (4.5%, central hypothesis).
Overall, these results for HIVST positivity are generally higher than the average overall positivity of
HIV testing services (excluding HIVST) in West Africa. For instance, in 2020 an estimated 1.9% of all HIV
tests performed were found to be positive in the region (95% credible intervals: 1.3 to 2.7%) [34].
Further, among 15-24 and 25-34 years old, which constitute more than 80% of our sample, overall
positivity was, respectively, 0.9% (0.7 to 1.3%) and 1.6% (1.2 to 2.2%). Collectively, these results provide
evidence that HIVST is a high-yield testing modality that can address the unmet HIV testing needs of key
populations and their partners.
Our linkage to confirmatory testing estimates were based on small numbers resulting in large
confidence intervals. Nevertheless, the overall proportion was clearly sub-optimal (44%, 95% confidence

interval from 33% to 55%). However, this estimate includes some individuals who did not adequately self-
interpreted their HIVST result as reactive. When considering only those who reported two lines and self-
interpreted their result as reactive, the linkage rate increased to 56% (36% to 74%). This percentage is
close to that was observed in a study conducted in Kenya on HIV testing of FSW male partners using
HIVST secondary distribution, where 65% of men with a reactive result had a confirmatory test [35].
Linkage to confirmatory testing happened relatively quickly after HIVST use: 53% did it in less than a week
and 91% in less than three months. Similar results were observed in a study in the general population in
Zambia[36], and a study among MSM in Nigeria [37]. For those who did confirmatory testing and were
confirmed HIV positive, initiation of antiretroviral treatment was almost systematic, showing good
linkage to care after confirmatory testing, as observed in many HIVST studies in sub-Saharan Africa [38–
40].

Previous analyses of ATLAS data showed that HIVST could reach people not reached by conventional
HIV testing approaches [41], particularly partners and clients of key populations and key population
members not self-identifying as such [25]. It is consistent with the finding that two-thirds of participants
who did confirmatory testing went to a general health facility rather than a community clinic dedicated to
key populations. In a study conducted in 2018 in Côte d'Ivoire among MSM, one-third of the participants
preferred community-based testing, one-third expressed no preference, and one-third preferred
undifferentiated HIV testing services (general population), mentioning the lack of discretion and
anonymity of community-based sites and the desire to avoid the gaze of others [42].

ATLAS' HIVST distribution strategy successfully reached people living with HIV in West Africa,
although linkage to confirmatory testing remained sub-optimal in these first years of HIVST
implementation. We showed that HIVST has the potential to reach more hidden populations and
constitutes a relevant complementary tool to existing screening services. To fully harness the potential of
self-tests, messaging around HIVST and its interpretation could be improved.
370

371 **Appendices**

372 Table S1. Eligibility and participation in phase 2 survey by sociodemographic characteristics, distribution channel, HIV testing history, the reported number of lines and
373 the self-interpreted HIV self-testing (HIVST) result

	Overall N= 2 615 (phase 1 participants)	Eligible for phase 2 completed the questionnaire N=78	Eligible for phase 2 but did not complete the questionnaire N= 42	Not eligible for Phase 2 N = 2495	p-value (Chi² test)
Country					0.9
Côte d'Ivoire	1 390 (53%)	39 (50%)	20 (48%)	1 331 (53%)	
Mali	984 (38%)	31 (40%)	18 (43%)	935 (37%)	
Senegal	241 (9.2%)	8 (10%)	4 (9.5%)	229 (9.2%)	
Sex and distribution channel					0.3
man MSM-based channels	997 (38%)	35 (45%)	14 (33%)	948 (38%)	
woman MSM-based channels	103 (3.9%)	5 (6.4%)	0 (0%)	98 (3.9%)	
man FSW-based channels	620 (24%)	22 (28%)	10 (24%)	588 (24%)	
woman FSW-based channels	685 (26%)	14 (18%)	15 (36%)	656 (26%)	
man other-based channels	137 (5.2%)	1 (1.3%)	2 (4.8%)	134 (5.4%)	
woman other-based channels	73 (2.8%)	1 (1.3%)	1 (2.4%)	71 (2.8%)	
Age group					0.5
24 years or younger	1 164 (45%)	27 (35%)	20 (48%)	1 117 (45%)	
25-34 years	1 063 (41%)	38 (49%)	16 (38%)	1 009 (40%)	
35 years or older	388 (15%)	13 (17%)	6 (14%)	369 (15%)	
Marital status					0.3
single	1 761 (67%)	54 (69%)	28 (67%)	1 679 (67%)	
living with partner / married	757 (29%)	18 (23%)	12 (29%)	727 (29%)	
divorced / separated / widowed	97 (3.7%)	6 (7.7%)	2 (4.8%)	89 (3.6%)	
Educational level					0.057
none / primary	503 (19%)	13 (17%)	10 (24%)	480 (19%)	
secondary	1 432 (55%)	50 (64%)	28 (67%)	1 354 (54%)	
higher	680 (26%)	15 (19%)	4 (9.5%)	661 (26%)	
First time testers					0.3
yes	1 078 (41%)	38 (49%)	20 (48%)	1 020 (41%)	
No	1 537 (59%)	40 (51%)	22 (52%)	1 475 (59%)	

	Overall N= 2 615 (phase 1 participants)	Eligible for phase 2 completed the questionnaire N=78	Eligible for phase 2 but did not complete the questionnaire N= 42	Not eligible for Phase 2 N = 2495	p-value (Chi² test)
Result and number line					<0.001
2 lines / reactive	50 (1.9%)	27 (35%)	20 (48%)	3 (0.1%)	
1 line / not reactive	2 292 (88%)	0 (0%)	0 (0%)	2 292 (92%)	
0 line / not reactive	3 (0.1%)	0 (0%)	0 (0%)	3 (0.1%)	
2 lines/ not reactive	35 (1.3%)	25 (32%)	9 (21%)	1 (<0.1%)	
1 line / reactive	10 (0.4%)	7 (9.0%)	3 (7.1%)	0 (0%)	
0-1 line/ invalid	4 (0.2%)	0 (0%)	0 (0%)	4 (0.2%)	
2 lines/ DK-R	29 (1.1%)	18 (23%)	9 (21%)	2 (<0.1%)	
1 line / DK-R	117 (4.5%)	0 (0%)	0 (0%)	117 (4.7%)	
0 line / DK-R	1 (<0.1%)	0 (0%)	0 (0%)	1 (<0.1%)	
DK-R / reactive	2 (<0.1%)	1 (1.3%)	1 (2.4%)	0 (0%)	
DK-R / not reactive	44 (1.7%)	0 (0%)	0 (0%)	44 (1.8%)	
DK-R / DK-R	28 (1.1%)	0 (0%)	0 (0%)	28 (1.1%)	

375   FSW: female sex workers, MSM: men having sex with men

377
378 Table S2. Main reason for not linking to confirmatory testing among phase 2 participants who did not link to confirmatory testing, by reported number of lines and self-interpreted HIVST result.

	Overall	2 lines / reactive	1 line / reactive	2 lines / non-reactive	2 lines / DK-R
My test was non-reactive	18 (41%)	6 (50%)	2 (33%)	5 (31%)	5 (50%)
I didn't know he should get a confirmatory test	10 (23%)	2 (17%)	2 (33%)	5 (31%)	1 (10%)
I didn't have time	8 (18%)	3 (25%)	0 (0%)	3 (19%)	2 (20%)
I feared that others would know the result	2 (4.5%)	0 (0%)	0 (0%)	1 (6.2%)	1 (10%)
I already knew the result before using HIVST	2 (4.5%)	1 (8.3%)	1 (17%)	0 (0%)	0 (0%)
I had no specific reason	2 (4.5%)	0 (0%)	1 (17%)	1 (6.2%)	0 (0%)
I didn't know where to take the test	1 (2.3%)	0 (0%)	0 (0%)	1 (6.2%)	0 (0%)
The testing site was too far away	1 (2.3%)	0 (0%)	0 (0%)	0 (0%)	1 (10%)
Total	44 (100%)	12 (27.3%)	6 (13.6%)	16 (36.4%)	10 (22.7%)

379   DK: don't know. R: refuse to answer

382 383 Table S3. Time between HIVST and confirmatory testing among phase 2 participants who did link to confirmatory testing, by reported number of lines and self-interpreted HIVST result.

	Overall	2 lines /reactive	1 line /reactive	2 lines /non-reactive	2 lines /DK-R	DK-R / reactive
less than a week	18 (53%)	12 (80%)	0 (0%)	0 (0%)	6 (75%)	0 (0%)
between 1 and 2 weeks	4 (12%)	1 (6.7%)	0 (0%)	2 (22%)	1 (12%)	0 (0%)
between 3 and 4 weeks	2 (5.9%)	1 (6.7%)	0 (0%)	0 (0%)	1 (12%)	0 (0%)
between one and two months	7 (21%)	1 (6.7%)	0 (0%)	5 (56%)	0 (0%)	1 (100%)
more than 3 months	3 (8.8%)	0 (0%)	1 (100%)	2 (22%)	0 (0%)	0 (0%)
Total	34 (100%)	15 (44.2%)	1 (2.9%)	9 (26.5%)	8 (23.5%)	1 (2.9%)

384 DK: don't know. R: refuse to answer

386 Table S4. Place of confirmatory testing among phase 2 participants who did link to confirmatory testing, by reported number of lines and self-interpreted HIVST result.

	Overall	2 lines /reactive	1 line /reactive	2 lines /non-reactive	2 lines /DK-R	DK-R / reactive
Health Center / Hospital / Clinic / Maternity	12 (35%)	3 (20%)	0 (0%)	6 (67%)	3 (38%)	0 (0%)
Community Clinic / KP-dedicated Health Center	22 (65%)	12 (80%)	1 (100%)	3 (33%)	5 (62%)	1 (100%)
Total	34 (100%)	15 (44.2%)	1 (2.9%)	9 (26.5%)	8 (23.5%)	1 (2.9%)

387 DK: don't know. R: refuse to answer
388

389 390 Table S5. Time between phase 1 and phase 2 interviews among phase 2 participants who did link to confirmatory testing, by reported number of lines and self-interpreted HIVST result.

	Overall	2 lines /reactive	1 line /reactive	2 lines /non-reactive	2 lines /DK-R	DK-R / reactive
between 4 and 6 months	67 (86%)	24 (89%)	5 (71%)	21 (84%)	17 (94%)	0 (0%)
less than 4 months	8 (10%)	3 (11%)	0 (0%)	4 (16%)	1 (5.6%)	0 (0%)
more than 6 months	3 (3.8%)	0 (0%)	2 (29%)	0 (0%)	0 (0%)	1 (100%)
Total	34 (100%)	15 (44.2%)	1 (2.9%)	9 (26.5%)	8 (23.5%)	1 (2.9%)

391 DK: don't know. R: refuse to answer

Acknowledgements

We wish to acknowledge the commitment and determination of all the ATLAS project teams, which made this research possible. We would also like to thank the interviewers for their professionalism in collecting this sensitive data. Finally, we are grateful to the participants who were kind enough to give us some of their time by agreeing to take part in the survey.

Data, scripts, code, and supplementary information availability

Data, scripts and code are available online on the Zenodo website.
(https://doi.org/ 10.5281/zenodo.7986077);

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of no financial conflicts of interest in relation to the content of the article. They declare no conflict of interest.

Funding

This work was supported by Unitaid (Grant Number: 2018-23 ATLAS) with additional funding from Agence Française pour le Développement (AFD). AKK benefits from an ANRS thesis allowance. MMG's research program is supported by a Canada Research Chair (Tier 2) in Population Health Modeling. The funding bodies were not involved in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

References

1. Das M, Chu PL, Santos G-M, Scheer S, Vittinghoff E, McFarland W, et al. Decreases in Community Viral Load Are Accompanied by Reductions in New HIV Infections in San Francisco. PLoS ONE. 2010;5:e11068.

2. Lima VD, Johnston K, Hogg RS, Levy AR, Harrigan PR, Anema A, et al. Expanded Access to Highly Active Antiretroviral Therapy: A Potentially Powerful Strategy to Curb the Growth of the HIV Epidemic. J Infect Dis. 2008;198:59–67.

3. Cohen MS, Chen YQ, McCauley M, Gamble T, Hosseinipour MC, Kumarasamy N, et al. Prevention of HIV-1 Infection with Early Antiretroviral Therapy. N Engl J Med. 2011;365:493–505.

4. Quinn TC, Wawer MJ, Sewankambo N, Serwadda D, Li C, Wabwire-Mangen F, et al. Viral Load and Heterosexual Transmission of Human Immunodeficiency Virus Type 1. N Engl J Med. 2000;342:921–9.

5. Hayes RJ, Donnell D, Floyd S, Mandla N, Bwalya J, Sabapathy K, et al. Effect of Universal Testing and Treatment on HIV Incidence — HPTN 071 (PopART). N Engl J Med. 2019;381:207–18.

6. Grinsztejn B, Hosseinipour MC, Ribaudo HJ, Swindells S, Eron J, Chen YQ, et al. Effects of early versus delayed initiation of antiretroviral treatment on clinical outcomes of HIV-1 infection: results from the phase 3 HPTN 052 randomised controlled trial. Lancet Infect Dis. 2014;14:281–90.

7. UNAIDS. REFERENCE UNAIDS DATA 2021 _ UNAIDS/JC3032E. 2021.

8. Unaids. Understanding Fast-Track: accelarating action to end the AIDS epidemic by 2023. Geneva: Unaids; 2015.

9. Johnson C, Baggaley R, Forsythe S, van Rooyen H, Ford N, Napierala Mavedzenge S, et al. Realizing the
Potential for HIV Self-Testing. AIDS Behav. 2014;18:391–5.

10. Njau B, Covin C, Lisasi E, Damian D, Mushi D, Boulle A, et al. A systematic review of qualitative
evidence on factors enabling and deterring uptake of HIV self-testing in Africa. BMC Public Health.
2019;19:1289.

11. Johnson CC, Kennedy C, Fonner V, Siegfried N, Figueroa C, Dalal S, et al. Examining the effects of HIV
self-testing compared to standard HIV testing services: a systematic review and meta-analysis. J Int AIDS
Soc. 2017;20:21594.

12. word Health Organisazion. WHO RECOMMENDS HIV SELFTESTING – EVIDENCE UPDATE AND
CONSIDERATIONS FOR SUCCESS. Geneva: Word Health Organization; 2019.

13. Choko AT, Jamil MS, MacPherson P, Corbett E, Chitembo L, Ingold H, et al. Measuring linkage to HIV
treatment services following HIV self-testing in low-income settings. J Int AIDS Soc. 2020;23.

14. World Health Organization. Guidelines on HIV self-testing and partner notification: supplement to
Consolidated guidelines on HIV testing services. 2016.

15. Hector J, Davies M-A, Dekker-Boersema J, Aly MM, Abdalad CCA, Langa EBR, et al. Acceptability and
performance of a directly assisted oral HIV self-testing intervention in adolescents in rural Mozambique.
PLOS ONE. 2018;13:e0195391.

16. Asiimwe S, Oloya J, Song X, Whalen CC. Accuracy of Un-supervised Versus Provider-Supervised Self-
administered HIV Testing in Uganda: A Randomized Implementation Trial. AIDS Behav. 2014;18:2477–84.

17. Figueroa C, Johnson C, Ford N, Sands A, Dalal S, Meurant R, et al. Reliability of HIV rapid diagnostic
tests for self-testing compared with testing by health-care workers: a systematic review and meta-
analysis. Lancet HIV. 2018;5:e277–90.

18. Chanda MM, Ortblad KF, Mwale M, Chongo S, Kanchele C, Kamungoma N, et al. HIV self-testing
among female sex workers in Zambia: A cluster randomized controlled trial. PLOS Med.
2017;14:e1002442.

19. Tonen-Wolyec S, Filali M, Mboup S, Bélec L. HIV self-testing in Africa: stakes and challenges.
Médecine Santé Trop. 2018;28:144–9.

20. Pant Pai N, Behlim T, Abrahams L, Vadnais C, Shivkumar S, Pillay S, et al. Will an Unsupervised Self-
Testing Strategy for HIV Work in Health Care Workers of South Africa? A Cross Sectional Pilot Feasibility
Study. PLoS ONE. 2013;8:e79772.

21. Brown AN, Djimeu EW, Cameron DB. A Review of the Evidence of Harm from Self-Tests. AIDS Behav.
2014;18:445–9.

22. UNAIDS. UNAIDS DATA 2022. Geneva: Joint United Nations Programme on
HIV/AIDS;UNAIDS/JC3063E. Geneva: Joint United Nations Programme on HIV/AIDS; 2022.

23. Rouveau N, Ky-Zerbo O, Boye S, Fotso AS, d'Elbée M, Maheu-Giroux M, et al. Describing, analysing
and understanding the effects of the introduction of HIV self-testing in West Africa through the ATLAS
programme in Côte d'Ivoire, Mali and Senegal. BMC Public Health. 2021;21:181.

24. Consolidated guidelines on HIV prevention, testing, treatment, service delivery and monitoring:
recommendations for a public health approach. Geneva, Switzerland: World Health Organization; 2021.

25. Kouassi AK, Simo Fotso A, N'Guessan KN, Geoffroy O, Younoussa S, Kanku Kabemba O, et al. Reaching
key and peripheral populations: a phone-based survey of HIV self-test users in West Africa. Durban:
poster #PEC004; 2021.

26. for the ATLAS Team, Rouveau N, Ky-Zerbo O, Boye S, Fotso AS, d'Elbée M, et al. Describing, analysing
and understanding the effects of the introduction of HIV self-testing in West Africa through the ATLAS
programme in Côte d'Ivoire, Mali and Senegal. BMC Public Health. 2021;21:181.

27. Simo Fotso A, Johnson C, Vautier A, Kouamé KB, Diop PM, Silhol R, et al. Routine programmatic data
show a positive population-level impact of HIV self-testing: the case of Côte d'Ivoire and implications for
implementation. AIDS. 2022;36:1871–9.

28. Ky-Zerbo O, Desclaux A, Boye S, Vautier A, Rouveau N, Kouadio BA, et al. Willingness to use and
distribute HIV self-test kits to clients and partners: a qualitative analysis of female sex workers' collective
opinion and attitude in Côte d'Ivoire, Mali, and Senegal. Women's Health. In press.

29. Boye S, Bouaré S, Ky-Zerbo O, Rouveau N, Simo Fotso A, d'Elbée M, et al. Challenges of HIV Self-Test
Distribution for Index Testing When HIV Status Disclosure Is Low: Preliminary Results of a Qualitative
Study in Bamako (Mali) as Part of the ATLAS Project. Front Public Health. 2021;9:653543.

30. d'Elbée M, Traore MM, Badiane K, Vautier A, Simo Fotso A, Kabemba OK, et al. Costs and Scale-Up
Costs of Integrating HIV Self-Testing Into Civil Society Organisation-Led Programmes for Key Populations
in Côte d'Ivoire, Senegal, and Mali. Front Public Health. 2021;9:653612.

31. R Core Team. R: A Language and Environment for Statistical Computing. 2021.

32. Asiimwe S, Oloya J, Song X, Whalen CC. Accuracy of Un-supervised Versus Provider-Supervised Self-
administered HIV Testing in Uganda: A Randomized Implementation Trial. AIDS Behav. 2014;18:2477–84.

33. Tippett Barr B, Lowrance D, Case Johnson C, Baggaley RC, Rogers J, Balachandra S, et al. Treatment-
adjusted prevalence to assess HIV testing programmes. Bull World Health Organ. 2021;99:874–82.

34. Giguère K, Eaton JW, Marsh K, Johnson LF, Johnson CC, Ehui E, et al. Trends in knowledge of HIV
status and efficiency of HIV testing services in sub-Saharan Africa, 2000–20: a modelling study using
survey and HIV testing programme data. Lancet HIV. 2021;8:e284–93.

35. Thirumurthy H, Masters SH, Mavedzenge SN, Maman S, Omanga E, Agot K. Promoting male partner
HIV testing and safer sexual decision making through secondary distribution of self-tests by HIV-negative
female sex workers and women receiving antenatal and post-partum care in Kenya: a cohort study.
Lancet HIV. 2016;3:e266–74.

36. Chipungu J, Bosomprah S, Zanolini A, Thimurthy H, Chilengi R, Sharma A, et al. Understanding linkage
to care with HIV self-test approach in Lusaka, Zambia - A mixed method approach. PLOS ONE.
2017;12:e0187998.

37. Tun W, Vu L, Dirisu O, Sekoni A, Shoyemi E, Njab J, et al. Uptake of HIV self-testing and linkage to
treatment among men who have sex with men (MSM) in Nigeria: A pilot programme using key opinion
leaders to reach MSM. J Int AIDS Soc. 2018;21:e25124.

38. Hlongwa M, Hlongwana K, Makhunga S, Choko A, Dzinamarira T, Conserve D, et al. Linkage to HIV
care following HIV self-testing among men: systematic review of quantitative and qualitative studies from
six countries in sub-Saharan Africa. preprint. In Review; 2022.

39. Choko AT, MacPherson P, Webb EL, Willey BA, Feasy H, Sambakunsi R, et al. Uptake, Accuracy, Safety,
and Linkage into Care over Two Years of Promoting Annual Self-Testing for HIV in Blantyre, Malawi: A
Community-Based Prospective Study. PLOS Med. 2015;12:e1001873.

40. Green KE, Vu BN, Phan HT, Tran MH, Ngo HV, Vo SH, et al. From conventional to disruptive: upturning
the HIV testing status quo among men who have sex with men in Vietnam. J Int AIDS Soc. 2018;21.

41. Ky-Zerbo O, Desclaux A, Kouadio AB, Rouveau N, Vautier A, Sow S, et al. Enthusiasm for Introducing
and Integrating HIV Self-Testing but Doubts About Users: A Baseline Qualitative Analysis of Key
Stakeholders' Attitudes and Perceptions in Côte d'Ivoire, Mali and Senegal. Front Public Health. 2021;9.

42. Inghels M, Kouassi AK, Niangoran S, Bekelynck A, Carilon S, Sika L, et al. Preferences and access to
community-based HIV testing sites among men who have sex with men (MSM) in Côte d'Ivoire. BMJ
Open. 2022;12:e052536.

517