HIV self-testing positivity rate and linkage to confirmatory testing and care: a telephone survey in Côte d'Ivoire, Mali, and Senegal

Arsène Kouassi Kra[1*], Arlette Simo Fotso[1,2], Nicolas Rouveau[1], Mathieu Maheu-Giroux[3], Marie-Claude Boily[4], Romain Silhol[4], Marc d'Elbée[1,5], Anthony Vautier[6] and Joseph Larmarange[1,2] on behalf of the ATLAS team

[1] Centre Population et Développement (Ceped), Université Paris Cité, Institut de Recherche pour le Développement (IRD), Inserm, France
[2] Institut National d'études Démographiques (Ined), France
[3] Department of Epidemiology and Biostatistics, School of Population and Global Health, McGill University, Montréal, QC, Canada
[4] MRC Centre for Global Infectious Disease Analysis, School of Public Health, Imperial College London, London, United Kingdom
[5] National Institute for Health and Medical Research UMR 1219, Research Institute for Sustainable Development EMR 271, Bordeaux Population Health Centre, University of Bordeaux, France
[6] Solidarité Thérapeutique et Initiatives pour la Santé (Solthis), Sénégal

*Corresponding author
Correspondence: arsene.kra.kouassi@ceped.org

ABSTRACT

HIV self-testing (HIVST) empowers individuals to decide when and where to test and with whom to share their results. From 2019 to 2022, the ATLAS program distributed ~ 400,000 HIVST kits in Côte d'Ivoire, Mali, and Senegal. It prioritised key populations, including female sex workers and men who have sex with men, and encouraged secondary distribution of HIVST to their partners, peers and clients.

To preserve the confidential nature of HIVST, use of kits and their results were not systematically tracked. Instead, an anonymous phone survey was carried out in two phases during 2021 to estimate HIVST positivity rates (phase 1) and linkage to confirmatory testing (phase 2). Initially, participants were recruited via leaflets from March to June and completed a sociobehavioural questionnaire. In the second phase (September-October), participants who had reported two lines or who reported a reactive result were recontacted to complete another questionnaire. Of the 2,615 initial participants, 89.7% reported consistent response between their interpretation and the number of lines on the HIVST (i.e., 1 for negative, 2 for reactive).

Overall HIV positivity rate was 2.5% (central hypothesis, low: 2.4%, high: 9.1%) based on self-interpreted results, and 4.5% (4.4% to 7.2%) based on the reported number of lines. Variations were observed according to country, distribution channel, sex and age group.

The second phase saw 78 out of 126 eligible participants complete the questionnaire. Of the 27 who reported a consistent reactive response in the first phase, 15 (56%, 95%CI: 36 to 74%) underwent confirmatory HIV testing, with 12 (80%) confirmed as HIV-positive, all of whom began antiretroviral treatment.

The confirmation rate of HIVST results was fast, with 53% doing so within a week and 91% within
three months of self-testing. Two-thirds (65%) went to a general public facility, and one-third to
a facility dedicated to key populations.
The ATLAS HIVST distribution strategy reached people living with HIV in West Africa. Linkage to
confirmatory testing following a reactive HIVST remained relatively low in these first years of
HIVST implementation. However, if confirmed HIV-positive, almost all initiated treatment. HIVST
constitutes a relevant complementary tool to existing screening services.
.

Keywords: AIDS; HIV; Self-Testing; Key Populations; MSM; sex-workers; phone-based survey; West Africa;
confirmatory testing; follow-up care; public health program evaluation.

55
56
57

Introduction

Early testing followed by successful linkage to antiretroviral treatment for those diagnosed with HIV can drastically reduce the risk of onward HIV transmission and mortality [1–6]. In 2021, according to the United Nations Program for HIV/AIDS (UNAIDS), 81% of the adult population living with HIV in West and Central Africa knew their status. Only 77% of them were on antiretroviral treatment[7], below the 95-95-95 UNAIDS targets for 2025 [8]. The 95-95-95 targets aim for 95% of people living with HIV to know their status, 95% of those diagnosed to receive treatment, and 95% of those on treatment to achieve viral suppression. Improving diagnosis coverage, especially among vulnerable key populations at high risk of HIV acquisition and transmission, is the necessary first step to achieve this goal.

HIV self-testing (HIVST) is the process by which a person who wants to know their HIV status collects their own sample (oral fluid or blood), performs the test, and then interprets the results themself, often in a private setting [9]. It is an innovative tool that empowers individuals and guarantees the confidentiality of the test result [10]. Individuals may decide when and where to test and with whom they want to share their result. It is a tool that is widely accepted by various populations, including key populations [11–18]. It has been shown to be effective in screening populations vulnerable to HIV acquisition and transmission that are often hardly reached through conventional approaches [19–21]. The World Health Organization (WHO) has recommended HIVST as a complementary testing approach since 2016 [22].

The HIV Self-Testing in Africa (STAR) carried in Eastern and Southern Africa and funded by Unitaid aimed to boost the global market for HIVST (https://www.psi.org/fr/project/star/). The project unfolded in three phases: Phase 1 ran from September 2015 to August 2017, Phase 2 spanned from August 2017 to July 2020, and Phase 3 took place between January 2020 and July 2021. Following the experience gained in Eastern and Southern Africa under the STAR project [11, 23–28], the Unitaid funding agency sought to stimulate HIVST in West Africa, where HIV epidemics are distinguished by their more concentrated and less generalised nature compared to those in Eastern and Southern Africa. In this region, the general population prevalences are relatively low to very low, and key populations (for example, female sex workers and men who have sex with men) are particularly affected and bear a disproportionate share of the HIV burden [29]. The ATLAS programme (*AutoTest de dépistage du VIH : Libre d'Accéder à la connaissance de son Statut*) aimed to promote, implement, and expand HIVST in Côte d'Ivoire, Mali, and Senegal [30] where the national HIV prevalence in 2021 was 1.9% (1.7%-2.2%) , 0.8% (0.6%-1%) , and 0.3% (0.3%-0.4%) respectively [31].

To preserve the anonymity and confidentiality of HIVST and not impede their use, ATLAS decided, in line with WHO recommendations, not to track the use and outcomes of distributed HIVST kits systematically. Such tracking can be logistically challenging and costly and could limit the distribution, redistribution and use of HIVST [32]. Without systematic tracking, it is challenging to obtain information on the users of the HIVST, their results and on linkage to confirmatory testing and treatment, which are crucial indicators to assess program effectiveness and impact. For instance, the positivity rate can reflect the yield of new individuals diagnosed with HIV and if the testing modality is indeed reaching those in need. Diagnosed individuals must seek confirmatory testing and be linked to care to maximise health benefits and decrease onward transmission.

We conducted an innovative survey by setting up an anonymous and free telephone platform in Côte d'Ivoire, Mali and Senegal while preserving anonymity and encouraging voluntary participation. In the second phase (September-October), participants who had reported two lines or a self-interpreted HIVST result as reactive were recontacted to complete another questionnaire. Here we present the HIV test positivity rates from the phase 1 questionnaire and the links with confirmatory tests and care.

Materials and Methods

ATLAS program description

ATLAS HIVST distribution was integrated into existing testing policies, programmes and activities in each country; 397,367 HIVST kits were freely distributed between July 2019 and February 2022 as part of the three countries' national AIDS strategies. At the time of ATLAS's implementation in 2019, only small-scale HIVST pilot programs had been previously conducted in Senegal and Côte d'Ivoire, whereas Mali had no previous

experience with HIVST. In Senegal, for instance, the first pilot survey took place between April 2017 and June
2018 [33].
The design of the different delivery channels and the priority populations were developed with country
stakeholders including national AIDS programs/councils, international institutions including the WHO,
international and national non-governmental organisations involved in local HIV programs, and civil society
and community leaders. ATLAS HIVST distribution was organised through eight different operational delivery
channels (Figure 1), i.e. five facility-based approaches (delivery of HIVST kits through public or community-
based health facilities) and three community-based approaches involving outreach activities engaging female
sex workers (FSW), men who have sex with men (MSM), and people who use drugs (PWUD) [30]. Peer
educators conducted these outreach activities through group activities (e.g. talks, discussion groups, night
visits, social events, or parties) and face-to-face activities (e.g. home visits). Outreach activities represented
the majority (~85%) of ATLAS's distribution volume.
119

121 **Figure 1.** ATLAS delivery channels (adapted from [30]). FSW=female sex workers, MSM=men who have
122 sex with men, PLHIV=people living with HIV PMTCT=prevention of mother-to-child transmission,
123 PWUD=people who use drugs, STI=sexually transmitted infection.

124
ATLAS activities relied on both primary and secondary distribution. HIVST kits were distributed by peer
educators and healthcare professionals to primary contacts for their personal use (primary distribution). With
secondary distribution, primary contacts were provided HIVST kits and invited to redistribute them to their
peers, partners, and clients. These secondary contacts were often members of key populations that can be
more difficult to engage in HIV prevention, along with other peripheral vulnerable populations. This chain-
referral distribution of HIVST implies that end-users were not limited to primary contacts.
Only oral self-testing (OraQuick HIV Self-Test®) has been distributed through ATLAS. OraSure Technologies,
the manufacturer of the OraQuick test, accompanies each HIVST kit with a user manual for result interpretation
(Figure 2). OraQuick HIVST results should be interpreted as follow: "reactive" ("positive") if two lines (C & T)
are visible (even barely), "non-reactive" ("negative") if only the C (control) line is visible, and "invalid" if no line
is visible or if only the T (test) line is visible. To be noted, the French version of the HIVST instructions distributed
by ATLAS used the wording "reactive" / "non-reactive" instead of "positive" / "negative" to qualify the HIVST
result, following WHO vocabulary in their HIVST guidelines[20] as an HIVST is triage test and does not provide
a definitive HIV-positive diagnosis. The questionnaire of the survey also used "reactive" / "non-reactive"
wording (https://doi.org/10.5281/zenodo.10210464).

OraQuick®
HIV SELF-TEST

INTERPRETING RESULTS 🔦 **Read test results in a well-lit area**

HIV POSITIVE RESULT

Two complete lines, even if the line is faint, means you may be HIV POSITIVE and you need to seek additional testing by a trained professional to confirm an HIV diagnosis.

As soon as possible . . .
Visit your nearest HIV Testing Centre or Health Facility

HIV NEGATIVE RESULT
IF READ BEFORE 20 MINUTES, RESULT MAY NOT BE CORRECT

ONE LINE next to the "C" and NO line next to the "T", your result is HIV NEGATIVE.

Seek regular testing. If you may have been exposed to HIV, test again in 3 months.

INVALID RESULT

If there is no line next to the "C" (even when there is a line next to the "T"), the test line or control line are not complete (all the way across the window), or a red background makes it impossible to read the test, the test is not working and should be repeated. **You will need to obtain another test.**

The test did not work properly. Visit your nearest HIV Testing Centre or Health Facility to test again.

NOT SURE OF RESULT
You do not know your result or you are unsure of your result. Visit your nearest HIV Testing Centre or Health Facility to test again.

DISPOSE
Remove the test stick, put the cap on the test tube, place in the disposal bag provided and throw away all contents in the normal trash.

142 **Figure 2.** Guidelines for interpreting HIVST result, extracted from the English version of the manufacturer
143 instructions for use (OraQuick HIV Self-Test®). To be noted, the French version distributed by ATLAS was
144 using the wording "Reactive" / "Non-reactive" instead of "Positive" / "Négative" to qualify the HIVST
145 result.

146
In addition to the manufacturer's instructions, locally adapted brochures and explanatory videos in French
and local languages have been developed to help users perform the test, interpret the result and know what
actions should be taken following a non-reactive, a reactive or indeterminate result (for example :
https://youtu.be/laCCjVEKZto in English or https://youtu.be/1xzitLD309U in French). They also encouraged

people with a reactive HIVST to seek confirmatory HIV testing and care. Individuals with a non-reactive test
were invited to retest after 3 months if still exposed to HIV. Existing toll-free hotlines in each counntry were
strengthened and trained on HIVST, to offer information about HIV, prevention, testing, use and interpretation
of HIVST and counseling.

Study design and data collection
The ATLAS program embedded multiple research activities, from qualitative studies to economic analyses,
which have been described in detail elsewhere [17, 30, 34–37].
The program included a voluntary anonymous phone survey. Between mid-March and mid-June 2021,
dedicated survey flyers were distributed with the HIVST kits inviting self-test users in each country to call a toll-
free number to complete a questionnaire (phase 1). All calls from the three countries, over the same period,
were rerouted to a telephone platform located in Abidjan and operated by Ipsos Côte d'Ivoire, which was
selected following an international call for tenders.
A pilot survey was initially conducted without offering financial compensation to the participants [38].
Following its results, we decided to introduce a reward as a token of appreciation for the time participants
dedicated to the survey. Consequently, completion of the questionnaire was rewarded with 2,000 XOF
(approximately 3.40 USD) of phone communication credit. This reward was mentioned on the survey flyers. In
order to participate in the survey, participants had to be of legal age to use an HIVST on their own without
parental permission (16 years in Côte d'Ivoire, 18 years in Mali, and 15 years in Senegal) and had to have used
an HIVST provided to them through the ATLAS project.
As the survey was anonymous, there was a risk that some HIVST users may participate more than once or
that individuals who have never used HIVST tried to participate to receive the financial incentive. To limit these
risks, several measures were taken: (i) the leaflet distributed with the HIVST kits had a unique 9-digit number
generated by the research team that was requested prior to participation in the survey, (ii) the same unique
number could not be used twice, (iii) the financial incentive was only paid out once the questionnaire was fully
completed (however individuals could refuse to answer any particular question), (iv) the same telephone
number could not be used twice to receive the telephone credit. These unique 9-digit numbers were generated
non-sequentially and were grouped by country, delivery channel and implementing partner. Thus, any unique
number could indirectly identify the delivery channel where the HIVST kit was initially dispensed.
The time when participants received their HIVST kit was not collected. However, as a survey leaflet was
mandatory to participate, we could estimate that all participants received their HIVST kit during the survey
period (i.e. between mid-March and mid-June 2021).
The phase 1 questionnaire, which lasted 20 to 30 minutes, collected information on sociodemographic
characteristics of HIVST users (including age, sex, marital status, education level), testing history (having ever
tested for HIV before using HIVST and date of last HIV test), sexual and preventive behaviours, HIVST use and
difficulties encountered [39]. Specifically, each participant was asked about the number of lines that appeared
when reading the HIVST result and their self-interpretation of it (reactive or non-reactive).
In total, 2,615 participants were recruited for phase 1[39]. Those who reported two lines or a reactive
result (n=126) were asked for their consent to be called back a few months later to participate in a
complementary survey and, if consented, to provide a phone contact (n=120). As some individuals may delay
their decision to undergo a confirmatory test by several weeks/months after using an HIV self-test, we chose
a minimum of 3-month gap between our two surveys to potentially get an estimate of the maximum number
of participants who eventually underwent confirmatory testing.
From September 27th to October 22nd, 2021, 96 were successfully recontacted and invited to complete a
5-minute questionnaire (phase 2). Among those, 89 accepted to participate in phase 2 and 78 fully completed
phase 2 questionnaire. Phase 2 questionnaire asked the participants if they had undergone a confirmatory test
following their HIVST, reasons for not linking to confirmatory testing (if not linked), time between HIVST and
confirmatory testing (if linked), type of facility for confirmatory testing, confirmation test result, linkage to
antiretroviral treatment (if confirmed HIV-positive).
199
The interviews were conducted in either French, English, Bambara, or Wolof. On-the-fly translation into
other local languages was also available. Compensation of XOF 2,000 (≈3.40 USD) in the form of telephone
credit was given to participants who completed the phase 2 questionnaire. Unlike in phase 1, it was not a
financial incentive as participants were informed about it only at the end of the interview. Interviews were not

audio-recorded. Questionnaires' responses were captured on a computer and stored in a database managed
by PAC-CI, an Ivorian research institute with expertise in clinical research.
At the end of the survey, collected telephone numbers (for appointments and rewards) were deleted from
the database. All procedures have been described in a publicly available data management plan
(https://dmp.opidor.fr/plans/3354/export.pdf). The complete project protocol, including the data
management plan (required by the ethics committees), was written in French. Both phase 1 and phase 2
questionnaires have been made available online and a link is now provided
(https://doi.org/10.5281/zenodo.10210464).

Data analysis
Following a previously published analysis [39], and due to small numbers of participants in certain
distribution channels, distribution channels (Figure 1) were grouped in three categories: FSW-based channels
(outreach or facility-based), MSM-based channels (outreach or facility-based) and other channels (PWUD-
based channels, index testing, STI consultations).
Based on phase 1 participants' self-reports, we distinguished between those who provided consistent
response, i.e. the reported number of visible lines was consistent with the reported self-interpretation (2
visible lines reported as reactive, one line reported as non-reactive, or no/one line and interpreted as invalid),
those who provided inconsistent response, i.e. the number of visible lines was inconsistent with the self-
interpretation of the result, or those who returned only a partial response (refusal to answer or answered "I
don't know" to one or both questions).
Due to inconsistent responses, we separately considered the self-interpreted results and the reported
number of lines on the HIVST to estimate HIVST positivity rates. For each source, we excluded "*don't know*"and
refusals (DK-R) from both the numerator and the denominator (central hypothese). We also computed the
highest positity (assuming all DK-R as reactive) and and lowest positivity (assuming all DK-R as non-reactive).
We conducted two binomial logistic regression analyses to examine the positivity rate in our central
scenario, based on self-interpreted results on one hand, and based on the reported number of lines on the
other. We deemed it important to stratify the positivity rates by country, sex, distribution channel, and age
group (15-24, 25-34, and 35+).
As the key population profile of participants should differ substantially by distribution channel (women
from the FSW-based channel are more likely FSW while those from the MSM-based channel are more likely
female partners of MSM; men from the MSM-based channel are more likely MSM while those from the FSW-
based channel are more likely partners or clients of FSW, see Figure 1), we decided to combine distribution
channel and sex into a single combined variable.
We described the selection of eligible participants for phase 2 questionnaires and corresponding
participation rates. To assess any participation bias, characteristics of phase 2 participants (country, sex and
distribution channel, age group, marital status, educational level, and first-time testers, i.e. whether they ever
tested for HIV before using HIVST) were compared with individuals eligible for phase 2 but who did not
participate and with phase 1 participants not eligible for phase 2. Bivariate comparison was done using chi-
squared tests, and multivariate comparison using a multinomial logistic regression model and then computing
likelihood-ratio tests.
Among phase 2 eligible participants who completed their questionnaire, linkage to confirmatory testing,
the proportion being confirmed HIV positive, and the proportion who initiated treatment were described,
stratified by the reported number of lines and self-interpreted HIVST result in phase 1 questionnaire.
We also described (i) for those who did not link to confirmatory testing, the main reported reason; and (ii)
for those who did link to confirmatory testing, the type of facility attended for confirmation and the time
between HIVST and confirmatory testing.
A dedicated anonymised dataset and the corresponding R script are available on Zenodo
(https://doi.org/10.5281/zenodo.10255772) to allow replication of the analysis. All analyses have been
performed using R version 4.3.1 [40]. All the descriptive tables were generated using the *tbl_summary()*
function from the *gtsummary* package [41]. Confidence intervals (95% confidence interval, 95%CI) were
computed using Wilson's method with Yate's continuity correction (*prop.test()* function in the 'stats' package).
Multinomial models were computed with *multinom()* from the 'nnet' package and likelihood-ratio tests with
Anova() from 'car'.
256

Ethics

ATLAS research protocol (version 3.0, October 8 2020) has been approved by the WHO Ethical Research Committee (January 12, 2021, reference: ERC 0003181), the National Ethics Committee for Life Sciences and Health of Côte d'Ivoire (November 27, 2020, reference: 191-20/MSHP/CNESVS-km, IRB:000111917), the Ethics Committee of the Faculty of Medicine and Pharmacy of the University of Bamako, Mali (November 16, 2020, reference: 2020/254/CE/FMPOS/FAPH), and the National Ethics Committee for Health Research of Senegal (January 26, 2021, protocol SEN19/32, n°8 MSAS/CNERS/Sec).

The full research protocol was written in French (https://hal.science/ATLAS_ADVIH/hal-04121482v1). The peer-reviewed protocol has been published in English elsewhere [30].

Results

HIVST results

Of the 2,615 participants recruited in phase 1, 2,346 (89.7%) reported a self-interpreted HIVST result consistent with their reported number of visibles lines on the HIVST: 2 292 (88.0%) reported one line self-interpreted as non-reactive, 50 (1.9%) two lines self-interpreted as reactive, and 4 (0.2%) no/one line self-interpreted as invalid (table 1). In contrast, 48 (1.8%) reported inconsistent response: 10 (0.4%) one line self-interpreted as reactive, 35 (1.3%) two lines self-interpreted as non-reactive/ and 3 (0.1%) no line self-interpreted as non-reactive. Finally, 221 (8.5%) reported a partial result: 147 (5.6%) reported 0, 1 or 2 lines but did not know how to interpret the result or refused to answer, 46 (1.7%) self-interpreted their result but did not know or refused to report the number of lines, and 28 (1.1%) did not know or refused to answer to both questions.

Table 1. Reported self-interpreted HIV self-test (HIVST) result, reported number of lines on the HIVST, and positivity rates according to different hypotheses among participants of the first phase of the survey in Côte d'Ivoire, Mali, and Senegal (2021).

Phase 1 participants	Formula	2615 (100%)
Consistent response (C)	$C = C_1 + C_2 + C_3$	**2346 (89.7%)**
2 lines / reactive[†]	C_1	50 (1.9%)
1 line / non-reactive	C_2	2 292 (88%)
0-1 line/ invalid	C_3	4 (0.2%)
Inconsistent response (I)	$I = I_1 + I_2 + I_3 + I_4 + I_5$	**48 (1.8%)**
1 line / reactive[†]	I_1	10 (0.4%)
0 line / reactive[†]	I_2	0 (0%)
2 lines/ non-reactive[†]	I_3	35 (1.3%)
0 line / non-reactive	I_4	3 (0.1%)
2 lines/ invalid[†]	I_5	0 (0%)
Partial response (P)	$P = P_1 + P_2 + P_3 + P_4 + P_5 + P_6 + P_7$	**221 (8.5%)**
0 line / DK-R	P_1	1 (<0.1%)
1 line / DK-R	P_2	117 (4.5%)
2 lines/ DK-R[†]	P_3	29 (1.1%)
DK-R / reactive[†]	P_4	2 (<0.1%)
DK-R / non-reactive	P_5	44 (1.7%)
DK-R / invalid	P_6	0 (0%)
DK-R / DK-R	P_7	28 (1.1%)
Positivity Rate		
Based on self-interpreted test results		
Low hypothesis (DK-R as not reactive)	$(C_1 + I_1 + I_2 + P_4) / n$	62 / 2615 (2.4 %)
Central hypothesis (DK-R excluded)	$(C_1 + I_1 + I_2 + P_4) / (C + I + P_4 + P_5 + P_6)$	62 / 2440 (2.5 %)
High hypothesis (DK-R as reactive)	$(C_1 + I_1 + I_2 + P_1 + P_2 + P_3 + P_4 + P_7) / n$	237 / 2615 (9.1%)
Based on the reported number of lines		
Low hypothesis (DK-R as 1 line)	$(C_1 + I_3 + I_5 + P_3) / n$	114 / 2615 (4.4 %)
Central hypothesis mid (DK-R excluded)	$(C_1 + I_3 + I_5 + P_3) / (C + I + P_1 + P_2 + P_3)$	114 / 2541 (4.5 %)
High hypothesis (DK-R as 2 lines)	$(C_1 + I_3 + I_5 + P_3 + P_4 + P_5 + P_6 + P_7) / (C + I + P_1 + P_2 + P_3)$	188 / 2615 (7.2 %)

†: Eligible for phase 2 Survey

DK: don't know. R: refused to answer

HIVST positivity rates

Based on self-interpreted HIVST results, the overall positivity rate was 2.4% when DK-R were considered non-reactive (low hypothesis, Figure 3, Table S2). Rate was similar at 2.5% by excluding DK-R from the numerator and the denominator (central hypothesis). Considering DK-R as reactive (high hypothesis) increased the positivity rate to 9.1%. Estimates based on the reported number of visible lines on the HIVST were 4.4%, 4.5% and 7.2%, respectively, for the low, central, and high hypotheses. Positivity rates ranged from 1.8% to 9.8% in Côte d'Ivoire, 3.5% to 7.8% in Mali, and 1.2% to 15.0% in Senegal depending on the hypothesis (e.g., low or high; Figure S4, Table S2).

Positivity rates (central hypothesis based on the number of lines) were higher among participants recruited
through community-based distribution channels. It was 4.8% for men and 4.9% for women in the MSM-based
channels, and 4.6% for men and 4.2% for women in the FSW-based channels. compared to 3.1% for men and
2.9% for women in the other distribution channels (PWUD-based and facility-based).
When analysing positivity rates by age group (Table S3), for participants aged between 15 to 24 years old,
the rates ranged from 2.2% to 7.4% based on the reported self-interpreted result and from 3.1% to 5.9% based
on the reported number of lines. Among those aged 25 to 34 years old, it fluctuated between 2.7% and 9.5%
based on the reported self-interpreted result and from 4.9% to 7.8% based on the reported number of lines.
Lastly, for individuals 35 years old or older, the rate layed between 1.8% and 12% based on the reported self-
interpreted result and between 4.9% and 9.3% based on the reported number of lines.

Participation in phase 2
During phase 1, 126 individuals reported two lines or self-interpreted their result as reactive and were
therefore eligible for phase 2 (Table 1). Among them, 6 refused to be recontacted after phase 1 (Figure 4).
Among the 120 (95%) who agreed to be recontacted, 24 (20%) were unreachable at the time of the phase 2
survey, and 96 (80%) were successfully recontacted. Among the latest, 89 (93%) accepted to participate in the
phase 2 survey. Ten dropped out before the end of the interview, and 1 disconnected and was unreachable
afterwards. As a result, 78 participants completed phase 2 questionnaire. Of the 78 participants, 39 (50%) were
from Côte d'Ivoire, 31 (40%) from Mali, and 8 (10%) from Senegal (Table S2). Participation rates were 54%
(27/50) for participants who reported a consistent response (2 lines and reactive), 71% (32/45) for those with
an inconsistent response (either 2 lines & non-reactive, or 1 line & reactive), and 66% (19/31) for those
reporting a partial response (2 lines & DK-R or DK-R & reactive).
The participants who completed the phase 2 questionnaire had similar sociodemographic characteristics
(e.g. country, sex, distribution channel, age group, marital status) compared to those eligible for phase 2, but
that did not complete it, and to phase 1 participants not eligible for phase 2 (Table S4). For most participants
(86%), phase 2 questionnaire was completed between 4 and 6 months after phase 1 questionnaire (Table S5).

a. Positivity rates based on self-interpreted HIVST result

b. Positivity rates based on the reported number of lines

Figure 3. Positivity rates and 95%CI based on self-interpreted HIVST results or the reported number of visible lines, by distribution channel, sex and country, among participants of the first survey phase in Côte d'Ivoire, Mali, and Senegal (2021). The asterix indicate that there was no participant in that distribution channel. FSW=female sex worker, MSM=men who have sex with men. MSM-based channels include facility-based and outreach. FSW-based channels include facility-based and outreach. Other channels include PWUD-based channels, index testing and STI consultations.

321　**Figure 4**. Flow chart of the participant selection process for the 2nd phase of the survey in Côte
322　d'Ivoire (CI), Mali (ML), and Senegal (SN) (2021).
323

Linkage to confirmatory testing and care

Overall, 34 of the 78 who completed the phase 2 questionnaire (44%) reported having performed confirmatory testing. Linkage was higher for those who reported 2 lines and correctly self-interpreted their result as reactive (56%,95%CI: 36-74%), than for those who reported two lines but did not know or refused to report their test interpretation (44%, (95%CI: 22-69%) and those who reported 2 lines but incorrectly self-interpreted the result as non-reactive (36%, 95%CI: 19-57%) (Table 3). Finally, among the 8 participants who reported none/one line or did not know how many lines and incorrectly self-interpreted the result as reactive, only 2 linked to confirmatory testing.

The main reason given for not linking to confirmatory testing was that "*their HIVST was non-reactive*" (18/44, 41%, and although 8 of these 18 reported a reactive result in phase 1 questionnaire), followed by "*not knowing that a confirmation test was required*" (10/44, 23%), and "*not having time*" (8/44, 18%) (Table S6).

When participants were linked to confirmatory testing, it was usually shortly after performing their HIVST: 53% linked in less than one week and 91% in less than 3 months (Table S5). Most participants (65%) performed their confirmatory testing in a general public facility (health centre, hospital, clinic or maternity) wheras 35% chose a community-based clinic or health centre dedicated to key populations (Table S7).

Among the 34 that linked to confirmatory testing, 19 (56%, 95%CI: 38-72%) were confirmed HIV-positive, and 18 (95%, 95%CI; 72-100%) initiated antiretroviral treatment. Of the 18 participants who initiated ART, 11 (72%) underwent their confirmation test less than a week after their self-test, 2 (11%) did so between 1 and 2 weeks, 1 (5.6%) between 3 and 4 weeks, 1 (5.6%) waited between 1 and 2 months, and 1 (5.6%) proceeded with the test three months later. Among the 27 who reported a consistent reactive response in the phase 1 questionnaire, 15 (56%, 95%CI: 36-74%) linked to confirmatory test, 12 (80%) were confirmed HIV-positive and all started treatment (100%).

349 **Table 2**. Linkage to confirmatory testing, proportion being confirmed HIV positive and treatment initiation, by reported number of lines and self-interpreted HIVST result
350 among eligible participants of the second phase of the survey who completed their questionnaire in Côte d'Ivoire, Mali, and Senegal (2021).

Reported number of lines/ self-interpreted HIVST result	Completed phase 2	Linked to confirmatory testing		Confirmed HIV positive		Initiated ART	
	n	n (%)	95%CI	n (%)	95%CI	n (%)	95%CI
Overall	78	34 (44%)	33% to 55%	19 (56%)	38% to 72%	18 (95%)	72% to 100%
2 lines / reactive	27	15 (56%)	36% to 74%	12 (80%)	51% to 95%	12 (100%)	70% to 100%
1 line / reactive	7	1 (14%)	1% to 58%	0 (0%)	0% to 80%		
2 lines / non-reactive	25	9 (36%)	19% to 57%	3 (33%)	9% to 69%	3 (100%)	31% to 100%
2 lines / DK-R	18	8 (44%)	22% to 69%	4 (50%)	22% to 78%	3 (75%)	22% to 99%
DK-R / reactive	1	1(100%)	5% to 100%	0 (0%)	0% to 95%		

351 DK: don't know. R: refuse to answer. CI: confidence interval.
352

Discussion

Our study shows that the strategy implemented by the ATLAS program, through primary and secondary
distribution of HIVST kits and dedicated channels, achieved HIV positivity rates of 2.5% (central hypothesis,
low: 2.4%, high: 9.1%) based on self-interpreted results, and 4.5% (central hypothesis, low: 4.4%, high:
7.2%) based on the reported number of lines. The proportion of participants with a reactive HIVST that
sought confirmatory testing was 44% (95% CI: 33%-55%). Of those who underwent confirmatory testing,
56% (95% CI: 38%-72%) were found to be HIV-positive and, among them, 95% (95% CI: 72%-100%) initiated
treatment. Among the participants who confirmed their reactive HIVST with a traditional facility-based HIV
test, 65% did so within a week and 91% within three months.

According to our estimates, HIVST positivity rates were 2.0% (central hypothesis, low: 1.8%, high: 9.8%)
based on self-interpreted results, and 3.9% (central hypothesis, low: 3.8%, high: 5.4%) based on the
reported number of lines in Côte d'Ivoire. In Mali, these rates were 3.6% (central hypothesis, low: 3.5%,
high: 6.7%) and 5% (central hypothesis, low: 4.9%, high: 7.8%), while in Senegal, they were 1.4% (central
hypothesis, low: 1.2%, high: 15%) and 6.0% (central hypothesis, low: 5.4%, high: 14.9%). Overall, these
results for HIVST positivity are generally higher than the average overall positivity of HIV testing services
(excluding HIVST) in West Africa. For instance, in 2020 an estimated 1.9% of all HIV tests performed were
found to be positive in the region (95% credible intervals: 1.3 to 2.7%) [42]. Further, among 15-24 and 25-
34 years old, which constitute more than 80% of our sample, overall positivity was 0.9% (0.7 to 1.3%) and
1.6% (1.2 to 2.2%), respectively. Collectively, these results provide evidence that HIVST is a high-yield
testing modality that can address the unmet HIV testing needs of key populations and their partners.
Our results are in line with data collected by ATLAS implementing partners. Between 2020 and 2021,
these ATLAS partners collected spontaneous feedback from HIVST users. This unpublished data collection
was non-systematic and varied from one partner to another. Among 4 463 documented feedbacks, HIVST
was reactive for 188 cases (4.2%), consistent with our estimates based on the reported number of visible
lines (4.5%).

In 2021, a study based on the UNAIDS-supported *Shiny90* mathematical model [43] estimated, using
data from 184 population surveys and reports from national HIV screening programs from 40 sub-Saharan
African countries, that the positivity rates for conventional HIV testing were 1.4% in Côte d'Ivoire, 2.2% in
Mali, and 1.0% in Senegal. Our estimates for HIVST were higher than these estimates for convential testing.
It is important to interpret HIV positivity rates while considering the treatment-adjusted prevalence
(i.e., removing those on treatment from the numerator and denominator of HIV prevalence), a more
reliable indicator for evaluating the effectiveness and positivity rates of targeted screening programs [44].
In West Africa, the treatment-adjusted prevalence remained relatively low in 2021: 0.6% in Côte d'Ivoire,
0.7% in Mali, and 0.06% in Senegal, according to UNAIDS data (https://aidsinfo.unaids.org/). Our positivy
rates in each country are higher than the treatment-adjusted prevalence, suggesting that the ATLAS HIVST
distribution strategy successfully reached a hard-to-reach population and at positivity levels at least as high
as with passive surveillance.

In our study, 2.0% of the participants reported an inconsistent response between the number of visible
lines and their self-interpretation of the result and 6.0% reported a number of lines but didn't know how
to interpret it or refused to answer, suggesting potential issues in interpreting the number of visible lines
on HIVST kits. In the context of the ATLAS program, the distribution strategy combining primary and
secondary approaches has led many HIVST users to perform their HIVST without receiving advice from a
healthcare professional or a trained peer educator. Although the HIVST is not designed to require
supervision, it is essential to have received information on its use before proceeding with the test. A study
conducted within the framework of the ATLAS program demonstrated that the manufacturer's instructions
alone were insufficient in a multilingual context with low literacy levels. The use of additional aid, such as
a demonstration video or a toll-free helpline proved to be necessary [45]. Similarly, a study carried out in
China in 2018 on the unsupervised use of HIVST among 27 MSM found that only 5 (or 19%) made no errors,
and 44% received an invalid test result due to various mistakes made [46]. However, the lack of supervision
is likely insufficient to explain the inconsistencies observed [23]. Some inconsistencies may result from a

misunderstanding of the terms "*reactive*" and "*non-reactive*", particularly considering that HIVST was a new tool in our context and that traditional terms used to describe conventional HIV testing are "*positive*" and "*negative*". This possible misunderstanding of the terms is also highlighted by the fact that 8 participants reported a "*reactive*" result in phase 1 questionnaire and then in phase 2 that their test was "*non-reactive*" as the main reason for not linking to confirmatory testing. Specific qualitative interviews or focus groups discussion with HIVST users could help better understand how they perceive different terms.

Linkage to confirmatory testing following a reactive test was 44% (95% confidence interval from 33% to 55%). However, this estimate includes some individuals who did not adequately self-interpreted their HIVST result as reactive. When considering only those who reported two lines and self-interpreted their result as reactive, the linkage rate increased to 56% (36% to 74%). This percentage is closer to that was observed in a study conducted in Kenya on HIV testing of FSW male partners using HIVST secondary distribution, where 65% of men with a reactive result had a confirmatory test [47]. Our estimates were based on small numbers resulting in large confidence intervals, but are still showing a low rate.

Linkage to confirmatory testing happened relatively quickly after HIVST use: 53% did it in less than a week and 91% in less than three months. Similar results were observed in a study in the general population in Zambia[48], and a study among MSM in Nigeria [49].

The main reasons given for not linking to confirmatory testing suggest potential misinterpretation of the result or misunderstanding about the need to perform a confirmatory HIV test, highlighting the need to improve messaging around HIVST, in particular when HIV self-testing policies will be scaled-up. For those who did confirmatory testing and were confirmed HIV positive, initiation of antiretroviral treatment was almost systematic, showing good linkage to care after confirmatory testing, as observed in many HIVST studies in sub-Saharan Africa [50–52].

Previous analyses of ATLAS data showed that HIVST could reach people not reached by conventional HIV testing approaches [53], particularly partners and clients of key populations and key population members not self-identifying as such [54]. It is consistent with the finding that two-thirds of participants who did confirmatory testing went to a general health facility rather than a community clinic dedicated to key populations. In a study conducted in 2018 in Côte d'Ivoire among MSM, one-third of the participants preferred community-based testing, one-third expressed no preference, and one-third preferred undifferentiated HIV testing services (general population), mentioning the lack of discretion and anonymity of community-based sites and the desire to avoid the gaze of others [55].

The implementation of a telephone survey, aimed at gathering information from HIVST users while preserving anonymity and without interfering with secondary distribution, has proven to be very useful to evaluate the ATLAS program. However, its high cost makes it difficult to integrate it into national strategies for assessing the impact of HIVST. Nevertheless, other impact evaluation methods, such as data triangulation [36] and modelling [37], may prove more suitable for routine monitoring of HIVST's impacts.

A previous analysis of this survey among ATLAS HIVST users showed that HIVST secondary distribution was feasible and acceptable [39]: participants reported that they appreciated the ease of use of HIVST, its discretion and the fact that they are autonomous in carrying out the test. Finally, HIVST appeared as a relevant additional approach for those usually distant from community activities and HIV testing services, and has the potential to reach, beyond key populations, partners, clients, and other groups vulnerable to HIV.

ATLAS' HIVST distribution strategy successfully reached people living with HIV in West Africa, although linkage to confirmatory testing following a reactive HIVST remained relatively low in these first years of HIVST implementation, and sub-optimal in the perspective of achieving UNAIDS 95-95-95 targets. However, among participants who confirmed their reactive self-test result with a traditional facility-based HIV test, a substantial proportion quickly proceeded with this confirmation (more than half in less than a week and the vast majority in less than three months). Furthermore, if individuals were confirmed HIV-positive, almost all began antiretroviral treatment. We showed that HIVST has the potential to reach more

460   hidden populations and constitutes a relevant complementary tool to existing screening services. To fully
461   harness the potential of self-tests, messaging around HIVST and its interpretation could be improved.
462

Appendices

Table S1. Positivity rates based on the reported number of visible lines or on self-interpreted HIVST results and associated factors (binomial logistic regression), among participants of the first survey phase in Côte d'Ivoire, Mali, and Senegal (2021)

	Positivity rate based on the reported number of visible lines (central hypothesis)			Positivity rate based on self-reported HIVST results (central hypothesis)		
	OR	95% CI	p-value	OR	95% CI	p-value
(Intercept)	0.04	0.02, 0.08	<0.001	0.03	0.01, 0.07	<0.001
Sex and distribution channel			0.4			0.10
Man : MSM-based channels	—	—		—	—	
Woman : MSM-based channels	1.05	0.36, 2.52	>0.9	0.25	0.01, 1.22	0.2
Man : FSW-based channels	0.90	0.54, 1.48	0.7	0.42	0.19, 0.86	0.023
Woman : FSW-based channels	0.66	0.38, 1.11	0.12	0.51	0.26, 0.98	0.048
Man: Other delivery channels	0.46	0.13, 1.30	0.2	0.26	0.01, 1.43	0.2
Woman : Other delivery channels	0.38	0.06, 1.44	0.2	0.40	0.02, 2.30	0.4
Country			0.2			0.10
Côte d'Ivoire	—	—		—	—	
Mali	1.33	0.88, 2.02	0.2	1.66	0.97, 2.90	0.067
Senegal	1.79	0.84, 3.58	0.11	0.70	0.15, 2.34	0.6
Age group			0.2			0.5
15-24 years or less	—	—		—	—	
25-34 years	1.44	0.94, 2.20	0.092	1.39	0.80, 2.43	0.2
35 years or more	1.61	0.84, 3.00	0.14	1.19	0.43, 2.93	0.7
Marital status			0.2			0.5
single	—	—		—	—	
divorced / separated / widowed	1.55	0.62, 3.50	0.3	2.02	0.54, 6.00	0.2
living with partner / married	0.76	0.46, 1.21	0.3	1.11	0.59, 2.02	0.7
Educational level			0.13			<0.001
none / primary	—	—		—	—	
secondary	1.01	0.62, 1.71	>0.9	0.98	0.52, 1.95	>0.9
higher	0.62	0.33, 1.18	0.14	0.23	0.07, 0.62	0.006

FSW: female sex workers, MSM: men having sex with men.

468 **Table S2**. Positivity rates based on self-interpreted HIVST results or the reported number of visible lines, by distribution channel, sex and country, among participants of
469 the first survey phase in Côte d'Ivoire, Mali, and Senegal (2021). FSW-based channels include facility-based and outreach. Other channels include PWUD-based channels,
470 index testing and STI consultations.

			MSM-based channels		FSW-based channels		Others delivery channels		Total
			Man	Woman	Man	Woman	Man	Woman	
Positivity rate based on self-reported HIVST results	**Low**	Côte d'Ivoire	2.5% (16/650)	1.4% (1/73)	1.5% (5/339)	1.2% (3/245)	0% (0/60)	0% (0/23) †	1.8% (25/1 390)
		Mali	4.6% (14/306)	0% (0/29)	1.9% (5/269)	3.9% (14/360)	9.1% (1/11) †	0% (0/9) †	3.5% (34/984)
		Senegal	4.9% (2/41)	0% (0/1) †	0% (0/12) †	0% (0/80)	0% (0/66)	2.4% (1/41)	1.2% (3/241)
		Overall	3.2% (32/997)	1.0% (1/103)	1.6% (10/620)	2.5% (17/685)	0.7% (1/137)	1.4% (1/73)	2.4% (62/2,615)
	Central	Côte d'Ivoire	2.7% (16/597)	1.4% (1/71)	1.6% (5/311)	1.4% (3/221)	0% (0/58)	0% (0/21) †	2.0% (25/1,279)
		Mali	4.7% (14/301)	0% (0/29)	1.9% (5/257)	4.1% (14/345)	9.1% (1/11) †	0% (0/9) †	3.6% (34/952)
		Senegal	6.1% (2/33)	0% (0/1) †	0% (0/11) †	0% (0/65)	0% (0/61)	2.6% (1/38)	1.4% (3/209)
		Overall	3.4% (32/931)	1.0% (1/101)	1.7% (10/579)	2.7% (17/631)	0.8% (1/130)	1.5% (1/68)	2.5% (62/2,440)
	High	Côte d'Ivoire	10.6% (69/650)	4.1% (3/73)	9.7% (33/339)	11% (27/245)	3.3% (2/60)	8.7% (2/23)	9.8% (136/1,390)
		Mali	6.2% (19/306)	0% (0/29)	6.3% (17/269)	8.1% (29/360)	9.1% (1/11) †	0% (0/9) †	6.7% (66/984)
		Senegal	24.0% (10/41)	0.0% (0/1) †	8.3% (1/12) †	19.0% (15/80)	7.6% (5/66)	9.8% (4/41)	15.0% (35/241)
		Overall	9.8% (98/997)	2.9% (3/103)	8.2% (51/620)	10.0% (71/685)	5.8% (8/137)	8.2% (6/73)	9.1% (237/2,615)
Positivity rate based on the reported number of visible lines	**Low**	Côte d'Ivoire	4.2% (27/650)	5.5% (4/73)	4.7% (16/339)	2.0% (5/245)	0% (0/60)	4.3% (1/23)	3.8% (53/1 390)
		Mali	4.9% (15/306)	3.4% (1/29)	4.5% (12/269)	5.3% (19/360)	9.1% (1/11) †	0% (0/9) †	4.9% (48/984)
		Senegal	12.2% (5/41)	0% (0/1) †	0% (0/12) †	5.0% (4/80)	4.5% (3/66)	2.4% (1/41)	5.4% (13/241)
		Overall	4.7% (47/997)	4.9% (5/103)	4.5% (28/620)	4.1% (28/685)	2.9% (4/137)	2.7% (2/73)	4.4% (114/2,615)
	Central	Côte d'Ivoire	4.2% (27/641)	5.5% (4/73)	4.8% (16/331)	2.1% (5/241)	0% (0/60)	4.5% (1/22) †	3.9% (53/1,368)
		Mali	5.0% (15/298)	3.4% (1/29)	4.5% (12/264)	5.5% (19/344)	9.1% (1/11) †	0% (0/9) †	5.0% (48/955)
		Senegal	13.2% (5/38)	0% (0/1) †	0% (0/10) †	5.3% (4/75)	5.3% (3/57)	2.7% (1/37)	6.0% (13/218)
		Overall	4.8% (47/977)	4.9% (5/103)	4.6% (28/605)	4.2% (28/660)	3.1% (4/128)	2.9% (2/68)	4.5% (114/2,541)
	High	Côte d'Ivoire	5.5% (36/650)	5.5% (4/73)	7.1% (24/339)	3.7% (9/245)	0% (0/60)	8.7% (2/23) †	5.4% (75/1,390)
		Mali	7.5% (23/306)	3.4% (1/29)	6.3% (17/269)	9.7% (35/360)	9.1% (1/11) †	0% (0/9) †	7.8% (77/984)
		Senegal	19.5% (8/41)	0% (0/1) †	16.7% (2/12) †	11.2% (9/80)	18.2% (12/66)	12.2% (5/41)	14.9% (36/241)
		Overall	6.7% (67/997)	4.9% (5/103)	6.9% (43/620)	7.7% (53/685)	9.5% (13/137)	9.6% (7/73)	7.2% (188/2,615)

471 DK: don't know. R: refusals. FSW: female sex workers, MSM: men having sex with men, PR: positivity rate.
472 †: indicated cells with less than 25 participants.
473 Low hypothesis: DK-R as non-reactive or 1 line. Central hypothesis: DK-R excluded from the numerator and the denominator. High hypothesis: DK-R as reactive or 2 lines.

Table S3. Positivity rates based on self-interpreted HIVST results or the reported number of visible lines, by age group and country, among participants of the first survey phase in Côte d'Ivoire, Mali, and Senegal (2021).

			15-24 years	25-34 years old	35 years or more	Total
Positivity rate based on self-reported HIVST results	Low	Côte d'Ivoire	1.7% (11/645)	2.0% (11/553)	1.6% (3/192)	1.8% (25/1 390)
		Mali	3.3% (15/455)	3.9% (16/415)	2.6% (3/114)	3.5% (34/984)
		Senegal	0.0% (0/64)	2.1% (2/95)	1.2% (1/82)	1.2% (3/241)
		Overall	2.2% (26/1 164)	2.7% (29/1 063)	1.8% (7/388)	2.4% (62/2 615)
	Central	Côte d'Ivoire	1.8% (11/604)	2.2% (11/506)	1.8% (3/169)	2.0% (25/1 279)
		Mali	3.4% (15/439)	4.0% (16/403)	2.7% (3/110)	3.6% (34/952)
		Senegal	0.0% (0/56)	2.4% (2/82)	1.4% (1/71)	1.4% (3/209)
		Overall	2.4% (26/1 099)	2.9% (29/991)	2.0% (7/350)	2.5% (62/2 440)
	High	Côte d'Ivoire	8.1% (52/645)	10.0% (58/553)	14.0% (26/192)	9.8% (136/1 390)
		Mali	6.8% (31/455)	6.7% (28/415)	6.1% (7/114)	6.7% (66/984)
		Senegal	13.0% (8/64)	16.0% (15/95)	15.0% (12/82)	15.0% (35/241)
		Overall	7.8% (91/1 164)	9.5% (101/1 063)	12.0% (45/388)	9.1% (237/2 615)
Positivity rate based on the reported number of visible lines	Low	Côte d'Ivoire	3.1% (20/645)	4.5% (25/553)	4.2% (8/192)	3.8% (53/1 390)
		Mali	4.8% (22/455)	4.8% (20/415)	5.3% (6/114)	4.9% (48/984)
		Senegal	1.6% (1/64)	7.4% (7/95)	6.1% (5/82)	5.4% (13/241)
		Overall	3.7% (43/1 164)	4.9% (52/1 063)	4.9% (19/388)	4.4% (114/2 615)
	Central	Côte d'Ivoire	3.1% (20/637)	4.6% (25/546)	4.3% (8/185)	3.9% (53/1 368)
		Mali	4.9% (22/447)	5.0% (20/401)	5.6% (6/107)	5.0% (48/955)
		Senegal	1.9% (1/54)	8.2% (7/85)	6.3% (5/79)	6.0% (13/218)
		Overall	3.8% (43/1 138)	5.0% (52/1 032)	5.1% (19/371)	4.5% (114/2 541)
	High	Côte d'Ivoire	4.3% (28/645)	5.8% (32/553)	7.8% (15/192)	5.4% (75/1 390)
		Mali	6.6% (30/455)	8.2% (34/415)	11.0% (13/114)	7.8% (77/984)
		Senegal	17.0% (11/64)	18.0% (17/95)	9.8% (8/82)	15.0% (36/241)
		Overall	5.9% (69/1 164)	7.8% (83/1 063)	9.3% (36/388)	7.2% (188/2 615)

Table S4. Eligibility and participation in phase 2 survey by sociodemographic characteristics, distribution channel, and HIV testing history (bivariate and multinomial regression model). FSW-based channels include facility-based and outreach. Other channels include PWUD-based channels, index testing and STI consultations.

	completed phase 2 questionnaire N = 78	eligible for phase 2 but did not complete the questionnaire N = 48	phase 1 participants not eligible for phase 2 N = 2,489	bivariate p-value (Chi² test)	multivariate p-value (multinomial model)	Overall N = 2,615 (phase 1 participants)
Country				0.9	0.8	
Côte d'Ivoire	39 (50%)	23 (48%)	1,328 (53%)			1,390 (53%)
Mali	31 (40%)	20 (42%)	933 (37%)			984 (38%)
Senegal	8 (10%)	5 (10%)	228 (9.2%)			241 (9.2%)
Sex and distribution channel				0.3	0.06	
Man: MSM-based channels	35 (45%)	18 (38%)	944 (38%)			997 (38%)
Woman: MSM-based channels	5 (6.4%)	0 (0%)	98 (3.9%)			103 (3.9%)
Man: FSW-based channels	22 (28%)	10 (21%)	588 (24%)			620 (24%)
Woman: FSW-based channels	14 (18%)	16 (33%)	655 (26%)			685 (26%)
Man: Other delivery channels	1 (1.3%)	3 (6.3%)	133 (5.3%)			137 (5.2%)
Woman: Other delivery channels	1 (1.3%)	1 (2.1%)	71 (2.9%)			73 (2.8%)
Age group				0.5	0.11	
15-24 years or less	27 (35%)	21 (44%)	1,116 (45%)			1,164 (45%)
25-34 years	38 (49%)	20 (42%)	1,005 (40%)			1,063 (41%)
35 years or more	13 (17%)	7 (15%)	368 (15%)			388 (15%)
Marital status				0.3	0.5	
single	54 (69%)	32 (67%)	1,675 (67%)			1,761 (67%)
divorced / separated / widowed	6 (7.7%)	2 (4.2%)	89 (3.6%)			97 (3.7%)
living with partner / married	18 (23%)	14 (29%)	725 (29%)			757 (29%)
Educational level				0.079	0.09	
none / primary	13 (17%)	13 (27%)	477 (19%)			503 (19%)
secondary	50 (64%)	29 (60%)	1,353 (54%)			1,432 (55%)
higher	15 (19%)	6 (13%)	659 (26%)			680 (26%)
First-time tester				0.2	0.228	
no	40 (51%)	25 (52%)	1,472 (59%)			1,537 (59%)
yes	38 (49%)	23 (48%)	1,017 (41%)			1,078 (41%)

481 FSW: female sex workers, MSM: men having sex with men.
482

485 **Table S5.** Time between HIVST and confirmatory testing among phase 2 participants who did link to confirmatory testing, by reported number of lines and self-interpreted
486 HIVST result

	Overall	2 lines /reactive	1 line /reactive	2 lines /non-reactive	2 lines /DK-R	DK-R / reactive
less than a week	18 (53%)	12 (80%)	0 (0%)	0 (0%)	6 (75%)	0 (0%)
between 1 and 2 weeks	4 (12%)	1 (6.7%)	0 (0%)	2 (22%)	1 (12%)	0 (0%)
between 3 and 4 weeks	2 (5.9%)	1 (6.7%)	0 (0%)	0 (0%)	1 (12%)	0 (0%)
between 1 and 2 months	7 (21%)	1 (6.7%)	0 (0%)	5 (56%)	0 (0%)	1 (100%)
more than 3 months	3 (8.8%)	0 (0%)	1 (100%)	2 (22%)	0 (0%)	0 (0%)
Total	34 (100%)	15 (44.2%)	1 (2.9%)	9 (26.5%)	8 (23.5%)	1 (2.9%)

487 DK: don't know. R: refuse to answer

489 **Table S6**. Main reason for not linking to confirmatory testing among phase 2 participants who did not link to confirmatory testing, by reported number of lines and self-
490 interpreted HIVST result.

	Overall	2 lines / reactive	1 line / reactive	2 lines / non-reactive	2 lines / DK-R
My test was non-reactive	18 (41%)	6 (50%)	2 (33%)	5 (31%)	5 (50%)
I didn't know we should get a confirmatory test	10 (23%)	2 (17%)	2 (33%)	5 (31%)	1 (10%)
I didn't have time	8 (18%)	3 (25%)	0 (0%)	3 (19%)	2 (20%)
I feared that others would know the result	2 (4.5%)	0 (0%)	0 (0%)	1 (6.2%)	1 (10%)
I already knew the result before using HIVST	2 (4.5%)	1 (8.3%)	1 (17%)	0 (0%)	0 (0%)
I had no specific reason	2 (4.5%)	0 (0%)	1 (17%)	1 (6.2%)	0 (0%)
I didn't know where to take the test	1 (2.3%)	0 (0%)	0 (0%)	1 (6.2%)	0 (0%)
The testing site was too far away	1 (2.3%)	0 (0%)	0 (0%)	0 (0%)	1 (10%)

491 DK: don't know. R: refuse to answer

494 **Table S7**. Place of confirmatory testing among phase 2 participants who did link to confirmatory testing, by reported number of lines and self-interpreted HIVST result.

	Overall	2 lines /reactive	1 line /reactive	2 lines /non-reactive	2 lines /DK-R	DK-R / reactive
Health Center / Hospital / Clinic / Maternity	12 (35%)	3 (20%)	0 (0%)	6 (67%)	3 (38%)	0 (0%)
Community Clinic / KP-dedicated Health Center	22 (65%)	12 (80%)	1 (100%)	3 (33%)	5 (62%)	1 (100%)
Total	34 (100%)	15 (44.2%)	1 (2.9%)	9 (26.5%)	8 (23.5%)	1 (2.9%)

495 DK: don't know. R: refuse to answer

497 **Table S8**. Time between phase 1 and phase 2 interviews among phase 2 participants who did link to confirmatory testing, by reported number of lines and self-interpreted
498 HIVST result.

	Overall	2 lines /reactive	1 line /reactive	2 lines /non-reactive	2 lines /DK-R	DK-R / reactive
less than 4 months	8 (10%)	3 (11%)	0 (0%)	4 (16%)	1 (5.6%)	0 (0%)
between 4 and 6 months	67 (86%)	24 (89%)	5 (71%)	21 (84%)	17 (94%)	0 (0%)
more than 6 months	3 (3.8%)	0 (0%)	2 (29%)	0 (0%)	0 (0%)	1 (100%)
Total	34 (100%)	15 (44.2%)	1 (2.9%)	9 (26.5%)	8 (23.5%)	1 (2.9%)

499 DK: don't know. R: refuse to answer

Acknowledgements

502   We wish to acknowledge the commitment and determination of all the ATLAS program teams, which
503   made this research possible. We would also like to thank the interviewers for their professionalism in
504   collecting this sensitive data. Finally, we are grateful to the participants who were kind enough to give us
505   some of their time by agreeing to take part in the survey.

Data, scripts, code, and supplementary information availability

507   Data, scripts and code are available online(https://doi.org/10.5281/zenodo.10255772) as well as the
508   survey questionnaires (https://doi.org/10.5281/zenodo.10210464).

Conflict of interest disclosure

510   The authors declare that they comply with the PCI rule of no financial conflicts of interest in relation to
511   the content of the article. They declare no conflict of interest.

Funding

This work was supported by Unitaid (Grant Number: 2018-23 ATLAS) with additional funding from
Agence Française pour le Développement (AFD). AKK benefits from an ANRS thesis allowance. MMG's
research program is supported by a Canada Research Chair (Tier 2) in Population Health Modeling. The
funding bodies were not involved in the design of the study and collection, analysis, and interpretation of
data and in writing the manuscript.

MCB and RS acknowledge funding from the MRC Centre for Global Infectious Disease Analysis (reference
MR/R015600/1), jointly funded by the UK Medical Research Council (MRC) and the UK Foreign,
Commonwealth & Development Office (FCDO), under the MRC/FCDO Concordat agreement and is also
part of the EDCTP2 programme supported by the European Union. For the purpose of open access, the
author has applied a Creative Commons Attribution (CC BY) license to any Author Accepted Manuscript
version arising.
524

References

1. Das M, Chu PL, Santos G-M, Scheer S, Vittinghoff E, McFarland W, et al. Decreases in Community
Viral Load Are Accompanied by Reductions in New HIV Infections in San Francisco. PLoS ONE.
2010;5:e11068.

2. Lima VD, Johnston K, Hogg RS, Levy AR, Harrigan PR, Anema A, et al. Expanded Access to Highly
Active Antiretroviral Therapy: A Potentially Powerful Strategy to Curb the Growth of the HIV
Epidemic. J Infect Dis. 2008;198:59–67.

3. Cohen MS, Chen YQ, McCauley M, Gamble T, Hosseinipour MC, Kumarasamy N, et al. Prevention
of HIV-1 Infection with Early Antiretroviral Therapy. N Engl J Med. 2011;365:493–505.

4. Quinn TC, Wawer MJ, Sewankambo N, Serwadda D, Li C, Wabwire-Mangen F, et al. Viral Load
and Heterosexual Transmission of Human Immunodeficiency Virus Type 1. N Engl J Med.
2000;342:921–9.

5. Hayes RJ, Donnell D, Floyd S, Mandla N, Bwalya J, Sabapathy K, et al. Effect of Universal Testing
and Treatment on HIV Incidence — HPTN 071 (PopART). N Engl J Med. 2019;381:207–18.

6. Grinsztejn B, Hosseinipour MC, Ribaudo HJ, Swindells S, Eron J, Chen YQ, et al. Effects of early
versus delayed initiation of antiretroviral treatment on clinical outcomes of HIV-1 infection: results
from the phase 3 HPTN 052 randomised controlled trial. Lancet Infect Dis. 2014;14:281–90.

7. UNAIDS. UNAIDS data 2021.Geneva: Joint United Nations Programme on
HIV/AIDS;UNAIDS/JC3032E. Geneva; 2021.

8. Joint United Nations Programme on HIV/AIDS. Fast-Track - Ending the AIDS epidemic by 2030.
Geneva; 2014.

9. Johnson C, Baggaley R, Forsythe S, van Rooyen H, Ford N, Napierala Mavedzenge S, et al. Realizing
the Potential for HIV Self-Testing. AIDS Behav. 2014;18:391–5.

10. Njau B, Covin C, Lisasi E, Damian D, Mushi D, Boulle A, et al. A systematic review of qualitative
evidence on factors enabling and deterring uptake of HIV self-testing in Africa. BMC Public Health.
2019;19:1289.

11. Hector J, Davies M-A, Dekker-Boersema J, Aly MM, Abdalad CCA, Langa EBR, et al.
Acceptability and performance of a directly assisted oral HIV self-testing intervention in adolescents in
rural Mozambique. PloS One. 2018;13:e0195391.

12. Grésenguet G, Longo J de D, Tonen-Wolyec S, Bouassa R-SM, Belec L. Acceptability and Usability
Evaluation of Finger-Stick Whole Blood HIV Self-Test as An HIV Screening Tool Adapted to The
General Public in The Central African Republic. Open AIDS J. 2017;11:101–18.

13. Harichund C, Moshabela M, Kunene P, Abdool Karim Q. Acceptability of HIV self-testing among
men and women in KwaZulu-Natal, South Africa. AIDS Care. 2019;31:186–92.

14. Kalibala S, Tun W, Cherutich P, Nganga A, Oweya E, Oluoch P. Factors Associated with
Acceptability of HIV Self-Testing Among Health Care Workers in Kenya. AIDS Behav. 2014;18:405–
14.

15. Kelvin EA, Cheruvillil S, Christian S, Mantell JE, Milford C, Rambally-Greener L, et al. Choice in
HIV testing: the acceptability and anticipated use of a self-administered at-home oral HIV test among
South Africans. Afr J AIDS Res. 2016;15:99–108.

16. Martínez Pérez G, Steele SJ, Govender I, Arellano G, Mkwamba A, Hadebe M, et al. Supervised
oral HIV self-testing is accurate in rural KwaZulu-Natal, South Africa. Trop Med Int Health.
2016;21:759–67.

17. Ky-Zerbo O, Desclaux A, Boye S, Vautier A, Rouveau N, Kouadio BA, et al. Willingness to use
and distribute HIV self-test kits to clients and partners: A qualitative analysis of female sex workers'
collective opinion and attitude in Côte d'Ivoire, Mali, and Senegal. Womens Health.
2022;18:174550572210922.

18. Kurth AE, Cleland CM, Chhun N, Sidle JE, Were E, Naanyu V, et al. Accuracy and Acceptability
of Oral Fluid HIV Self-Testing in a General Adult Population in Kenya. AIDS Behav. 2016;20:870–9.

19. Johnson CC, Kennedy C, Fonner V, Siegfried N, Figueroa C, Dalal S, et al. Examining the effects
of HIV self-testing compared to standard HIV testing services: a systematic review and meta-analysis.
J Int AIDS Soc. 2017;20:21594.

20. World Health Organization. RECOMMENDS HIV SELFTESTING – EVIDENCE UPDATE AND
CONSIDERATIONS FOR SUCCESS. Geneva: Word Health Organization; 2019.

21. Choko AT, Jamil MS, MacPherson P, Corbett E, Chitembo L, Ingold H, et al. Measuring linkage to
HIV treatment services following HIV self-testing in low-income settings. J Int AIDS Soc. 2020;23.

22. World Health Organization. Guidelines on HIV self-testing and partner notification: supplement to
Consolidated guidelines on HIV testing services. https://apps.who.int/iris/handle/10665/251655. 2016.

23. Asiimwe S, Oloya J, Song X, Whalen CC. Accuracy of Un-supervised Versus Provider-Supervised
Self-administered HIV Testing in Uganda: A Randomized Implementation Trial. AIDS Behav.
2014;18:2477–84.

24. Figueroa C, Johnson C, Ford N, Sands A, Dalal S, Meurant R, et al. Reliability of HIV rapid
diagnostic tests for self-testing compared with testing by health-care workers: a systematic review and
meta-analysis. Lancet HIV. 2018;5:e277–90.

25. Chanda MM, Ortblad KF, Mwale M, Chongo S, Kanchele C, Kamungoma N, et al. HIV self-testing
among female sex workers in Zambia: A cluster randomized controlled trial. PLOS Med.
2017;14:e1002442.

26. Tonen-Wolyec S, Filali M, Mboup S, Bélec L. HIV self-testing in Africa: stakes and challenges.
Médecine Santé Trop. 2018;28:144–9.

27. Pant Pai N, Behlim T, Abrahams L, Vadnais C, Shivkumar S, Pillay S, et al. Will an Unsupervised
Self-Testing Strategy for HIV Work in Health Care Workers of South Africa? A Cross Sectional Pilot
Feasibility Study. PLoS ONE. 2013;8:e79772.

28. Brown AN, Djimeu EW, Cameron DB. A Review of the Evidence of Harm from Self-Tests. AIDS
Behav. 2014;18:445–9.

29. Dwyer-Lindgren L, Cork MA, Sligar A, Steuben KM, Wilson KF, Provost NR, et al. Mapping HIV
prevalence in sub-Saharan Africa between 2000 and 2017. Nature. 2019;570:189–93.

30. Rouveau N, Ky-Zerbo O, Boye S, Fotso AS, d'Elbée M, Maheu-Giroux M, et al. Describing,
analysing and understanding the effects of the introduction of HIV self-testing in West Africa through
the ATLAS programme in Côte d'Ivoire, Mali and Senegal. BMC Public Health. 2021;21:181.

31. UNAIDS. UNAIDS DATA 2022. Geneva: Joint United Nations Programme on
HIV/AIDS;UNAIDS/JC3063E.https://www.unaids.org/sites/default/files/media_asset/data-book-
2022_en.pdf. Geneva: Joint United Nations Programme on HIV/AIDS; 2022.

32. World Health Organization. Consolidated guidelines on HIV prevention, testing, treatment, service
delivery and monitoring: recommendations for a public health approach. 2021 update. Geneva: World
Health Organization; 2021.

33. Lyons CE, Coly K, Bowring AL, Liestman B, Diouf D, Wong VJ, et al. Use and Acceptability of
HIV Self-Testing Among First-Time Testers at Risk for HIV in Senegal. AIDS Behav. 2019;23:130–
41.

34. Boye S, Bouaré S, Ky-Zerbo O, Rouveau N, Simo Fotso A, d'Elbée M, et al. Challenges of HIV
Self-Test Distribution for Index Testing When HIV Status Disclosure Is Low: Preliminary Results of a
Qualitative Study in Bamako (Mali) as Part of the ATLAS Project. Front Public Health. 2021;9:653543.

35. d'Elbée M, Traore MM, Badiane K, Vautier A, Simo Fotso A, Kabemba OK, et al. Costs and Scale-
Up Costs of Integrating HIV Self-Testing Into Civil Society Organisation-Led Programmes for Key
Populations in Côte d'Ivoire, Senegal, and Mali. Front Public Health. 2021;9:653612.

36. Simo Fotso A, Johnson C, Vautier A, Kouamé KB, Diop PM, Silhol R, et al. Routine programmatic
data show a positive population-level impact of HIV self-testing: the case of Côte d'Ivoire and
implications for implementation. AIDS. 2022;36:1871–9.

37. Silhol R, Maheu-Giroux M, Soni N, Fotso AS, Rouveau N, Vautier A, et al. Assessing the potential
population-level impacts of HIV self-testing distribution among key populations in Côte d'Ivoire, Mali,
and Senegal: a mathematical modelling analysis. preprint. HIV/AIDS; 2023.

38. Simo Fotso A, Kra AK, Maheu-Giroux M, Boye S, d'Elbée M, Ky-zerbo O, et al. Is it possible to
recruit HIV self-test users for an anonymous phone-based survey using passive recruitment without
financial incentives? Lessons learned from a pilot study in Côte d'Ivoire. Pilot Feasibility Stud.
2022;8:4.

39. Kra AK, Fosto AS, N'guessan KN, Geoffroy O, Younoussa S, Kabemba OK, et al. Can HIV self-
testing reach first-time testers? A telephone survey among self-test end users in Côte d'Ivoire, Mali,
and Senegal. BMC Infect Dis. 2023;22:972.

40. R Core Team. R: A Language and Environment for Statistical Computing. 2021.

41. Sjoberg D D, Whiting K, Curry M, Lavery J A, Larmarange J. Reproducible Summary Tables with
the gtsummary Package. R J. 2021;13:570.

42. Giguère K, Eaton JW, Marsh K, Johnson LF, Johnson CC, Ehui E, et al. Trends in knowledge of
HIV status and efficiency of HIV testing services in sub-Saharan Africa, 2000–20: a modelling study
using survey and HIV testing programme data. Lancet HIV. 2021;8:e284–93.

43. Maheu-Giroux M, Marsh K, Doyle CM, Godin A, Lanièce Delaunay C, Johnson LF, et al. National
HIV testing and diagnosis coverage in sub-Saharan Africa: a new modeling tool for estimating the 'first
90' from program and survey data. AIDS. 2019;33 Supplement 3:S255–69.

44. Tippett Barr B, Lowrance D, Case Johnson C, Baggaley RC, Rogers J, Balachandra S, et al.
Treatment-adjusted prevalence to assess HIV testing programmes. Bull World Health Organ.
2021;99:874–82.

45. Vautier A. La notice d'utilisation du fabricant suffit-elle dans un contexte multilingue et de faible
alphabétisation ? L'exemple de l'autodépistage du VIH en Afrique de l'Ouest. 2020.

46. Wei C, Yan L, Li J, Su X, Lippman S, Yan H. Which user errors matter during HIV self-testing? A
qualitative participant observation study of men who have sex with men (MSM) in China. BMC Public
Health. 2018;18:1108.

47. Thirumurthy H, Masters SH, Mavedzenge SN, Maman S, Omanga E, Agot K. Promoting male
partner HIV testing and safer sexual decision making through secondary distribution of self-tests by
HIV-negative female sex workers and women receiving antenatal and post-partum care in Kenya: a
cohort study. Lancet HIV. 2016;3:e266–74.

48. Chipungu J, Bosomprah S, Zanolini A, Thimurthy H, Chilengi R, Sharma A, et al. Understanding
linkage to care with HIV self-test approach in Lusaka, Zambia - A mixed method approach. PLOS
ONE. 2017;12:e0187998.

49. Tun W, Vu L, Dirisu O, Sekoni A, Shoyemi E, Njab J, et al. Uptake of HIV self-testing and linkage
to treatment among men who have sex with men (MSM) in Nigeria: A pilot programme using key
opinion leaders to reach MSM. J Int AIDS Soc. 2018;21:e25124.

50. Hlongwa M, Hlongwana K, Makhunga S, Choko A, Dzinamarira T, Conserve D, et al. Linkage to
HIV care following HIV self-testing among men: systematic review of quantitative and qualitative
studies from six countries in sub-Saharan Africa. preprint. In Review; 2022.

51. Choko AT, MacPherson P, Webb EL, Willey BA, Feasy H, Sambakunsi R, et al. Uptake, Accuracy,
Safety, and Linkage into Care over Two Years of Promoting Annual Self-Testing for HIV in Blantyre,
Malawi: A Community-Based Prospective Study. PLOS Med. 2015;12:e1001873.

52. Green KE, Vu BN, Phan HT, Tran MH, Ngo HV, Vo SH, et al. From conventional to disruptive:
upturning the HIV testing status quo among men who have sex with men in Vietnam. J Int AIDS Soc.
2018;21 Suppl 5:e25127.

53. Ky-Zerbo O, Desclaux A, Kouadio AB, Rouveau N, Vautier A, Sow S, et al. Enthusiasm for
Introducing and Integrating HIV Self-Testing but Doubts About Users: A Baseline Qualitative Analysis
of Key Stakeholders' Attitudes and Perceptions in Côte d'Ivoire, Mali and Senegal. Front Public Health.
2021;9.

54. Kouassi AK, Fosto AS, N'Guessan NK, Geoffroy O, Younoussa S, Kabemba OK, et al. Reaching
key and peripheral populations: a phone-based survey of HIV self-test users in West Africa.Durban:
poster #PEC004.https://hal.science/hal-04121478. 2021.

55. Inghels M, Kouassi AK, Niangoran S, Bekelynck A, Carilon S, Sika L, et al. Preferences and access
to community-based HIV testing sites among men who have sex with men (MSM) in Côte d'Ivoire.
BMJ Open. 2022;12:e052536.