Socioeconomic inequity in access to care among older people in Japan

Shohei Okamoto[1*], Atsuhiro Yamada[2], Erika Kobayashi[1], Jersey Liang[3, 4]

[1]Research Team for Social Participation and Healthy Aging, Tokyo Metropolitan Institute for Geriatrics and Gerontology, Tokyo, Japan
[2]Faculty of Economics, Keio University, Tokyo, Japan
[3]Department of Health Management and Policy, School of Public Health, University of Michigan, Michigan, USA
[4]Department of Healthcare Management and Healthy Aging Research Center, Chang Gung University, Taiwan

*Corresponding author: Dr Shohei Okamoto, Research Team for Social Participation and Healthy Aging, Tokyo Metropolitan Institute for Geriatrics and Gerontology: 35-2 Sakae-cho, Itabashi-ku, Tokyo, Japan 1730015 (E-mail: shohei@z2.keio.jp)

Declarations
Ethics approval and consent to participate
This study was approved by the institutional review board of the Tokyo Metropolitan Geriatric Hospital and Institute of Gerontology (R21-008). All participants provided their informed consent to participate in the study.

Consent for publication
Consent for publication was not required because all information collected through the surveys was deidentified.

Availability of data and materials
The data used for this study are partially accessible through the webpage of the Centre for Social Research and Data Archives, Institute of Social Sciences of The University of Tokyo: https://csrda.iss.u-tokyo.ac.jp/english/infrastructure/

Competing interests
The authors declare no competing interests associated with this study.

Funding
This study received funding from the World Health Organization Centre for Health Development (WHO Kobe Centre: K21003), which also provided technical support in reviewing an earlier draft of this manuscript. Also, the study was supported by Grants-in-Aid for Scientific Research (No. 20H00091 and 23H00063) and by the Open Research Areas (ORA) program by the Japan Society for the Promotion of Science (IN-CARE).

Authors' contributions
SO conceptualised the study, conducted statistical analysis, and prepared a first draft, which was refined by AY, EK, and JL. SO, AY, EK, and JL contributed to the design of the survey and data collection. All authors have reviewed and agreed to the final draft of the submitted paper.

Acknowledgements
We are grateful for the helpful comments from Drs Megumi Rosenberg and Shinichi Tomioka at the World Health Organization Centre for Health Development.

Abstract

Background
Equity in access to long-term care (LTC) enables older people to maintain their well-being after they undergo a decline in their intrinsic capacity.

Methods
We used data from Wave 6 (2002) through Wave 10 (2021) of the National Survey of the Japanese Elderly to assess gradients by income and education in access to medical care and LTC among Japanese individuals aged 60 years and above. Specifically, we assessed self-reported unmet needs for medical care and LTC, and public LTC use, and estimated the concentration indices (CI) to measure the degree of inequality and inequity. We standardised public LTC use by need and non-need variables. We analysed data derived from up to 1,775 person-wave observations from 1,370 individuals.

Findings
The pooled incidence across waves of forgone medical care, self-reported unmet support for activities of daily living (ADL) or instrumental ADL (IADL), and those not certified for LTC services even with ADL or IADL limitations were 4.6%, 15.5%, and 62.5%, respectively. Public LTC use demonstrated pro-higher education and pro-rich distribution, whereas the gaps decreased for need-predicted use. Based on the CI estimates, no explicit inequality was found for forgone medical care. However, we observed inequity in standardised LTC use across education, indicating pro-higher education inequality, particularly among women and those aged ≥80 years.

Conclusion
Improving the understanding of available resources and strengthening the functions of health centres and communities are required to detect the needs of citizens and facilitate their access to necessary care.

Background
Universal health coverage (UHC) means everyone, everywhere can access nationally determined sets of needed essential health services, including promotion, prevention, treatment, rehabilitation, and palliative care without incurring financial hardship (1). Older people may require more care to maintain their functional ability (i.e. the interaction of intrinsic capacity with environmental characteristics), human dignity, and well-being because their intrinsic capacity (i.e. a combination of their physical and mental capacities) declines (2). Ageing population is a global trend (3); therefore, ensuring equitable access to necessary care, including long-term care (LTC) for older people, is essential to achieve progress toward UHC.

LTC includes numerous services required by people dependent on assistance with the basic activities of daily living (ADL) (4). Among the OECD countries, the total LTC expenditure per country's gross domestic product ranges from 0.1% to 4.1%, in addition to differences in care providers (e.g. nursing homes, hospitals, home care, households, and social providers) (5). To date, only six countries have adopted national social LTC insurance systems, namely the Netherlands, Israel, Germany, Luxembourg, South Korea, and Japan (6). Japan is the first Asian country, which has adopted a public universal LTC insurance in 2000, principally covering all citizens aged ≥40 years (7). To be eligible for public LTC benefits, people requiring care undergo an objective test to evaluate their level of ADL dependence and an assessment by their attending physician and are classified into independence, support-need-levels 1 to 2, and LTC need-levels 1 to 5. The types and amounts of available benefits are determined based on this classification. The users bear charges for LTC services at a 10% co-payment rate (or 20% or 30% depending on one's income level). In addition, several financial protection policies are available to mitigate excessive out-of-pocket payments and the burden of insurance premiums for low-income individuals. Further details on LTC insurance in Japan have been described previously (7).

Socioeconomic inequity in access to care and unmet LTC needs
Despite evidence of socioeconomic inequity in access to healthcare (8-10), there is limited evidence of such inequity in LTC. Based on the Grossman model (11), socioeconomic inequity in access to healthcare can arise from various channels, such as disparities in health literacy, income, time preference, and available time for producing health. However, inequitable access to LTC can be driven by different channels from healthcare, including due to substitution between formal and informal care. Those unable to access the necessary care experience unmet needs, which could lead to negative health outcomes (12, 13); or ensuring access to LTC can have a protective effect for increased medical care expenditures, such as one for emergency care use (14). Women, lower education, and worse economic status predict unmet needs for personal assistance in ADL among older adults (15, 16). Inequity in LTC access exists in South Korea and the US, suggesting the importance of extending insurance coverage and subsidising low-income individuals (17-19). Formal care among people without dementia is pro-rich, and poor people are more likely to experience unmet care needs in England (20). Furthermore, findings from Spain and the Survey of Health, Aging, and Retirement in Europe suggest that formal services are concentrated among richer people, whereas poorer people tend to use intensive informal care (21, 22). Studies in the Netherlands also find a pro-poor gradient in home care use (23, 24). However, a Japanese study found that formal LTC use was generally equitable across income groups (25).

LTC services may still exhibit economic inequity despite the availability of universal insurance or benefits, as suggested by studies in South Korea and European countries (17, 18, 22). Nevertheless, income inequity in LTC service use among those eligible for LTC benefits has not been observed in Japan (25), potentially owing to its low co-payment rate and financial protection policies. However, as mentioned before, socioeconomic inequity in care access can arise from financial and non-financial causes. Under the social insurance scheme, users need to contract with providers to purchase services at a (quasi) market, which would require a certain level of knowledge about LTC services and their own preferences in service use. Furthermore, in Japan, individuals are required to submit an application to be eligible for public LTC services, which may serve as a hindrance to accessing care for non-financial reasons. Therefore, we aimed to expand the literature in assessing

socioeconomic inequity in LTC access using both financial and non-financial indicators under the public universal LTC insurance system.

Methods
Data
Data were obtained from the National Survey of the Japanese Elderly (26), comprising Japanese adults aged ≥60 years. The survey was first conducted in 1987 and was followed up with participants every 3 to 6 years, adding new samples to complement sample size declines caused by deaths and a loss to follow-up. The sample was extracted from the Basic Resident Registration System using a stratified two-stage random sampling method based on a combination of regional blocks and population. From waves 1 to 10, 7,892 individuals responded in one or more waves. To evaluate access to LTC services, we analysed data collected from wave 6 (2002) to wave 10 (2021), after public LTC insurance introduction in 2000.

We restricted our analysis to participants who responded to interviews, excluding mail, proxy, and non-responders' surveys because LTC-need-related information and self-reported unmet needs were only available for these people. Thus, we obtained 10,743 person-wave observations from 5,471 unique, non-institutionalised individuals from Wave 6 to Wave 10. We then focused on those with LTC needs, leading to 1,975 person-wave observations by 1,512 individuals. Finally, we excluded those with missing information, resulting in the final sample size of at most 1,775 person-wave observations by 1,370 individuals.

Access to medical care: Forgone care

To assess the inequality in medical care access, we used self-reported forgone medical care (10, 27). In the survey, for the question, 'During the past 3 months, how often did you reduce the dose of medication or did not see a physician even though it was necessary?', each respondent selected one of the following options: 1. most times; 2. sometimes; 3. hardly/none; and 4. they did not need to consult a physician or consume medicine. Respondents who selected 'most times' or 'sometimes' were regarded as forgoing needed care. We excluded those who did not need to consult a physician or consume medicine. To assess inequality in medical care access, we analysed the data obtained in waves 8, 9, and 10, because the questionnaire was administered only in these waves.

Access to LTC

(1) Self-reported unmet needs

Respondents with LTC needs were questioned if someone else (e.g. family members and LTC workers) had helped them with their activities within the past 3 months. They rated the frequency of receiving support as follows: 1. almost always; 2. sometimes; 3. occasionally; 4. never; and 5. did not need support. Respondents who selected 'sometimes', 'occasionally', or 'never' were categorised as having experienced an unmet need for LTC. This operational definition is reasonable because people who cannot obtain necessary care whenever they need them experience unmet need.

To define the LTC needs operationally, we used questionnaires on difficulties in performing basic and instrumental activities of daily living (ADL and IADL). Similar to the Katz index (28), ADL was measured using the following six items: bathing, dressing, feeding, transferring, outing, and toileting. IADL was measured using the following four items: shopping for personal items, using a telephone, riding the bus or subway alone, and performing light tasks around the house (29). Each item was rated on a five-point Likert scale, ranging from 0 (never difficult) to 4 (unable to do at all). Respondents experiencing difficulties with at least one of the items (i.e. 2. moderately difficult to 5. unable to do it at all) were defined as having LTC needs.

(2) Utilisation-based measurement

Considering health disparities across socioeconomic statuses (30, 31), the care need differs across groups, leading to dissimilar levels of care utilisation. Thus, higher utilisation among the poor with worse health status does not imply pro-poor inequity in care access based on the vertical equity principle. Therefore, standardised utilisation must be used to assess inequity in access to care (32).

We standardised LTC use by the LTC need explained by demographics, health status, and morbidity status (32), formalised as a non-linear model with repeated measures (i.e. multilevel mixed-effects logistic regression) as follows:

$$y_{it} = \frac{\exp\left(\alpha + \sum_j \beta_j x_{jit} + \sum_k \gamma_k z_{kit} + u_t\right)}{1 + \exp\left(\alpha + \sum_j \beta_j x_{jit} + \sum_k \gamma_k z_{kit} + u_t\right)}$$

where y_{it} denotes a binary variable corresponding to 1 if respondent *i* in year *t* is considered eligible for public LTC services at any level, and 0 otherwise. The *j*th need variables (x) to predict LTC use and *k*th control variables (z) to avoid omitted-variables bias, with parameters β and γ, are included. α is a constant, and u is the random effects. Accepting that the variance of need-standardised use may depend on the formulation of the z variables in the standardisation procedure, need-standardised LTC use ($\hat{y}_{it}^{ST}$) is defined as:

$$\hat{y}_{it}^{ST} = y_{it} - \frac{\exp\left(\hat{\alpha} + \sum_j \hat{\beta}_j x_{jit} + \sum_k \hat{r}_k \bar{z}_k + u_t\right)}{1 + \exp\left(\hat{\alpha} + \sum_j \hat{\beta}_j x_{jit} + \sum_k \hat{r}_k \bar{z}_k + u_t\right)} + \frac{1}{n}\sum_i \frac{\exp\left(\hat{\alpha} + \sum_j \hat{\beta}_j x_{jit} + \sum_k \hat{r}_k \bar{z}_k + u_t\right)}{1 + \exp\left(\hat{\alpha} + \sum_j \hat{\beta}_j x_{jit} + \sum_k \hat{r}_k \bar{z}_k + u_t\right)}$$

where n denotes the sample size, and the z variables are set to their means ($\bar{z}$). The need variables directly influencing LTC needs are proxied by the demographic and morbidity characteristics (32), including the age, sex, self-rated health, the number of chronic conditions (ranging from 0 to 6), the degree of difficulty in performing each ADL (0–24) and IADL (0–16), and cognitive functioning (0–9). The non-need control variables include the marital status, the number of co-resident members, residential area, the population size of the residential area, and year dummies. These non-need variables may affect the health status through social support and relationships (33); however, they are expected to affect formal and informal care availability, generating potential heterogeneity across group behaviour in the application and utilisation of formal LTC services. The need and non-need variables are defined in the Appendix Table A-1.

Empirical strategy
Concentration index
To measure inequality and horizontal inequity in the access to medical and LTC services across socioeconomics groups, we estimated the concentration indices (CI) frequently used to evaluate health disparities (32). CI in each year is defined as twice the area between the concentration curve and the 45° line as follows (32):

$$C = \frac{2cov(h_i, R_i)}{\bar{h}} = \frac{1}{n}\sum\left\{\frac{h_i}{\bar{h}}(2R_i - 1)\right\}$$

where h_i denotes the health variable of interest to measure inequity, and $\bar{h}$ is its mean. R_i is the rank variable in which the health gradients are measured. For binary outcomes, the concentration index (W) was rescaled to consider its dependence on the variable means and to satisfy the mirror condition, i.e. the absolute value of a measured inequality is symmetric when computed over either attainments or shortfalls as follows (34):

$$W = \frac{1}{n}\sum\left\{\frac{(a^{max} - a^{min})a_i}{(a^{max} - \bar{a})(\bar{a} - a^{min})}(2R_i - 1)\right\}$$

The CI is positive when access is more frequent among higher socioeconomic groups and negative if otherwise. We used the years of education and income as the rank variables to measure inequality and inequity. For income, we used the couple's gross annual income equalised by the marital status (i.e. divided by the square root of 2 if married) and residualised by employment status (See Appendix B). To calculate the descriptive differences in LTC utilisation across education and income groups, but not the CI, both indicators were divided into five categories. The category for education included lowest (<6), lower-middle (6–9), middle (9–12), higher-middle (12–15), and higher (16+). Despite different pre- and post-World War II educational systems, this categorisation roughly corresponded to the distinctions between elementary, secondary, high school, and university or higher. Income was divided into 20th percentile groups.

Demographic factors, such as sex and age, can influence help-seeking behaviours (35); thus, we evaluated inequities according to the sex and age. Considering that the average age of the sample was approximately 81, we categorised the participants into those aged <80 years and ≥80 years. This categorisation is reasonable because people can become drastically dependent on ADL around the age of 80 years (36).

To partially adjust for potential selection bias, we adopted two types of weights, namely, cross-sectional and longitudinal. Cross-sectional weights were estimated by logistic regression as probabilities of responding to baseline surveys predicted by the age, sex, the geographic area of residence, and the municipal/population category of a residential area. Longitudinal weights were estimated as response probabilities in each wave, estimated by the age, sex, employment status, marital status, education, self-rated health, the geographic area and population category of a residential area at baseline or the closest. These approaches were similar to multiple imputations based on random missing data (37, 38).

Results
Descriptive statistics
Table 1 summarises the incidence of forgone medical care and unmet LTC needs in each wave. The average incidence for medical care was 4.6% across the waves as follows: 4.5%, 6.0%, and 3.8% in 2012, 2017, and 2021, respectively. The average incidence for ADL or IADL support was 41.6%, ranging from 37.9% to 44.6% from 2002 to 2021. Approximately, 62.0% of the participants with ADL or IADL limitations were not certified for LTC services (55.3–71.8%).

Table 2 defines the pooled descriptive statistics between Wave 6 (2002) and Wave 10 (2021) for those with LTC needs. Of all participants, 38.0% were certified for public LTC use. The average age was approximately 80.8 years, and the majority of participants were women (72.6%).

Appendix Table A-2 presents the regression results to predict LTC use by need and non-need variables. Both needs (e.g. chronic conditions and ADL and IADL limitations) and non-needs (e.g. the number of co-resident members) were associated with the certification for public LTC use.

Table 3 summarises the mean probabilities of non-standardised, need-predicted, and need-standardised LTC use by education and income, pooling the data between waves 6 and 10. Remarkably, the actual distribution of public LTC was pro-higher-education and pro-rich, whereas the gaps decreased for need-predicted use. Thus, the participants with lower education and income used public LTC services less than expected by 3% to 7% points, whereas their counterparts with middle and higher levels used services more than expected by 3% to 26% points. Despite standardisation, higher education and income groups used more services than the lowest groups.

Socioeconomic inequality and inequity in access to medical care and LTC
Table 4 summarises CIs for medical care and LTC use by education and income in each wave. Figure 1 presents visualised horizontal inequities in LTC use by education and income. We observed no explicit inequality for forgone medical care. However, inequity was observed in LTC use across education in 2006 and 2012, indicating pro-higher-education inequality: CIs with standard errors were

0.12 (0.03) and 0.09 (0.03). Meanwhile, the inequity was less obvious for income, showing that concentration curves crossed 45-degree lines in most waves and pro-rich inequality was evident only in 2012. For non-need-standardised self-reported unmet needs for LTC support, both education and income inequality were evident in 2002.

Heterogeneity by sex and age
Standardised LTC use was pro-higher-education and pro-rich for both men and women (Appendix Table A-3). CI-based inequities in each wave were particularly evident among women, corresponding to the results of our primary analysis (Appendix Table A-4). Regarding the heterogeneity by age, we observed similar pro-higher education, particularly among those aged 80 years (Appendix A-5 and A-6).

Discussion
This study aimed to assess unmet need and socioeconomic inequity in LTC access and found three main findings. First, more than one-third of the participants underwent unmet needs for LTC. Second, LTC access was pro-higher-education, which was more evident than income inequity. Third, pro-higher-educational inequity was evident among women and the oldest-old population, despite being descriptively similar among their male and younger (i.e. 60–80 years) counterparts.

Japan has the highest UHC achievement (39); nonetheless, challenges exist with certain proportions of people experiencing unmet medical and LTC needs. People with higher education may be better at accessing the necessary services, considering health literacy is related to education (40) and predicts a lack of understanding and use of health services (41). For medical care, older people might not experience unmet needs frequently due to the absence of gatekeeping and low copayment rates.

Educational inequity in LTC use was more pronounced among women and the old. This may be explained by different help-seeking behaviours by men and women despite similar health needs. Women are more likely to seek and use care than men (42); therefore, disparities could be more apparent when a specific group forgoes it.

Our findings generate policy implications for enhancing access to care. First, barriers to accessing LTC or relevant services may exist because people need to make an application by themselves. In Japan's policies for care for older people, community comprehensive support centres located in each municipality play an important role in care management and facilitating the use of necessary care. Nevertheless, reaching out to these centres still requires care seeking from people with some difficulties themselves or their family members For some older people, a decline in intrinsic capacity and care needs may be detected by health check-ups or their attending physicians. However, poor health is associated with social isolation (43); therefore, people with health issues may be unable to access the necessary care because of physical barriers (e.g. the lack of transportation) or health-literacy-related issues (e.g. the lack of knowledge about how to seek care). Therefore, it is necessary to enhance access through any of these channels, such as education, to improve an understanding of available resources, strengthening the functions of health centres and communities to detect the needs of citizens, and facilitating their access to necessary care.

This study had four limitations. First, the sample size was small because we focused only on individuals with LTC needs. Therefore, we may not obtain significant results for some analyses or may be unable to generalise our findings. Second, our measure of LTC certification was self-reported and did not include information on the number of services used. Further studies should address these limitations by using extensive administrative data. Third, we could not distinguish between unmet needs for formal and informal LTC because of data restrictions. Depending on the policy context in each country, the availability and service coverage of formal long-term care services largely vary. Therefore, assessing the unmet needs for both formal and informal care comprehensively is required to design the service coverage of formal care. Fourth, there were non-responders, particularly in Wave 10. Thus, we may have missed vulnerable populations during the state of emergency in large cities, by which we cannot conclude that inequity was absent during the pandemic.

Conclusions

In conclusion, approximately 42% of individuals with LTC needs experienced an unmet need for ADL or IADL support, and approximately 62% were not certified for public LTC use even with ADL or IADL limitations. Inequity in care access was driven by education rather than income. Even if the services are universally available, financial and non-financial supports are indispensable to ensuring all people can access care whenever and wherever they need them.

References

1. World Health Organization, World Bank. Tracking universal health coverage: 2021 global monitoring report. 2021.
2. World Health Organization. World report on ageing and health. 2015.
3. United Nations, Department of Economics and Social Affairs, Population Division. World population prospects 2022. New York: 2022.
4. OECD. Long-term Care for Older People. Paris: OECD Publishing; 2005.
5. OECD. Ageing and long-term care. Health at a Glance 2021: OECD Indicators. Paris: OECD Publishing; 2021.
6. Fischer J. The social long-term care insurance model: comparing actor configurations across countries and time. Journal of International and Comparative Social Policy. 2022;38(2):93-110. Epub 2022/08/23.
7. Okamoto S, Komamura K. Towards universal health coverage in the context of population ageing: a narrative review on the implications from the long-term care system in Japan. Arch Public Health. 2022;80(1):210. Epub 20220921.
8. Pulok MH, van Gool K, Hajizadeh M, Allin S, Hall J. Measuring horizontal inequity in healthcare utilisation: a review of methodological developments and debates. Eur J Health Econ. 2020;21(2):171-80. Epub 20190921.
9. Albert-Ballestar S, Garcia-Altes A. Measuring health inequalities: a systematic review of widely used indicators and topics. Int J Equity Health. 2021;20(1):73. Epub 20210310.
10. Rahman MM, Rosenberg M, Flores G, Parsell N, Akter S, Alam MA, et al. A systematic review and meta-analysis of unmet needs for healthcare and long-term care among older people. Health Econ Rev. 2022;12(1):60. Epub 20221209.
11. Grossman M. The Human Capital Model. In: Culyer AJ, Newhouse JP, editors. Handbook of Health Economics: Elsevier; 2000. p. 347-408.
12. Zhen Z, Feng Q, Gu D. The Impacts of Unmet Needs for Long-Term Care on Mortality Among Older Adults in China. Journal of Disability Policy Studies. 2013;25(4):243-51.
13. LaPlante MP, Kaye HS, Kang T, Harrington C. Unmet need for personal assistance services: estimating the shortfall in hours of help and adverse consequences. J Gerontol B Psychol Sci Soc Sci. 2004;59(2):S98-S108.
14. Rapp T, Chauvin P, Sirven N. Are public subsidies effective to reduce emergency care? Evidence from the PLASA study. Soc Sci Med. 2015;138:31-7. Epub 20150523.
15. Zhu H. Unmet needs in long-term care and their associated factors among the oldest old in China. BMC Geriatr. 2015;15:46. Epub 20150412.
16. Liu Y-H, Chang H-J, Huang C-C. The Unmet Activities of Daily Living (ADL) Needs of Dependent Elders and their Related Factors: An Approach from Both an Individual- and Area-Level Perspective. International Journal of Gerontology. 2012;6(3):163-8.
17. Kim H, Kwon S, Yoon NH, Hyun KR. Utilization of long-term care services under the public long-term care insurance program in Korea: Implications of a subsidy policy. Health Policy. 2013;111(2):166-74. Epub 20130524.
18. Park JM. The Determinants of Long-Term Care Utilization and Equity of Access to Care among Older Adults in Dong-Ku of Incheon Metropolitan City, South Korea. Asia Pacific Journal of Public Health. 2005;17(2):104-9.
19. Park JM. Equity of access to long-term care among the American elderly. Journal of Public Health. 2003;11(2):121-30.
20. Hu B, Read S, Wittenberg R, Brimblecombe N, Rodrigues R, Banerjee S, et al. Socioeconomic inequality of long-term care for older people with and without dementia in England. Ageing and Society. 2022:1-21.
21. Garcia-Gomez P, Hernandez-Quevedo C, Jimenez-Rubio D, Oliva-Moreno J. Inequity in long-term care use and unmet need: two sides of the same coin. J Health Econ. 2015;39:147-58. Epub 20141122.
22. Lera J, Pascual-Saez M, Cantarero-Prieto D. Socioeconomic Inequality in the Use of Long-Term Care among European Older Adults: An Empirical Approach Using the SHARE Survey. Int J Environ Res Public Health. 2020;18(1). Epub 20201222.

23. Tenand M, Bakx P, van Doorslaer E. Equal long-term care for equal needs with universal and comprehensive coverage? An assessment using Dutch administrative data. Health Econ. 2020;29(4):435-51. Epub 20200120.

24. Tenand M, Bakx P, van Doorslaer E. Eligibility or use? Disentangling the sources of horizontal inequity in home care receipt in the Netherlands. Health Econ. 2020;29(10):1161-79. Epub 20200708.

25. Endo H, Yamada A. The equity of the long-term care insurance in Japan. Japanese journal of health economics & policy. 2007;19(2):147-67.

26. JAHEAD/NSJE Project Group. Japanese Aging and Health Dynamics Study: National Survey of the Japanese Elderly. https://www2.tmig.or.jp/jahead/eng/index.html (Accessed: 14 May 2020).

27. Gibson G, Grignon M, Hurley J, Wang L. Here comes the SUN: Self-assessed unmet need, worsening health outcomes, and health care inequity. Health Econ. 2019;28(6):727-35. Epub 2019/04/26.

28. Katz S, Downs TD, Cash HR, Grotz RC. Progress in development of the index of ADL. Gerontologist. 1970;10(1):20-30.

29. Lawton MP, Brody EM. Assessment of Older People: Self-Maintaining and Instrumental Activities of Daily Living. The Gerontologist. 1969;9(3 Part 1):179-86.

30. Adler NE, Ostrove JM. Socioeconomic status and health: What we know and what we don't. Ann N Y Acad Sci. 1999;896(1):3-15.

31. Feinstein JS. The Relationship between Socioeconomic Status and Health: A Review of the Literature. The Milbank Quarterly. 1993;71(2).

32. O'Donnell O, Van Doorslaer E, Wagstaff A, Lindelow M. Analyzing Health Equity Using Household Survey Data : A Guide to Techniques and Their Implementation. Washington DC: World Bank; 2008.

33. Reblin M, Uchino BN. Social and emotional support and its implication for health. Curr Opin Psychiatry. 2008;21(2):201-5.

34. Wagstaff A. The bounds of the concentration index when the variable of interest is binary, with an application to immunization inequality. Health Econ. 2005;14(4):429-32.

35. Mackenzie CS, Gekoski WL, Knox VJ. Age, gender, and the underutilization of mental health services: the influence of help-seeking attitudes. Aging Ment Health. 2006;10(6):574-82.

36. Ishizaki T, Kai I, Kobayashi Y, Matsuyama Y, Imanaka Y. The effect of aging on functional decline among older Japanese living in a community: a 5-year longitudinal data analysis. Aging Clin Exp Res. 2004;16(3):233-9.

37. Jones AM, Koolman X, Rice N. Health-related non-response in the British Household Panel Survey and European Community Household Panel: using inverse-probability-weighted estimators in non-linear models. Journal of the Royal Statistical Society: Series A (Statistics in Society). 2006;169(3):543-69.

38. Hofler M, Pfister H, Lieb R, Wittchen HU. The use of weights to account for non-response and drop-out. Soc Psychiatry Psychiatr Epidemiol. 2005;40(4):291-9.

39. World Health Organization. Global Health Observatory n.d. [cited 2022 13 April]; Available from: https://www.who.int/gho/ncd/risk_factors/physical_activity/en/.

40. Paasche-Orlow MK, Parker RM, Gazmararian JA, Nielsen-Bohlman LT, Rudd RR. The prevalence of limited health literacy. J Gen Intern Med. 2005;20(2):175-84.

41. Berkman ND, Sheridan SL, Donahue KE, Halpern DJ, Crotty K. Low health literacy and health outcomes: an updated systematic review. Ann Intern Med. 2011;155(2):97-107.

42. Bertakis KD, Azari R, Helms LJ, Callahan EJ, Robbins JA. Gender differences in the utilization of health care services. J Fam Pract. 2000;49(2):147-52. Epub 2000/03/16.

43. Coyle CE, Dugan E. Social isolation, loneliness and health among older adults. J Aging Health. 2012;24(8):1346-63. Epub 20120923.

Table 1. Incidence of unmet need for medical and long-term care

		2002	2006	2012	2017	2021	Total
Forgone medical care	%			4.5%	6.0%	3.8%	4.6%
	Total, N			2,044	1,264	1,542	4,850
Unmet need for ADL or IADL support	%	44.6%	40.4%	41.4%	37.9%	41.9%	41.6%
	Total, N	442	385	471	320	157	1,775

Non-certified for LTC services even with ADL or IADL limitations	%	71.8%	60.2%	59.1%	55.3%	58.1%	62.0%
	Total, N	442	385	471	320	157	1,775

Note. These are weighted by both cross-sectional and longitudinal weights.

Table 2. Pooled descriptive statistics, Wave 6 (2002) to Wave 10 (2021)

Stats	% or mean	Standard Deviation		%
Certified for long-term care use	38.0%		Residential area (Geographic region of Japan)	
Standardised long-term care use	0.38	0.44	Hokkaido	3.6%
Years of education	9.28	2.58	Tohoku	10.1%
Couple's income (10,000JPY)	175.21	152.43	Kanto	18.8%
Age	80.80	6.28	Hokuriku	7.1%
Women	72.6%		Tozan	6.9%
Employment status: Working	5.1%		Tokai	8.0%
SRH: Very good	11.5%		Kinki	13.2%
SRH: Good	39.8%		Chugoku	9.7%
SRH: Fair	35.6%		Shikoku	4.3%
SRH: Bad	11.1%		Kyushu	18.3%
SRH: Very bad	2.0%		Municipal/population category: Government-designated	13.0%
N of chronic conditions	2.02	0.99	Population size: 200K+	20.0%
ADL score (0–24)	2.77	3.87	Population size: 100K–200K	16.4%
IADL score (0–16)	4.79	4.22	Population size: <100K	28.1%
Memory test (0–9)	1.89	1.71	Population size: Town and villages	22.5%
N of household members	1.94	1.69		
Marital status: Single	55.1%			

Note: SRH stands for self-rated health; ADL stands for activities of daily living; IADL stands for instrumental activities of daily living; government-designated cities have populations greater than 500,000 and have been designated by the Cabinet of Japan; these descriptive statistics are calculated for 1,775 respondents with long-term care needs (i.e. ADL or IADL limitations), weighted by both cross-sectional and longitudinal weights; for couples' income, the sample size is 1,375; individuals whose marital status is single include those divorced and widowed.

Table 3. Non-standardised and standardised long-term care use by education and income: Mean probabilities, Pooled between Wave 6 (2002) and Wave 10 (2021)

		Actual	Need-predicted	Difference	Standardised
Education	Lowest: <6	0.30	0.37	-0.07	0.31
	Lower middle: 6–9	0.37	0.39	-0.03	0.35
	Middle: 9–12	0.46	0.38	0.08	0.45
	Higher middle: 12–15	0.42	0.39	0.03	0.40
	Higher: 16+	0.70	0.44	0.26	0.64
Income (Quintile)	Poorest 20%	0.34	0.37	-0.03	0.34
	2nd poorest 20%	0.35	0.38	-0.02	0.35
	Middle	0.42	0.38	0.04	0.42
	2nd richest 20%	0.49	0.38	0.10	0.48
	Richest 20%	0.42	0.37	0.05	0.42

Table 4. Concentration indices for medical and long-term care use

			2002	2006	2012	2017	2021
Forgone medical care	Education	Concentration index			-0.02	0.11	-0.02
		SE			0.07	0.07	0.08
		P-value			0.73	0.11	0.80
		N			2,044	1,264	1,542
	Income	Concentration index			-0.09	-0.00	0.04
		SE			0.08	0.08	0.09
		P-value			0.26	0.99	0.63
		N			1,744	1,103	1,224
Unmet need for long-term care	Education	Concentration index	-0.13	0.03	-0.00	-0.04	0.05
		SE	0.06	0.06	0.06	0.07	0.09
		P-value	<0.05	0.60	0.99	0.55	0.58
		N	442	385	471	320	157
	Income	Concentration index	-0.14	-0.16	-0.07	-0.03	0.07
		SE	0.06	0.07	0.07	0.08	0.12
		P-value	<0.05	<0.05	0.30	0.71	0.54
		N	376	300	381	256	118
Non-standardised long-term care use	Education	Concentration index	0.11	0.22	0.12	-0.00	0.16
		SE	0.06	0.06	0.06	0.07	0.10
		P-value	0.08	<0.01	<0.05	0.96	0.09
		N	442	385	471	320	157
	Income	Concentration index	0.07	0.08	0.14	0.07	0.08
		SE	0.07	0.07	0.07	0.08	0.11
		P-value	0.34	0.25	<0.05	0.42	0.49
		N	376	300	381	256	118
Standardised long-term care use	Education	Concentration index	0.07	0.12	0.09	-0.00	0.08
		SE	0.04	0.03	0.03	0.04	0.05
		P-value	0.05	<0.01	<0.01	0.94	0.13
		N	442	385	471	320	157
	Income	Concentration index	0.04	0.05	0.08	0.03	0.06
		SE	0.04	0.04	0.04	0.04	0.06
		P-value	0.32	0.17	<0.05	0.43	0.33
		N	376	300	381	256	118

Note: Standard errors (SE) are adjusted for clusters in each respondent; Wagstaff indices are presented; and estimates are weighted by both cross-sectional and longitudinal weights.

Figure 1. Concentration curves on standardised long-term care use by education and income

Note: The 95% confidence interval (CI; grey area) is calculated based on standard errors (SE) adjusted for clusters for each respondent; estimates are weighted by both cross-sectional and longitudinal weights. L(p) is the concentration curve for need-standardised long-term care (LTC) use.

Supplementary material for
Socioeconomic inequity in access to care among older people in Japan

Contents

Appendix A: Tables

Appendix B: Income Residualisation

Appendix A

Appendix Table A-1. A list of need and non-need variables

Classification	Variable	Description
Need	Age	Respondent's age
	Gender	Respondent's gender
	Self-rated health	1: Very bad, 2: Bad, 3: Fair, 4: Good, 5: Very good
	Chronic conditions	The number of chronic conditions respondents suffer from, including chronic respiratory diseases, hypertension, heart diseases, diabetes, stroke, and cancer (0–6)
	Activities of daily living (ADL)	A sum of the magnitudes of difficulties in performing bathing, dressing, feeding, transferring, outing, and toileting, ranging from 1: Never difficult to 5: Unable to do at all (0–24)
	Instrumental activities of daily living (IADL)	A sum of the magnitudes of difficulties in shopping for personal items, using a telephone, riding the bus or subway alone, and performing light tasks around the house, ranging from 1: Never difficult to 5: Unable to do at all (0–16)
	Cognitive functioning	The number of misreporting of cognitive functioning assessments comprising respondent's home address, interview date, interview day, mother's maiden name, name of the current prime minister, name of the previous prime minister, a simple calculation, respondent's birthday, and respondent's age (0–9)
Non-need	Marital status	1 if single (including unmarried, divorced, and widowed); 0 otherwise
	Co-residence	The number of family members living together with a respondent
	Residential area (Geographic region of Japan)	Hokkaido, Tohoku, Kanto, Hokuriku, Tozan, Tokai, Kinki, Chugoku, Shikoku, and Kyushu
	Municipal/population category of residential area	Government-designated, 200K+, 100K-200K, <100K, and Towns and villages
	Year	Wave 6 (2002), Wave 7 (2006), Wave 8 (2012), Wave 9 (2017), Wave 10 (2021)

Appendix Table A-2. Regression results: Prediction of long-term care use by need and non-need variables

Variables	beta (SE)	Variables	beta (SE)
Age: 60–64	-1.44 (1.27)	Geographic region: Hokkaido	-0.06 (0.32)
Age: 65–69	-0.35 (0.39)	Geographic region: Tohoku	0.13 (0.25)
Age: 70–74	-0.41 (0.26)	Geographic region: Kanto	Ref.
Age: 75–79	-0.30 (0.19)	Geographic region: Hokuriku	-0.67* (0.30)
Age: 80–84	-0.13 (0.14)	Geographic region: Tozan	0.00 (0.30)
Age: 85+	Ref.	Geographic region: Tokai	0.04 (0.27)
Women	-0.05 (0.16)	Geographic region: Kinki	0.20 (0.23)
SRH: Very bad	Ref.	Geographic region: Chugoku	-0.09 (0.24)
SRH: Bad	0.20 (0.21)	Geographic region: Shikoku	0.41 (0.35)
SRH: Fair	-0.32 (0.22)	Geographic region: Kyushu	-0.09 (0.22)
SRH: Good	-0.53# (0.27)	Municipal/population category: Government-designated	Ref.
SRH: Very good	-0.56 (0.43)	Municipal/population category: 200K+	-0.20 (0.22)
ln(Chronic conditions)	0.40** (0.14)	Municipal/population category: 100–200K	-0.27 (0.23)
ln(ADL difficulties)	0.44** (0.07)	Municipal/population category: <100K	-0.66** (0.22)
ln(IADL difficulties)	0.49** (0.09)	Municipal/population category: Towns and villages	-0.34 (0.24)
ln(Cognitive functioning)	-0.10 (0.08)	Wave 6 (2002)	Ref.
Marital status: Single	0.27# (0.16)	Wave 7 (2006)	0.46* (0.17)
ln(N of co-resident members)	-0.32** (0.09)	Wave 8 (2012)	0.46** (0.17)
Variance(_cons[id])	0.00 (0.00)	Wave 9 (2016)	0.64** (0.20)
Constant	-1.95** (0.46)	Wave 10 (2021)	0.67** (0.23)
Observations		1,855	
Number of unique individuals		1,431	

Note: Robust standard errors (SE) are in parentheses; estimates are weighted by both cross-sectional and longitudinal weights; ADL stands for activities of daily living; SRH stands for self-rated health; ** p<0.01, * p<0.05, # p<0.1

Table A-3. Gender-specific non-standardised and standardised long-term care use by education and income: Mean probabilities, Pooled between Wave 6 (2002) and Wave 10 (2021)

Gender			Actual	Need-predicted	Difference	Standardised
Men	Education	Lowest: <6	0.38	0.42	-0.04	0.33
		Lower middle: 6–9	0.34	0.40	-0.06	0.32
		Middle: 9–12	0.42	0.41	0.01	0.39
		Higher middle: 12–15	0.38	0.39	-0.02	0.36
		Higher: 16+	0.70	0.45	0.25	0.62
	Income (Quintile)	Poorest 20%	0.36	0.39	-0.02	0.35
		2nd poorest 20%	0.30	0.39	-0.10	0.28
		Middle	0.43	0.39	0.04	0.41
		2nd richest 20%	0.41	0.41	-0.01	0.37
		Richest 20%	0.45	0.40	0.05	0.42
Women	Education	Lowest: <6	0.29	0.36	-0.07	0.30
		Lower middle: 6–9	0.38	0.39	-0.01	0.36
		Middle: 9–12	0.47	0.37	0.10	0.47
		Higher middle: 12–15	0.45	0.38	0.07	0.45
		Higher: 16+	0.69	0.38	0.32	0.69
	Income (Quintile)	Poorest 20%	0.33	0.37	-0.04	0.34
		2nd poorest 20%	0.40	0.36	0.04	0.41
		Middle	0.42	0.37	0.05	0.42
		2nd richest 20%	0.53	0.37	0.15	0.53
		Richest 20%	0.38	0.33	0.05	0.43

Table A-4. Concentration indices for long-term care use by gender

Gender			2002	2006	2012	2017	2021
Men	Education	Concentration index	0.02	0.10	0.06	-0.02	0.15
		SE	0.10	0.06	0.06	0.07	0.08
		P-value	0.85	0.10	0.36	0.79	0.05
		N	130	103	135	114	57
	Income	Concentration index	-0.11	0.12	0.10	0.00	-0.06
		SE	0.09	0.07	0.06	0.07	0.08
		P-value	0.21	0.09	0.12	0.94	0.49
		N	120	79	119	99	46
Women	Education	Concentration index	0.10	0.14	0.11	0.02	0.02
		SE	0.04	0.04	0.04	0.05	0.07
		P-value	<0.01	<0.01	<0.01	0.64	0.76
		N	312	282	336	206	100
	Income	Concentration index	0.11	0.03	0.08	0.06	0.11
		SE	0.04	0.04	0.05	0.05	0.08
		P-value	<0.01	0.42	0.07	0.31	0.18
		N	256	221	262	157	72

Note: Standard errors (SE) are adjusted for clusters in each respondent; Wagstaff indices are presented; and estimates are weighted by both cross-sectional and longitudinal weights.

Table A-5. Age-specific non-standardised and standardised long-term care use by education and income: Mean probabilities, Pooled between Wave 6 (2002) and Wave 10 (2021)

Age			Actual	Need-predicted	Difference	Standardised
60–79	Education	Lowest: <6	0.27	0.31	-0.04	0.33
		Lower middle: 6–9	0.32	0.37	-0.05	0.32
		Middle: 9–12	0.36	0.34	0.02	0.40
		Higher middle: 12–15	0.32	0.40	-0.07	0.30
		Higher: 16+	0.59	0.39	0.20	0.57
	Income (Quintile)	Poorest 20%	0.33	0.35	-0.02	0.36
		2nd poorest 20%	0.37	0.38	-0.01	0.36
		Middle	0.46	0.33	0.12	0.50
		2nd richest 20%	0.30	0.34	-0.04	0.33
		Richest 20%	0.32	0.32	0.00	0.37
80+	Education	Lowest: <6	0.31	0.39	-0.07	0.30
		Lower middle: 6–9	0.39	0.40	-0.02	0.36
		Middle: 9–12	0.50	0.40	0.10	0.47
		Higher middle: 12–15	0.47	0.38	0.09	0.46
		Higher: 16+	0.78	0.48	0.30	0.67
	Income (Quintile)	Poorest 20%	0.34	0.38	-0.04	0.33
		2nd poorest 20%	0.13	0.29	-0.16	0.21
		Middle	0.41	0.39	0.02	0.40
		2nd richest 20%	0.53	0.40	0.13	0.51
		Richest 20%	0.47	0.39	0.08	0.45

Appendix Table A-6. Concentration indices for long-term care use by age

Age			2002	2006	2012	2017	2021
60–79	Education	Concentration index	0.01	0.15	0.02	-0.13	0.08
		SE	0.05	0.06	0.07	0.18	0.07
		P-value	0.76	<0.05	0.75	0.48	0.29
		N	207	118	102	33	58
	Income	Concentration index	0.02	0.00	0.02	0.02	0.04
		SE	0.05	0.06	0.08	0.16	0.09
		P-value	0.65	0.98	0.78	0.89	0.65
		N	184	98	80	28	43
80+	Education	Concentration index	0.13	0.12	0.13	0.03	0.06
		SE	0.05	0.04	0.04	0.04	0.08
		P-value	<0.05	<0.01	<0.01	0.37	0.43
		N	235	267	369	287	99
	Income	Concentration index	0.07	0.09	0.11	0.04	0.09
		SE	0.07	0.05	0.04	0.04	0.07
		P-value	0.35	0.07	<0.01	0.37	0.19
		N	192	202	301	228	75

Note: Standard errors (SE) are adjusted for clusters in each respondent; Wagstaff indices are presented; and estimates are weighted by both cross-sectional and longitudinal weights.

Figure A-1. Concentration curves on standardised long-term care use by non-residualised income

Note: The 95% confidence interval (CI) is calculated based on standard errors (SE) adjusted for clusters for each respondent; estimates are weighted by both cross-sectional and longitudinal weights. The sample sizes in each wave were 376, 300, 382, 256, and 118, respectively, and L(p) is the concentration curve for long-term care (LTC).

Appendix B: Income Residualisation

Employed and unemployed individuals are non-comparable because of large differences: theoretically, people decumulate their savings to smooth their consumption levels, particularly after retirement (i.e. life-cycle hypothesis) (1); Thus, a low income does not always suggest being deprived among older people. Instead of using limited information about savings and consumption from the National Survey of the Japanese Elderly, we residualised the income to partially adjust for the differences between workers and non-workers by fixed-effects ordinary least squares as follows:

$$e_{i,t} = Y_{i,t} - \tilde{Y}_{i,t}$$
$$\ln(Y_{i,t}) = c + \delta_i Work_{i,t} + \theta_i \pi_{i,t} + \mu_i + \epsilon_{it}$$

where $e_{i,t}$ is the residualised income of individual i in year t, defined as the predicted income ($\tilde{Y}_{i,t}$) subtracted from the actual income ($Y_{i,t}$). The income is predicted by the employment status ($Work_{i,t}$; =1 if in paid work; =0 if otherwise) and other independent variables ($\pi_{i,t}$), including the age, sex, marital status, residential area, municipal or population category of residential area, year-fixed effects, and individual fixed effects (μ_i). c is a constant, ϵ is the stochastic disturbance, and δ and θ are the estimated coefficients of the independent variables. This formalisation enabled controlling for potential imbalances in the income arising from the employment status, demographic factors, price differences across areas and years, and time-invariant individual heterogeneity (Appendix Table B-1).

Appendix Table B-1. Regression result: Prediction of couple's income

Variables	beta (SE)
Age: 60–64	Ref.
Age: 65–69	-0.02 (0.02)
Age: 70–74	-0.01 (0.03)
Age: 75–79	-0.04 (0.05)
Age: 80–84	-0.04 (0.05)
Age: 85+	-0.05 (0.08)
Employment status: Currently working	0.20** (0.02)
Marital status: Single	-0.06* (0.02)
Individual FE	Yes
Residential area FE	Yes
City-scale FE	Yes
Year FE	Yes
Constant	5.13** (0.03)
Observations	17,031
Number of unique individuals	4,586

Note: Robust standard errors (SE) are in parentheses; estimates are weighted by both cross-sectional and longitudinal weights; FE denotes fixed effects; ** $p<0.01$, * $p<0.05$, # $p<0.1$

References for Appendices

1.   Ando A, Modigliani F. The "Life Cycle" Hypothesis of Saving: Aggregate Implications and Tests. The American Economic Review. 1963;53(1):55-84.