3   **An ecological examination of early adolescent e-cigarette use: A machine learning**

4   **approach to understanding a health epidemic**

Alejandro L. Vázquez [1*], Cynthia M. Navarro Flores [1], Byron H. Garcia [2], Tyson S. Barrett [3],

Melanie M. Domenech Rodríguez [4]

[1] Department of Psychology, University of Tennessee, Knoxville, Knoxville, Tennessee, United

States

[2] Department of Psychology, Arizona State University, Tempe, Arizona, United States

[3] Highmark Health, Pittsburg, Pennsylvania, United States

[4] Department of Psychology, Utah State University, Logan, Utah, United States

* Correspondence author

Email: avazquez4004@gmail.com

Abstract

E-cigarette use among adolescents is a national health epidemic spreading faster than researchers can amass evidence for risk and protective factors and long-term consequences associated with use. New technologies, such as machine learning, may asset prevention programs in identifying at-risk youth and potential targets for intervention before adolescents enter developmental periods were e-cigarette use escalates. The current study utilized machine learning algorithms to explore a wide array of individual and socioecological variables in relation to patterns of lifetime e-cigarette use during early adolescence (i.e., exclusive, or with tobacco). Extant data was utilized from 14,346 students middle school students (M_{age} = 12.5, SD = 1.1; 6th and 8th grades) who participated in the Utah Prevention Needs Assessment survey. Students self-reported their substance use behaviors and related risk and protective factors. Machine learning algorithms examined 112 individual and socioecological as potential classifiers of lifetime e-cigarette use outcomes. Algorithms achieved outstanding classification for lifetime exclusive (AUC = .926) and dual use (AUC = .944) on a validation test set. Six high value classifiers were identified that varied in importance by outcome: Lifetime alcohol or marijuana use, perception of e-cigarette availability and risk, school suspension(s), and perceived risk of smoking marijuana regularly. Specific classifiers were important for lifetime exclusive (parent's attitudes regarding student vaping, best friend[s] tried alcohol or marijuana) and dual use (best friend[s] smoked cigarettes, lifetime inhalant use). Our findings provide specific targets for the adaptation of existing substance use prevention programs to address early adolescent e-cigarette use.

Keywords: e-cigarette use, vaping, machine learning, early adolescence, ecological

Introduction

E-cigarette use among adolescents is a national epidemic [1]. The popularity of these

devices has spread faster than health researchers could amass evidence for E-cigarette deleterious

effects. As the prevalence of traditional cigarette smoking among U.S. adolescents has declined,

e-cigarette use, or vaping, has become the most commonly used form of nicotine uptake among

youth in the U.S. [2]. Adolescents' decisions to engage in e-cigarette use may be understood

through ecological framework that accounts for complex interactions between spheres of

influence [3]. Research is underway to identify individual and socioecological risk-factors

associated with e-cigarette use [4–11]. However, this literature has prominently focused on high

school samples resulting in a dearth of knowledge regarding e-cigarette risk-factors during early

adolescence. Identifying factors associated with the emergence of e-cigarette use during early

adolescence may facilitate intervention prior to developmental periods where use escalates (i.e.,

middle to late adolescence [12]). These efforts may be bolstered by new methodologies that

allow researchers to efficiently explore the importance of a wide range of variables in relation to

e-cigarette use [13]. In this study we use machine learning algorithms to simultaneously consider

a large number of individual and socioecological factors in relation to patterns of e-cigarette

usage among middle school students [7].

Researchers have documented complex relationships between individual (e.g., academic

performance, substance use, perceptions of use) and socioecological (e.g., access,

advertisements, peer and parental factors) influences implicated in e-cigarette use during middle

to late adolescence [7–9]. Less is known about e-cigarette use risk-factors during early

adolescence. The early adolescent e-cigarette literature has predominately focused on the

prevalence and reasons for use, or factors associated with susceptibility rather than initiation

[12,14–17]. Studies examining adolescent e-cigarette use have also had a narrow focus when

considering potential individual and socioecological influences on adolescent e-cigarette use. For

example, studies commonly focus on specific adolescent attitudes (e.g., perceived danger of e-

cigarettes and tobacco), substance use behaviors (e.g., alcohol, tobacco, marijuana), aspects of

the environment (e.g., access, advertisements), and social influences (e.g., peer and parental e-

cigarette or cigarette use [7–9,12,16]). Research is needed to examine these risk-factors in

conjunction with a broader array of influences traditionally associated with early substance use

(e.g., anti-social behavior; parenting practices; school involvement, performance, environment;

community attachment, norms, drug use, delinquency; [18–22]).

77   Studies examining adolescent e-cigarette use have had a narrow focus when considering

potential individual and socioecological influences on adolescent e-cigarette use. For example,

studies commonly focus on specific adolescent attitudes (e.g., perceived danger of e-cigarettes

and tobacco), substance use behaviors (e.g., alcohol, tobacco, marijuana), aspects of the

environment (e.g., access, advertisements), and social influences (e.g., peer and parental e-

cigarette or cigarette use; [7–9,12,16]). Research is needed to examine these risk-factors in

conjunction with a broader array of influences traditionally associated with adolescent substance

use (e.g., anti-social behavior; parenting practices; school involvement, performance,

environment; community attachment, norms, drug use, delinquency; [18–22]).

86   It is also important to consider patterns of use when identifying correlates of early

adolescent vaping. Research suggests risk-factors vary between youth who have utilized e-

cigarettes exclusively and those who have used them in combination with tobacco, with dual use

being associated with greater behavioral problems (i.e., lifetime use; $M = 14.6$ years, $SD = 0.7$)

and substance use (i.e., lifetime alcohol, marijuana, drug use prescription drug misuse; 9th and

12th graders; [7,23]). Exclusive use may represent adolescent using e-cigarettes as a "safer"

alternative to traditional tobacco cigarettes [16]. Dual use may be associated with tobacco

cessation or recreational use in conjunction with other substances [16,23]. Research has yet to

determine whether differences in risk-factors for exclusive or dual e-cigarette use exist during

early adolescence.

Methodological challenges may explain the limited number of studies examining a broad

array of correlates of e-cigarette use during early adolescence. For example, lifetime e-cigarette

is a low base rate behavior during early adolescence relative to later developmental periods [23].

Prevalence rates are even lower when researchers examine exclusive and dual e-cigarette use

relative to general lifetime use [7] Furthermore, limitations associated with traditional statistical

methodologies may pose a barrier to examining the broad array of potential factors implicated in

early adolescent e-cigarette use (e.g., statistical power issues, multicellularity; familywise error

rate[13]). These limitations can be addressed with large datasets, however meeting statistical

assumptions for multicollinearity and reducing family-wise error may limit the number of

potential risk-factors that can be simultaneously considered in relation to early adolescent e-

cigarette use.

Machine learning may facilitate the examination of factors associated with early

adolescent e-cigarette use. Machine learning provides an efficient method of simultaneously

examining large numbers of variables representing youth individual and socioecological factors

to determine their *importance* in classifying substance use [13]. Within the context of machine

learning, variable importance refers to the relative ability for variables to reduce the error in

models' predictions of group membership (e.g., exclusive e-cigarette user or non-user) compared

to other covariates in the model [24]. Elastic net, random forest, k-nearest neighbors, and neural

networks are examples of common algorithms capable of producing highly accurate

classification models for lifetime substance use [13,25]. Each of these algorithms approach

classification with contrasting linear (i.e., elastic net) and nonlinear (i.e., random forest, k-nearest

neighbors, neural networks) methods, providing the opportunity to identify the algorithm that

best performs the classification task for each outcome [24].

Identifying high value correlates of e-cigarette initiation during early adolescence could

improving our ability to identify at-risk youth prior to developmental period were the prevalence

and frequency of vaping escalates (i.e., middle to late adolescence; [12]). While machine

learning may provide an additional tool for informing substance use prevention effort [26], few

studies have utilized machine learning to identify factors associated with patterns of early e-

cigarette use (i.e., lifetime exclusive use, dual use with tobacco). Prior applications of machine

learning have predominately focused on unstructured data (i.e., pictures, text) to classify e-

cigarette use [27,28]. While a recent study trained machine learning algorithms on survey data

collected on older teens (M_{age} =15.36 years old; SD = 1.85), this research had a narrow focus on

tobacco related substance use predictors that precludes a broader understanding of factors

associated with early vaping initiation (i.e., LASSO and Random Forest; [29]). Thus, our aim

was to explore a wide array of factors using machine learning to identify important classifiers of

lifetime exclusive e-cigarette and dual use within a sample of middle school students. These

analyses can help quantify the relative importance of predictors and established the extent to

which e-cigarette use can be classified by individual and socioecological factors.

Materials and method

The current study utilized data from the Utah Student Health and Risk Prevention

(SHARP) survey project, which has been collecting and disseminating information on substance

use prevalence and related behaviors since 2007 (Utah Department of Human Services [UDHS];

[30]). SHARP was developed as a collaboration between multiple state agencies with the

purpose of assessing risk and protective factors for problem behaviors among Utah middle and

high school students. Students complete the Utah Prevention Needs Assessment (PNA) survey

biannually, during the spring of odd numbered years, as a part of the SHARP survey project. The

PNA survey gathers statewide data on substance use and individual/socioecological factors that

influence the use of alcohol, tobacco, and other drugs. PNA surveys are implemented and used to

inform statewide prevention policy and programming across the United States. Surveys are

completed in schools and are self-administered using paper and pencil. The present study used

data collected during the Spring of 2017 (i.e., March-June 2017) as part of a PNA survey in

Utah. Parents provided written consent for their child to participate in the survey. Parents of

youths that did not consent to their child participating in the PNA were not administered the

survey. Student also provided verbal assent prior to participating in the PNA survey.

Participation in the survey was voluntary and students could opt to participate in an alternative

activity or discontinue at any time. The Utah State University Institutional Review Board

approved secondary analyses of the 2017 Utah PNA survey data as non-human subjects research

as participants could not be re-identified (protocol #10108). Previous research has utilized

similar statewide school-based samples to identify factors associated with e-cigarette use among

adolescents across the United States (e.g., Hawaii, Texas, Connecticut, New Jersey; [6–9,17,31]).

The current study focused on 14,346 middle school students (i.e., 6th and 8th grade) that

participated in the 2017 Utah PNA survey. Participants were approximately 12 years old on

average ($M = 12.5$; $SD = 1.1$), were relatively balanced on sex (girls; n = 7,532, 52.5%) and 6th

grade (n = 7,473, 52.1%), and were predominantly White (9,491, 71%). Nearly a third of

students attended school within Salt Lake County (n = 4,173, 29.1%) in Utah. Youths in this

sample reported a 9.4% (n = 1,343) prevalence of lifetime e-cigarette use and 5.4% (n = 784)

tobacco use. Students largely reported abstaining from both tobacco and e-cigarette use (91%; n

= 13,003) and reported greater lifetime exclusive e-cigarette use (5.5%; n = 791) relative to

exclusive tobacco use (1.6%; n = 232). Within the sample, 3.8% (n = 552) of students reported

dual lifetime use of tobacco and e-cigarettes. See Table 1 for sample demographic information

by outcomes.

Measures

Individual and socioecological variables

The measures utilized in the current study have been traditionally used and reported by

the SHARP survey project as individual items [30]. Variables examined used in the current study

have been identified as being theoretically and/or empirically important factors in the substance

use literature. We decided to examine individual items to provide a nuanced understanding of e-

cigarette use risk factors [13]. Variables included a wide array of factors including *individual*

(i.e., antisocial behaviors/attitudes, rebelliousness, academic performance, perceived risk of drug

use, intentions for adulthood substance use, lifetime substance use), *community* (i.e., attachment,

prosocial involvement and reward, drug use consequences and antisocial behavior, perceived

availability of substances), *school* (i.e., learning environment perceptions, enjoyment,

commitment, benefits of learning, truancy), *home* (i.e., parenting practices, family history of

substance abuse, rewards for prosocial behavior, parental attitudes regarding antisocial behavior

and drug use, relationship quality with parents), and *social* (i.e., best friends engaged in

antisocial behavior, tried alcohol or drugs, exhibited prosocial behavior; social rewards for

antisocial and prosocial behaviors) influences. See supplementary Table S1 for all items

examined in the current study.

Outcome

185   Students reported whether they ever tried electronic cigarettes or e-cigarettes (i.e., *yes* or

186   *no)*. They also reported whether they had ever tried tobacco cigarettes, even just a puff (i.e., *yes*

187   or *no*). Two dichotomous outcome variables were created from these items to represent lifetime

188   exclusive e-cigarette use and dual use (i.e., tobacco and e-cigarette). The comparison group for

189   each outcome were students who did not utilize either substance.

Analytic plan

In our sample, 47% ($n = 6,744$) of participants were missing at least one covariate. Prior

to imputation, data was randomly resampled into training (70%; $n = 9,657$ e-cigarette; $n = 9,490$

dual) and testing sets (30%; $n = 4,237$ e-cigarette; $n = 4,065$ dual). We then used mode

imputation, wherein missing values were replaced with the mode for each variable to address

missingness independently for training and testing sets. Mode imputation is commonly utilized

within the context of machine learning for classification task [25]. As algorithms can struggle to

predict low base rate outcomes, a method known as down sampling was used to randomly

resample and reduce the negative class (i.e., those that did not use e-cigarettes or tobacco) until it

was equal to the positive class within the training set [24]. Thus, rates of lifetime use and non-

use were equal for each outcome within the resampled training sets. The training sets were $n =$

1,108 for exclusive e-cigarette use and $n = 774$ for dual use.

Four dissimilar machine learning algorithms-elastic net, random forest, neural networks,

and k-nearest neighbors–were then fitted to the training set to create classification models for

each outcome [32]. Each classification algorithm drew information from 112 variables

representing student individual and socioecological factors. Five-fold cross-validation was used

to identify variables that improve classification across random subsets of data within the training

set [25]. Model performance was assessed on test sets using the Area Under (AUC) of the

Receiving Operator Characteristic (ROC) curve, which represents the ability of a model to

classify outcomes across all possible cut points [24]. The top performing classification algorithm

on the test set was selected for each outcome (i.e., AUC; sensitivity, specificity; [24]). Variable

importance figures reflect results from the best performing algorithms for each outcome. High

value classifiers were then identified through visual inspection of relative importance figures.

Variables that demonstrate large increase in relative importance over subsequent covariates were

said to be high value classifiers [25]. High value classifiers were examined using a

crosstabulation visualization to determine the nature of the relationship between each variable

and the corresponding outcome [13].

Results

Patterns of lifetime e-cigarette use differed by demographic variables within the current

sample. Chi-square test of independence suggest e-cigarette usage was significantly ($p < .001$)

associated with student gender, grade, and race/ethnicity. Boys, 8th graders, Latinxs, Native

Americans, and mixed-race students reported the greatest proportion of use across outcomes.

Exclusive and dual use were generally associated with a greater proportion of lifetime use across

substances relative to no-users. See Table 1 for demographic variables by outcomes.

Exclusive use

Algorithmic performance on the exclusive e-cigarette use classification task ranged from

excellent to outstanding (AUC = .837 - .926) on the test set. See supplemental Fig S1 for ROCs

for classification algorithms. Elastic net was the best performing algorithm in classifying

exclusive e-cigarette use (AUC = .926, sensitivity = .857, specificity = .848). Elastic net

identified perceived availability of e-cigarettes, lifetime alcohol use, parents' attitudes regarding

their use of vape products, school suspension, perceived risk of e-cigarette use, lifetime

marijuana use, best friend(s) tried alcohol, best friend(s) used marijuana, and perceived risk of

smoking marijuana regularly as the best discriminators between lifetime exclusive e-cigarette

users and non-users. See Fig 1 for variable importance. Visual inspection of cross-tabulation

mosaics suggests that perceived availability of e-cigarettes (i.e., *sort of hard*, v*ery easy, sort of*

easy), lifetime substance use (i.e., alcohol, marijuana), school suspensions (i.e., 1 or more), lower

levels of perceived risk associated with e-cigarette use (i.e., *none* to *moderate*), best friend(s)

tried alcohol or used marijuana (i.e., 1 or more), and less perceived risk associated with smoking

marijuana regularly were all associated with a greater proportion of lifetime e-cigarette use.

Students who reported that their parents would view their use of vape products as "*very wrong*"

had the lowest proportion of use relative to other levels of approval (i.e., *wrong* to *not wrong at*

all). See supplemental Fig S3-11 for cross-tabulation visualizations.

Fig 1. Top 20 Variables with the Highest Relative Importance in Classifying Lifetime E-

Cigarette Use. Results represent validation on a separate test dataset.

Dual use

Algorithmic performance on the dual tobacco and e-cigarette use classification task also

ranged from excellent to outstanding (AUC = .862 - .944) on the test set. See supplemental Fig

S2 for ROCs for classification algorithms. Elastic net and random forest had the same AUC

score (.944). However, elastic net (sensitivity = .824, specificity = .947) outperformed random

forest (sensitivity = .818, specificity = .939) based on sensitivity and specificity. Elastic net

identified lifetime alcohol use, lifetime marijuana use, perceived availability of e-cigarettes, best

friend(s) cigarette use, perceived risk of e-cigarette use, lifetime inhalants use, school

suspension, and perceived risk of smoking marijuana regularly as the best discriminators

between lifetime dual users and non-users. See Fig 2 for variable importance. Visual inspection

of cross-tabulation mosaics suggests that lifetime substance use (i.e., alcohol, marijuana,

inhalants), higher levels of perceived availability of e-cigarettes (i.e., *very easy, sort of easy*),

best friend(s) that have smoked cigarettes (i.e., 1 or more), school suspensions (i.e., 1 or more),

lower levels of perceived risk associated with e-cigarette use and using marijuana regularly (i.e.,

none to *moderate*) were all associated with a greater proportion of lifetime dual use. See

supplemental Fig S12-19 for cross-tabulation visualizations.

Fig 2. Top 20 Variables with the Highest Relative Importance in Classifying Lifetime Dual

Use. Results represent validation on a separate test dataset.

Discussion

The current study expands the literature through the simultaneous exploration of

established correlates of e-cigarette initiation and traditional factors associated with substance

use in relation to early adolescent vaping. Algorithms utilizing information regarding student

individual characteristics and socioecological context demonstrated high levels of classification

accuracy for both lifetime exclusive and dual e-cigarette use. While the order of importance of

classifiers differed by outcome, algorithms consistently identified six high value classifiers

across usage groups: lifetime alcohol or marijuana use, perception of e-cigarette availability and

risk, school suspension(s), and perceived risk of smoking marijuana regularly. Several high value

classifiers differed between youth who reported lifetime exclusive (i.e., parent's attitudes

regarding their use of vaping products, best friend[s] tried alcohol, best friend[s] used marijuana)

and dual e-cigarette use (i.e., best friend[s] smoked cigarettes, lifetime inhalants use). These

findings highlight important commonalities and difference in risk profiles between lifetime

exclusive and dual e-cigarette users.

Research using high school samples have documented higher rates of life substance use

among dual versus exclusive e-cigarette users [7]. Within our sample, rates of substance use

were generally higher among exclusive and dual users relative to those who abstained from both.

Consistent with prior research, the greatest portions of substance use were found among youth

who had reported dual use [23]. However, only lifetime alcohol and tobacco were found to be

important classifiers of both e-cigarette use outcomes among middle school students, which is

consistent with prior findings in high school samples [7]. Our findings also extend prior work

through the identification of inhalant use as a novel risk factor specifically related to lifetime

dual use during early adolescence. It is possible that dual users may access a wide variety of

substances recreationally and may utilize inhalants as they are easy to access within the home

[23,33]. Further research is needed to understand the relationship between lifetime inhalant and

dual use.

Our findings confirm that availability of e-cigarettes is an important influence on early

adolescent vaping [16], despite a Utah state law that restrict the sale of these products to

individuals under the age of 20. Accessibility was especially important to exclusive e-cigarette

use, which is concerning as this may translate to future traditional cigarette use among

adolescents who may have otherwise abstained from tobacco use [8]. Consistent with prior

research, students who reported lower perceived danger of using e-cigarettes reported a greater

proportion of lifetime use [12]. Our findings suggest the need to consider perceptions regarding

the danger of other substances, such as marijuana, when assessing risk for early adolescent e-

cigarette use. While recent findings have highlighted the importance of school-based factors in

assessing risk for e-cigarette using among highs school samples (i.e., truancy and poor academic

performance; [10]), our findings suggest that student suspensions were the most relevant aspect

of school in relation to early adolescent substance use. It is possible school suspensions may be

associated with an increased risk for e-cigarette use as a potential proxy for rule breaking

behaviors or through greater unsupervised time outside of school [7,34]. Further research is

needed to elucidate the relationship between school suspensions and early adolescent e-cigarettes

use.

It is important to mention that factors traditionally associated with substance use such as

adolescent and peer delinquency, community substance use norms, and school involvement were

not important predictors of lifetime exclusive and dual e-cigarette use within the current sample.

These findings may signal potential differences between factors underlying e-cigarette and other

forms of substance use. Additionally, factors identified by prior research as relevant predictors of

vaping (e.g., parenting practices, perceived risk of smoking tobacco) were not relevant correlates

of patterns of e-cigarette use within the current sample [7,12]. It is possible that when competing

against other variables within a machine learning approach, these important predictors are truly

of lesser importance relative to high value classifiers identified in the current study.

Implications

Our findings support addressing early adolescent vaping through prevention programs

aiming to address substance use prevention broadly (i.e., alcohol, tobacco, marijuana, inhalants,

e-cigarette). There is ample evidence for the efficacy of prevention programs whether focused on

a single substance or multiple ones [35]. Our findings provide specific structural and behavioral

targets that may inform the adaptation of programs seeking to prevent different patterns of early

adolescent e-cigarette use. Substance use prevention programs may benefit from adding

components that equip parents to effectively communicate disapproval regarding their child's

use of e-cigarette and teach youth skills to resist peer substance use influences [36,37]. Targeted

prevention programs are already supported by research documenting perception of risk

associated with vaping as a consistent predictor of e-cigarette use [7,8] and our findings highlight

the importance of also considering perceptions of danger regarding marijuana use during early

adolescence. Specific to programming, youth's perceptions of risk do not align with research

evidence providing an important point of content for preventive interventions [38]. Altering

youth's perception of e-cigarette accessibility, however, may require intervention at a broader

social level (e.g., public media campaigns). Alternatively, decreasing accessibility to e-cigarettes

may be achieved by actions external to youths such as strong enforcement of laws regarding

possession and/or consumption for underage users and/or those selling e-cigarette products to

them, or by way of increasing prices for goods associated with e-cigarette use.

Machine learning appears to be a promising screening tool for the identification of risk

factors that can accelerate the development of the e-cigarette knowledge base needed to curb the

rapid spread of vaping among adolescents nationally. Algorithms were able to efficiently explore

a wide range of factors in association with early adolescent e-cigarette use, which confirmed

findings from later developmental stages and identified several novel risk factors (i.e., inhalant

use, perceived risk of marijuana, school suspensions). An important consideration in this

research is that the tools utilized to identify e-cigarette use classifiers are publicly available. R

offers open access statistical packages for machine learning. Additionally, there are substantial

training materials available for free online. These tools can provide an accessible and replicable

method of generating and disseminating scientific knowledge regarding e-cigarette use classifiers

nationally. We encourage researchers to apply machine learning algorithms to their data to draw

new insight regarding factors contributing to a variety of e-cigarette use outcomes among

adolescents. Examining cross-sectional markers of e-cigarette use could also identify important

variables that can be examined longitudinally in prospective research. Machine learning

algorithms have many exciting applications when applied to longitudinal data, including

identifying context specific predictors of service use, specific targets for substance use

prevention programs, and ensure that important factors are not excluded from causal models

examining mechanism underpinning early vaping initiation.

Limitations

Results from algorithms used in the current study do not necessarily imply causal

mechanisms explaining patterns of lifetime e-cigarette use but rather identify factors that are

strong correlates of group membership (i.e., use or non-use). Longitudinal research is needed to

establish causal links. The current study examined lifetime substance use that may range from

experimentation to habitual use. Future research may consider using machine learning as a

method of identifying youth at-risk for habitual e-cigarette use. Although a large number of the

Utah adolescent population was captured, the PNA survey does not include students in private

schools, correctional facilities, or treatment centers. Additionally, students who were not in

attendance, declined participation, or did not return parental consent forms are not represented.

Furthermore, findings may not generalize to students in other states. Further research is needed

to replicate our findings in different contexts and developmental periods.

Conclusions

The current study utilized a machine learning approach to efficiently explore and identify

high value correlates of early adolescent lifetime e-cigarette use. This approach identified several

shared risk factors for exclusive and dual e-cigarette use such as lifetime use of specific

substances (i.e., alcohol, marijuana), perception of e-cigarette availability and risk, school

suspension(s), and perceived risk of smoking marijuana regularly. Several differences were also

identified between youth who reported lifetime exclusive (i.e., parent's attitudes regarding their

use of vaping products, best friend[s] tried alcohol or used marijuana) and dual use (i.e., best

friend[s] smoked cigarettes, lifetime inhalants use) relative to non-users. This information

provides a first step towards identifying youth at-risk for e-cigarette use during early

adolescence. Further research is needed to examine high value classifiers identified by the

current study using explanatory models and longitudinal data to understand mechanism

underlying their importance in accounting for differences in risk profiles between e-cigarette

usage groups during early adolescence.

Reporting

Funding

378   This research was funded by a collaboration between multiple state agencies in Utah (i.e.,

379   Department of Health, Department of Human Services, and the State Board of Education).

Disclosure Statement

381   The authors have no conflicts of interest associated with the publication of this

382   manuscript.

Acknowledgements

384   This research was funded by a collaboration between multiple state agencies in Utah (i.e.,

385   Department of Health, Department of Human Services, and the State Board of Education).

Data Availability

387   Data used in the current study can be requested from the Utah Department of Human and

388   Health Services.

Data Deposition

390   We are not authorized to share the Utah Prevention Needs Assessment data used in the

391   current study as it is owed and managed by the Utah Department of Human and Health Services.

References

1.   Office of the Surgeon General. Surgeon General's Advisory on E-cigaretteUse Among

Youth. 2019. Available: https://www.cdc.gov/tobacco/basic_information/e-

cigarettes/surgeon-general-advisory/index.html

2.   Centers for Disease Control and Prevention. Tobacco product use among middle and high

school students—United States, 2011–2017. Morb Mortal Wkly Rep. 2018;63: 629–633.

doi:http://dx.doi.org/ 10.15585/mmwr.mm6722a3

3.   Bronfenbrenner U. Toward an experimental ecology of human development. Am Psychol.

1977;32: 513–531. doi:http://doi.org/10.1037/0003-066X.32.7.513

4.   Barrington-Trimis JL, Berhane K, Unger JB, Cruz TB, Urman R, Chou P, et al. The E-

cigarette Social Environment, E-cigarette Use, and Susceptibility to Cigarette Smoking.

2016 [cited 26 Jun 2019]. doi:10.1016/j.jadohealth.2016.03.019

5.   Bold KW, Morean ME, Kong G, Simon P, Camenga DR, Cavallo DA, et al. Early age of

e-cigarette use onset mediates the association between impulsivity and e-cigarette use

frequency in youth. Drug Alcohol Depend. 2017;181: 146–151.

doi:10.1016/j.drugalcdep.2017.09.025

6.   Giovenco DP, Casseus M, Duncan DT, Coups EJ, Lewis MJ, Delnevo CD. Association

Between Electronic Cigarette Marketing Near Schools and E-cigarette Use Among Youth.

J Adolesc Heal. 2016;59: 627–634. doi:10.1016/j.jadohealth.2016.08.007

7.   Wills TA, Knight R, Williams RJ, Pagano I, Sargent JD. Risk factors for exclusive e-

cigarette use and dual e-cigarette use and tobacco use in adolescents. Pediatrics. 2015;135:

e43–e51. doi:10.1542/peds.2014-0760

8.   Wills TA, Sargent JD, Knight R, Pagano I, Gibbons FX. E-cigarette Use and Willingness

to Smoke in a Sample of Adolescent Nonsmokers. 2016;2: 1–16.

doi:10.14440/jbm.2015.54.A

9.   Simon P, Camenga DR, Morean ME, Kong G, Bold KW, Cavallo DA, et al.

Socioeconomic status and adolescent e-cigarette use: The mediating role of e-cigarette

advertisement exposure. Prev Med (Baltim). 2018;112: 193–198.

doi:10.1016/j.ypmed.2018.04.019

10.   McCabe SE, West BT, Veliz P, Boyd CJ. E-cigarette Use, Cigarette Smoking, Dual Use,

and Problem Behaviors Among U.S. Adolescents: Results From a National Survey. J

Adolesc Heal. 2017;61: 155–162. doi:10.1016/j.jadohealth.2017.02.004

11.   Rocheleau GC, Vito AG, Intravia J. Peers, Perceptions, and E-Cigarettes: A Social

Learning Approach to Explaining E-Cigarette Use Among Youth. J Drug Issues. 2020;50:

472–489. doi:10.1177/0022042620921351

12.   Kwon E, Seo DC, Lin HC, Chen Z. Predictors of youth e-cigarette use susceptibility in a

U.S. nationally representative sample. Addict Behav. 2018;82: 79–85.

doi:10.1016/j.addbeh.2018.02.026

13.   Vázquez AL, Domenech Rodríguez MM, Barrett TS, Schwartz S, Amador Buenabad NG,

Bustos Gamiño MN, et al. Innovative Identification of Substance Use Predictors: Machine

Learning in a National Sample of Mexican Children. Prev Sci. 2020;21: 171–181.

doi:10.1007/s11121-020-01089-4

14.   Carey FR, Rogers SM, Cohn EA, Harrell ☐ MB, Wilkinson A V, Perry CL.

Understanding susceptibility to e-cigarettes: A comprehensive model of risk factors that

influence the transition from non-susceptible to susceptible among e-cigarette naïve

adolescents. Addict Behav. 2019;91: 68–74. doi:10.1016/j.addbeh.2018.09.002

15.   Cullen KA, Gentzke AS, Sawdey MD, Chang JT, Anic GM, Wang TW, et al. e-Cigarette

457   Use among Youth in the United States, 2019. JAMA - J Am Med Assoc. 2019;322: 2095–

458   2103. doi:10.1001/jama.2019.18387

459   16.   Kong G, Morean ME, Cavallo DA, Camenga DR, Krishnan-Sarin S. Reasons for

460   electronic cigarette experimentation and discontinuation among adolescents and young

461   adults. Nicotine Tob Res. 2015;17: 847–854. doi:10.1093/ntr/ntu257

462   17.   Krishnan-Sarin S, Morean ME, Camenga DR, Cavallo DA, Kong G. E-cigarette use

463   among high school and middle school adolescents in Connecticut. Nicotine Tob Res.

464   2015;17: 810–818. doi:10.1093/ntr/ntu243

465   18.   Eitle DJ, Eitle TM. School and county characteristics as predictors of school rates of drug,

466   alcohol, and tobacco offenses. J Health Soc Behav. 2004;45: 408–421.

467   doi:10.1177/002214650404500404

468   19.   Evans WD, Powers A, Hersey J, Renaud J. The influence of social environment and social

469   image on adolescent smoking. Heal Psychol. 2006;25: 26–33.

470   doi:https://doi.org/10.1037/0278-6133.25.1.26

471   20.   Mason MJ, Zaharakis NM, Rusby JC, Westling E, Light JM, Mennis J, et al. A

472   longitudinal study predicting adolescent tobacco, alcohol, and cannabis use by behavioral

473   characteristics of close friends. Psychol Addict Behav. 2017;31: 712–720.

474   doi:https://doi.org/10.1037/adb0000299

475   21.   Novak SP, Clayton RR. The influence of school environment and self-regulation on

476   transitions between stages of cigarette smoking: A multilevel analysis. Heal Psychol.

477   2001;20: 196–207. doi:10.1037/0278-6133.20.3.196

22.   Voelkl KE, Frone MR. Predictors of substance use at school among high school students.

J Educ Psychol. 2000;92: 583–592. doi:10.1037/0022-0663.92.3.583

23.   Curran KA, Burk T, Pitt PD, Middleman AB. Trends and Substance Use Associations

With E-Cigarette Use in US Adolescents. Clin Pediatr (Phila). 2018;57: 1191–1198.

doi:10.1177/0009922818769405

24.   Kuhn M, Johnson K. Applied Predictive Modeling with Applications in R. 2013.

Available:

http://appliedpredictivemodeling.com/s/Applied_Predictive_Modeling_in_R.pdf

25.   Hastie T, Tibshirani R, Friedman J. The Elements of Statistical Learning. Second. New

York: Springer; 2013. doi:10.1007/978-1-4419-9863-7_941

26.   Barenholtz E, Fitzgerald ND, Hahn WE. Machine-learning approaches to substance-abuse

research: emerging trends and their implications. Curr Opin Psychiatry. 2020;33: 334–

342. doi:10.1097/YCO.0000000000000611

27.   Ketonen V, Malik A. Characterizing vaping posts on instagram by using unsupervised

machine learning. Int J Med Inform. 2020;141: 104223.

doi:10.1016/j.ijmedinf.2020.104223

28.   Malik A, Khan MI, Karbasian H, Nieminen M, Ammad-Ud-Din M, Khan SA. Modeling

Public Sentiments About JUUL Flavors on Twitter Through Machine Learning. Nicotine

Tob Res. 2021; 1–11. doi:10.1093/ntr/ntab098

29.   Choi J, Jung H-T, Ferrell A, Woo S, Haddad L. Machine Learning-Based Nicotine

Addiction Prediction Models for Youth E-Cigarette and Waterpipe (Hookah) Users. J Clin

Med. 2021;10: 972. doi:10.3390/jcm10050972

30.   Utah Department of Human Services. Student Health and Risk Prevention (SHARP)

survey reports. 2019.

31.   Temple JR, Shorey RC, Lu Y, Torres E, Stuart GL, Le VD. E-cigarette use of young

adults motivations and associations with combustible cigarette alcohol, marijuana, and

other illicit drugs. Am J Addict. 2017;26: 343–348. doi:10.1111/ajad.12530

32.   James G, Witten D, Hastie T, Tibshirani R. An Introduction to Statistical Learning.

Current Medicinal Chemistry. Springer; 2013.

33.   Medina-Mora M´a E, Real T. Epidemiology of inhalant use. Curr Opin Psychiatry.

2008;21: 247–251. doi:https://doi.org/10.1097/yco.0b013e3282fc9875

34.   Lee KTH, Vandell DL. Out-of-School Time and Adolescent Substance Use. J Adolesc

Heal. 2015;57: 523–529. doi:10.1016/j.jadohealth.2015.07.003

35.   Das JK, Salam RA, Arshad A, Finkelstein Y, Bhutta ZA. Interventions for Adolescent

Substance Abuse: An Overview of Systematic Reviews. J Adolesc Heal. 2016;59: S61–

S75. doi:10.1016/j.jadohealth.2016.06.021

36.   Marsiglia FF, Kulis S, Yabiku ST, Nieri TA, Coleman E. When to Intervene: Elementary

School, Middle School or Both? Effects of keepin' It REAL on Substance Use

Trajectories of Mexican Heritage Youth. Prev Sci. 2011;12: 48–62. doi:10.1007/s11121-

010-0189-y

37.   Patterson G. The next generation of PMTO models. Behav Ther. 2005;28: 25–32.

38.   Cheeta S, Halil A, Kenny M, Sheehan E, Zamyadi R, Williams AL, et al. Does perception

of drug-related harm change with age? A cross-sectional online survey of young and older

people. BMJ Open. 2018;8. doi:10.1136/bmjopen-2017-021109

Table 1. Student Demographics by Lifetime E-Cigarette Use Groups.					
	Total	**Neither**	**Exclusive Use**	**Dual use**	*p* value[a]
Sample	*n* = 14346	*n* = 13003	*n* = 791	*n* = 552	
Age *M*(*SD*)	12.5 (1.1)	12.5 (1.1)	13.1 (1.0)	13.1 (1.0)	
Sex					< .001
Boy	6766 (47.2%)	6066 (89.7%)	427 (6.3%)	273 (4%)	
Girl	7532 (52.5%)	6894 (91.5%)	361 (4.8%)	277 (3.7%)	
Grade					< .001
6th	7473 (52.1%)	7110 (95.1%)	217 (2.9%)	146 (2%)	
8th	6873 (47.9%)	5893 (85.7%)	574 (8.4%)	406 (5.9%)	
Race/ethnicity					< .001
White	10191 (71%)	9491 (93.1%)	415 (4.1%)	285 (2.8%)	
Native American	289 (2%)	247 (85.5%)	20 (6.9%)	22 (7.6%)	
Asian	238 (1.7%)	221 (92.9%)	12 (5%)	5 (2.1%)	
Black	210 (1.5%)	186 (88.6%)	11 (5.2%)	13 (6.2%)	
Latinx	1888 (13.2%)	1554 (82.3%)	210 (11.1%)	124 (6.6%)	
Pacific Islander	210 (1.5%)	191 (91%)	13 (6.2%)	6 (2.9%)	
Mixed Race	1320 (9.2%)	1113 (84.3%)	110 (8.3%)	97 (7.3%)	
Substance use[b]					
Alcohol					< .001
Yes	1329 (9.3%)	600 (45.1%)	353 (26.6%)	376 (28.3%)	
No	12982 (90.5%)	12371 (95.3%)	436 (3.4%)	175 (1.3%)	
Marijuana					< .001
Yes	602 (4.2%)	124 (20.6%)	201 (33.4%)	277 (46%)	
No	13671 (95.3%)	12812 (93.7%)	585 (4.3%)	274 (2%)	
Inhalants					< .001
Yes	669 (4.7%)	414 (61.9%)	105 (15.7%)	150 (22.4%)	
No	13579 (94.7%)	12499 (92%)	679 (5%)	401 (3%)	
Prescription drugs					< .001
Yes	546 (3.8%)	332 (60.8%)	81 (14.8%)	133 (24.4%)	
No	13562 (94.5%)	12468 (91.9%)	690 (5.1%)	404 (3%)	
Hallucinogens					< .001
Yes	99 (0.7%)	22 (22.2%)	25 (25.3%)	52 (52.5%)	
No	14128 (98.5%)	12871 (91.1%)	760 (5.4%)	497 (3.5%)	
Synthetic marijuana					< .001
Yes	86 (0.6%)	12 (14%)	21 (24.4%)	53 (61.6%)	
No	14199 (99%)	12940 (91.1%)	766 (5.4%)	493 (3.5%)	

525   Variable frequency is displayed by column for the total and row for usage groups. [a] Chi-square
526   test of independence. [b] Lifetime use.

Figure 1

Figure 2