3   **An ecological examination of early adolescent e-cigarette use: A machine learning**

4   **approach to understanding a health epidemic**

Alejandro L. Vázquez [1*], Cynthia M. Navarro Flores [1], Byron H. Garcia [2], Tyson S. Barrett [3],

Melanie M. Domenech Rodríguez [4]

[1] Department of Psychology, University of Tennessee, Knoxville, Knoxville, Tennessee, United

States

[2] Department of Psychology, Arizona State University, Tempe, Arizona, United States

[3] Highmark Health, Pittsburg, Pennsylvania, United States

[4] Department of Psychology, Utah State University, Logan, Utah, United States

* Correspondence author

Email: avazquez4004@gmail.com

Abstract

E-cigarette use among adolescents is a national health epidemic spreading faster than researchers can amass evidence for risk and protective factors and long-term consequences associated with use. New technologies, such as machine learning, may asset prevention programs in identifying at-risk youth and potential targets for intervention before adolescents enter developmental periods where e-cigarette use escalates. The current study utilized machine learning algorithms to explore a wide array of individual and socioecological variables in relation to patterns of lifetime e-cigarette use during early adolescence (i.e., exclusive, or with tobacco). Extant data was used from 14,346 students middle school students (M_{age} = 12.5, SD = 1.1; 6th and 8th grades) who participated in the Utah Prevention Needs Assessment survey. Students self-reported their substance use behaviors and related risk and protective factors. Machine learning algorithms examined 112 individual and socioecological factors as potential classifiers of lifetime e-cigarette use outcomes. The elastic net algorithm achieved outstanding classification for lifetime exclusive (AUC = .926) and dual use (AUC = .944) on a validation test set. Six high value classifiers were identified that varied in importance by outcome: Lifetime alcohol or marijuana use, perception of e-cigarette availability and risk, school suspension(s), and perceived risk of smoking marijuana regularly. Specific classifiers were important for lifetime exclusive (parent's attitudes regarding student vaping, best friend[s] tried alcohol or marijuana) and dual use (best friend[s] smoked cigarettes, lifetime inhalant use). Our findings provide specific targets for the adaptation of existing substance use prevention programs to address early adolescent e-cigarette use.

Keywords: e-cigarette use, vaping, machine learning, early adolescence, ecological

Introduction

E-cigarette use among adolescents is a national epidemic [1]. The popularity of these devices has spread faster than health researchers could amass evidence for the potential deleterious effects of e-cigarette use. As the prevalence of traditional cigarette smoking among U.S. adolescents has declined, e-cigarette use, or vaping, has become the most commonly used form of nicotine uptake among youth in the U.S. [2]. Adolescents' decisions to engage in e-cigarette use may be understood through an ecological framework that accounts for complex interactions between spheres of influence [3]. Research is underway to identify individual and socioecological risk-factors associated with e-cigarette use [4–11]. However, this literature has prominently focused on high school samples resulting in a dearth of knowledge regarding e-cigarette risk-factors during early adolescence. Identifying factors associated with the emergence of e-cigarette use during early adolescence may facilitate intervention prior to developmental periods where use escalates (i.e., middle to late adolescence [12]). These efforts may be bolstered by new methodologies that allow researchers to efficiently explore the importance of a wide range of variables in relation to e-cigarette use [13]. In this study we use machine learning algorithms to simultaneously consider a large number of individual and socioecological factors in relation to patterns of e-cigarette usage among middle school students [7].

The use of the e-cigarettes has been touted as a healthier alternative to tobacco cigarettes, despite their delivery of nicotine and other potentially harmful chemicals [14]. A major concern of nicotine consumption during early adolescence is the possible alteration of function in the brain's reward systems at a sensitive developmental period, in ways that can increase risk for other substance use, mood disorders, and difficulties with concentration and learning [14]. In addition to nicotine-related risks, other carcinogenic agents found in chemicals in the e-liquid as

well as those produced in the vaporizing product or even associated with the e-cigarette materials

(i.e., nickel, chromium, cadmium; [14]). Chemicals used to flavor e-liquid have also been found

to have sufficiently high toxicity to warrant medical concerns [15] or even cause death [16]. The

possible harms of e-cigarettes go well beyond exposure to nicotine.

Researchers have documented complex relationships between individual (e.g., academic

performance, substance use, perceptions of use) and socioecological (e.g., access,

advertisements, peer and parental factors) influences implicated in e-cigarette use during middle

to late adolescence [7–9]. Less is known about e-cigarette use risk-factors during early

adolescence. The early adolescent e-cigarette literature has predominately focused on the

prevalence and reasons for use, or factors associated with susceptibility rather than initiation

[12,17–20]. Studies examining adolescent e-cigarette use have also had a narrow focus when

considering potential individual and socioecological influences on adolescent e-cigarette use. For

example, studies commonly focus on specific adolescent attitudes (e.g., perceived danger of e-

cigarettes and tobacco), substance use behaviors (e.g., alcohol, tobacco, marijuana), aspects of

the environment (e.g., access, advertisements), and social influences (e.g., peer and parental e-

cigarette or cigarette use [7–9,12,19]). Research is needed to examine these risk-factors in

conjunction with a broader array of influences traditionally associated with early substance use

(e.g., anti-social behavior; parenting practices; school involvement, performance, environment;

community attachment, norms, drug use, delinquency; [21–25]).

It is also important to consider patterns of use when identifying correlates of early

adolescent vaping. Research suggests risk-factors vary between youth who have utilized e-

cigarettes exclusively and those who have used them in combination with tobacco, with dual use

being associated with greater behavioral problems (i.e., lifetime use; $M = 14.6$ years, $SD = 0.7$)

and substance use (i.e., lifetime alcohol, marijuana, drug use prescription drug misuse; 9th and

12th graders; [7,26]). Exclusive use may represent adolescent using e-cigarettes as a "safer"

alternative to traditional tobacco cigarettes [19]. Dual use may be associated with tobacco

cessation or recreational use in conjunction with other substances [19,26]. Research has yet to

determine whether differences in risk-factors for exclusive or dual e-cigarette use exist during

early adolescence.

Methodological challenges may explain the limited number of studies examining a broad

array of correlates of e-cigarette use during early adolescence. For example, lifetime e-cigarette

is a low base rate behavior during early adolescence relative to later developmental periods [26].

Prevalence rates are even lower when researchers examine exclusive and dual e-cigarette use

relative to general lifetime use [7]. Furthermore, limitations associated with traditional statistical

methodologies may pose a barrier to examining the broad array of potential factors implicated in

early adolescent e-cigarette use (e.g., statistical power issues, multicellularity; familywise error

rate[13]). These limitations can be addressed with large datasets, however meeting statistical

assumptions for multicollinearity and reducing family-wise error may limit the number of

potential risk-factors that can be simultaneously considered in relation to early adolescent e-

cigarette use.

Machine learning may facilitate the examination of factors associated with early

adolescent e-cigarette use. Machine learning provides an efficient method of simultaneously

examining large numbers of variables representing youth individual and socioecological factors

to determine their *importance* in classifying substance use [13]. Within the context of machine

learning, variable importance refers to the relative ability for variables to reduce the error in

models' predictions of group membership (e.g., exclusive e-cigarette user or non-user) compared

to other covariates in the model [27]. Elastic net, random forest, k-nearest neighbors, and neural

networks are examples of common algorithms that are capable of provided suprior accuracy in

classifying lifetime substance use relative to traditional logistic regression [13,28]. Each of these

algorithms approach classification with contrasting linear (i.e., elastic net) and nonlinear (i.e.,

random forest, k-nearest neighbors, neural networks) methods, providing the opportunity to

identify the algorithm that best performs the classification task for each outcome [27].

120   Identifying high value correlates of e-cigarette initiation during early adolescence could

improving our ability to identify at-risk youth prior to developmental periods were the

prevalence and frequency of vaping escalates (i.e., middle to late adolescence; [12]). While

machine learning may provide an additional tool for informing substance use prevention effort

[29], few studies have utilized machine learning to identify factors associated with patterns of

early e-cigarette use (i.e., lifetime exclusive use, dual use with tobacco) or determined which

method provides the best classification accuracy. Prior applications of machine learning have

predominately focused on unstructured data (i.e., pictures, text) to classify e-cigarette use

[30,31]. While a recent study trained machine learning algorithms on survey data collected on

older teens (M_{age} =15.36 years old; SD = 1.85), this research had a narrow focus on tobacco

related substance use predictors that precludes a broader understanding of factors associated with

early vaping initiation (i.e., LASSO and Random Forest; [32]). Thus, our aim was to (a) explore

a wide array of factors using machine learning to identify important classifiers of lifetime

exclusive e-cigarette and dual use within a sample of middle school students, (b) and identify the

algorithm that best performs the classification task. These analyses can help quantify the relative

importance of predictors and established the extent to which e-cigarette use can be classified by

individual and socioecological factors.

Materials and method

The current study utilized data from the Utah Student Health and Risk Prevention

(SHARP) survey project, which has been collecting and disseminating information on substance

use prevalence and related behaviors since 2007 (Utah Department of Human Services [UDHS];

[33]). SHARP was developed as a collaboration between multiple state agencies with the

purpose of assessing risk and protective factors for problem behaviors among Utah middle and

high school students. Students complete the Utah Prevention Needs Assessment (PNA) survey

biannually, during the spring of odd numbered years, as a part of the SHARP survey project. The

PNA survey gathers statewide data on substance use and individual/socioecological factors that

influence the use of alcohol, tobacco, and other drugs. PNA surveys are implemented and used to

inform statewide prevention policy and programming across the United States. Surveys are

completed in schools and are self-administered using paper and pencil. The present study used

data collected during the Spring of 2017 (i.e., March-June 2017) as part of a PNA survey in

Utah. Parents provided written consent for their child to participate in the survey. Parents of

youths that did not consent to their child participating in the PNA were not administered the

survey. Student also provided verbal assent prior to participating in the PNA survey.

Participation in the survey was voluntary and students could opt to participate in an alternative

activity or discontinue at any time. The Utah State University Institutional Review Board

approved secondary analyses of the 2017 Utah PNA survey data as non-human subjects research

as participants could not be re-identified (protocol #10108). Previous research has utilized

similar statewide school-based samples to identify factors associated with e-cigarette use among

adolescents across the U.S. (e.g., Hawaii, Texas, Connecticut, New Jersey; [6–9,20,34]).

The current study focused on 14,346 middle school students (i.e., 6th and 8th grade) that

participated in the 2017 Utah PNA survey. Participants were approximately 12 years old on

average (M = 12.5; SD = 1.1), were relatively balanced on sex (girls; n = 7,532, 52.5%) and 6th

grade (n = 7,473, 52.1%), and were predominantly White (9,491, 71%). Nearly a third of

students attended school within Salt Lake County (n = 4,173, 29.1%) in Utah. Youths in this

sample reported a 9.4% (n = 1,343) prevalence of lifetime e-cigarette use and 5.4% (n = 784)

tobacco use. Students largely reported abstaining from both tobacco and e-cigarette use (91%; n

= 13,003) and reported greater lifetime exclusive e-cigarette use (5.5%; n = 791) relative to

exclusive tobacco use (1.6%; n = 232). Within the sample, 3.8% (n = 552) of students reported

dual lifetime use of tobacco and e-cigarettes. See Table 1 for sample demographic information

by outcomes.

170   Measures

171   Individual and socioecological variables

The measures utilized in the current study have been traditionally used and reported by

the SHARP survey project as individual items [33]. Variables examined in the current study have

been identified as being theoretically and/or empirically important factors in the substance use

literature. We decided to examine individual items to provide a nuanced understanding of e-

cigarette use risk-factors [13]. Variables included a wide array of factors representing *individual*

(i.e., antisocial behaviors/attitudes, rebelliousness, academic performance, perceived risk of drug

use, intentions for adulthood substance use, lifetime substance use), *community* (i.e., attachment,

prosocial involvement and reward, drug use consequences and antisocial behavior, perceived

availability of substances), *school* (i.e., learning environment perceptions, enjoyment,

commitment, benefits of learning, truancy), *home* (i.e., parenting practices, family history of

substance abuse, rewards for prosocial behavior, parental attitudes regarding antisocial behavior

and drug use, relationship quality with parents), and *social* (i.e., best friends engaged in

antisocial behavior, tried alcohol or drugs, exhibited prosocial behavior; social rewards for

antisocial and prosocial behaviors) influences. See supplementary Table S1 for all items

examined in the current study.

Outcome

188   Students reported whether they ever tried electronic cigarettes or e-cigarettes (i.e., *yes* or

189   *no)*. They also reported whether they had ever tried tobacco cigarettes, even just a puff (i.e., *yes*

190   or *no*). Two dichotomous outcome variables were created from these items to represent lifetime

191   exclusive e-cigarette use and dual use (i.e., tobacco and e-cigarette). The comparison group for

192   each outcome were students who did not use either substance.

Analytic plan

In our sample, 47% ($n = 6,744$) of participants were missing at least one covariate. Prior

to imputation, data was randomly resampled into training (70%; $n = 9,657$ e-cigarette; $n = 9,490$

dual) and testing sets (30%; $n = 4,237$ e-cigarette; $n = 4,065$ dual). We then used mode

imputation, wherein missing values were replaced with the mode for each variable to address

missingness independently for training and testing sets. Mode imputation is commonly utilized

within the context of machine learning for classification task [28]. As algorithms can struggle to

predict low base rate outcomes, a method known as down sampling was used to randomly

resample and reduce the negative class (i.e., those that did not use e-cigarettes or tobacco) until it

was equal to the positive class within the training set [27]. Thus, rates of lifetime use and non-

use were equal for each outcome within the resampled training sets. The training sets were $n =$

1,108 for exclusive e-cigarette use and $n = 774$ for dual use.

Five dissimilar machine learning algorithms-elastic net, random forest, neural networks,

k-nearest neighbors, and logistic regression–were then fitted to the training set to create

classification models for each outcome [35]. Each classification algorithm drew information

from 112 variables representing student individual and socioecological factors. 5-fold cross-

validation was used to identify variables that improved classification accuracy across random

subsets of data within the training set [28]. Model performance was assessed on a test set using

the Area Under (AUC) of the Receiving Operator Characteristic (ROC) curve, which represents

the ability of a model to classify outcomes across all possible cut points [27]. The top performing

classification algorithm on the test set was selected for each outcome (i.e., AUC; sensitivity,

specificity; [27]). Variable importance figures reflect results from the best performing algorithms

for each outcome. High value classifiers were then identified through visual inspection of the

relative importance figures. Variables that demonstrate large increase in relative importance over

subsequent covariates were said to be high value classifiers [28]. High value classifiers were

examined using a crosstabulation visualization to determine the nature of the relationship

between each variable and the corresponding outcome [13].

Results

Patterns of lifetime e-cigarette use differed by demographic variables within the current

sample. Chi-square test of independence suggest e-cigarette usage was significantly ($p < .001$)

associated with student gender, grade, and race/ethnicity. Boys, 8[th] graders, Latinxs, Native

Americans, and mixed-race students reported the greatest proportion of use across outcomes.

Exclusive and dual use were generally associated with a greater proportion of lifetime use across

substances relative to no-users. See Table 1 for demographic variables by outcomes.

Exclusive use

Algorithmic performance on the exclusive e-cigarette use classification task ranged from

good to outstanding (AUC = . 787 - .926) on the test set. See supplemental Fig S1 for ROCs for

classification algorithms. Elastic net was the best performing algorithm in classifying exclusive

e-cigarette use (AUC = .926, sensitivity = .857, specificity = .848). In contrast, logistic

regression was the worst performing algorithm in classifying lifetime e-cigarette use (AUC =

.787, sensitivity = .768, specificity = .806). Elastic net identified perceived availability of e-

cigarettes, lifetime alcohol use, parents' attitudes regarding their use of vape products, school

suspension, perceived risk of e-cigarette use, lifetime marijuana use, best friend(s) tried alcohol,

best friend(s) used marijuana, and perceived risk of smoking marijuana regularly as the best

discriminators between lifetime exclusive e-cigarette users and non-users. See Fig 1 for variable

importance. Visual inspection of cross-tabulation mosaics suggests that perceived availability of

e-cigarettes (i.e., *sort of hard*, v*ery easy, sort of easy*), lifetime substance use (i.e., alcohol,

marijuana), school suspensions (i.e., 1 or more), lower levels of perceived risk associated with e-

cigarette use (i.e., *none* to *moderate*), best friend(s) tried alcohol or used marijuana (i.e., 1 or

more), and less perceived risk associated with smoking marijuana regularly were all associated

with a greater proportion of lifetime e-cigarette use. Students who reported that their parents

would view their use of vape products as "*very wrong*" had the lowest proportion of use relative

to other levels of approval (i.e., *wrong* to *not wrong at all*). See supplemental Fig S3-11 for

cross-tabulation visualizations.

Dual use

Algorithmic performance on the dual tobacco and e-cigarette use classification task also

ranged from excellent to outstanding (AUC = . 725 - .944) on the test set. See supplemental Fig

S2 for ROCs for classification algorithms. Elastic net and random forest had the same AUC

score (.944). However, elastic net (sensitivity = .824, specificity = .947) outperformed random

forest (sensitivity = .818, specificity = .939) based on sensitivity and specificity. Logistic

regression was the worst performing algorithm in classifying lifetime dual use (AUC = .725,

sensitivity = .630, specificity = .779). Elastic net identified lifetime alcohol use, lifetime

marijuana use, perceived availability of e-cigarettes, best friend(s) cigarette use, perceived risk of

e-cigarette use, lifetime inhalants use, school suspension, and perceived risk of smoking

marijuana regularly as the best discriminators between lifetime dual users and non-users. See Fig

2 for variable importance. Visual inspection of cross-tabulation mosaics suggests that lifetime

substance use (i.e., alcohol, marijuana, inhalants), higher levels of perceived availability of e-

cigarettes (i.e., v*ery easy, sort of easy*), best friend(s) that have smoked cigarettes (i.e., 1 or

more), school suspensions (i.e., 1 or more), lower levels of perceived risk associated with e-

cigarette use and using marijuana regularly (i.e., *none* to *moderate*) were all associated with a

greater proportion of lifetime dual use. See supplemental Fig S12-19 for cross-tabulation

visualizations.

Discussion

The current study expands the literature through the simultaneous exploration of

established correlates of e-cigarette initiation and traditional factors associated with substance

use in relation to early adolescent vaping. Algorithms utilizing information regarding student

individual characteristics and socioecological context demonstrated high levels of classification

accuracy for both lifetime exclusive and dual e-cigarette use. Elastic net generally outperformed

other algorithm in classification accuracy. While the order of importance of classifiers differed

by outcome, elastic net consistently identified six high value classifiers across usage groups:

lifetime alcohol or marijuana use, perception of e-cigarette availability and risk, school

suspension(s), and perceived risk of smoking marijuana regularly. Several high value classifiers

differed between youth who reported lifetime exclusive (i.e., parent's attitudes regarding their

use of vaping products, best friend[s] tried alcohol, best friend[s] used marijuana) and dual e-

cigarette use (i.e., best friend[s] smoked cigarettes, lifetime inhalants use). These findings

highlight important commonalities and difference in risk profiles between lifetime exclusive and

dual e-cigarette users.

Research using high school samples have documented higher rates of life substance use

among dual versus exclusive e-cigarette users [7]. Within our sample, rates of substance use

were generally higher among exclusive and dual users relative to those who abstained from both.

Consistent with prior research, the greatest portions of substance use were found among youth

who had reported dual use [26]. However, only lifetime alcohol and tobacco were found to be

important classifiers of both e-cigarette use outcomes among middle school students, which is

consistent with prior findings in high school samples [7]. Our findings also extend prior work

through the identification of inhalant use as a novel risk factor specifically related to lifetime

dual use during early adolescence. It is possible that dual users may access a wide variety of

substances recreationally and may utilize inhalants as they are easy to access within the home

[26,36]. Further research is needed to understand the relationship between lifetime inhalant and

dual use.

Our findings confirm that availability of e-cigarettes is an important influence on early

adolescent vaping [19], despite a Utah state law that restrict the sale of these products to

individuals under the age of 20. Accessibility was especially important to exclusive e-cigarette

use, which is concerning as this may translate to future traditional cigarette use among

adolescents who may have otherwise abstained from tobacco use [8]. Consistent with prior

research, students who reported lower perceived danger of using e-cigarettes reported a greater

proportion of lifetime use [12]. Our findings suggest a need to consider perceptions regarding the

danger of other substances, such as marijuana, when assessing risk for early adolescent e-

cigarette use. While recent findings have highlighted the importance of school-based factors in

assessing risk for e-cigarette using among high school samples (i.e., truancy and poor academic

performance; [10]), our findings suggest that student suspensions were the most relevant aspect

of school in relation to early adolescent e-cigarette use. It is possible school suspensions may be

associated with an increased risk for e-cigarette use as a potential proxy for rule breaking

behaviors or through greater unsupervised time outside of school [7,37]. Further research is

needed to elucidate the relationship between school suspensions and early adolescent e-cigarettes

use.

It is important to mention that factors traditionally associated with substance use such as

adolescent and peer delinquency, community substance use norms, and school involvement were

not important predictors of lifetime exclusive and dual e-cigarette use within the current sample.

These findings may signal potential differences between factors underlying e-cigarette and other

forms of substance use. Additionally, factors identified by prior research as relevant predictors of

vaping (e.g., parenting practices, perceived risk of smoking tobacco) were not relevant correlates

of patterns of e-cigarette use within the current sample [7,12]. It is possible that when competing

against other variables within a machine learning approach, these important predictors are truly

of lesser importance relative to high value classifiers identified in the current study.

Implications

Our findings support addressing early adolescent vaping through prevention programs

aiming to address substance use prevention broadly (i.e., alcohol, tobacco, marijuana, inhalants,

e-cigarette). There is ample evidence for the efficacy of prevention programs whether focused on

a single substance or multiple ones [38]. Our findings provide specific structural and behavioral

targets that may inform the adaptation of programs seeking to prevent different patterns of early

adolescent e-cigarette use. Substance use prevention programs may benefit from adding

components that equip parents to effectively communicate disapproval regarding their child's

use of e-cigarette and teach youth skills to resist peer substance use influences [39,40]. Targeted

prevention programs are already supported by research documenting perception of risk

associated with vaping as a consistent predictor of e-cigarette use [7,8] and our findings highlight

the importance of also considering perceptions of danger regarding marijuana use during early

adolescence. Specific to programming, youth's perceptions of risk do not align with research

evidence providing an important point of content for preventive interventions [41]. Altering

youth's perception of e-cigarette accessibility, however, may require intervention at a broader

social level (e.g., public media campaigns). Alternatively, decreasing accessibility to e-cigarettes

may be achieved by actions external to youths such as strong enforcement of laws regarding

possession and/or consumption for underage users and/or those selling e-cigarette products to

them, or by way of increasing prices for goods associated with e-cigarette use.

Machine learning appears to be a promising screening tool for the identification of risk

factors that can accelerate the development of the e-cigarette knowledge base needed to curb the

rapid spread of vaping among adolescents nationally. Algorithms were able to efficiently explore

a wide range of factors in association with early adolescent e-cigarette use, which confirmed

findings from later developmental stages and identified several novel risk factors (i.e., inhalant

use, perceived risk of marijuana, school suspensions). An important consideration in this

research is that the tools utilized to identify e-cigarette use classifiers are publicly available. R

offers open access statistical packages for machine learning. Additionally, there are substantial

training materials available for free online. These tools can provide an accessible and replicable

method of generating and disseminating scientific knowledge regarding e-cigarette use classifiers

nationally. We encourage researchers to apply machine learning algorithms to their data to draw

new insight regarding factors contributing to a variety of e-cigarette use outcomes among

adolescents. Examining cross-sectional markers of e-cigarette use could also identify important

variables that can be examined longitudinally in prospective research. Machine learning

algorithms have many exciting applications when applied to longitudinal data, including

identifying context specific predictors of service use, specific targets for substance use

prevention programs, and ensure that important factors are not excluded from causal models

examining mechanism underpinning early vaping initiation.

Limitations

Results from algorithms used in the current study do not necessarily imply causal

mechanisms explaining patterns of lifetime e-cigarette use but rather identify factors that are

strong correlates of group membership (i.e., use or non-use). Longitudinal research is needed to

establish causal links. The current study examined lifetime substance use that may range from

experimentation to habitual use. Future research may consider using machine learning as a

method of identifying youth at-risk for habitual e-cigarette use. Although a large number of the

Utah adolescent population was captured, the PNA survey does not include students in private

schools, correctional facilities, or treatment centers. Additionally, students who were not in

attendance, declined participation, or did not return parental consent forms are not represented.

Furthermore, findings may not generalize to students in other states. Further research is needed

to replicate our findings in different contexts and developmental periods.

Conclusions

The current study utilized a machine learning approach to efficiently explore and identify high value correlates of early adolescent lifetime e-cigarette use. This approach identified several shared risk factors for exclusive and dual e-cigarette use such as lifetime use of specific substances (i.e., alcohol, marijuana), perception of e-cigarette availability and risk, school suspension(s), and perceived risk of smoking marijuana regularly. Several differences were also identified between youth who reported lifetime exclusive (i.e., parent's attitudes regarding their use of vaping products, best friend[s] tried alcohol or used marijuana) and dual use (i.e., best friend[s] smoked cigarettes, lifetime inhalants use) relative to non-users. This information provides a first step towards identifying youth at-risk for e-cigarette use during early adolescence. Further research is needed to examine high value classifiers identified by the current study using explanatory models and longitudinal data to understand mechanism underlying their importance in accounting for differences in risk profiles between e-cigarette usage groups during early adolescence.

Reporting

Funding

This research was funded by a collaboration between multiple state agencies in Utah (i.e., Department of Health, Department of Human Services, and the State Board of Education).

Disclosure Statement

The authors have no conflicts of interest associated with the publication of this manuscript.

Acknowledgements

This research was funded by a collaboration between multiple state agencies in Utah (i.e., Department of Health, Department of Human Services, and the State Board of Education).

Data Availability

Data used in the current study can be requested from the Utah Department of Human and Health Services.

Data Deposition

We are not authorized to share the Utah Prevention Needs Assessment data used in the current study as it is owed and managed by the Utah Department of Human and Health Services.

References

1. Office of the Surgeon General. Surgeon General's Advisory on E-cigaretteUse Among Youth. 2019. Available: https://www.cdc.gov/tobacco/basic_information/e-cigarettes/surgeon-general-advisory/index.html

2. Centers for Disease Control and Prevention. Tobacco product use among middle and high school students—United States, 2011–2017. Morb Mortal Wkly Rep. 2018;63: 629–633. doi:http://dx.doi.org/ 10.15585/mmwr.mm6722a3

3. Bronfenbrenner U. Toward an experimental ecology of human development. Am Psychol. 1977;32: 513–531. doi:http://doi.org/10.1037/0003-066X.32.7.513

4. Barrington-Trimis JL, Berhane K, Unger JB, Cruz TB, Urman R, Chou P, et al. The E-cigarette Social Environment, E-cigarette Use, and Susceptibility to Cigarette Smoking. 2016 [cited 26 Jun 2019]. doi:10.1016/j.jadohealth.2016.03.019

5. Bold KW, Morean ME, Kong G, Simon P, Camenga DR, Cavallo DA, et al. Early age of e-cigarette use onset mediates the association between impulsivity and e-cigarette use frequency in youth. Drug Alcohol Depend. 2017;181: 146–151. doi:10.1016/j.drugalcdep.2017.09.025

6. Giovenco DP, Casseus M, Duncan DT, Coups EJ, Lewis MJ, Delnevo CD. Association Between Electronic Cigarette Marketing Near Schools and E-cigarette Use Among Youth. J Adolesc Heal. 2016;59: 627–634. doi:10.1016/j.jadohealth.2016.08.007

7. Wills TA, Knight R, Williams RJ, Pagano I, Sargent JD. Risk factors for exclusive e-cigarette use and dual e-cigarette use and tobacco use in adolescents. Pediatrics. 2015;135: e43–e51. doi:10.1542/peds.2014-0760

8. Wills TA, Sargent JD, Knight R, Pagano I, Gibbons FX. E-cigarette Use and Willingness

to Smoke in a Sample of Adolescent Nonsmokers. 2016;2: 1–16.

doi:10.14440/jbm.2015.54.A

9.   Simon P, Camenga DR, Morean ME, Kong G, Bold KW, Cavallo DA, et al.

Socioeconomic status and adolescent e-cigarette use: The mediating role of e-cigarette

advertisement exposure. Prev Med (Baltim). 2018;112: 193–198.

doi:10.1016/j.ypmed.2018.04.019

10.   McCabe SE, West BT, Veliz P, Boyd CJ. E-cigarette Use, Cigarette Smoking, Dual Use,

and Problem Behaviors Among U.S. Adolescents: Results From a National Survey. J

Adolesc Heal. 2017;61: 155–162. doi:10.1016/j.jadohealth.2017.02.004

11.   Rocheleau GC, Vito AG, Intravia J. Peers, Perceptions, and E-Cigarettes: A Social

Learning Approach to Explaining E-Cigarette Use Among Youth. J Drug Issues. 2020;50:

472–489. doi:10.1177/0022042620921351

12.   Kwon E, Seo DC, Lin HC, Chen Z. Predictors of youth e-cigarette use susceptibility in a

U.S. nationally representative sample. Addict Behav. 2018;82: 79–85.

doi:10.1016/j.addbeh.2018.02.026

13.   Vázquez AL, Domenech Rodríguez MM, Barrett TS, Schwartz S, Amador Buenabad NG,

Bustos Gamiño MN, et al. Innovative Identification of Substance Use Predictors: Machine

Learning in a National Sample of Mexican Children. Prev Sci. 2020;21: 171–181.

doi:10.1007/s11121-020-01089-4

14.   National Institute on Drug Abuse. What are electronic cigarettes? 2018.

15.   Tierney PA, Karpinski CD, Brown JE, Luo W, Pankow JF. Flavour chemicals in

electronic cigarette fluids. Tob Control. 2016;25: e10–e15. doi:10.1136/tobaccocontrol-

2014-052175

16.   Centers for Disease Control and Prevention. Outbreak of lung injury associated with the

458   use of e-cigarette , or vaping, products. 2020.

459   17.   Carey FR, Rogers SM, Cohn EA, Harrell ☐ MB, Wilkinson A V, Perry CL.

460   Understanding susceptibility to e-cigarettes: A comprehensive model of risk factors that

461   influence the transition from non-susceptible to susceptible among e-cigarette naïve

462   adolescents. Addict Behav. 2019;91: 68–74. doi:10.1016/j.addbeh.2018.09.002

463   18.   Cullen KA, Gentzke AS, Sawdey MD, Chang JT, Anic GM, Wang TW, et al. e-Cigarette

464   Use among Youth in the United States, 2019. JAMA - J Am Med Assoc. 2019;322: 2095–

465   2103. doi:10.1001/jama.2019.18387

466   19.   Kong G, Morean ME, Cavallo DA, Camenga DR, Krishnan-Sarin S. Reasons for

467   electronic cigarette experimentation and discontinuation among adolescents and young

468   adults. Nicotine Tob Res. 2015;17: 847–854. doi:10.1093/ntr/ntu257

469   20.   Krishnan-Sarin S, Morean ME, Camenga DR, Cavallo DA, Kong G. E-cigarette use

470   among high school and middle school adolescents in Connecticut. Nicotine Tob Res.

471   2015;17: 810–818. doi:10.1093/ntr/ntu243

472   21.   Eitle DJ, Eitle TM. School and county characteristics as predictors of school rates of drug,

473   alcohol, and tobacco offenses. J Health Soc Behav. 2004;45: 408–421.

474   doi:10.1177/002214650404500404

475   22.   Evans WD, Powers A, Hersey J, Renaud J. The influence of social environment and social

476   image on adolescent smoking. Heal Psychol. 2006;25: 26–33.

477   doi:https://doi.org/10.1037/0278-6133.25.1.26

478   23.   Mason MJ, Zaharakis NM, Rusby JC, Westling E, Light JM, Mennis J, et al. A

479   longitudinal study predicting adolescent tobacco, alcohol, and cannabis use by behavioral

characteristics of close friends. Psychol Addict Behav. 2017;31: 712–720.

doi:https://doi.org/10.1037/adb0000299

24. Novak SP, Clayton RR. The influence of school environment and self-regulation on

transitions between stages of cigarette smoking: A multilevel analysis. Heal Psychol.

2001;20: 196–207. doi:10.1037/0278-6133.20.3.196

25. Voelkl KE, Frone MR. Predictors of substance use at school among high school students.

J Educ Psychol. 2000;92: 583–592. doi:10.1037/0022-0663.92.3.583

26. Curran KA, Burk T, Pitt PD, Middleman AB. Trends and Substance Use Associations

With E-Cigarette Use in US Adolescents. Clin Pediatr (Phila). 2018;57: 1191–1198.

doi:10.1177/0009922818769405

27. Kuhn M, Johnson K. Applied Predictive Modeling with Applications in R. 2013.

Available:

http://appliedpredictivemodeling.com/s/Applied_Predictive_Modeling_in_R.pdf

28. Hastie T, Tibshirani R, Friedman J. The Elements of Statistical Learning. Second. New

York: Springer; 2013. doi:10.1007/978-1-4419-9863-7_941

29. Barenholtz E, Fitzgerald ND, Hahn WE. Machine-learning approaches to substance-abuse

research: emerging trends and their implications. Curr Opin Psychiatry. 2020;33: 334–

342. doi:10.1097/YCO.0000000000000611

30. Ketonen V, Malik A. Characterizing vaping posts on instagram by using unsupervised

machine learning. Int J Med Inform. 2020;141: 104223.

doi:10.1016/j.ijmedinf.2020.104223

31. Malik A, Khan MI, Karbasian H, Nieminen M, Ammad-Ud-Din M, Khan SA. Modeling

Public Sentiments About JUUL Flavors on Twitter Through Machine Learning. Nicotine

Tob Res. 2021; 1–11. doi:10.1093/ntr/ntab098

32.   Choi J, Jung H-T, Ferrell A, Woo S, Haddad L. Machine Learning-Based Nicotine

Addiction Prediction Models for Youth E-Cigarette and Waterpipe (Hookah) Users. J Clin

Med. 2021;10: 972. doi:10.3390/jcm10050972

33.   Utah Department of Human Services. Student Health and Risk Prevention (SHARP)

survey reports. 2019.

34.   Temple JR, Shorey RC, Lu Y, Torres E, Stuart GL, Le VD. E-cigarette use of young

adults motivations and associations with combustible cigarette alcohol, marijuana, and

other illicit drugs. Am J Addict. 2017;26: 343–348. doi:10.1111/ajad.12530

35.   James G, Witten D, Hastie T, Tibshirani R. An Introduction to Statistical Learning.

Current Medicinal Chemistry. Springer; 2013.

36.   Medina-Mora M´a E, Real T. Epidemiology of inhalant use. Curr Opin Psychiatry.

2008;21: 247–251. doi:https://doi.org/10.1097/yco.0b013e3282fc9875

37.   Lee KTH, Vandell DL. Out-of-School Time and Adolescent Substance Use. J Adolesc

Heal. 2015;57: 523–529. doi:10.1016/j.jadohealth.2015.07.003

38.   Das JK, Salam RA, Arshad A, Finkelstein Y, Bhutta ZA. Interventions for Adolescent

Substance Abuse: An Overview of Systematic Reviews. J Adolesc Heal. 2016;59: S61–

S75. doi:10.1016/j.jadohealth.2016.06.021

39.   Marsiglia FF, Kulis S, Yabiku ST, Nieri TA, Coleman E. When to Intervene: Elementary

School, Middle School or Both? Effects of keepin' It REAL on Substance Use

Trajectories of Mexican Heritage Youth. Prev Sci. 2011;12: 48–62. doi:10.1007/s11121-

010-0189-y

40.   Patterson G. The next generation of PMTO models. Behav Ther. 2005;28: 25–32.

41.   Cheeta S, Halil A, Kenny M, Sheehan E, Zamyadi R, Williams AL, et al. Does perception

527   of drug-related harm change with age? A cross-sectional online survey of young and older

528   people. BMJ Open. 2018;8. doi:10.1136/bmjopen-2017-021109

Table 1. Student Demographics by Lifetime E-Cigarette Use Groups.					
	Total	**Neither**	**Exclusive Use**	**Dual use**	***p* value[a]**
Sample	*n* = 14346	*n* = 13003	*n* = 791	*n* = 552	
Age *M*(*SD*)	12.5 (1.1)	12.5 (1.1)	13.1 (1.0)	13.1 (1.0)	
Sex					< .001
Boy	6766 (47.2%)	6066 (89.7%)	427 (6.3%)	273 (4%)	
Girl	7532 (52.5%)	6894 (91.5%)	361 (4.8%)	277 (3.7%)	
Grade					< .001
6th	7473 (52.1%)	7110 (95.1%)	217 (2.9%)	146 (2%)	
8th	6873 (47.9%)	5893 (85.7%)	574 (8.4%)	406 (5.9%)	
Race/ethnicity					< .001
White	10191 (71%)	9491 (93.1%)	415 (4.1%)	285 (2.8%)	
Native American	289 (2%)	247 (85.5%)	20 (6.9%)	22 (7.6%)	
Asian	238 (1.7%)	221 (92.9%)	12 (5%)	5 (2.1%)	
Black	210 (1.5%)	186 (88.6%)	11 (5.2%)	13 (6.2%)	
Latinx	1888 (13.2%)	1554 (82.3%)	210 (11.1%)	124 (6.6%)	
Pacific Islander	210 (1.5%)	191 (91%)	13 (6.2%)	6 (2.9%)	
Mixed Race	1320 (9.2%)	1113 (84.3%)	110 (8.3%)	97 (7.3%)	
Substance use[b]					
Alcohol					< .001
Yes	1329 (9.3%)	600 (45.1%)	353 (26.6%)	376 (28.3%)	
No	12982 (90.5%)	12371 (95.3%)	436 (3.4%)	175 (1.3%)	
Marijuana					< .001
Yes	602 (4.2%)	124 (20.6%)	201 (33.4%)	277 (46%)	
No	13671 (95.3%)	12812 (93.7%)	585 (4.3%)	274 (2%)	
Inhalants					< .001
Yes	669 (4.7%)	414 (61.9%)	105 (15.7%)	150 (22.4%)	
No	13579 (94.7%)	12499 (92%)	679 (5%)	401 (3%)	
Prescription drugs					< .001
Yes	546 (3.8%)	332 (60.8%)	81 (14.8%)	133 (24.4%)	
No	13562 (94.5%)	12468 (91.9%)	690 (5.1%)	404 (3%)	
Hallucinogens					< .001
Yes	99 (0.7%)	22 (22.2%)	25 (25.3%)	52 (52.5%)	
No	14128 (98.5%)	12871 (91.1%)	760 (5.4%)	497 (3.5%)	
Synthetic marijuana					< .001
Yes	86 (0.6%)	12 (14%)	21 (24.4%)	53 (61.6%)	
No	14199 (99%)	12940 (91.1%)	766 (5.4%)	493 (3.5%)	

549   Variable frequency is displayed by column for the total and row for usage groups. [a] Chi-square
550   test of independence. [b] Lifetime use.

551   **Fig 1. Top 20 Variables with the Highest Relative Importance in Classifying Lifetime E-**

552   **Cigarette Use.** Results represent validation on a separate test dataset.

558   **Fig 2. Top 20 Variables with the Highest Relative Importance in Classifying Lifetime Dual**

559   **Use.** Results represent validation on a separate test dataset.

560