Real-world Effectiveness of BNT162b2 in Children and Adolescents in Preventing Infection and Severe Diseases with SARS-CoV-2 During the Delta and Omicron Periods: Findings from Trial Emulation Using an EHR-based Cohort

Qiong Wu, PhD[1]^, Jiayi Tong[1]^, Bingyu Zhang[1], Dazheng Zhang[1], Jie Xu, PhD[2], Yishan Shen[1], Lu Li[1], L. Charles Bailey, MD, PhD[3], Jiang Bian PhD[2], Dimitri A. Christakis, MD, MPH[4], Megan L. Fitzgerald, PhD[5], Kathryn Hirabayashi[3], Ravi Jhaveri, MD[6], Alka Khaitan, MD[7], Tianchen Lyu[2], Suchitra Rao, MBBS, MSCS[8], Hanieh Razzaghi[3], Hayden T. Schwenk[9], Fei Wang, PhD[10], Margot I. Witvliet, PhD[11], Eric J. Tchetgen Tchetgen, PhD[12], Jeffrey S. Morris, PhD[1]*, Christopher B. Forrest, MD, PhD[10]*, and Yong Chen, PhD[1]*

1. Department of Biostatistics, Epidemiology, and Informatics, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, USA
2. Department of Health Outcomes Biomedical Informatics, University of Florida, Gainesville, FL, USA
3. Applied Clinical Research Center, The Children's Hospital of Philadelphia, Philadelphia, PA, USA
4. Center for Child Health, Behavior, and Development, Seattle Children's Research Institute, Seattle, WA, USA
5. Department of Medicine, Grossman School of Medicine, New York University, New York, NY, USA
6. Division of Pediatric Infectious Diseases, Ann & Robert H. Lurie Children's Hospital of Chicago, Chicago, IL, USA
7. Department of Pediatrics, Ryan White Center for Pediatric Infectious Diseases and Global Health, Indiana University School of Medicine, IN, USA
8. Department of Pediatrics, University of Colorado School of Medicine and Children's Hospital Colorado, Aurora, CO, USA
9. Department of Pediatrics, Stanford School of Medicine, Stanford, CA, USA
10. Department of Population Health Sciences, Weill Cornell Medicine, New York, NY, USA
11. Department of Sociology, Social Work and Criminal Justice, Lamar University, Beaumont, TX, USA
12. Department of Statistics and Data Science, The Wharton School, The University of Pennsylvania, PA, USA

^: Co-first authors
*: Co-senior authors

Corresponding authors:
Yong Chen, PhD, ychen123@pennmedicine.upenn.edu
Christopher B. Forrest, MD, PhD, forrestc@chop.edu

Authorship has been determined according to ICMJE recommendations.

ABSTRACT

BACKGROUND
The current understanding of the long-term effectiveness of the BNT162b2 vaccine for a range of outcomes across diverse U.S. pediatric populations is limited. In this study, we assessed the effectiveness of BNT162b2 against various strains of the SARS-CoV-2 virus using data from a national collaboration of pediatric health systems (PEDSnet).

METHODS
We emulated three target trials to assess the real-world effectiveness of BNT162b2: adolescents aged 12 to 20 years during the Delta variant period (Target trial 1), children aged 5 to 11 years (Target trial 2) and adolescents aged 12 to 20 years during the Omicron variant period (Target trial 3). The outcomes included documented infection, COVID-19 illness severity, admission to an intensive care unit (ICU), and two cardiac-related outcomes, myocarditis and pericarditis. In the U.S., immunization records are often captured and stored across multiple disconnected sources, resulting in incomplete vaccination records in patients' electronic health records (EHR). We implemented a novel trial emulation pipeline accounting for possible misclassification bias in vaccine documentation in EHRs. The effectiveness of the BNT162b2 vaccine was estimated from the Poisson regression model with confounders balanced via propensity score stratification.

RESULTS
During the Delta period, the BNT162b2 vaccine demonstrated an overall effectiveness 98.4% (95% CI, 98.1 to 98.7) against documented infection among adolescents, with no significant waning after receipt of the first dose. During the Omicron period, the overall effectiveness was estimated to be 74.3% (95% CI, 72.2 to 76.2) in preventing documented infection among children, which was higher against moderate or severe COVID-19 (75.5%; 95% CI, 69.0 to 81.0) and ICU admission with COVID-19 (84.9%; 95% CI, 64.8 to 93.5). In the adolescent population, the overall effectiveness against documented Omicron infection was 85.5% (95% CI, 83.8 to 87.1), with effectiveness of 84.8% (95% CI, 77.3 to 89.9) against moderate or severe COVID-19, and 91.5% (95% CI, 69.5 to 97.6) against ICU admission with COVID-19. The effectiveness of the BNT162b2 vaccine against the Omicron variant declined after 4 months following the first dose and then stabilized with higher levels of uncertainty. Across all three cohorts, the risk of cardiac outcomes was approximately 65% to 85% lower in the vaccinated group than that of the unvaccinated group accounting for possible misclassification bias.

CONCLUSIONS
This study suggests BNT162b2 was effective among children and adolescents in Delta and Omicron periods for a range of COVID-19-related outcomes and is associated with a lower risk for cardiac complications. Waning effectiveness over time suggests that revaccination may be needed in the future.

INTRODUCTION

The Food and Drug Administration (FDA) expanded the emergency use authorization of the BNT162b2 mRNA COVID-19 vaccine (Pfizer-BioNTech) to 12-15-year-olds on May 10, 2021, and to 5-11-year-olds on October 29, 2021. As of April 5, 2023, the Centers for Disease Control and Prevention (CDC) reports indicate that 40% of U.S. children aged 5-to-11-year-olds and 72% of adolescents aged 12-to-18-year-olds had received at least one dose of the vaccine. The emergence of the Omicron variant (B.1.1.529) and its subvariants in early 2022 led to a new surge in COVID-19 cases worldwide[1]. The randomized trials of the BNT162b2 vaccine which demonstrated high efficacy of 2 doses against COVID-19 (100% and 91% among those aged 12-15 and 5-11 years, respectively) were conducted before the emergence of the Omicron variant[2,3].

Several observational studies have been conducted to investigate the effectiveness of vaccination in real-world settings[4–8]. However, prior studies have had limited follow-up periods, covering the Delta variant or earlier subvariants of Omicron periods only. Studies evaluating the Omicron variant have only assessed the short-term effects of the vaccine, with only one study involving children evaluating the effect beyond 3 months[9]. There is limited information on the long-term durability of vaccine protection during the Omicron period. Few existing studies on U.S. pediatric populations have covered both hospitalized patients and those with mild or asymptomatic conditions. Furthermore, while studies have acknowledged limitations due to misclassification in vaccination status in real-world effectiveness studies, none have rigorously evaluated the impacts of such misclassification nor accounted for the potential bias it may introduce.

To address these gaps in our knowledgebase of the pediatric effectiveness of SARS-CoV-2 vaccination, we designed this study to assess the real-world effectiveness of BNT162b2 among children and adolescents during the Delta and Omicron variant-predominant periods using electronic health record (EHR) data from a national network of U.S. pediatric medical centers. Our study used a trial emulation design and adjusted for misclassification issues in vaccination status and has several attractive features that strengthen credibility of inferences. First, it is the largest study to date in the U.S. estimating vaccine effectiveness in children and adolescents, covering a broad spectrum of the U.S. pediatric population. Second, the study examined the effectiveness against infection over a longer follow-up period than any previous study, enabling evaluation of the durability of vaccine protection. Third, the study included a diverse representation of U.S. pediatric populations from primary care, specialty care, emergency department, testing centers, and inpatient settings. Fourth, the study was the first to account for the incomplete capture of vaccination status by health systems in the U.S. Finally, besides infection and severe disease endpoints, we also studied the effect of vaccination on the incidence of myocarditis and pericarditis to assess the effect of vaccination relative to this known potential risk.

METHODS

DATA SOURCES

This study used EHR data from PEDSnet[10], which is a national collaboration of pediatric health systems that share EHR data, conduct research, and improve outcomes together. Participating institutions in this study included: Children's Hospital of Philadelphia, Cincinnati Children's Hospital Medical Center, Children's Hospital Colorado, Ann & Robert H. Lurie Children's Hospital of Chicago, Nationwide Children's Hospital, Nemours Children's Health System (inclusive of the Delaware and Florida health system), Seattle Children's Hospital, and Stanford Children's Health. Data were extracted from the PEDSnet COVID-19 Database Version Week 141[11].

SPECIFICATION AND EMULATION OF TARGET TRIALS

Target trials are hypothetical randomized controlled trials (RCT) that guide the design of observational studies to assess the real-world effectiveness of treatments[12]. We designed and conducted emulation of three target trials to investigate the effectiveness of the BNT162b2 vaccine in preventing infection with various strains of the SARS-CoV-2 virus in children and adolescents in the United States (Table 1). The three target trials focused on documented SARS-CoV-2 infection and outcomes in:

- *Target trial 1* (Delta study in adolescents): adolescents aged 12 to 20 years during the period when the Delta variant was prevalent from July 1, 2021, to November 30, 2021.
- *Target trial 2* (Omicron study in children): children aged 5 to 11 years during the period when the Omicron variant was prevalent from January 1, 2022, to November 30, 2022.
- *Target trial 3* (Omicron study in adolescents): adolescents aged 12 to 20 years during the period when the Omicron variant was prevalent from January 1, 2022, to November 30, 2022.

The design of target trials and trial emulation procedures in real-world data are summarized in Table 1.

Eligibility criteria included age of 5 to 11 years for children or 12 to 20 years for adolescents at the start of the study period and no previous COVID-19 vaccination or documented SARS-CoV-2 infection. Additionally, participants were required to have a prior encounter (including telephone or telehealth encounters) within 18 months of cohort entrance to ensure that they had an ongoing interaction with the health system.

The intervention of interest was vaccination, in comparison with no receipt of any type of COVID-19 vaccine. Since the BNT162b2 vaccine covered more than 85% of documented vaccinations among children and adolescents in the PEDSnet database, in this study we focused primarily on studying the effectiveness of this vaccine, although the supplementary appendix reports a sensitivity analysis investigating all types of reported COVID-19 assessed in the U.S., with 85.6% BNT162b2, 1.2% mRNA-1273, and 12.4% unspecified COVID-19 vaccine.

In the emulation of the target trials, the cohort entrance date for the intervention group was defined as the date of the first dose of the BNT162b2 vaccine, while for the comparator group, it was a randomly selected date from visits, chosen to ensure the distribution of index dates for the control group matched the

distribution of index dates for the vaccination group to control for time effects. The risk period for the study began 28 days after the index date such that infections within 28 days were excluded.

Randomized trials achieve balance across potential confounders by randomly allocating the treatment to intervention and comparator groups. In our trial emulation, we balanced the intervention and comparator groups by adjusting for a large number of measured confounders using propensity score stratification[13]. We built the propensity score model based on demographic factors including age, sex, race/ethnicity, clinical factors including obesity status, a chronic condition indicator as defined by the Pediatric Medical Complexity Algorithm (PMCA)[14], and a list of pre-existing chronic conditions, and healthcare utilization factors including the number of inpatients, outpatients, ED visits, unique mediations, and the number of negative COVID-19 tests prior to the cohort entry. We stratified the patients into propensity score quintiles based on these factors. We refer to Table S1 in the Supplementary Appendix for detailed definitions of study variables.

The four COVID-19 outcomes of interest were: documented SARS-CoV-2 infection, mild COVID-19, moderate/severe COVID-19, and ICU admission with COVID-19. We did not evaluate death from COVID-19 as it was too rare among children and adolescents to study quantitatively. In the emulation of target trials, SARS-CoV-2 infections were defined by and occurrence of positive polymerase-chain-reaction (PCR), serology, or antigen tests or diagnoses of COVID-19, post-acute sequelae of SARS-CoV-2 (PASC), or multisystem inflammatory syndrome in children (MIS-C) regardless of the presence of symptoms. Classification of mild, moderate, or severe COVID-19 infections was defined based on the severity classification of COVID-19 in Forrest et al. (2022)[15]. ICU admission with COVID-19 was defined by any ICU visit 7 days prior to 13 days after documented SARS-CoV-2 infection. Additionally, we also considered the clinical outcomes of myocarditis or pericarditis incidence to evaluate the effect of the BNT162b2 vaccine in terms of cardiac outcomes.

STATISTICAL ANALYSIS

We evaluated covariate balancing after propensity score stratification by plotting the standardized mean differences (SMD) between variable values for the vaccinated and unvaccinated groups, with a difference of 0.1 or less indicating an acceptable balance. We used Poisson regression to estimate the relative risk between two treatment arms for the risk of each outcome while adjusting for different follow-up lengths among participants. Since immunization records are often captured and stored across multiple disconnected sources, resulting in incomplete vaccination records in patients' EHRs, we mitigated the potential bias arising from this measurement error by incorporating an integration method of the Poisson likelihood in a pre-specified range of misclassification rates. The vaccine effectiveness was defined as 100*(1-relative risk). The details of statistical methods are described in Section S1 of the Supplementary Appendix.

We conducted secondary analyses stratified by 2-month intervals since receipt of vaccination to investigate the durability of vaccine protection. Subgroup analyses were also performed to investigate differences in vaccine protection according to age groups (5-to-8, 9-to-11, 12-to-15, 16-to-20).

SENSITIVITY ANALYSES

Extensive sensitivity analyses were conducted to evaluate the robustness of the research findings; see Supplementary Appendix sections S3-10 for the impact of cohort design. In scenarios in which any categorical covariates were unbalanced (with a standardized mean difference>0.1), we included a sensitivity analysis excluding participants in that category. Since the proportion of patients entered with ED visits is relatively low in the vaccinated cohort compared to the unvaccinated, we conducted a sensitivity analysis excluding all participants who entered the cohort due to an ED visit. We studied the dose-response of BNT162b2 by defining a single-dose vaccination as one-dose 28 days prior to infection and two-dose vaccination as the receipt of a second dose 14 days prior to infection. To further evaluate the risk of cardiac outcomes associated with vaccination, we estimated the vaccine effects on myocarditis and pericarditis stratified by gender and age groups. Residual study bias from unmeasured and systematic sources can still exist in observational studies after controlling for measured confounding; thus, we conducted negative control outcome experiments, where the null hypothesis of no effect was believed to be true using 40 control outcomes specified by pediatric physicians. The empirical null distribution and calibrated effectiveness were reported as sensitivity results. We also reported the estimated vaccine effectiveness from all brands of COVID-19 vaccines.

MISSINGNESS IN VACCINE RECORDS

Vaccine status may be missing for individuals whose vaccine doses were administered by a site outside of the PEDSnet network care delivery sites. It is likely that patients recorded as vaccinated in the EHR are true positives, so specificity could be very high, but sensitivity would be reduced by undocumented vaccinations (false negatives). To account for potential bias from the misclassification of vaccination status, a range of possible sensitivities based on our prior study was pre-specified for each study. The sensitivity range was considered to be 0.8 to 1 for the study involving children and 0.7 to 0.9 for the studies involving adolescents. By accounting for misclassification and specifying a range of sensitivity, the study aimed to minimize the impact of bias caused by the misclassification in the estimation of the effectiveness of the COVID-19 vaccine among children and adolescents. The details of statistical methods are described in Section S1 of the Supplementary Appendix.

To further evaluate the robustness of statistical methods we used to account for misclassification, we varied the trial emulation procedure by considering various methods for bias correction, including the naive method (without adjusting for misclassification), using different ranges of misclassification rates, and using a fully Bayesian method[16]. To evaluate the impact of differential misclassification on effectiveness estimates, we conducted sensitivity analyses simulating vaccination status according to various differential misclassification scenarios. Results from these sensitivity analyses are summarized in section S11-13 of the Supplementary Appendix.

RESULTS

STUDY POPULATION

A total of 77,392 adolescents within the PEDSnet network were identified to study the effectiveness of vaccination against Delta infection and severe outcomes (see Table 2A for baseline characteristics).
111,539 children and 56,080 adolescents were included in the cohort to study the effectiveness of vaccination against the Omicron infections (see Table 2B for baseline characteristics). The vaccinated and

unvaccinated groups had a slightly unbalanced distribution of testing rates before cohort entry across all three cohorts. After propensity-score stratification, all covariates were well balanced between vaccinated and unvaccinated groups with an SMD smaller than 0.1 in the Omicron study involving children (Figure S1) and involving adolescents (Figure S2). In the study evaluating vaccine effectiveness for adolescents during the Delta period, one site remained unbalanced after propensity-score stratification with an SMD larger than 0.1, and thus a sensitivity analysis was conducted by excluding participants from this site which gave consistent results with the primary analysis (see Figure S3 and Section S4 in Supplementary Appendix).

VACCINE EFFECTIVENESS

Table 3A summarizes the estimated vaccine effectiveness in three target-trial emulations and Figure 2 shows the durability of protection. The vaccine effectiveness was estimated to be 98.4% (95% CI, 98.1 to 98.7) among adolescents in the Delta period, 74.3% (95% CI, 72.2 to 76.2) against documented infection among children in the Omicron period, and 85.5% (95% CI, 83.8 to 87.1) among adolescents in the Omicron period. During the Delta period, the vaccine effectiveness against documented infection remained stable throughout the follow-up period of the study. After 4 months following the first dose, vaccine effectiveness against documented infection with Omicron declined from 82.3% (95% CI, 77.9 to 85.8) to 70.6% (95% CI, 65.9 to 74.6) among children, and from 91.3% (95% CI, 87.6 to 94.0) to 82.9% (95% CI, 79.0 to 86.1) among adolescents. Although vaccine effectiveness against documented infection stabilized after this initial decline, the corresponding confidence intervals were much wider indicating higher levels of uncertainty.

VACCINE EFFECTIVENESS: SEVERE ILLNESS AND COMPLICATIONS

During the Delta period, the vaccine was observed to have high effectiveness against documented infection 98.4% (95% CI, 98.1, 98.7) and severe COVID-19 outcomes. The estimated relative risk reduction in percentage of the vaccine on myocarditis was 73.8% (95% CI, 43.8, 87.6) and pericarditis was 66.1% (95% CI, 14.8, 86.5). The estimated vaccine effectiveness against the Omicron variant in children was 73.5% (95% CI, 69.2 to 77.1) against mild COVID-19, 75.5% (95% CI, 69.0 to 81.0) against moderate or severe COVID-19, and 84.9% (95% CI, 64.8 to 93.5) against ICU admission with COVID-19. The estimated relative risk reduction in percentage of the vaccine on myocarditis was 71.4% (95% CI, 8.8, 91.0) and pericarditis was 69.1% (95% CI, -19.7, 92.0). In the Omicron study in adolescents, the vaccine effectiveness was estimated to be 87.0% (95% CI, 83.5 to 89.8) against mild COVID-19, 84.8% (95% CI, 77.3 to 89.9) against moderate or severe COVID-19, and 91.5% (95% CI, 69.5 to 97.6) against ICU admission with COVID-19. The estimated relative risk reduction in percentage of the vaccine on myocarditis was 82.0% (95% CI, 46.6, 94.0) and pericarditis was 83.7% (95% CI, 43.3, 95.3) in this cohort.

SENSITIVITY ANALYSES AND ADDRESSING MISCLASSIFICATION BIAS

Table S9 presents the proportion of vaccinated participants with single-dose and two-dose vaccination in three studies. Table 3B summarizes the single-dose and two-dose vaccine effectiveness indicating the two doses of the BNT162b2 vaccine are slightly more effective against the Omicron variant while much more effective against the Delta variant than one dose. Section S9 presents negative control experiments of three target-trial emulations using 40 negative control outcomes. After accounting for systematic error through calibration using negative control outcomes, our findings indicate a slight shift in point estimates accompanied by wider confidence intervals. This suggests the presence of a minor degree of systematic error, as well as additional uncertainty based on the information derived from the negative control outcomes.

Section S11 shows effectiveness estimated from the naive method and proposed trial emulation pipeline with different ranges of sensitivity. The comparison results indicated that the vaccine effectiveness was reasonably consistent across different sensitivity ranges, suggesting that our primary analysis was robust to changes in the range of sensitivity considered. Section S12 shows the comparative results to a fully Bayesian method indicating nearly identical results. Section S13 shows sensitivity analyses on differential misclassifications which demonstrates the trial emulation pipeline corrects the bias even when the non-differential misclassification assumption does not hold.

DISCUSSION

We emulated three target trials to evaluate the effectiveness of BNT162b2 vaccines for the prevention of documented COVID-19 infections and severe disease in a national network of pediatric health systems in the U.S. During the period of time where the Omicron variant was dominant, the BNT162b2 vaccine was associated with a moderate protection against infection and severe outcomes with an effectiveness of approximately 70% for children and 85% for adolescents. The estimated protection decreased by roughly 10% 4 months from the first dose and slightly waned over time. During the period of time where the Delta variant was dominant, the BNT162b2 vaccine in adolescents was associated with strong protection with effectiveness higher than 95% and with little evidence of waning during the follow-up period.

Our study has several strengths. First, we used a national network of academic medical centers that covered a diverse cohort being more representative of the general pediatric population, provided a robust sample size, and allowed for multiple subgroup analyses and detection of rare outcomes. Second, the richness of these EHR data allowed us to investigate the effectiveness against infection of different levels of severity as well as adjust for a broad set of confounders. Third, we conducted the negative control outcome experiments to assess the potential residual bias due to unmeasured confounders and other potential sources of systematic bias. These experiments revealed a small amount of systematic error but with excessive uncertainty across different negative control outcomes, leading to wider confidence intervals of our estimated effectiveness that honestly reflects the impacts of unmeasured confounding and other potential sources of residual biases[17]. Finally, to the best of our knowledge, this is the first target trial emulation evaluating COVID-19 vaccines against infection and severe outcomes that explicitly handle exposure misclassification.

Our study also has several potential limitations. First, effectiveness was investigated against documented infection in a cohort without previous infection, but potential bias from undocumented infections cannot be ruled out, especially if these occurred differentially in the vaccinated and unvaccinated cohorts. Our inclusion of previous negative COVID-19 tests as a confounder can adjust for the propensity to get tested which could partially adjust for this factor. Moreover, the increasing availability of at-home rapid antigen testing kits over time could have further reduced the testing frequency captured by EHR. This factor is expected to have less impact on severe outcomes that are associated with specific symptoms of the infection. Baseline confounders were balanced between vaccinated and unvaccinated groups, which should adjust for between-cohort differences in exposure risks and risks of severe infections. Second, as in any observational study, assignment to the vaccine group was non-random and the validity of the results could be impacted

by unmeasured confounders. To evaluate the impact of unmeasured confounders and residual bias, we conducted negative control experiments that confirmed the robustness of our results.

Third, in this study, patients who had received vaccinations prior to the start of the study period were excluded. Due to missing vaccine records, some patients who had previously been vaccinated may have still entered the cohort, particularly in the unvaccinated group. However, the trial emulation pipeline used in this study adjusted for potential bias resulting from unrecorded vaccinated patients which could also reduce the bias resulting from this issue. Fourth, in the Omicron study involving adolescents, the cohort included adolescents who had their first vaccine after January 1, 2022. Since the use of BNT162b2 vaccines was authorized in adolescents aged 12-15 years on May 10, 2021, this cohort may represent a population with late vaccines which reduces the generalizability of the findings. Finally, in the dose-response analyses, the effectiveness of a single dose of the BNT162b2 vaccine was estimated by comparing patients who had a one-dose record in EHR to those with no vaccination, which may potentially reveal the effectiveness of mixing of single- and two-dose due to the missingness of the second dose. However, as the second dose is always scheduled at the receipt of the first dose, we consider the proportion of missingness on the second dose should be low.

Although this study provides evidence of a slight waning of vaccine effectiveness 4 months following the first dose against Omicron infection and the effectiveness is stabilized after 4 months, waning can be impacted by vaccines during the follow-up period and other factors. Patients who got boosters during the follow-up period were not excluded from the study. A sensitivity analysis evaluating the durability of two-dose vaccine effectiveness considering the third dose as censoring didn't suggest a significantly different conclusion. A future study is warranted to investigate the effect of booster vaccination. Furthermore, despite the recognized risk of myocarditis associated with COVID-19 vaccines in young men and teen boys, the study reveals a lower relative risk of myocarditis in vaccinated groups which can be explained by its reduced likelihood of infection[18].

In summary, this study involving national pediatric cohorts in the U.S. estimates moderate effectiveness of the BNT162b2 vaccine for preventing infection and severe diseases of the SARS-CoV-2 Omicron variant, and high effectiveness against the Delta variant. This study reveals a low risk of cardiac-related outcomes among children and adolescents who were vaccinated. This study significantly contributes to our knowledge of the BNT162b2 vaccine in the U.S. pediatric population using a rigorously designed trial emulation pipeline accounting for the incomplete capture of vaccination status in EHR data in the U.S.

Figure 1. Selection of participants for the emulation of three target trials evaluating the effectiveness of the BNT162b2 vaccine in preventing infection with SARS-CoV-2 in (*Target trial 1*) adolescents aged 12-20 years during the period when the Delta variant was prevalent, (*Target trial 2*) children aged 5 to 11 years and (*Target trial 3*) adolescents aged 12 to 20 years during the period when the Omicron variant was prevalent.

Table 1. Specification and emulation of three target trials evaluating effectiveness of the BNT162b2 vaccine in preventing infection with Delta and Omicron variants in children and adolescents.

Protocol component	Target trial specification	Target trial emulation
Eligibility criteria	• Aged 12-20 at 2021-07-01 for studying Delta in adolescents (**Target trial 1**), aged 5-11 at 2022-01-01 for studying Omicron in children (**Target trial 2**), and aged 12-20 at 2022-01-01 for studying Omicron in adolescents (**Target trial 3**), or • No previously documented infection SARS-CoV-2 based on positive polymerase-chain-reaction (PCR), serology, or antigen tests or diagnoses of COVID-19, post-acute sequelae of SARS-CoV-2 (PASC), or multisystem inflammatory syndrome in children (MIS-C) • No previous COVID-19 vaccination • Known demographic information • User of the healthcare system of PEDSnet (defined as having at least one primary care visit in the past 18 months)	Same as for the target trial, except that, participants without any interaction with the healthcare system during the study period were excluded to ensure these persons had an ongoing relationship with the health systems.
Treatment strategies	(1) Receive one dose of BNT162b2 vaccine (2) No receipt of any COVID-19 vaccines	Same as for the target trial. We defined the vaccination status and date using immunization records ascertained from health system records, using either the presence of a CVX code designating an administered or patient-reported dose or of a source value containing the terms "COVID" or "sars" in the immunization table.
Treatment assignment	Participants are randomly assigned to be vaccinated within strata defined based on age, sex, race/ethnicity, obesity, PMCA, # of inpatient, outpatient, ED, unique mediations, negative tests 18 months to 7 days prior to cohort entry (0,1,2,>=3), hospital index, medical conditions. Individuals will be aware of the assigned treatment strategy.	We assumed random assignment after propensity score stratification using the same set of variables.
Outcomes	• Documented SARS-CoV-2 infection • Mild COVID-19 (COVID-19-related symptoms: abdominal pain, anorexia, cough, diarrhea, fever/chills, headache, loss of taste or smell, fatigue, myalgia, nasal congestion, nausea, sore throat, respiratory symptoms, and vomiting.) • Moderate or severe COVID-19 (Moderate COVID-19 including acute bronchitis, bronchiolitis, dehydration, gastroenteritis, pneumonia, and use of intravenous fluids during the first 6 hours of an ED visit, and severe COVID-19 including acute respiratory distress syndrome, acute kidney injury, acute liver failure, death, encephalopathy/encephalitis, myocarditis, pericarditis, respiratory failure, sepsis, shock, thromboembolism, evidence of ICU level care, invasive or noninvasive mechanical ventilation, or vasopressor/inotropic support.) • ICU admission with COVID-19	Same as for the target trial.
Follow-up	For each participant, follow-up starts 28 days after the day of vaccination (baseline) and ends on 2021-11-30 for the Delta study in adolescents (**Target trial 1**), and 2022-11-30 for Omicron studies in children and adolescents (**Target trials 2 and 3**).	Same as for the target trial.
Causal contrasts	Intention-to-treat effect.	Observational analogue.
Statistical analysis	Relative risk comparing the vaccinated and unvaccinated cohorts using Poisson regression to adjust for different lengths of follow-up. Subgroup analyses by baseline age and time since the receipt of the first dose of the BNT162b2 vaccine.	Same as for the target trial, except that, we adjust for the potential bias from the misclassified vaccination status by integrating the Poisson likelihood in a reasonable range of misclassification rates.

Table 2A. Baseline characteristics of adolescents 12 to 20 years of age in the study of the effectiveness of the BNT162b2 vaccine in preventing infection and severe diseases with SARS-CoV-2 during the period when the Delta variant was prevalent.

	Delta study in adolescents		
	Vaccinated (N=45,007)	**Unvaccinated (N=32,385)**	**Overall (N=77,392)**
Age			
Median [Q1, Q3]	14 [13, 16]	15 [13, 17]	15 [13, 16]
Distribution			
12	8922 (19.8%)	4926 (15.2%)	13848 (17.9%)
13	8099 (18.0%)	4864 (15.0%)	12963 (16.7%)
14	8037 (17.9%)	4923 (15.2%)	12960 (16.7%)
15	7311 (16.2%)	4749 (14.7%)	12060 (15.6%)
16	5450 (12.1%)	4546 (14.0%)	9996 (12.9%)
17	4076 (9.1%)	3991 (12.3%)	8067 (10.4%)
18	1674 (3.7%)	2075 (6.4%)	3749 (4.8%)
19	942 (2.1%)	1461 (4.5%)	2403 (3.1%)
20	496 (1.1%)	850 (2.6%)	1346 (1.7%)
Gender			
Female	23,589 (52.4%)	16,500 (50.9%)	40,089 (51.8%)
Male	21,416 (47.6%)	15,880 (49.0%)	37,296 (48.2%)
Ethnicity			
White	16,446 (50.8%)	17,964 (39.9%)	34,410 (44.5%)
Black/AA	6,019 (18.6%)	12,012 (26.7%)	18,031 (23.3%)
Hispanic	4,925 (15.2%)	9,629 (21.4%)	14,554 (18.8%)
Other/Unknown	4,995 (15.4%)	5,402 (12.0%)	10,397 (13.4%)
Hospital			
A	5,424 (12.1%)	7,385 (22.8%)	12,809 (16.6%)
B	12,884 (28.6%)	5,216 (16.1%)	18,100 (23.4%)
C	6,333 (14.1%)	3,457 (10.7%)	9,790 (12.6%)
D	1,723 (3.8%)	914 (2.8%)	2,637 (3.4%)
E	4,369 (9.7%)	6,063 (18.7%)	10,432 (13.5%)
F	12,831 (28.5%)	3,409 (10.5%)	16,240 (21.0%)
G	1,430 (3.2%)	1,457 (4.5%)	2,887 (3.7%)
H	13 (0.0%)	4,484 (13.8%)	4,497 (5.8%)
Entry time			
07/2021 - 09/2021	38,335 (85.2%)	24,509 (75.7%)	62,844 (81.2%)
10/2021 - 11/2021	6,672 (14.8%)	7,876 (24.3%)	14,548 (18.8%)
Obesity			
0	28,029 (62.3%)	22,479 (69.4%)	50,508 (65.3%)
1	16,978 (37.7%)	9,906 (30.6%)	26,884 (34.7%)
PMCA			
0	25,634 (57.0%)	19,916 (61.5%)	45,550 (58.9%)

1	10,915 (24.3%)	6,417 (19.8%)	17,332 (22.4%)
2	8,458 (18.8%)	6,052 (18.7%)	14,510 (18.7%)
Negative tests prior entry			
0	1,330 (3.0%)	2,888 (8.9%)	4,218 (5.5%)
1	34,272 (76.1%)	16,299 (50.3%)	50,571 (65.3%)
2	7,388 (16.4%)	9,739 (30.1%)	17,127 (22.1%)
>=3	2,017 (4.5%)	3,459 (10.7%)	5,476 (7.1%)

Table 2B. Baseline characteristics of children 5 to 11 and adolescents 12 to 20 years of age in the study of the effectiveness of the BNT162b2 vaccine in preventing infection and severe diseases with SARS-CoV-2 during periods when Omicron variant was prevalent.

	Omicron study in children			Omicron study in adolescents		
	Vaccinated (N=50,398)	Unvaccinated (N=61,141)	Overall (N=111,539)	Vaccinated (N=21,180)	Unvaccinated (N=34,900)	Overall (N=56,080)
Age						
Median [Q1, Q3]	8 [6, 10]	7 [6, 9]	8 [6, 10]	14 [13, 16]	15 [13, 17]	15 [13, 17]
Distribution						
5	8,165 (16.2%)	13,321 (21.8%)	21,486 (19.3%)			
6	7,447 (14.8%)	11,314 (18.5%)	18,761 (16.8%)			
7	7,090 (14.1%)	9,151 (15.0%)	16,241 (14.6%)			
8	7,028 (13.9%)	7,922 (13.0%)	14,950 (13.4%)			
9	6,773 (13.4%)	7,085 (11.6%)	13,858 (12.4%)			
10	7,011 (13.9%)	6,434 (10.5%)	13,445 (12.1%)			
11	6,884 (13.7%)	5,914 (9.7%)	12,798 (11.5%)			
12				4754 (22.4%)	5760 (16.5%)	10514 (18.7%)
13				3421 (16.2%)	5520 (15.8%)	8941 (15.9%)
14				3338 (15.8%)	5299 (15.2%)	8637 (15.4%)
15				3123 (14.7%)	5315 (15.2%)	8438 (15.0%)
16				2634 (12.4%)	4944 (14.2%)	7578 (13.5%)
17				2201 (10.4%)	3836 (11.0%)	6037 (10.8%)
18				986 (4.7%)	2114 (6.1%)	3100 (5.5%)
19				475 (2.2%)	1400 (4.0%)	1875 (3.3%)
20				248 (1.2%)	712 (2.0%)	960 (1.7%)
Gender						
Female	23,962 (47.5%)	28,669 (46.9%)	52,631 (47.2%)	11,402 (53.8%)	17,954 (51.4%)	29,356 (52.3%)
Male	26,436 (52.5%)	32,468 (53.1%)	58,904 (52.8%)	9,775 (46.2%)	16,939 (48.5%)	26,714 (47.6%)
Ethnicity						
White	14,399 (28.6%)	24,644 (40.3%)	39,043 (35.0%)	16,240 (46.5%)	6,836 (32.3%)	23,076 (41.1%)
Black/AA	13,711 (27.2%)	13,733 (22.5%)	27,444 (24.6%)	6,154 (17.6%)	6,157 (29.1%)	12,311 (22.0%)
Hispanic	12,119 (24.0%)	12,781 (20.9%)	24,900 (22.3%)	6,287 (18.0%)	3,784 (17.9%)	10,071 (18.0%)
Other/Unknown	10,169 (20.2%)	9,983 (16.3%)	20,152 (18.1%)	6,219 (17.8%)	4,403 (20.8%)	10,622 (18.9%)
Hospital						
A	5,019 (10.0%)	9,266 (15.2%)	14,285 (12.8%)	2,131 (10.1%)	5,183 (14.9%)	7,314 (13.0%)
B	15,229 (30.2%)	13,168 (21.5%)	28,397 (25.5%)	6,397 (30.2%)	6,556 (18.8%)	12,953 (23.1%)
C	5,482 (10.9%)	7,409 (12.1%)	12,891 (11.6%)	1,719 (8.1%)	4,075 (11.7%)	5,794 (10.3%)
D	4,766 (9.5%)	2,878 (4.7%)	7,644 (6.9%)	678 (3.2%)	1,337 (3.8%)	2,015 (3.6%)
E	5,843 (11.6%)	10,551 (17.3%)	16,394 (14.7%)	2,047 (9.7%)	4,263 (12.2%)	6,310 (11.3%)
F	9,786 (19.4%)	11,348 (18.6%)	21,134 (18.9%)	3,563 (16.8%)	5,186 (14.9%)	8,749 (15.6%)

G	1,250 (2.5%)	3,239 (5.3%)	4,489 (4.0%)	622 (2.9%)	2,353 (6.7%)	2,975 (5.3%)
H	3,023 (6.0%)	3,282 (5.4%)	6,305 (5.7%)	4,023 (19.0%)	5,947 (17.0%)	9,970 (17.8%)
Entry time						
01/2022 - 03/2022	37,970 (75.3%)	32,523 (53.2%)	70,493 (63.2%)	14,684 (69.3%)	1,9032 (54.5%)	33,716 (60.1%)
04/2022 - 06/2022	5,882 (11.7%)	11,919 (19.5%)	17,801 (16.0%)	3,344 (15.8%)	7,087 (20.3%)	10,431 (18.6%)
07/2022 - 09/2022	4,994 (9.9%)	10,329 (16.9%)	15,323 (13.7%)	2,206 (10.4%)	5,479 (15.7%)	7,685 (13.7%)
10/2022 - 11/2022	1,552 (3.1%)	6,370 (10.4%)	7,922 (7.1%)	946 (4.5%)	3,302 (9.5%)	4,248 (7.6%)
Obesity						
0	33,381 (66.2%)	42,165 (69.0%)	75,546 (67.7%)	13,832 (65.3%)	23,895 (68.5%)	37,727 (67.3%)
1	17,017 (33.8%)	18,976 (31.0%)	35,993 (32.3%)	7,348 (34.7%)	11,005 (31.5%)	18,353 (32.7%)
PMCA						
0	33,870 (67.2%)	40,976 (67.0%)	74,846 (67.1%)	13,482 (63.7%)	21,079 (60.4%)	34,561 (61.6%)
1	10,000 (19.8%)	11,189 (18.3%)	21,189 (19.0%)	4,382 (20.7%)	6,764 (19.4%)	11,146 (19.9%)
2	6,528 (13.0%)	8,976 (14.7%)	15,504 (13.9%)	3,316 (15.7%)	7,057 (20.2%)	10,373 (18.5%)
Negative tests prior entry						
0	2,337 (4.6%)	5,640 (9.2%)	7,977 (7.2%)	768 (3.6%)	2,966 (8.5%)	3,734 (6.7%)
1	34,077 (67.6%)	28,417 (46.5%)	62,494 (56.0%)	15,707 (74.2%)	17,303 (49.6%)	33,010 (58.9%)
2	10,514 (20.9%)	19,816 (32.4%)	30,330 (27.2%)	3,654 (17.3%)	11,012 (31.6%)	14,666 (26.2%)
>=3	3,470 (6.9%)	7,268 (11.9%)	10,738 (9.6%)	1,051 (5.0%)	3,619 (10.4%)	4,670 (8.3%)

Table 3A. Estimated effectiveness of the BNT162b2 vaccine in preventing infection and severe diseases with SARS-CoV-2 in children and adolescents.

	Vaccinated	Unvaccinated	Overall	Vaccine effectiveness (in %) and 95% CI
Delta study in adolescents				
Follow-up				
Total follow-up — no. of person-wk	644,162	398,906	1,043,068	
Median [Q1, Q3]	16 [12, 18]	13 [9, 17]	15 [10, 18]	
Incidence rate per 10,000 person-wk				
Documented infection	2.47	42.72	17.86	98.4 (98.1, 98.7)
Mild COVID-19	0.43	11.58	4.70	99.0 (98.5, 99.3)
Moderate or severe COVID-19	0.20	3.94	1.63	98.7 (97.4, 99.3)
ICU admission with COVID-19	0.02	0.43	0.17	99.0 (92.5, 99.9)
Myocarditis	0.26	0.75	0.45	73.8 (43.8, 87.6)*
Pericarditis	0.19	0.43	0.28	66.1 (14.8, 86.5)*
Age 12-15 yr				
Total follow-up — no. of person-wk	458,981	229,083	688,064	
Documented infection	2.37	44.87	16.52	99.0 (98.6, 99.3)
Age 16-21 yr				
Total follow-up — no. of person-wk	185,181	169,823	355,003	
Documented infection	2.70	39.81	20.45	97.0 (95.9, 97.8)
Omicron study in children				
Follow-up				
Total follow-up — no. of person-wk	1,925,686	1,911,599	3,837,285	
Median [Q1, Q3]	44 [35, 46]	36 [19, 44]	40 [25, 45]	
Incidence rate per 10,000 person-wk				
Documented infection	4.95	17.46	11.18	74.3 (72.2, 76.2)
Mild COVID-19	1.41	4.96	3.18	73.5 (69.2, 77.1)
Moderate or severe COVID-19	0.49	2.13	1.31	75.5 (69.0, 81.0)
ICU admission with COVID-19	0.04	0.26	0.15	84.9 (64.8, 93.5)
Myocarditis	0.02	0.07	0.04	71.4 (8.8, 91.0)*
Pericarditis	0.02	0.05	0.03	69.1 (-19.7, 92.0)*
Age 5-8 yr				
Total follow-up — no. of person-wk	1,101,418	1,254,819	2,356,236	
Documented infection	5.08	15.99	10.89	71.3 (68.2, 74.1)
Age 9-11 yr				
Total follow-up — no. of person-wk	824,268	656,780	1,481,049	
Documented infection	4.79	20.28	11.66	77.9 (75.1, 80.4)

Omicron study in adolescents				
Follow-up				
Total follow-up — no. of person-wk	772,176	1,113,561	1,885,736	
Median [Q1, Q3]	42 [30, 45]	37 [22, 44]	39 [25, 45]	
Incidence rate per 10,000 person-wk				
Documented infection	4.99	25.59	17.16	85.5 (83.8, 87.1)
Mild COVID-19	1.17	6.48	4.31	87.0 (83.5, 89.8)
Moderate or severe COVID-19	0.40	2.76	1.79	84.8 (77.3, 89.9)
ICU admission with COVID-19	0.04	0.43	0.27	91.5 (69.5, 97.6)
Myocarditis	0.05	0.25	0.17	82.0 (46.6, 94.0)*
Pericarditis	0.04	0.21	0.14	83.7 (43.3, 95.3)*
Age 12-15 yr				
Total follow-up — no. of person-wk	524,053	654,315	1,178,368	
Documented infection	5.15	25.57	16.49	85.8 (83.6, 87.7)
Age 16-21 yr				
Total follow-up — no. of person-wk	248,123	459,246	707,368	
Documented infection	4.63	25.63	18.26	85.9 (82.7, 88.5)

: vaccine effectiveness was defined as 100(1- relative risk) which represents the relative risk reduction in percentage in the vaccinated groups compared to the unvaccinated.

Table 3B. Estimated single-dose and two-dose vaccine effectiveness in preventing infection and severe diseases with SARS-CoV-2 in children and adolescents

Outcomes	Single-dose Vaccine effectiveness (95 % CI)	Two-dose Vaccine effectiveness (95 % CI)
Delta study in adolescents		
Documented infection	91.2 (88.0, 93.6)	98.6 (98.2, 98.9)
Mild COVID-19	94.7 (89.0, 97.4)	99.0 (98.4, 99.4)
Moderate or severe COVID-19	86.8 (68.7, 94.4)	99.0 (97.6, 99.6)
ICU admission with COVID-19	99.0 (-, -)	98.5 (85.3, 99.8)
Omicron study in children		
Documented infection	73.8 (69.4, 77.5)	72.3 (69.8, 74.6)
Mild COVID-19	67.2 (57.3, 74.8)	73.3 (68.4, 77.5)
Moderate or severe COVID-19	75.9 (60.9, 85.1)	72.7 (64.2, 79.2)
ICU admission with COVID-19	54.4 (0.0, 84.1)	90.7 (70.7, 97.0)
Omicron study in adolescents		
Documented infection	83.4 (80.7, 85.6)	85.4 (82.5, 87.9)
Mild COVID-19	84.8 (79.3, 88.8)	88.2 (82.7, 92.0)
Moderate or severe COVID-19	86.4 (75.5, 92.4)	81.5 (68.1, 89.3)
ICU admission with COVID-19	89.2 (52.4, 97.5)	89.2 (16.4, 98.6)

Figure 2. Stratified effectiveness of the BNT162b2 vaccine in preventing infection with SARS-CoV-2 in children and adolescents by 2-month intervals since receipt of the first dose

References

1. Flaxman S, Whittaker C, Semenova E, et al. Assessment of COVID-19 as the Underlying Cause of Death Among Children and Young People Aged 0 to 19 Years in the US. JAMA Netw Open 2023;6(1):e2253590.

2. Frenck RW, Klein NP, Kitchin N, et al. Safety, Immunogenicity, and Efficacy of the BNT162b2 Covid-19 Vaccine in Adolescents. New England Journal of Medicine 2021;385(3):239–50.

3. Walter EB, Talaat KR, Sabharwal C, et al. Evaluation of the BNT162b2 Covid-19 Vaccine in Children 5 to 11 Years of Age. New England Journal of Medicine 2022;386(1):35–46.

4. Chemaitelly H, AlMukdad S, Ayoub HH, et al. Covid-19 Vaccine Protection among Children and Adolescents in Qatar. New England Journal of Medicine 2022;387(20).

5. Cohen-Stavi CJ, Magen O, Barda N, et al. BNT162b2 Vaccine Effectiveness against Omicron in Children 5 to 11 Years of Age. New England Journal of Medicine 2022;387(3):227–36.

6. Florentino PTV, Alves FJO, Cerqueira-Silva T, et al. Vaccine effectiveness of CoronaVac against COVID-19 among children in Brazil during the Omicron period. Nat Commun 2022;13(1).

7. Olson SM, Newhams MM, Halasa NB, et al. Effectiveness of BNT162b2 Vaccine against Critical Covid-19 in Adolescents. New England Journal of Medicine 2022;386(8):713–23.

8. Sacco C, Del Manso M, Mateo-Urdiales A, et al. Effectiveness of BNT162b2 vaccine against SARS-CoV-2 infection and severe COVID-19 in children aged 5–11 years in Italy: a retrospective analysis of January–April, 2022. The Lancet 2022;400(10346):97–103.

9. Khan FL, Nguyen JL, Singh TG, et al. Estimated BNT162b2 Vaccine Effectiveness Against Infection With Delta and Omicron Variants Among US Children 5 to 11 Years of Age. JAMA Netw Open 2022;5(12):e2246915.

10. Home - PEDSnet [Internet]. [cited 2023 May 21];Available from: https://pedsnet.org/

11. Data_Models_Public/PEDSnet at master · PEDSnet/Data_Models_Public · GitHub [Internet]. [cited 2023 May 21];Available from: https://github.com/PEDSnet/Data_Models_Public/tree/master/PEDSnet

12. Hernán MA, Robins JM. Using Big Data to Emulate a Target Trial When a Randomized Trial Is Not Available. Am J Epidemiol [Internet] 2016 [cited 2023 May 21];183(8):758–64. Available from: https://pubmed.ncbi.nlm.nih.gov/26994063/

13. Suchard MA, Schuemie MJ, Krumholz HM, et al. Comprehensive comparative effectiveness and safety of first-line antihypertensive drug classes: a systematic, multinational, large-scale analysis. The Lancet 2019;394(10211):1816–26.

14. Simon TD, (COE4CCN) for the C of E on Q of CM for C with CN, Cawthon ML, et al. Development and Validation of the Pediatric Medical Complexity Algorithm (PMCA) Version 2.0. Hosp Pediatr [Internet] 2017 [cited 2023 Jun 7];7(7):373–7. Available from: /hospitalpediatrics/article/7/7/373/26493/Development-and-Validation-of-the-Pediatric

15. Forrest CB, Burrows EK, Mejias A, et al. Severity of Acute COVID-19 in Children <18 Years Old March 2020 to December 2021. Pediatrics 2022;149(4).

16. Johnson LT, Geyer CJ. Variable transformation to obtain geometric ergodicity in the random-walk Metropolis algorithm. The Annals of Statistics 2012;40(6).

17.   J. Schuemie M, Soledad Cepede M, A. Suchard M, et al. How Confident Are We About Observational Findings in Health Care: A Benchmark Study. Harv Data Sci Rev 2020;

18.   Block JP, Boehmer TK, Forrest CB, et al. Cardiac Complications After SARS-CoV-2 Infection and mRNA COVID-19 Vaccination — PCORnet, United States, January 2021–January 2022. MMWR Morb Mortal Wkly Rep [Internet] 2022 [cited 2023 Jun 8];71(14):517–23. Available from: https://www.cdc.gov/mmwr/volumes/71/wr/mm7114e1.htm