Assessment of Ultra Processed Foods consumption in Senegal: Validation of the Nova-UPF screener

Saliou Diombo Kébé[1]*, Adama Diouf[1], Papa Mamadou Dit Doudou Sylla[1,4], Kalidou Kane[1], Caroline dos Santos Costa[2], Fernanda Helena Marrocos Leite[2], Giovanna Calixto Andrade[2], Abdou Badiane[1], Jean-Claude Moubarac[3], Nicole Idohou-Dossou[1] and Carlos Augusto Monteiro[2]

[1] Laboratoire de Recherche en Nutrition et Alimentation Humaine (LARNAH), Faculté des Sciences et Techniques, Université Cheikh Anta Diop. BP 5005, Dakar – Fann, Sénégal; labonut@ucad.edu.sn

[2] Center for Epidemiological Research in Nutrition and Health, School of Public Health, University of Sao Paulo, Brazil nupens.usp@gmail.com

[3] Département de Nutrition, Faculté de médecine, Université de Montréal. Montréal, Canada; info-nutrition@umontreal.ca

[4] Laboratoire des sciences Biologiques, Agronomiques, Alimentaires et de Modélisation des Systèmes complexes (LABAAM), UFR des Sciences Agronomiques, de l'Aquaculture et des Technologies Alimentaires, Université Gaston Berger de Saint-Louis. BP 234 Saint-Louis, Sénégal, s2ata@ugb.edu.sn

* Correspondence: salioudiombo.kebe@ucad.edu.sn ;

Abstract: Ultra-processed foods (UPF), as proposed by the Nova food classification system, are linked to the development of several non-communicable chronic diseases and deaths from all causes. The Nova-UPF screener developed in Brazil is a simple and quick tool to assess and monitor the consumption of these food products. The aim of this study was to adapt and validate, against the 24-hour dietary recall, this short food-based screener to assess UPF consumption in the Senegalese context. The tool adaptation was undertaken using Delphi methodology with national experts and data from a food market survey. The validation study was conducted in the urban area of Dakar in a sample of 301 adults, using the dietary share of UPF on the day prior to the survey, expressed as a percentage of total energy intake obtained via 24-hour recall. Association between the Nova score and the dietary share of UPF was evaluated using linear regression models. The Pabak index was used to assess the agreement in participants' classification according to fifths of Nova score and fifths of dietary share of UPF. Results show a direct association (p-value<0.001) between intervals of the Nova-UPF score and the average dietary share of UPF. There was an almost perfect agreement in the distribution of individuals according to score's quintiles and UPF dietary share quintiles (Pabak index = 0.84). The study concluded that the score provided by the Nova-UPF screener adapted to the Senegalese context is a valid assessment of UPF consumption.

Keywords: Validation study; Questionnaire; Food consumption; Ultra-processed Foods; Senegal; Non-communicable disease; Nova classification, Diet Surveys, food classification systems

1. Introduction

The Nova food classification system divides all foods according to the nature, extent and purpose of the industrial processes they undergo. These are physical, biological and chemical techniques used after food has been extracted from nature and before it is consumed or made into dishes and meals [1]. Nova classifies all foods into four groups including Ultra-Processed Foods (UPF). They are defined as industrial formulations made from food derived substances with little or no whole foods to which flavours, colours and other cosmetic additives are added. Their poor nutritional profile and level of processing make them nutritionally unbalanced, hyperpalatable and addictive [2]. A meta-analysis involving many representative studies and several other epidemiological studies have showed the negative effect of high UPF consumption on the nutritional quality of diets [3] and the effective link between UPF consumption and the development of chronic non-communicable diseases (NCDs) such as type II diabetes, obesity, hypertension, cardiovascular diseases and all-cause mortality [4–8].

The consumption of these products is rapidly increasing in low- and middle-income countries (LMICs) and this evolution, strongly linked to the industrialization of food systems and economic development, represents today a major threat to public health [9–11]. Such as for Senegal, which is facing a rise in the prevalence of overweight and obesity due to lifestyle changes, including sedentarism and physical inactivity, and unhealthy eating habits. Prevalence of overweight and obesity in 2015 was 22.1% and 6.4%, respectively, among people aged 18 to 69 years. The prevalence of hypertension at national level was 29.8%, and 19.2% and 2.1% of adults had hypercholesterolemia and type II diabetes, respectively [12].

Senegalese public health system regularly conducts surveys to assess the state of malnutrition in all its forms, the dietary diversity and micronutrient intakes, but none specifically addresses UPF consumption. It is necessary to study the increasing consumption of UPF and their effects on health, but it is difficult for researchers and policy makers to monitor consumption trends of UPF. Assessing the dietary contribution of UPF using quantitative data, based on 24-hour recall or semi-quantitative food frequency data, is expensive, time consuming and complex to analyse. There is also a considerable lack of data on new dietary patterns and sales and/or consumption trends of ultra-processed foods.

The Nova-UPF screener developed and validated in Brazil is a short food-based questionnaire that specifically addresses the consumption of UPF. It asks about UPF consumption ("yes" or "no") in the previous day and assigns a score ranging from 0 to 23 according to a list of 23 predefined UPF subcategories. It is a simple and quick tool to administer, requires a low workload, is easy to use and its application will allow the monitoring and evaluation of UPF

consumption and possibly their impact on the development of diet-related NCDs [13–14]. The

aim of this study was to describe the adaptation of the Nova-UPF screener to the Senegalese

context and evaluate the performance of the score obtained by applying the adapted tool in

comparison with the daily energy contribution of UPF obtained from 24-hour dietary recalls.

2. Materials and Methods

2.1. Ethics

This study received approval (Protocol SEN21/17 - 00000079MSAS/CNERS/SP) from the

National Committee of Ethics for Research in Health (CNERS) of the Ministry of Health and

Social Action (MSAS). Free and informed consent was obtained from the participants before

starting the study.

2.2. Adaptation of the Nova-UPF Screener

The original version of the Nova-UPF screener was developed in Brazil and validated against

a full 24-hour recall [14]. To be applicable in our context, the original content (**table 1**) was

adapted and tested to reflect Senegalese dietary patterns by the Laboratoire de Recherche en

Nutrition et Alimentation Humaine (LARNAH) team and its partners, including researchers

who worked on the development of the original version (*Núcleo de Pesquisas Epidemiológicas*

em Nutrição e Saúde, NUPENS). The changes made to the original version of the Nova-UPF

screener were carried out through a collaborative process, led by the LARNAH research team

with key stakeholders in the domains of nutrition, epidemiology, public health and food

technology at national level. First, a literature review was performed on limited existing data,

especially on Euromonitor reports of countries in the same economic zone and with a similar

food environment [15–16], to carry out a first content adjustment. Then, two virtual workshops

and individual consultation sessions involving the aforementioned stakeholders were

organised. During the workshops, the DELPHI method, which aims to gather expert opinions

on a specific subject and to highlight convergences and consensus on the orientations to be

given to a project by subjecting these experts to successive waves of questions, was applied

[17].

Table 1: Original version of the Nova-UPF screener

N°1	Subcategories
1	Regular or Diet soda
2	Fruit juice or fruit drink in can or box
3	Fruit drink prepared from a powdered mix
4	Chocolate milk in can or box or prepared from a powdered mix
5	Tea or coffee in can or box or prepared from a powdered mix
6	Any type of flavored yogurt
7	Sausage, hamburger or nuggets
8	Ham, salami or bologna
9	Bun, roll or any type of packaged bread
10	Margarine
11	Branded mayonnaise, ketchup, or mustard
12	Branded salad dressing
13	Frozen French fries or from fast-food restaurants
14	Frozen pizza or from fast-food restaurants
15	Instant noodles or instant powdered soup
16	Frozen lasagna or any other frozen ready meal
17	Potato chips, crackers or any other type of packaged salty snacks
18	Cookies or biscuits with or without filling
19	Branded cake or muffin (not homemade or artisanal)
20	Cereal bar
21	Branded ice cream or ice lolly (not homemade or artisanal)
22	Chocolate bar or chocolate candies
23	Sugary breakfast cereals

Source: Caroline et al. (2021).

The first virtual workshop conducted in June 2021 with over 20 national stakeholders was an opportunity to present the Nova-UPF screener to the audience and give the main guidelines for adapting the original content of the Nova-UPF screener. These included finding resources to fill the data gap on UPF consumption in the country and to define a methodological approach to revise the proposed subcategories. In that order, we implemented a market survey to characterise the food supply in Dakar (capital of Senegal). The Dakar region concentrates most of the economic activity, flow of food products to the rest of the country and all major distribution chains [18]. More than 4,700 types of packaged food products, of which more than 70% was ultra-processed, were listed and classified into different subcategories.

Following the first virtual workshop, individual one-hour consultations were held with experts to directly obtain their opinions and suggestions on the content to be adapted. The results of these interviews and data from the market survey were used to develop a first adapted content proposal. The second workshop carried out in September 2021 XX with same stakeholders reviewed the suitability of selected UPF subcategories and examples of products included in

each subcategory to ensure that they were relevant to the Senegalese context. Following this second consultation with national experts, the final content of the Nova-UPF screener for Senegal was obtained.

Before implementing the tool, an apparent validity or "face-validity" was carried out. This was a pre-test to assess the tool's design, content, structure, and ease of understanding. The Nova-UPF screener in its original design is a self-administered questionnaire where the respondent reads the names of the subcategories and checks the answers. However, given the low level of literacy and linguistic diversity, the Nova-UPF screener for Senegal was administered by an interviewer who was responsible for explaining to the respondent, if necessary, the meaning of each UPF subcategory. In this phase, an electronic questionnaire was designed to allow interviewers to personally assess, after each question (subcategory): i) whether the wording of the requested subcategory was well understood and ii) whether the respondent had no difficulty in answering. Then, participants were asked: i) whether he/she had understood the meaning of each subcategory, ii) whether the proposed UPF subcategories corresponded to the food products to which they were often exposed, iii) whether the examples used were clear and well recognised. Based on these results, further minor adjustments to the phrasing of the questions were made. From the original version, several changes were made. These included rewording of subcategory names, the splitting or grouping of certain subcategories and the introduction of new subcategories. In **Figure 1**, we have a summary of the changes made from the original version. Like the original tool, the Nova-UPF screener for Senegal also presents 23 subcategories where foods are grouped within the same subcategory according to their composition or use. At the end of the adaptation phase and content validation (face-validity), a final version adapted for Senegal was obtained (see **Table 2**).

Original		Adapted
Chocolate milk in can or box or prepared from a powdered mix	→	Instant milk powder or instant chocolate powder
Tea or coffee in can or box or prepared from a powdered	→	Iced or flavoured tea, coffee milk or flavoured milk in a can or bottle or prepared from a powdered mixture
Any type of flavored yogurt	→	Any type of flavoured yoghurt or industrial curdled milk
Sausage, hamburger or nuggets Ham, salami or bologna	→	Processed meats or meat spreads or nuggets
Branded salad dressing	→	Bouillons, Dipping Sauce, vinaigrette or industrial salad dressing
Frozen French fries or from fast-food restaurants Frozen pizza or from fast-food restaurants	→	Frozen fries or frozen pizza
Frozen lasagna or any other frozen ready meal	→	Any frozen ready-to-eat meal
Cereal bar Chocolate bar or chocolate candies	→	Chocolate bars, chocolate candies, confectionery or chewy products

Three (3) new categories

Industrial jams or marmalades   Chocolate or hazelnut paste spreads   Spreadable cheese

137   Figure 1: Summary of changes to the original content

140   Table 2: Final adapted version

N°	Subcategories	Code
1	Soda, energizing or diet beverages	CAT 01
2	Flavoured drinks, concentrates and/or fruit nectars in a can or box	CAT 02
3	Fruit-flavoured drinks prepared from a powdered mixture	CAT 03
4	Instant milk powder or instant chocolate powder	CAT 04
5	Iced or flavoured tea, coffee milk or flavoured milk in a can or bottle or prepared from a powdered mixture	CAT 05
6	Any type of flavoured yoghurt or industrial curdled milk	CAT 06
7	Processed meats or meat spreads or nuggets	CAT 07
8	Bread, rusks, any other type of packaged industrial bread	CAT 08
9	Frozen fries or frozen pizza	CAT 09
10	Margarine	CAT 10
11	Industrial jams or marmalades	CAT 11
12	Chocolate or hazelnut paste spreads	CAT 12
13	Spreadable cheese	CAT 13
14	Industrial mayonnaise, ketchup or mustard	CAT 14
15	Bouillons, Dipping Sauce, vinaigrette or industrial salad dressing	CAT 15
16	Instant soup powder or instant noodles	CAT 16
17	Any frozen ready-to-eat meal	CAT 17
18	Chips, crackers, or any other type of packaged salty snack	CAT 18
19	Cookies or biscuits with or without fillings	CAT 19
20	Industrial cakes, muffins, or pastries	CAT 20
21	Sweetened breakfast cereals	CAT 21
22	Industrial ice cream or popsicles	CAT 22
23	Chocolate bars, chocolate candies, confectionery or chewy products	CAT 23

2.3. Validation of the Nova-UPF Screener

2.3.1. Design and selection of participants

The validation study was conducted in urban areas of Dakar, which concentrates nearly a

quarter of the national population and almost half of the country's urban population (49.6%)

[19]. For validation studies, higher numbers of subjects will provide better estimates of

reproducibility or validity and a sample size of at least 50 to100 subjects is recommended [20].

Based on this, a sample size of 300 adults aged 18 years and older was retained.

A two-stage random sampling design was used. First, a random selection of 25 urban census

districts (CD), spread across nine (9) of health districts in Dakar medical region. Second, 12

households were randomly selected in each CD (13 in one of them), by using a sampling step

defined according to the size of the CD. Finally, in each household, one adult was randomly

selected, taking into account the 50/50 balance between men and women. Overall, 301 adult

subjects were enrolled to participate in this study.

2.3.2. Data collection

2.3.2. Data collection

The data collection was carried out on December 2021, by experienced nutritionists and

investigators who were trained in the use of the Nova-UPF screener and quantitative 24h recall

by the research team of the LARNAH. All participants were informed about the purpose of the

study and were invited to join after agreeing to participate by signing the consent form. Then

sociodemographic and food consumption data (Nova-UPF screener and 24-hour recall) were

collected. The electronic questionnaire was deployed in tablets using the ODK application for

data collection, where the 23 subcategories were listed along with a descriptive text of the

category and illustrative images as shown in **Figure 2**.

Figure 2: Electronic version of the Nova-UPF screener for Senegal.

The Nova-UPF screener was administered first, for an average time of 05 minutes. After

completion of the tool, investigators conducted the 24-hour dietary recall, where participants

informed about all foods and the amount consumed on the day before.

The 24-hour dietary recall was conducted using the automated multiple-pass method [21]. First,

participants report, in a rapid and uninterrupted manner, all foods and beverages consumed.

Next, the interviewer asks, from an initial list of often forgotten foods, what other foods or

beverages the respondent might have omitted to report. The participant is then asked about the

type, time and place of each meal, and then provides details such as the method of preparation,

quantities and the addition of other foods (e.g., sugar, salt…). Finally, the interviewer lists all

the foods previously reported by the respondent, reviewing any omissions. Given the local food

habits, where people are used to eat the main meals together around the same bowl, the quantity
of food or meals consumed was estimated by direct weighing (if a replica was available in the
household), by non-standard measurements (e.g., slice, unit, tablespoon, bag, handful, ...) or by
using substitutes (water, dry millet, modelling paste) to get the weight that refer to the volume
occupied by the amount of food consumed.
181

2.4. Data analysis

Each participant's Nova-UPF score corresponds to the total number of subcategories that the
participant reports having consumed among the 23 listed in the tool. To estimate the dietary
share of UPF, each food reported in the 24-hour recall was first classified according to the
Nova classification system as 1) unprocessed or minimally processed, 2) processed culinary
ingredients, 3) processed foods or 4) ultra-processed foods [1–2]. Then, the real amount
consumed of each food was obtained using the database of food conversion factors and the
Senegalese standard recipe database, developed under the national food consumption survey
implemented by Consortium pour la Recherche Économique et Sociale (CRES) in
collaboration with LARNAH, the Ministry of Health and FAO-Senegal. These conversion
factors are coefficients that allow us to move from non-standard measurements or substitute
quantities to real quantities of food. The amount of each food (in grams) was converted to
calories using the 2019 West African Food Composition Table [22] and the CIQUAL 2020
Nutritional Food Composition Table [23]. Finally, for each person, we calculated the total
calories consumed in the day, the total calories from UPF and also the percentage of calories
from UPF.

To test the association between the score provided by the Nova-UPF screener and the dietary
share of UPF obtained from the 24-h recall, the evolution of percentage of calories from UPF
was examined according to the evolution of the score on a continuous basis and also by
comparing the quintiles of distribution of the two variables. In both cases, linear regression
models were used to estimate the association. The level of agreement between the quintiles for
the score and the percentage of calories from UPF was assessed by calculating the prevalence
and bias-adjusted Kappa index (Pabak) [24]. Values greater than 0.80 indicate near perfect
agreement; between 0.61 and 0.80, substantial agreement; between 0.41 and 0.60, moderate;
between 0.21 and 0.40, fair; and equal to or less than 0.20, slight [25]. Data analyses were
performed with the Stata® 16.1 software and the Pabak index calculated using R Studio
software.

3. Results

3.1. Socio-demographic characteristics

Sociodemographic characteristics of participants are described in **Table 3**. The mean age of subjects was 42 ± 14 years. All 301 volunteers in this study were aged between 20 and 68 years, and the most represented age group was those of 20-39 years. More than half of the individuals were married (61.5%) and had reached high school or university (52.8%). Regarding their occupation, 22.9% were in trade sector and 11% were unemployed, including 18.7% of women.

Table 3: Socio-demographic characteristics (n=301). Senegal (2021).

	Total (n= 301)	Men (n= 151)	Women (n= 150)
Mean age	42 ± 14	41 ± 14	43 ± 15
Age group			
20 – 39	49.5 (149)	55 (83)	44 (66)
40 – 59	34.2 (103)	31.1 (47)	37.3 (56)
60 – 68	16.3 (49)	13.9 (21)	18.7 (28)
Literacy	77.1 (232)	82.8 (125)	71.3 (107)
Level of Education			
Primary school	21.6 (65)	13.9 (21)	29.3 (44)
College / high school	24.2 (73)	26.5 (40)	22 (33)
University	28.6 (86)	25.8 (39)	31.4 (47)
Vocational school	16.3 (49)	21.2 (32)	11.3 (17)
Others	9.3 (28)	12.6 (19)	6 (9)
Marital status			
Single	23.6 (71)	31.1 (47)	16 (24)
Divorced	4.6 (14)	3.3 (5)	6 (9)
Married monogamous	51.5 (155)	55 (83)	48 (72)
Married polygamous	14 (42)	10.6 (16)	17.3 (26)
Widow(er)	6.3 (19)	0 (0)	12.7 (19)
Profession			
No occupation	11 (33)	3.3 (5)	18.7 (28)
Student	9.6 (29)	8.6 (13)	10.7 (16)
Manual worker	7 (21)	13.3 (20)	0.7 (1)
Trader	22.9 (69)	13.3 (20)	32.7 (49)
Housekeeper	11.6 (35)	2 (3)	21.3 (32)

The results are expressed as percentages with the effective (n) or mean $\pm$ SD.

3.2. UPF consumption and distribution of the Nova-UPF score

As described in **Figure 3**, the most consumed UPF subcategories were Bouillons, dipping sauce, vinaigrette or industrial salad dressings (32.5%); Industrial mayonnaise, ketchup or mustard (26.3%); Instant milk powder or instant chocolate powder (14.5%); Margarine (6.6%), and Chocolate bars, chocolate candies, confectionery or chewy products (4.9%). Overall, no products belonging to the subcategories Fruit-flavoured drinks prepared from a powdered mixture; Industrial jams or marmalades; and Industrial ice cream or popsicles were consumed by our study population.

Figure 3: Consumption frequencies of ultra-processed food subcategories.

The Nova scores ranged from zero (0) to eight (8) (**Figure 4**). The majority of participants got scores of 3 (28.6%), 2 (27.2%), 1 (20.3%), and 4 (13.3%). Only one individual (0.3%) had a score of zero and 10.3% of individuals had a score of 5 or higher (≥ 5).

235   Figure 4: Nova-UPF score distribution.

3.3. Dietary share of ultra-processed foods according to Nova score intervals

Overall, the average contribution of UPF to daily energy intake was 17.4% (CI= 18.5 - 21.3).

The results showed that the dietary share of UPF consumption increased linearly with the

increase in the intervals of the Nova-UPF score (p-value for linear trend <0001) (**Figure 5**).

Individuals in the first quintile had a mean energy contribution from UPF lower than 3%, while

the mean energy contribution from UPF for those in the fifth quintile was more than 30%.

Figure 5: Mean dietary share of ultra-processed foods obtained from the full 24-h recall

according to Nova-UPF score intervals. Senegal (n=301), 2021

3.4. Agreement between the Nova-UPF Screener and the 24-hour dietary recall

Table 4 shows the distribution of individuals according to quintiles of Nova-UPF score and

quintiles of energy contribution from ultra-processed foods. The Pabak index value was 0.84

(0.68 - 1). This indicates an almost perfect agreement (>0.80) between the two criteria, the

Nova-UPF score and the 24-hour dietary recall.

Table 4: Distribution of individuals by score quintiles and UPF energy contribution quintiles.

	1	**2**	**3**	**4**	**5**	**Total**
Q1 (≤ 2.18)	15.28	4.32	0.33	0.33	0.33	20.60
Q2 (2.19 – 10.89)	3.65	11.96	8.97	2.33	0.33	27.24
Q3 (10.90 – 26.80)	1.33	6.64	10.96	4.98	4.65	28.57
Q4 (26.81 – 41.53)	0.33	1.66	5.98	2.99	2.33	13.29
Q5 (≥ 41.54)	0.00	2.66	2.33	2.66	2.66	10.30
Total	20.60	27.24	28.57	13.29	10.30	100.00

Pabak index (Kappa adjusted for prevalence bias) = 0.84

4. Discussion

Results of this study show a positive association (P<0.001) between the score obtained with

the Nova-UPF screener for Senegal and the dietary share of UPF obtained from the full 24-

hour dietary recall. There was near perfect agreement between the score quintiles distribution

of individuals and the UPF energy contribution quintiles.

The average percentage of calories provided by UPF (17.4%) was similar to that reported in

France [26] and in Brazilian adults [27], and to other studies in Colombia [28] and Spain [29].

This reflects that consumption of these products occurs regardless of differences in dietary

habits and economic development between countries. However, UPF consumption in the

Senegalese context is still much lower than the rates reported in the USA [30], Canada [31]

and Great Britain [7], where the availability of these products in the food supply is much higher

[9].

The most consumed UPF category was "Bouillons, dipping sauce, vinaigrette or industrial

salad dressings". This is due to the presence of bouillons in this category, which is widely

consumed by the majority (90%) of the Senegalese households' [32], to enhance the taste of

prepared dishes. The same happens for the category of salty sauces (Industrial mayonnaise,

ketchup or mustard). The frequency of use of foods belonging to the categories "Instant milk

powder or instant chocolate powder" and "Margarine" is mainly due to the habit of people to

consume instant milk powder with coffee in the morning and bread coated with margarine, for
breakfast. The consumption of "Chocolate bars, chocolate candies, confectionery or chewy
products" is mostly due to snacking behaviour between meals, especially among younger
people.

The absence of consumption of 'industrial ice cream or popsicles' could be attributed to the data
collection period, which took place in winter, whereas these products are generally more
consumed during hot weather (summer). Despite their high availability on the market, it
appeared that 'Fruit-flavoured drinks prepared from a powdered mixture' and 'Industrial jams
or marmalades' were not consumed by our study population. This may be due to the fact that
we have not caught profiles of people who consume these products, or because their
consumption is more common in areas other than those covered by our study (e.g., rural or
peri-urban). Applying the Nova tool at national level will give us a clearer assessment on
consumption of all categories, and possibly allow us to adjust the adapted content further on.
The scores obtained and their distribution were very similar to results from the validation tool
in Brazil. Indeed, in both studies, the most common scores were 1, 2, 3, and 4 [14]. However,
the average percentage of energy contribution according to the score quintiles was lower for
Senegal. It varied between 2.8% for the first quintile and 30.4% for the fifth quintile, while the
Brazilian average contribution varied between 19.3% and 43.9%, respectively [14]. When
considering the distribution of individuals according to Nova score quintiles and quintiles of
energy contribution of UPF, the agreement between the Nova-UPF screener for Senegal and
the 24h recall is higher in the first three quintiles while the original Brazilian tool highlights
more the extreme quintiles (first and fifth). These differences could be explained by the
difference in dietary patterns and food culture between Senegal and Brazil, and the sample
characteristics, such as age, sex, and education level, which can influence the consumption of
UPF in quantity and variety.

Beyond the significant association, the concordance expressed through the Pabak index is
better for the adapted version in Senegal compared to the original one [14]. Adaptation from a
predefined original content allows for a more precise and exhaustive adjustment of the tool's
content. This suggests good results when considering that the Nova-UPF screener can be
adapted in other countries, or its content can be updated in the future, according to changes in
dietary habits.

The sample size of this study is unrepresentative, but this has no impact on the results because
this study compares two methods to estimate UPF consumption in the same population. It was
conducted on the basis of probability sampling with a balanced sex-ratio population whose

socio-demographic characteristics (level of education, age, profession, etc.) cover the entire concerned population. This is the first study on food consumption in Senegal using the Nova food classification and gives a fairly good idea of what the situation could be at the national level.

5. Conclusions

The Nova-UPF screener for Senegal is valid for measuring and monitoring the consumption of ultra-processed foods at the populational level and over time. The score obtained with this tool accurately reflects the level of energy intake from these targeted foods. However, it will be necessary to perform this tool on a larger sample to better appreciate its applicability in nutritional surveillance and monitoring system.

Funding: "This work is funded through the Innovative Methods and Metrics for Agriculture and Nutrition Action (IMMANA) programme (Grant No. IMMANA 3.06), led by the London School of Hygiene & Tropical Medicine (LSHTM)"

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki, and approved by "Comité National d'Éthique pour la Recherche en Santé" of the Ministry of Health and Social Action (Protocol SEN21/17 – approved 14 May 2021).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study

Acknowledgments: Thanks to. Fousseni Sama, Kodjovi AMAVI (statisticians), and Ms. Mariam Sy (nutritionist) for their technical support during the implementation of this study. We thank all the administrative and technical staff of LARNAH, the people involved in data collection and the participants for allowing us to carry out this work.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

1. Monteiro, C.A.; Cannon, G.; Moubarac, J-C.; Levy, R.B.; Louzada, M.L.C.; Jaime, P.C. The UN Decade of Nutrition, the NOVA food classification and the trouble with ultra-processing. Public Health Nutr 2018, 21, 5–17.

2. Monteiro, C.A.; Cannon, G.; Levy, R.B.; Moubarac, J-C.; Louzada, M.L.C.; Rauber, F.; Khandpur, N.; Cediel, G.; Neri, D.; Martinez-Steele, E. et al. Ultra-processed foods: what they are and how to identify them. Public Health Nutr 2019, 22, 936–41.

3. Martini, D.; Godos, J.; Bonaccio, M.; Vitaglione, P.; Grosso, G. Ultra-Processed Foods and Nutritional Dietary Profile: A Meta-Analysis of Nationally Representative Samples. Nutrients 2021, 13, 3390.

4. Rauber, F.; Chang, K.; Vamos, EP.; Louzada, M.L.C.; Monteiro, C.A.; Millett, C.; Levy, R.B. Ultra-processed food consumption and risk of obesity: a prospective cohort study of UK Biobank. Eur J Nutr 2021, 60, 2169–80.

5. Llavero-Valero, M.; Escalada-San Martín, J.; Martínez-González, M.A.; Basterra-Gortari, F.J.; de la Fuente-Arrillaga, C.; Bes-Rastrollo, M. Ultra-processed foods and type-2 diabetes risk in the SUN project: A prospective cohort study. Clinical Nutrition 2021, 40, 2817–2824.

6. Machado, P.P.; Steele, E.M.; Levy, R.B.; da Costa Louzada, M.L.; Rangan, A.; Woods, J.; Gill, T.; Scrinis, G.; Monteiro, C.A. Ultra-processed food consumption and obesity in the Australian adult population. Nutr. Diabetes 2020, 10, 39.

7. Pagliai, G.; Dinu, M.; Madarena, M.P.; Bonaccio, M.; Iacoviello, L.; Sofi, F. Consumption of ultra-processed foods and health status: a systematic review and meta-analysis. Br J Nutr 2021, 125, 308–318

8. Suksatan, W.; Moradi, S.; Naeini, F.; Bagheri, R.; Mohammadi, H.; Talebi, S.; Mehrabani, S.; Kermani, M.H.; Suzuki, K. Ultra-Processed Food Consumption and Adult Mortality Risk: A Systematic Review and Dose–Response Meta-Analysis of 207,291 Participants. Nutrients 2021, 14, 174.

9. Vandevijvere, S.; Jaacks, L.M.; Monteiro, C.A.; Moubarac, J-C.; Girling-Butcher, M.; Lee, A.C.; Pan, A.; Bentham, J.; Swinburn, B. Global trends in ultraprocessed food and drink product sales and their association with adult body mass index trajectories. Obesity Reviews 2019, 20, 10–9.

10. Baker, P.; Machado, P.; Santos, T.; Sievert, K.; Backholer, K.; Hadjikakou, M.; Russell, C.; Huse, O.; Bell, C.; Scrinis, G.; et al. Ultra-processed foods and the nutrition transition: Global, regional and national trends, food systems transformations and political economy drivers. Obesity Reviews 2020, 21, e13126.

11. Moodie, R.; Bennett, E.; Kwong, E.J.L.; Santos, T.M.; Pratiwi, L.; Williams, J.; Baker, P. Ultra-Processed Profits: The Political Economy of Countering the Global Spread of Ultra-Processed Foods – A Synthesis Review on the Market and Political Practices of Transnational Food Corporations and Strategic Public Health Responses. Int J Health Policy Manag 2021, 10, 968-82.

12. Agence Nationale de la Statistique et de la Démographie (ANSD). Enquête nationale sur les facteurs de risques des maladies non transmissibles (STEPS). Sénégal : Ministère de la Santé et de l'Action Sociale ; 2015.

13. Sattamini, I.F. Instrumentos de avaliação da qualidade de dietas: desenvolvimento, adaptação e validação no Brasil. Doutorado em Nutrição em Saúde Pública, Universidade de São Paulo, São Paulo, 2019.

14. Costa, C.S.; Faria, F.R.; Gabe, K.T.; Sattamini, I.F.; Khandpur, N.; Leite, F.H.M.; Steele, E.M.; Louzada, M.L.C.; Levy, R.B.; Monteiro, C.A. Nova score for the consumption of ultra-processed foods: description and performance evaluation in Brazil. Rev. saúde pública 2021, 55, 13.

15. Euromonitor International. Packaged Food in Ghana. Passport, 2019.

16. Euromonitor International. Packaged Food in Côte d'Ivoire. Passport, 2019.

17. De Villiers, M.R.; De Villiers, P J.; Kent, AP. The Delphi technique in health sciences education research. Med Teach 2005, 27(7), 639–43.

18. Agence Nationale de la Statistique et de la Démographie (ANSD). Situation économique et sociale régionale 2019. Sénégal : ANSD ; 2021.

19. Agence Nationale de la Statistique et de la Démographie (ANSD). Rapport définitif Recensement Général de la Population et de l'Habitat, de l'Agriculture et de l'Elevage (RGPHAE). Sénégal : ANSD ; 2013.

20. Cade J., Thompson, R., Burley, V., & Warm, D. Development, validation and utilisation of food-frequency questionnaires – a review. Public Health Nutrition 2002, 5(4), 567–587.

21. Moshfegh, A.J.; Rhodes, D.G.; Baer, D.J.; Murayi, T.; Clemens, J.C.; Rumpler, W.V.; Paul, D.R.; Sebastian, R.S.; Kuczynski, K.J.; Ingwersen, L.A. The US Department of Agriculture Automated Multiple-Pass Method reduces bias in the collection of energy intakes. Am. J. Clin. Nutr. 2008, 88, 324–32.

22. Vincent, A.; Grande, F.; Compaoré, E.; Amponsah Annor, G.; Addy, P.A.; Aburime, L.C.;
Ahmed, D.; Bih Loh, A.M.; Dahdouh Cabia, S.; Deflache, N.; Dembélé, F.M.; Dieudonné,
B.; Edwige, O.B.; Ene-Obong, H.N.; Fanou Fogny, N.; Ferreira, M.; Omaghomi Jemide,
J.; Kouebou, P.C.; Muller, C.; Nájera Espinosa, S.; Ouattara, F.; Rittenschober, D.;
Schönfeldt, H.; Stadlmayr, B.; van Deventer, M.; Razikou Yiagnigni, A.; Charrondière,
U.R. FAO/INFOODS Food Composition Table for Western Africa (2019) User Guide &
Condensed Food Composition Table. Rome, FAO ; 2020.

23. CIQUAL Table de composition nutritionnelle des aliments 2020. Available online:
https://ciqual.anses.fr. (accessed on 24 October 2022)

24. Byrt, T.; Bishop, J.; Carlin, J.B. Bias, prevalence and kappa. Journal of Clinical
Epidemiology 1993, 46, 423–9.

25. Landis, J.R.; Koch, G.G. The Measurement of Observer Agreement for Categorical Data.
Biometrics 1977, 33, 159.

26. Srour, B.; Fezeu, L.K.; Kesse-Guyot, E.; Allès, B.; Méjean, C.; Andrianasolo, R.M.;
Chazelas, E.; Deschasaux, M.; Hercberg, S.; Galan, P.; Monteiro, C.A.; Julia, C.; Touvier,
M. Ultra-processed food intake and risk of cardiovascular disease: prospective cohort
study (NutriNet-Santé). BMJ 2019, 365, l1451.

27. Cattafesta, M.; Petarli, G.B.; Zandonade, E.; Bezerra, O.M.P.A.; Abreu, S.M.R.; Salaroli,
L.B. Energy contribution of NOVA food groups and the nutritional profile of the Brazilian
rural workers' diets. PLoS ONE 2020, 15, e0240756.

28. Khandpur, N.; Cediel, G.; Obando, D.A.; Jaime, P.C.; Parra, D.C. Sociodemographic
factors associated with the consumption of ultra-processed foods in Colombia. Rev. saúde
pública 2020, 54, 19.

29. Sandoval-Insausti, H.; Jiménez-Onsurbe, M.; Donat-Vargas, C.; Rey-García, J.; Banegas,
J.R.; Rodríguez-Artalejo, F.; Guallar-Castillón, P. Ultra-Processed Food Consumption Is
Associated with Abdominal Obesity: A Prospective Cohort Study in Older Adults.
Nutrients 2020, 12, 2368.

30. Yang, Q.; Zhang, Z.; Steele, E.M.; Moore, L.V.; Jackson, S.L. Ultra-Processed Foods and
Excess Heart Age Among U.S. Adults. American Journal of Preventive Medicine 2020,
59, e197–e206.

31. Nardocci, M.; Polsky, J.Y.; Moubarac, J.-C. Consumption of ultra-processed foods is
associated with obesity, diabetes and hypertension in Canadian adults. Can J Public Health
2021, 112, 421–429.

431