TITLE: Predictors of mortality among low birth weight neonates after hospital discharge
in a low-resource setting: A case study in Uganda.

Authors

Ronald Nsubuga[1,3]*, Joseph Rujumba[1,3], Saleh Nyende[1], Stevens Kisaka[2], Richard
Idro[1,3], Jolly Nankunda[1,3]

[1]Department of Paediatrics and Child Health, College of Health Sciences, Makerere
University, Kampala, Uganda.

[2]School Public Health, Makerere University, Kampala, Uganda

[3]Department of Paediatrics and Child Health, Mulago National Referral Hospital,
Kampala, Uganda

*Ronald Nsubuga

Email: ronald.nsubuga@mak.ac.ug | rnsubuga@dr.com

All authors contributed equally to this work.

Abstract

Background: Most of neonatal deaths occur among low birth weight infants. However, in
resource-limited settings, these infants are commonly discharged early which further
exposes them to mortality. This study aimed at determining the proportion and
predictors of mortality among low birth weight neonates that are discharged from
Mulago National Referral Hospital in Uganda.

Methods: This was a prospective cohort study of 220 low birth weight neonates
discharged from a Special Care Baby Unit at Mulago National Referral Hospital. These
were followed up to 28 completed days of life, or death, whichever occurred first.
Proportions were used to express mortality. To determine the predictors of mortality,
Cox hazards regression was performed.

Results: Of the 220 enrolled participants, 216 (98.1%) completed the follow-up. The mean gestational age of study participants was 34 ±3 weeks. The median weight at discharge was 1,650g (IQR: 1,315g -1,922g) and 46.1% were small for gestational age. During follow-up, 14/216 (6.5%) of neonates died. Mortality was highest (7/34, 20.6%) among neonates with discharge weights less than 1,200g. The causes of death included presumed neonatal sepsis (10/14, 71.4%), suspected aspiration pneumonia (2/14, 14.3%), and suspected cot death (2/14, 14.3%). The median time to death after discharge was 11 days (range 3-16 days). The predictors of mortality were a discharge weight of less than 1,200g (adj HR: 23.47, p <0.001), a 5-minute Apgar score of less than 7 (adj HR: 4.25, p = 0.016), and a diagnosis of neonatal sepsis during admission (adj HR: 7.93, p = 0.009).

Conclusion: Post-discharge mortality among low birth weight neonates in Uganda is high. A discharge weight of less than 1,200g may be considered unsafe among neonates. Caregiver education about neonatal danger signs, and measures to prevent sepsis, aspiration, and cot death should be emphasized before discharge and during follow-up visits.

Keywords: *Mortality, low birth weight, neonate, post-discharge.*

Introduction

Worldwide, an estimated 20 million infants are born with low birth weight (LBW) annually. The majority of these births (96.5%) occur in low and middle-income countries (LMICs)(1). Low birth weight contributes 60-80% of all neonatal deaths worldwide. Preterm birth, which is commonly associated with LBW is the most common direct cause of neonatal mortality(3). Low birth weight neonates are at increased odds of death compared to their heavier counterparts(4) and LBW is considered the single most important predictor of infant mortality (5).

The World Health Organization (WHO) developed discharge criteria for LBW infants(6). However, this has been difficult to adhere to in resource-limited settings like Uganda with an over-stretched healthcare system and an estimated 10% of births have LBW(2).

In Uganda, many neonatal units lack proper Kangaroo Mother Care (KMC) facilities(7,8), yet this is known to improve the survival of LBW infants (9).

Previous studies have assessed the morbidity and mortality of LBW infants after early discharge. Nonetheless, these mainly focused on Very Low Birth Weight (VLBW) infants and without the description of post-discharge neonatal mortality (10–12). Post-discharge outcomes of LBW neonates are largely unknown.

In our setting, LBW infants are discharged from the Special Care Baby Unit (SCBU) once stable (off antibiotics and oxygen, with stable body temperature in a cot). In addition, they should be able to tolerate 10ml per feed of expressed breastmilk by cup or tube, regardless of body weight or weight growth velocity. These babies are followed up weekly until they attain a body weight of 2500g. This practice may serve to reduce pressure on available limited resources, but its safety has not been assessed. This study aimed at determining the proportion of LBW neonates that die after discharge from the SCBU, and the predictors of mortality.

Methods and materials

Study setting

This was a prospective cohort study conducted at the SCBU of Mulago National Referral Hospital from 8th November 2018 to 28th January 2019. This SCBU was a level II neonatal unit. On average 4,600 babies are admitted to the unit annually, and 50% of these are low birth weight neonates.

The SCBU had an official bed capacity of 50 with 13 baby cots, 22 incubators, 11 radiant warmers, and 4 beds, but it often houses over 100 babies. It had term and preterm baby sections but with no KMC wing. The unit was run by 4 Paediatricians, 4-6 Paediatrics residents, 2 intern doctors on a rotational basis, and 18 nurses (working in 3 shifts a day).

The care offered included the provision of warmth (incubators and radiant warmers), phototherapy, and bubble Continuous Positive Airway Pressure (CPAP). Admitted neonates may also receive intravenous fluids (two hourly boluses), intravenous

antibiotics, and nasogastric feeding. Vital signs were intermittently assessed and
caregivers feed their babies every two hours under the supervision of the nursing team.

Low birth weight infants are discharged from the SCBU once they are deemed
physiologically stable (normal axillary temperature, off oxygen, off intravenous fluids,
and apnea-free), tolerating at least 10ml per enteral feed, and gaining weight. The
caregiver must demonstrate knowledge and competence in the provision of care to the
infant before discharge. On average, 138 LBW neonates are discharged monthly.

After discharge, LBW infants were followed up in the clinic every week until a body
weight of 2500g is attained. During clinic visits, growth monitoring, drug refills (Iron and
multivitamins), nasogastric tube replacement, and adjustment of feeds were done.
Unwell infants were identified and readmitted.

Study population
94 All neonates discharged from the SCBU with a birth weight of <2500g. Those who were
95 neither reachable physically nor by telephone were excluded from the study.

Sample size
97 The sample size was estimated using an OpenEpi formula for frequency in a population
98 (13). Since no similar studies were addressing post-discharge mortality among LBW
99 neonates, we assumed a frequency of 50%.

100 The calculated sample size (n) was 220 with a precision level of 5%.

Data collection procedure
All neonates with birth weight <2500g were identified by a research assistant at
discharge. Their parents were then introduced to the study. For neonates who met the
inclusion criteria, the research assistant obtained written informed consent from the
caregiver/ guardian. Recruitment was conducted daily and consecutively until the
desired sample size was achieved.

Recruited neonates received a study identification number and their details were
captured in a logbook. Each parent provided 3 phone contacts for purposes of follow-

up. Phone contacts of the principal investigator (PI) and research assistants were
shared with parents for them to call in case of any concerns.

Data were captured at discharge and every clinic visit. Study participants were followed
up for 28 days. Parents of study participants were contacted by telephone once a week
and significant events (death, illness) were registered. Those who missed their
scheduled follow-up visits were contacted by phone and requested to bring the neonate
for review on the next clinic day. A neonate who missed two consecutive clinic visits and
whose parents were not reachable by phone throughout the study period was
considered lost to follow-up.

Data management and analysis

Captured data were entered into Epidata version 3.1, checked for accuracy and
completeness, and exported to STATA Version 14 for analysis. Continuous variables
were summarized using the mean and standard deviation, and median and interquartile
ranges for skewed data. Categorical variables were summarized using frequencies and
proportions.

The proportion of LBW neonates that died after discharge from the SCU was expressed
as a percentage. The numerator was the number of babies that died within the first 28
days of life and the denominator was the total number of LBW neonates recruited into
the study. Chi-square and Fisher's exact tests were used to compare the proportion of
LBW neonates that died after discharge from the SCU by their baseline characteristics.

Student-t and Wilcoxon rank sum tests were used to compare means and median
respectively of the continuous variables of participants that died and those that were
alive after 28 completed days of life. To determine the predictors of mortality, relative
death rates (Hazard Ratios) were obtained using simple Cox proportional hazards
regression.

Variables that achieved a level of significance of ≤0.05 were then included in a
multivariable Cox proportional hazards model. Backward elimination was used to build a
model of best fit. The level of significance was set at ≤0.05 and a confidence interval of

95% was used. Kaplan Meier curves were used to compare survival across gender,
gestation age at birth, and birth weight.

Results

Of the 243 LBW neonates identified for potential recruitment into the study, 5 did not
consent to the study, and 18 did not have phone contact. Therefore, a total of 220
(representing 90.5%) neonates were enrolled in the study. Of these, 216 completed
follow-up. The mean gestational age at birth was 34 ±3 weeks, and the median weight
at discharge was 1650g (IQR 1315-1922 grams). All four neonates who were lost to
follow-up had discharge weight >2000g.

The median length of hospital stay of study participants was 5 days (IQR 3-8 days). The
majority of the study participants (134/216, 62%) were fully breastfeeding, and the rest
were breastfeeding and receiving top-up feeds through a nasogastric tube. Five study
participants had confirmed neonatal sepsis during admission, and they had completed
treatment before discharge (**Table 1** shows the details of the characteristics of the study
participants)

Table 1. Baseline characteristics of study participants

Variable	Frequency N= 216	Percentage (%)	p-value
Gender			
Male	93	43.0	0.988[c]
Female	123	57.0	
Birth type			
Singleton	143	66.2	0.777[c]
Twin	73	33.8	
Gestational age at birth (weeks)*			
<30	7	3.7	0.069[f]
30- 32	40	20.9	
33- 34	43	22.5	
35- <37	82	42.9	

≥37	19	10.0	
Post-menstrual age at discharge (weeks)			
<30	1	0.5	**0.027**[f]
30-32	33	17.3	
33-34	43	22.5	
35- <37	91	47.6	
≥37	23	12.1	
Birth weight (grams)			
<1200	18	8.3	**0.001**[f]
1200-1499	36	16.7	
1500-1799	56	25.9	
1800-1999	36	16.7	
2000- 1499	70	32.4	
Discharge weight (grams)			
<1200	34	15.7	**0.001**[f]
1200-1499	49	22.7	
1500-1799	46	21.3	
1800-1999	45	20.8	
≥2000	42	19.5	
SGA*			
No	103	53.9	0.16[f]
Yes	88	46.1	
5- minute Apgar score**			
<7	30	15.2	**0.009**[f]
≥7	167	84.8	
Neonatal sepsis during admission			
No	211	97.7	**0.035**[f]
Yes	5	2.3	
Maternal age (years)			
<20	21	11.3	0.407[f]
20-24	54	28.5	
25-29	53	28.0	

≥30	61	32.2	
Residence			
Rural	14	7.4	0.278[f]
Urban	175	92.6	
Maternal level of education			
Never gone to school and Primary	55	29.2	0.853[f]
Secondary	110	58.2	
Tertiary	24	12.6	
Maternal monthly income (UShs)			
0-50,000 (0-13.5$)	35	18.9	0.925[f]
>50,000-200,000 (13.5-54$)	35	18.9	
>200,000-500,000 (54-135$)	56	29.6	
>500,000 (>135$)	63	32.6	
Preceding birth interval (months)**			
0-24	32	24.0	0.34[f]
25-48	54	40.6	
>48	47	35.4	

Note: p-values are based on Pearson's Chi-square test (c) and Fisher's exact test (f).

During follow-up, 14/216 (6.4%) neonates died. Of these, 8 occurred from home, 5

occurred at a health facility, and 1 occurred on the way to the hospital. The

circumstances surrounding each death were established by contacting the primary

caregiver by phone. The likely causes of death included neonatal sepsis (71.4%),

aspiration pneumonia (14.3%), and cot death (14.3%). The median time to death after

discharge was 11 days (range 3-16 days).

The proportion of neonates that died was highest among those with discharge weight

<1200g (20.6%) and gestational age at birth <30 weeks (28.6%).

Figure 1 shows that the overall survival probability was 93.5% and the greatest drop in

survival probability occurred in the first week after discharge. The median survival time

had not been reached by the time the study closed meaning that survival was sustained

above 50% throughout the study period. Survival was compared according to birth

weight. **Figure 2** shows that survival probability was highest among study participants
with birth weight >2,000g. Notably, survival dropped with reducing birth weight.

The neonates who died had a lower median body weight at birth, discharge, and first
follow-up visit, compared to those who survived. In both groups, the median discharge
weight was lower than the birth weight. Among those who died, the median body weight
at the first follow-up visit was lower than the median weight at discharge as shown in
Figure 3.

Predictors of mortality were assessed at bivariate and multivariate levels of analysis.
Discharge weight <1200g was associated with a 23.5-fold increase in the risk of death
compared to babies of discharge weight greater than 1,500g (p<0.001). The neonates
who posted a 5-minute Apgar of <7 were 4.42 times more likely to die compared to
those with a 5-minute Apgar of 7 and above (p=0.016). Similarly, having a diagnosis of
neonatal sepsis at admission was significantly associated with mortality compared with
those that did not have such a diagnosis (adj HR = 7.93, p = 0.009). Notably, none of
the maternal sociodemographic characteristics was significantly associated with
mortality. The details of the unadjusted and adjusted hazard ratios are shown in **Table**
2.

Table 2. Multiple Cox regression for predictors of mortality among 216 LBW neonates
after discharge from the SCBU.

Variable	Total	Died after discharge		Unadjusted	HRp-value	Adjusted HR	p-value
	N= 216	No	Yes	(95% CI)		(95% CI)	
		n= 202 (%)	n= 14 (%)				
Discharge weight (grams)							
≥1500	133	130 (97.8)	3 (2.2)	Ref		Ref	
1200-1499	49	45 (91.8)	4 (8.2)	4.55	0.05	5.2	0.08
<1200	34	27 (79.4)	7 (20.6)	13.5	**<0.001**	23.47	**<0.001**
5- minute Apgar score							
≥7	167	160 (95.8)	7 (4.2)	Ref		Ref	
<7	30	25 (83.3)	5 (16.7)	4.42	**0.011**	4.25	**0.016**
Neonatal sepsis during admission							
No	211	199 (94.3)	12 (5.7)	Ref		Ref	

| Yes | 5 | 3 (60) | 2 (40) | 8.92 | **0.004** | 7.93 | **0.009** |

Discussion

In this study, we observed a post-discharge neonatal mortality of 6.5%. Mortality was highest among neonates delivered before 30 weeks of gestation (28.6%), those with birth weight <1500g (18.5%), and discharge weight <1200g (20.6%). Most of the deaths occurred at home and were due to possible neonatal sepsis.

The post-discharge mortality we observed at the referral hospital was lower than what was previously reported in two other similar settings. In Bangladesh, a study by Yasmin et al, 2001 found a mortality of 13.3% (4), while in Malawi, Blencove et al, 2009 found a mortality of 12.4% (14). The discrepancies may be explained by the differences in engagement between the healthcare facilities and the caregivers post-discharge. The level of mortality was however comparable to what Vazirinejad et al, 2012 found (5.4%) in Iran(15), and what Kibona et al found in Tanzania (16).

In all these studies, however, mortality was similarly higher among neonates with birth weights <1500g and those with gestational age at birth <30 weeks. The lower mortality observed in our study and Tanzania may be due to advances in neonatal care over the last 10 years since the other two studies were carried out. The higher mortality among those with birth weight <1500g, gestational age at birth <30 weeks, or discharge weight <1200g may be explained by immature oromotor coordination, poor homeostatic control among neonates with lower birth weight and lower gestational age at birth(17), and increased risk of sepsis in the same age group.

In our study, 3 in 5 LBW neonates who died had possible neonatal sepsis. This is comparable to the findings by Abdallah et al (10). It is not surprising because neonatal sepsis accounts for about 1 in 5 neonatal deaths in Uganda (18) and it is a significant contributor to neonatal deaths in most low and middle-income countries(19). Studies have shown that low birth weight is a risk factor for neonatal sepsis(20–22). Disease progression is fast and deaths occur rapidly. Indeed, our study found that the majority of the deaths occurred at home before parents had taken action to save their infants.

These findings are similar to what was found in Malawi (14). In part, this could reflect
inadequacies by parents to recognize danger signs in neonates and seek care promptly
as has been documented in Southwestern Uganda(23). On the other hand, this may be
explained by the fact that caregivers who think an infant is likely to die may be less likely
to seek urgent care, especially if the infant is fragile and small(24).

Predictors of mortality in this study were; discharge weight <1200g, 5- minute Apgar
score <7, and a diagnosis of neonatal sepsis during admission. Low birth weight
neonates with discharge weight <1200g were 23 times more likely to die compared to
heavier neonates. These results are comparable to what Abdallah et al, and Blencowe
et al found (10,14); that the risk of post-discharge mortality increases with decreasing
weight at discharge. This could be because smaller neonates, especially those with
weight <1500g are more likely to be preterm(1), and have not yet attained oromotor
maturity and appropriate homeostatic control. These neonates are therefore at greater
risk of complications like hypothermia, hypoglycemia, and aspiration. They are also
more likely to have immature immune systems (22). To compound this, they are likely to
have been born before the maximal transplacental transfer of antibodies from the
mother, which occurs in the last 8 weeks of gestation (25). They are therefore at
increased risk of infections, hence mortality.

Low birth weight neonates with an Apgar score <7 at 5 minutes were 4 times more likely
to die compared to those with an Apgar score ≥7. These results are in keeping with
what Abdullah et al found in Bangladesh, what a systematic review by Ehrestein found,
and what Cnattingius et al found in Sweden; that an Apgar score of <7 at 5 minutes was
associated with an increased risk of mortality among LBW neonates (26–28). This may
be due to complications of perinatal asphyxia like intraventricular hemorrhage, acute
kidney injury, and hypoxic ischaemic encephalopathy. The association between
prematurity and low Apgar score (29) may also explain the increased risk of mortality
among these neonates.

Neonates who had sepsis during admission were 8 times more likely to die after
discharge compared to those without this diagnosis. Neonatal sepsis is one of the

major causes of neonatal mortality worldwide (30). The increased likelihood of post-discharge mortality among LBW neonates treated for neonatal sepsis during admission could be a reflection of the persistence of some modifiable and non-modifiable risk factors of neonatal sepsis, like poor caregiver hygiene practices and prematurity. These neonates may therefore be at risk of re-infection and death even after discharge.

Strengths and limitations of the study

This was the first study in Uganda to assess mortality and its predictors among LBW neonates following discharge from the SCBU. The prospective cohort study design used for this study allowed us to establish a temporal association between mortality and its predictors. The study was carried out at a tertiary care facility that receives many referrals. This may have caused selection bias. We relied on clinical symptoms identified and reported by caregivers to ascertain the possible causes of death.

Conclusion

Neonatal mortality of 6.5% among low birth weight infants discharged from the National Referral Hospital is high. The predictors of mortality include discharge weight <1200g, a 5-minute Apgar score of <7, and a diagnosis of neonatal sepsis during admission. Most of the deaths occurred from home, in the first week after discharge, and were due to possible neonatal sepsis.

Improving survival of LBW neonates especially those with Apgar score <7 at 5 minutes, discharge weight <1200g, and /or have had sepsis during hospitalization requires intensive follow-up and care. There is a need to amend the guidelines for the discharge of LBW neonates so that they are discharged with weights >1200. Caregiver education on feeding, prevention of neonatal sepsis, early recognition of danger signs in neonates, and prompt care-seeking should be intensified.

Abbreviations

ANC: Antenatal care

BPD: Bronchopulmonary dysplasia

CPAP: Continuous Positive Airway Pressure

ELBW: Extremely low birth weight

HIV: Human Immunodeficiency Virus

IMR: Infant Mortality Rate

IVH: Intraventricular hemorrhage

KMC: Kangaroo Mother Care

LBW: Low birth weight

LMIC: Low and middle-income countries

LNMP: Last Normal Menstruation Period

NEC: Necrotising enterocolitis

NICU: Neonatal Intensive Care Unit

NMR: Neonatal Mortality Rate

MNRH: Mulago National Referral Hospital

SCBU: Special Care Baby Unit

SDGs: Sustainable Development Goals

SGA: Small for Gestational Age

SSA: Sub-Saharan Africa

VLBW: Very low birth weight

Acknowledgments

The authors are grateful to all the research assistants) and the study participants. We are also grateful to the staff at the Special Care Baby Unit, at Mulago National Referral Hospital.

Author contributions

All authors made a substantial contribution to the work, whether that was in the conception, study design, implementation, data collection, analysis, and interpretation, or all these areas: manuscript drafting, review, and revision. All authors read and approved the final manuscript.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Availability of data and materials

Available on request

Declarations

Ethical approval and consent to participate

This study protocol was subjected to approval by the Makerere University School of Medicine Research Ethics Committee. The investigators ensured that caregivers (parents or guardians) of study participants were given full and adequate oral and written information about the nature, purpose, possible risks, and benefits of the study. They were given adequate opportunities to ask questions and allowed time to consider the information provided voluntarily. Written consent was obtained from caregivers before conducting this study. Participants in this database were identified by their unique enrollment numbers. Only the PI had access to the participant identification list, which included their unique codes, full names, phone numbers and address.

Consent for publication

298   Not applicable.

300   **Competing interests**

301   The authors declare that they have no competing interests.

References

1.   Eds CP Howson, MV Kinney JLWHOG. Born too soon. Born Too Soon, Glob
Action Rep Preterm Birth World Heal Organ Geneva. 2012;13(5):1–126.

2.   Uganda Bureau of Statistics, ICF, DHS Program. Uganda Demographic and
Health Survey. 2016.

3.   WHO. WHO | Causes of child mortality. WHO [Internet]. 2016 [cited 2018 May
23]; Available from: http://www.who.int/gho/child_health/mortality/causes/en/

4.   Yasmin S, Osrin D, Paul E, Costello A. Neonatal mortality of low-birth-weight
infants in Bangladesh. Bull World Health Organ. 2001;79(7):608–14.

5.   Ryan CA, Ryan F, Keane E, Hegarty H. Trend analysis and socio-economic
differentials in infant mortality in the Southern Health Board, Ireland (1988-1997).
Ir Med J. 2000;93(7):204–6.

6.   WHO. Kangaroo mother care A practical guide. WHO Reprod Heal Res. 2003;

7.   Aliganyira P, Kerber K, Davy K, Gamache N, Sengendo NH, Bergh AM. Helping
small babies survive: An evaluation of facility-based Kangaroo Mother Care
implementation progress in Uganda. Pan Afr Med J. 2014 Sep 16;19.

8.   Bergh A-M, Davy K, Otai CD, Nalongo AK, Sengendo NH, Aliganyira P.
Evaluation of kangaroo mother care services in Uganda [Internet]. 2012. Available
from: http://www.mchip.net/

9.   Conde-Agudelo A, Díaz-Rossello JL. Kangaroo mother care to reduce morbidity
and mortality in low birthweight infants. Vol. 2014, Cochrane Database of

Systematic Reviews. John Wiley and Sons Ltd; 2014.

10.   Abdallah Y, Namiiro F, Nankunda J, Mugalu J, Vaucher Y. Mortality among very
325   low birth weight infants after hospital discharge in a low resource setting. BMC
326   Pediatr. 2018;18(239).

327   11.   De Jesus LC, Pappas A, Shankaran S, Kendrick D, Das A, Higgins RD, et al. Risk
328   Factors for Post-Neonatal Intensive Care Unit Discharge Mortality among
329   Extremely Low Birth Weight Infants. J Pediatr. 2012;161(1):70–4.

330   12.   Kugelman A, Reichman B, Chistyakov I, Boyko V, Levitski O, Lerner-Geva L, et
331   al. Postdischarge Infant Mortality Among Very Low Birth Weight Infants: A
332   Population-Based Study. Pediatrics. 2007;120(4):e788–94.

333   13.   Kevin M S. OpenEpi - Open Source Statistics for Public Health.

334   14.   Blencowe H, Kerac M, Molyneux E. Safety, effectiveness and barriers to follow-up
335   using an 'early discharge' kangaroo care policy in a resource poor setting. J Trop
336   Pediatr. 2009;55(4):244–8.

337   15.   Vazirinejad R, Masoodpour N, Puyanfar A. Survival rate of low and very low birth
338   weight neonates in an Iranian community. Iran J Public Health. 2012;41(2):87–93.

339   16.   Kibona SE, Sungwa EE, Dika HI, Laisser RM, Mhada T. Immediate Seven Day
340   Outcomes and Risk Factors of Low Birth Weight Neonates at Referral Hospitals in
341   Mwanza City , Tanzania in October 2020. Afr J Health Sci. 2021;34(4):451–63.

342   17.   Miller MJ, Kiatchoosakun P. Relationship between respiratory control and feeding
343   in the developing infant. Semin Neonatol. 2004 Jun;9(3):221–7.

344   18.   UNICEF. Uganda Maternal and Newborn Health Disparities [Internet]. 2015.
345   Available   from: https://data.unicef.org/wp-
346   content/uploads/country_profiles/Uganda/country profile_UGA.pdf

19.   Lawn JE, Blencowe H, Oza S, You D, Lee AC, Waiswa P, et al. Every Newborn:

progress, priorities, and potential beyond survival. Lancet. 2014 Jul
12;384(9938):189–205.

20. Leal YA, Álvarez-Nemegyei J, Velázquez JR, Rosado-Quiab U, Diego-Rodríguez
N, Paz-Baeza E, et al. Risk factors and prognosis for neonatal sepsis in
southeastern Mexico: analysis of a four-year historic cohort follow-up. BMC
Pregnancy Childbirth. 2012;12:48.

21. Adegoke S, Bankole K, Olugbemiga A, Tinuade O. Intraventricular hemorrhage in
newborns weighing <1500 g: Epidemiology and short-term clinical outcome in a
resource-poor setting. Ann Trop Med Public Heal. 2014;7(1):48.

22. Adatara P, Afaya A, Salia SM, Afaya RA, Kuug AK, Agbinku E, et al. Risk Factors
for Neonatal Sepsis: A Retrospective Case-Control Study among Neonates Who
Were Delivered by Caesarean Section at the Trauma and Specialist Hospital,
Winneba, Ghana. Biomed Res Int. 2018;2018.

23. Sandberg J, Pettersson KO, Asp G, Kabakyenga J, Agardh A. Inadequate
knowledge of neonatal danger signs among recently delivered women in
southwestern rural Uganda: A community survey. PLoS One. 2014;9(5).

24. O'Leary M, Edmond K, Floyd S, Newton S, Thomas G, Thomas SL. A cohort
study of low birth weight and health outcomes in the first year of life, Ghana. Bull
World Health Organ. 2017;95(8):574–83.

25. Fouda GG, Martinez DR, Swamy GK, Permar SR. The Impact of IgG
transplacental transfer on early life immunity. ImmunoHorizons. 2018 Jan
1;2(1):14–25.

26. Abdullah A, Hort K, Butu Y, Simpson L. Risk factors associated with neonatal
deaths: A matched case-control study in Indonesia. Glob Health Action.
2016;9(1).

27. Ehrenstein V. Clinical Epidemiology Association of Apgar scores with death and
neurologic disability. Clin Epidemiol. 2009;1:1–45.

28. Cnattingius S, Johansson S, Razaz N. Apgar Score and Risk of Neonatal Death among Preterm Infants. N Engl J Med. 2020;383(1):49–57.

29. Svenvik M, Brudin L, Blomberg M. Preterm Birth: A Prominent Risk Factor for Low Apgar Scores. Biomed Res Int. 2015 Aug 27;2015:1–8.

30. Perin J, Mulick A, Yeung D, Villavicencio F, Lopez G, Strong KL, et al. Global, regional, and national causes of under-5 mortality in 2000–19: an updated systematic analysis with implications for the Sustainable Development Goals. Lancet Child Adolesc Heal. 2022;6(2):106–15.

Figure 1. Kaplan Meier survival estimates for study participants (neonates) followed up over a period of 28 days.

Figure 2. Kaplan Meier survival estimates by birth weight categories among the 216 neonates

Figure 3. A box plot showing the variation of body weight at birth, discharge, and first follow-up visit among the 202 neonates who survived and the 14 who died.