TITLE: Predictors of mortality among low birth weight neonates after hospital discharge

in a low-resource setting: A case study in Uganda.

Authors

Ronald Nsubuga[1,3]*, Joseph Rujumba[1,3], Saleh Nyende[1], Stevens Kisaka[2], Richard

Idro[1,3], Jolly Nankunda[1,3]

[1]Department of Paediatrics and Child Health, College of Health Sciences, Makerere

University, Kampala, Uganda.

[2]School Public Health, Makerere University, Kampala, Uganda

[3]Department of Paediatrics and Child Health, Mulago National Referral Hospital,

Kampala, Uganda

*Ronald Nsubuga

Email: ronald.nsubuga@mak.ac.ug | rnsubuga@dr.com

All authors contributed equally to this work.

Abstract

Background: Most of neonatal deaths occur among low birth weight infants. However, in

resource-limited settings, these infants are commonly discharged early which further

exposes them to mortality. This study aimed at determining the proportion and predictors

of mortality among low birth weight neonates that are discharged from Mulago National

Referral Hospital in Uganda.

Methods: This was a prospective cohort study of 220 low birth weight neonates

discharged from a Special Care Baby Unit at Mulago National Referral Hospital. These

were followed up to 28 completed days of life, or death, whichever occurred first.

Proportions were used to express mortality. To determine the predictors of mortality, Cox

hazards regression was performed.

Results: Of the 220 enrolled participants, 216 (98.1%) completed the follow-up. The mean gestational age of study participants was 34 ±3 weeks. The median weight at discharge was 1,650g (IQR: 1,315g -1,922g) and 46.1% were small for gestational age. During follow-up, 14/216 (6.5%) of neonates died. Mortality was highest (7/34, 20.6%) among neonates with discharge weights less than 1,200g. The causes of death included presumed neonatal sepsis (10/14, 71.4%), suspected aspiration pneumonia (2/14, 14.3%), and suspected cot death (2/14, 14.3%). The median time to death after discharge was 11 days (range 3-16 days). The predictors of mortality were a discharge weight of less than 1,200g (adj HR: 23.47, p <0.001), a 5-minute Apgar score of less than 7 (adj HR: 4.25, p = 0.016), and a diagnosis of neonatal sepsis during admission (adj HR: 7.93, p = 0.009).

Conclusion: Post-discharge mortality among low birth weight neonates in Uganda is high. A discharge weight of less than 1,200g may be considered unsafe among neonates. Caregiver education about neonatal danger signs, and measures to prevent sepsis, aspiration, and cot death should be emphasized before discharge and during follow-up visits.

Keywords: *Mortality, low birth weight, neonate, post-discharge.*

Introduction

Worldwide, an estimated 20 million infants are born with low birth weight (LBW) annually. The majority of these births (96.5%) occur in low and middle-income countries (LMICs)(1). Low birth weight contributes 60-80% of all neonatal deaths worldwide. Preterm birth, which is commonly associated with LBW is the most common direct cause of neonatal mortality(3). Low birth weight neonates are at increased odds of death compared to their heavier counterparts(4) and LBW is considered the single most important predictor of infant mortality (5).

The World Health Organization (WHO) developed discharge criteria for LBW infants(6). However, this has been difficult to adhere to in resource-limited settings like Uganda with an over-stretched healthcare system and an estimated 10% of births have LBW(2). In

Uganda, many neonatal units lack proper Kangaroo Mother Care (KMC) facilities(7,8), yet this is known to improve the survival of LBW infants (9).

Previous studies have assessed the morbidity and mortality of LBW infants after early discharge. Nonetheless, these mainly focused on Very Low Birth Weight (VLBW) infants and without the description of post-discharge neonatal mortality (10–12). Post-discharge outcomes of LBW neonates are largely unknown.

In our setting, LBW infants are discharged from the Special Care Baby Unit (SCBU) once stable (off antibiotics and oxygen, with stable body temperature in a cot). In addition, they should be able to tolerate 10ml per feed of expressed breastmilk by cup or tube, regardless of body weight or weight growth velocity. These babies are followed up weekly until they attain a body weight of 2500g. This practice may serve to reduce pressure on available limited resources, but its safety has not been assessed. This study aimed at determining the proportion of LBW neonates that die after discharge from the SCBU, and the predictors of mortality.

Methods and materials

Study setting

This was a prospective cohort study conducted at the SCBU of Mulago National Referral Hospital from November 2018 to January 2019. This SCBU was a level II neonatal unit. On average 4,600 babies are admitted to the unit annually, and 50% of these are low birth weight neonates.

The SCBU had an official bed capacity of 50 with 13 baby cots, 22 incubators, 11 radiant warmers, and 4 beds, but it often houses over 100 babies. It had term and preterm baby sections but with no KMC wing. The unit was run by 4 Paediatricians, 4-6 Paediatrics residents, 2 intern doctors on a rotational basis, and 18 nurses (working in 3 shifts a day).

The care offered included the provision of warmth (incubators and radiant warmers), phototherapy, and bubble Continuous Positive Airway Pressure (CPAP). Admitted neonates may also receive intravenous fluids (two hourly boluses), intravenous

antibiotics, and nasogastric feeding. Vital signs were intermittently assessed and caregivers feed their babies every two hours under the supervision of the nursing team.

Low birth weight infants are discharged from the SCBU once they are deemed physiologically stable (normal axillary temperature, off oxygen, off intravenous fluids, and apnea-free), tolerating at least 10ml per enteral feed, and gaining weight. The caregiver must demonstrate knowledge and competence in the provision of care to the infant before discharge. On average, 138 LBW neonates are discharged monthly.

After discharge, LBW infants were followed up in the clinic every week until a body weight of 2500g is attained. During clinic visits, growth monitoring, drug refills (Iron and multivitamins), nasogastric tube replacement, and adjustment of feeds were done. Unwell infants were identified and readmitted.

Study population

All neonates discharged from the SCBU with a birth weight of <2500g. Those who were neither reachable physically nor by telephone were excluded from the study.

Sample size

The sample size was estimated using an OpenEpi formula for frequency in a population (13). Since no similar studies were addressing post-discharge mortality among LBW neonates, we assumed a frequency of 50%.

The calculated sample size (n) was 220 with a precision level of 5%.

Data collection procedure

All neonates with birth weight <2500g were identified by a research assistant at discharge. Their parents were then introduced to the study. For neonates who met the inclusion criteria, the research assistant obtained written informed consent. Recruitment was conducted daily and consecutively until the desired sample size was achieved.

Recruited neonates received a study identification number and their details were captured in a logbook. Each parent provided 3 phone contacts for purposes of follow-up. Phone contacts of the principal investigator (PI) and research assistants were shared with parents for them to call in case of any concerns.

Data were captured at discharge and every clinic visit. Study participants were followed
up for 28 days. Parents of study participants were contacted by telephone once a week
and significant events (death, illness) were registered. Those who missed their scheduled
follow-up visits were contacted by phone and requested to bring the neonate for review
on the next clinic day. A neonate who missed two consecutive clinic visits and whose
parents were not reachable by phone throughout the study period was considered lost to
follow-up.

116 Data management and analysis

Captured data were entered into Epidata version 3.1, checked for accuracy and
completeness, and exported to STATA Version 14 for analysis. Continuous variables
were summarized using the mean and standard deviation, and median and interquartile
ranges for skewed data. Categorical variables were summarized using frequencies and
proportions.

The proportion of LBW neonates that died after discharge from the SCU was expressed
as a percentage. The numerator was the number of babies that died within the first 28
days of life and the denominator was the total number of LBW neonates recruited into the
study. Chi-square and Fisher's exact tests were used to compare the proportion of LBW
neonates that died after discharge from the SCU by their baseline characteristics.

Student-t and Wilcoxon rank sum tests were used to compare means and median
respectively of the continuous variables of participants that died and those that were alive
after 28 completed days of life. To determine the predictors of mortality, relative death
rates (Hazard Ratios) were obtained using simple Cox proportional hazards regression.

Variables that achieved a level of significance of ≤0.05 were then included in a
multivariable Cox proportional hazards model. Backward elimination was used to build a
model of best fit. The level of significance was set at ≤0.05 and a confidence interval of
95% was used. Kaplan Meier curves were used to compare survival across gender,
gestation age at birth, and birth weight.

Results

Of the 243 LBW neonates identified for potential recruitment into the study, 5 did not consent to the study, and 18 did not have phone contact. Therefore, a total of 220 (representing 90.5%) neonates were enrolled in the study. Of these, 216 completed follow-up. The mean gestational age at birth was 34 ±3 weeks, and the median weight at discharge was 1650g (IQR 1315-1922 grams). All four neonates who were lost to follow-up had discharge weight >2000g.

The median length of hospital stay of study participants was 5 days (IQR 3-8 days). The majority of the study participants (134/216, 62%) were fully breastfeeding, and the rest were breastfeeding and receiving top-up feeds through a nasogastric tube. Five study participants had confirmed neonatal sepsis during admission, and they had completed treatment before discharge (**Table 1** shows the details of the characteristics of the study participants)

Table 1. Baseline characteristics of study participants

Variable	Frequency N= 216	Percentage (%)	p-value
Gender			
Male	93	43.0	0.988[c]
Female	123	57.0	
Birth type			
Singleton	143	66.2	0.777[c]
Twin	73	33.8	
Gestational age at birth (weeks)*			
<30	7	3.7	0.069[f]
30- 32	40	20.9	
33- 34	43	22.5	
35- <37	82	42.9	
≥37	19	10.0	
Post-menstrual age at discharge (weeks)			
<30	1	0.5	**0.027[f]**

30-32	33	17.3	
33-34	43	22.5	
35- <37	91	47.6	
≥37	23	12.1	
Birth weight (grams)			
<1200	18	8.3	**0.001**[f]
1200-1499	36	16.7	
1500-1799	56	25.9	
1800-1999	36	16.7	
2000- 1499	70	32.4	
Discharge weight (grams)			
<1200	34	15.7	**0.001**[f]
1200-1499	49	22.7	
1500-1799	46	21.3	
1800-1999	45	20.8	
≥2000	42	19.5	
SGA*			
No	103	53.9	0.16[f]
Yes	88	46.1	
5- minute Apgar score**			
<7	30	15.2	**0.009**[f]
≥7	167	84.8	
Neonatal sepsis during admission			
No	211	97.7	**0.035**[f]
Yes	5	2.3	
Maternal age (years)			
<20	21	11.3	0.407[f]
20-24	54	28.5	
25-29	53	28.0	
≥30	61	32.2	
Residence			
Rural	14	7.4	0.278[f]

Urban	175	92.6
Maternal level of education		
Never gone to school and Primary	55	29.2	0.853[f]
Secondary	110	58.2
Tertiary	24	12.6
Maternal monthly income (UShs)		
0-50,000 (0-13.5$)	35	18.9	0.925[f]
>50,000-200,000 (13.5-54$)	35	18.9
>200,000-500,000 (54-135$)	56	29.6
>500,000 (>135$)	63	32.6
Preceding birth interval** (months)		
0-24	32	24.0	0.34[f]
25-48	54	40.6
>48	47	35.4

Note: *p*- values are based on Pearson's Chi-square test (c) and Fisher's exact test (f).

During follow-up, 14/216 (6.4%) neonates died. Of these, 8 occurred from home, 5

occurred at a health facility, and 1 occurred on the way to the hospital. The circumstances

surrounding each death were established by contacting the primary caregiver by phone.

The likely causes of death included neonatal sepsis (71.4%), aspiration pneumonia

(14.3%), and cot death (14.3%). The median time to death after discharge was 11 days

(range 3-16 days).

The proportion of neonates that died was highest among those with discharge weight

<1200g (20.6%) and gestational age at birth <30 weeks (28.6%).

Figure 1 shows that the overall survival probability was 93.5% and the greatest drop in

survival probability occurred in the first week after discharge. The median survival time

had not been reached by the time the study closed meaning that survival was sustained

above 50% throughout the study period.

Figure 1. Kaplan Meier survival estimates for study participants (neonates) followed up over a period of 28 days.

Survival was compared according to birth weight. **Figure 2** shows that survival probability was highest among study participants with birth weight >2,000g. Notably, survival dropped with reducing birth weight.

Figure 2. Kaplan Meier survival estimates by birth weight categories among the 216 neonates

The neonates who died had a lower median body weight at birth, discharge, and first
follow-up visit, compared to those who survived. In both groups, the median discharge
weight was lower than the birth weight. Among those who died, the median body weight
at the first follow-up visit was lower than the median weight at discharge as shown in
Figure 3.

Figure 3. A box plot showing the variation of body weight at birth, discharge, and first
follow-up visit among the 202 neonates who survived and the 14 who died.

Predictors of mortality were assessed at bivariate and multivariate levels of analysis.
Discharge weight <1200g was associated with a 23.5-fold increase in the risk of death
compared to babies of discharge weight greater than 1,500g (p<0.001). The neonates
who posted a 5-minute Apgar of <7 were 4.42 times more likely to die compared to those
with a 5-minute Apgar of 7 and above (p=0.016). Similarly, having a diagnosis of neonatal
sepsis at admission was significantly associated with mortality compared with those that
did not have such a diagnosis (adj HR = 7.93, p = 0.009). Notably, none of the maternal
sociodemographic characteristics was significantly associated with mortality. The details
of the unadjusted and adjusted hazard ratios are shown in **Table 2**.

Table 2. Multiple Cox regression for predictors of mortality among 216 LBW neonates
after discharge from the SCBU.

Variable	Total	Died after discharge		Unadjusted	HR*p*-value	Adjusted HR	*p*-value
	N= 216	No	Yes	(95% CI)		(95% CI)	
		n= 202 (%)	n= 14 (%)				
Discharge weight (grams)							
≥1500	133	130 (97.8)	3 (2.2)	Ref		Ref	
1200-1499	49	45 (91.8)	4 (8.2)	4.55	0.05	5.2	0.08
<1200	34	27 (79.4)	7 (20.6)	13.5	<0.001	23.47	<0.001
5- minute Apgar score							
≥7	167	160 (95.8)	7 (4.2)	Ref		Ref	
<7	30	25 (83.3)	5 (16.7)	4.42	0.011	4.25	0.016
Neonatal sepsis during admission							
No	211	199 (94.3)	12 (5.7)	Ref		Ref	
Yes	5	3 (60)	2 (40)	8.92	0.004	7.93	0.009

Discussion

In this study, we observed a post-discharge neonatal mortality of 6.5%. Mortality was
highest among neonates delivered before 30 weeks of gestation (28.6%), those with birth
weight <1500g (18.5%), and discharge weight <1200g (20.6%). Most of the deaths
occurred at home and were due to possible neonatal sepsis.

The post-discharge mortality we observed at the referral hospital was lower than what
was previously reported in two other similar settings. In Bangladesh, a study by Yasmin
et al, 2001 found a mortality of 13.3% (4), while in Malawi, Blencove et al, 2009 found a
mortality of 12.4% (14). The discrepancies may be explained by the differences in
engagement between the healthcare facilities and the caregivers post-discharge. The
level of mortality was however comparable to what Vazirinejad et al, 2012 found (5.4%)
in Iran(15), and what Kibona et al found in Tanzania (16).

In all these studies, however, mortality was similarly higher among neonates with birth
weights <1500g and those with gestational age at birth <30 weeks. The lower mortality
observed in our study and Tanzania may be due to advances in neonatal care over the

last 10 years since the other two studies were carried out. The higher mortality among
those with birth weight <1500g, gestational age at birth <30 weeks, or discharge weight
<1200g may be explained by immature oromotor coordination, poor homeostatic control
among neonates with lower birth weight and lower gestational age at birth(17), and
increased risk of sepsis in the same age group.

In our study, 3 in 5 LBW neonates who died had possible neonatal sepsis. This is
comparable to the findings by Abdallah et al (10). It is not surprising because neonatal
sepsis accounts for about 1 in 5 neonatal deaths in Uganda (18) and it is a significant
contributor to neonatal deaths in most low and middle-income countries(19). Studies have
shown that low birth weight is a risk factor for neonatal sepsis(20–22). Disease
progression is fast and deaths occur rapidly. Indeed, our study found that the majority of
the deaths occurred at home before parents had taken action to save their infants. These
findings are similar to what was found in Malawi (14). In part, this could reflect
inadequacies by parents to recognize danger signs in neonates and seek care promptly
as has been documented in Southwestern Uganda(23). On the other hand, this may be
explained by the fact that caregivers who think an infant is likely to die may be less likely
to seek urgent care, especially if the infant is fragile and small(24).

Predictors of mortality in this study were; discharge weight <1200g, 5- minute Apgar score
<7, and a diagnosis of neonatal sepsis during admission. Low birth weight neonates with
discharge weight <1200g were 23 times more likely to die compared to heavier neonates.
These results are comparable to what Abdallah et al, and Blencowe et al found (10,14);
that the risk of post-discharge mortality increases with decreasing weight at discharge.
This could be because smaller neonates, especially those with weight <1500g are more
likely to be preterm(1), and have not yet attained oromotor maturity and appropriate
homeostatic control. These neonates are therefore at greater risk of complications like
hypothermia, hypoglycemia, and aspiration. They are also more likely to have immature
immune systems (22). To compound this, they are likely to have been born before the
maximal transplacental transfer of antibodies from the mother, which occurs in the last 8
weeks of gestation (25). They are therefore at increased risk of infections, hence
mortality.

Low birth weight neonates with an Apgar score <7 at 5 minutes were 4 times more likely
to die compared to those with an Apgar score ≥7. These results are in keeping with what
Abdullah et al found in Bangladesh, what a systematic review by Ehrestein found, and
what Cnattingius et al found in Sweden; that an Apgar score of <7 at 5 minutes was
associated with an increased risk of mortality among LBW neonates (26–28). This may
be due to complications of perinatal asphyxia like intraventricular hemorrhage, acute
kidney injury, and hypoxic ischaemic encephalopathy. The association between
prematurity and low Apgar score (29) may also explain the increased risk of mortality
among these neonates.

Neonates who had sepsis during admission were 8 times more likely to die after discharge
compared to those without this diagnosis. Neonatal sepsis is one of the major causes of
neonatal mortality worldwide (30). The increased likelihood of post-discharge mortality
among LBW neonates treated for neonatal sepsis during admission could be a reflection
of the persistence of some modifiable and non-modifiable risk factors of neonatal sepsis,
like poor caregiver hygiene practices and prematurity. These neonates may therefore be
at risk of re-infection and death even after discharge.

Strengths and limitations of the study

This was the first study in Uganda to assess mortality and its predictors among LBW
neonates following discharge from the SCBU. The prospective cohort study design used
for this study allowed us to establish a temporal association between mortality and its
predictors. The study was carried out at a tertiary care facility that receives many referrals.
This may have caused selection bias. We relied on clinical symptoms identified and
reported by caregivers to ascertain the possible causes of death.

Conclusion

Neonatal mortality of 6.5% among low birth weight infants discharged from the National
Referral Hospital is high. The predictors of mortality include discharge weight <1200g, a
5-minute Apgar score of <7, and a diagnosis of neonatal sepsis during admission. Most
of the deaths occurred from home, in the first week after discharge, and were due to
possible neonatal sepsis.

Improving survival of LBW neonates especially those with Apgar score <7 at 5 minutes,

discharge weight <1200g, and /or have had sepsis during hospitalization requires

intensive follow-up and care. There is a need to amend the guidelines for the discharge

of LBW neonates so that they are discharged with weights >1200. Caregiver education

on feeding, prevention of neonatal sepsis, early recognition of danger signs in neonates,

and prompt care-seeking should be intensified.

Abbreviations

ANC:	Antenatal care
BPD:	Bronchopulmonary dysplasia
CPAP:	Continuous Positive Airway Pressure
ELBW:	Extremely low birth weight
HIV:	Human Immunodeficiency Virus
IMR:	Infant Mortality Rate
IVH:	Intraventricular hemorrhage
KMC:	Kangaroo Mother Care
LBW:	Low birth weight
LMIC:	Low and middle-income countries
LNMP:	Last Normal Menstruation Period
NEC:	Necrotising enterocolitis
NICU:	Neonatal Intensive Care Unit
NMR:	Neonatal Mortality Rate
MNRH:	Mulago National Referral Hospital
SCBU:	Special Care Baby Unit
SDGs:	Sustainable Development Goals
SGA:	Small for Gestational Age
SSA:	Sub-Saharan Africa
VLBW:	Very low birth weight

Acknowledgments

The authors are grateful to all the research assistants) and the study participants. We are also grateful to the staff at the Special Care Baby Unit, at Mulago National Referral Hospital.

Author contributions

All authors made a substantial contribution to the work, whether that was in the conception, study design, implementation, data collection, analysis, and interpretation, or all these areas: manuscript drafting, review, and revision. All authors read and approved the final manuscript.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Availability of data and materials

Available on request

Declarations

Ethical approval and consent to participate

This study protocol was subjected to approval by the Makerere University School of Medicine Research Ethics Committee. The investigators ensured that participants were given full and adequate oral and written information about the nature, purpose, possible risks, and benefits of the study. They were given adequate opportunities to ask questions and allowed time to consider the information provided voluntarily. The subject's signed and dated informed consent was obtained before conducting this study. Participants in this database were identified by their unique enrollment numbers. Only the PI had access

299 to the participant identification list, which included their unique codes, full names, and the
300 latest known address.

Consent for publication

302 Not applicable.

Competing interests

304 The authors declare that they have no competing interests.

References

1. Eds CP Howson, MV Kinney JLWHOG. Born too soon. Born Too Soon, Glob Action
Rep Preterm Birth World Heal Organ Geneva. 2012;13(5):1–126.

2. Uganda Bureau of Statistics, ICF, DHS Program. Uganda Demographic and Health
Survey. 2016.

3. WHO. WHO | Causes of child mortality. WHO [Internet]. 2016 [cited 2018 May 23];
Available from: http://www.who.int/gho/child_health/mortality/causes/en/

4. Yasmin S, Osrin D, Paul E, Costello A. Neonatal mortality of low-birth-weight infants
in Bangladesh. Bull World Health Organ. 2001;79(7):608–14.

5. Ryan CA, Ryan F, Keane E, Hegarty H. Trend analysis and socio-economic
differentials in infant mortality in the Southern Health Board, Ireland (1988-1997). Ir
Med J. 2000;93(7):204–6.

6. WHO. Kangaroo mother care A practical guide. WHO Reprod Heal Res. 2003;

7. Aliganyira P, Kerber K, Davy K, Gamache N, Sengendo NH, Bergh AM. Helping
small babies survive: An evaluation of facility-based Kangaroo Mother Care
implementation progress in Uganda. Pan Afr Med J. 2014 Sep 16;19.

8. Bergh A-M, Davy K, Otai CD, Nalongo AK, Sengendo NH, Aliganyira P. Evaluation
of kangaroo mother care services in Uganda [Internet]. 2012. Available from:
http://www.mchip.net/

9.   Conde-Agudelo A, Díaz-Rossello JL. Kangaroo mother care to reduce morbidity
and mortality in low birthweight infants. Vol. 2014, Cochrane Database of
Systematic Reviews. John Wiley and Sons Ltd; 2014.

10.   Abdallah Y, Namiiro F, Nankunda J, Mugalu J, Vaucher Y. Mortality among very
low birth weight infants after hospital discharge in a low resource setting. BMC
Pediatr. 2018;18(239).

11.   De Jesus LC, Pappas A, Shankaran S, Kendrick D, Das A, Higgins RD, et al. Risk
Factors for Post-Neonatal Intensive Care Unit Discharge Mortality among
Extremely Low Birth Weight Infants. J Pediatr. 2012;161(1):70–4.

12.   Kugelman A, Reichman B, Chistyakov I, Boyko V, Levitski O, Lerner-Geva L, et al.
Postdischarge Infant Mortality Among Very Low Birth Weight Infants: A Population-
Based Study. Pediatrics. 2007;120(4):e788–94.

13.   Kevin M S. OpenEpi - Open Source Statistics for Public Health.

14.   Blencowe H, Kerac M, Molyneux E. Safety, effectiveness and barriers to follow-up
using an 'early discharge' kangaroo care policy in a resource poor setting. J Trop
Pediatr. 2009;55(4):244–8.

15.   Vazirinejad R, Masoodpour N, Puyanfar A. Survival rate of low and very low birth
weight neonates in an Iranian community. Iran J Public Health. 2012;41(2):87–93.

16.   Kibona SE, Sungwa EE, Dika HI, Laisser RM, Mhada T. Immediate Seven Day
Outcomes and Risk Factors of Low Birth Weight Neonates at Referral Hospitals in
Mwanza City , Tanzania in October 2020. Afr J Health Sci. 2021;34(4):451–63.

17.   Miller MJ, Kiatchoosakun P. Relationship between respiratory control and feeding
in the developing infant. Semin Neonatol. 2004 Jun;9(3):221–7.

18.   UNICEF. Uganda Maternal and Newborn Health Disparities [Internet]. 2015.
Available   from: https://data.unicef.org/wp-
content/uploads/country_profiles/Uganda/country profile_UGA.pdf

19.   Lawn JE, Blencowe H, Oza S, You D, Lee AC, Waiswa P, et al. Every Newborn:
progress, priorities, and potential beyond survival. Lancet. 2014 Jul
12;384(9938):189–205.

20.   Leal YA, Álvarez-Nemegyei J, Velázquez JR, Rosado-Quiab U, Diego-Rodríguez
N, Paz-Baeza E, et al. Risk factors and prognosis for neonatal sepsis in
southeastern Mexico: analysis of a four-year historic cohort follow-up. BMC
Pregnancy Childbirth. 2012;12:48.

21.   Adegoke S, Bankole K, Olugbemiga A, Tinuade O. Intraventricular hemorrhage in
newborns weighing <1500 g: Epidemiology and short-term clinical outcome in a
resource-poor setting. Ann Trop Med Public Heal. 2014;7(1):48.

22.   Adatara P, Afaya A, Salia SM, Afaya RA, Kuug AK, Agbinku E, et al. Risk Factors
for Neonatal Sepsis: A Retrospective Case-Control Study among Neonates Who
Were Delivered by Caesarean Section at the Trauma and Specialist Hospital,
Winneba, Ghana. Biomed Res Int. 2018;2018.

23.   Sandberg J, Pettersson KO, Asp G, Kabakyenga J, Agardh A. Inadequate
knowledge of neonatal danger signs among recently delivered women in
southwestern rural Uganda: A community survey. PLoS One. 2014;9(5).

24.   O'Leary M, Edmond K, Floyd S, Newton S, Thomas G, Thomas SL. A cohort study
of low birth weight and health outcomes in the first year of life, Ghana. Bull World
Health Organ. 2017;95(8):574–83.

25.   Fouda GG, Martinez DR, Swamy GK, Permar SR. The Impact of IgG transplacental
transfer on early life immunity. ImmunoHorizons. 2018 Jan 1;2(1):14–25.

26.   Abdullah A, Hort K, Butu Y, Simpson L. Risk factors associated with neonatal
deaths: A matched case-control study in Indonesia. Glob Health Action. 2016;9(1).

27.   Ehrenstein V. Clinical Epidemiology Association of Apgar scores with death and
neurologic disability. Clin Epidemiol. 2009;1:1–45.

28.   Cnattingius S, Johansson S, Razaz N. Apgar Score and Risk of Neonatal Death
among Preterm Infants. N Engl J Med. 2020;383(1):49–57.

29.   Svenvik M, Brudin L, Blomberg M. Preterm Birth: A Prominent Risk Factor for Low
Apgar Scores. Biomed Res Int. 2015 Aug 27;2015:1–8.

30.   Perin J, Mulick A, Yeung D, Villavicencio F, Lopez G, Strong KL, et al. Global,
regional, and national causes of under-5 mortality in 2000–19: an updated
systematic analysis with implications for the Sustainable Development Goals.
Lancet Child Adolesc Heal. 2022;6(2):106–15.

386