Preventable suicides involving medicines before the covid-19 pandemic: a systematic case series of coroners' reports in England and Wales (pre-print)

Grace Anthony[1]; Jeffrey K. Aronson[2]; Richard Brittain[3]; Carl Heneghan[2]; Georgia C. Richards[2]*

[1]Oxford University Medical School, John Radcliffe Hospital, Headington, University of Oxford, Oxford, OX3 9DU, UK.
[2]Centre for Evidence-Based Medicine, Nuffield Department of Primary Care Health Sciences, University of Oxford, Radcliffe Observatory Quarter, Woodstock Road, Oxford, OX2 6GG, UK.
[3]St Pancras Coroner's Court, London, N1C 4PP, UK.

*Corresponding author: Dr Georgia C. Richards; Centre for Evidence-Based Medicine, University of Oxford, email: georgia.richards@phc.ox.ac.uk

Word count: 2969

Abstract

Background: Over 5000 suicides are registered in England and Wales each year. Coroners' Prevention of Future Deaths reports (PFDs) share concerns to promote actions to reduce the risks of similar deaths.

Aims: To systematically review coroners' PFDs involving suicides in which a medicine caused or contributed to the death, to identify lessons for suicide prevention.

Methods: 3037 PFDs were screened for eligibility between July 2013 and December 2019. Following data extraction, descriptive statistics and content analysis were performed to assess coroners' concerns, the recipients, and the response rates to coroners.

Results: There were 734 suicide-related PFDs, with 100 reporting a medicine. Opioids (40%) were the most common class involved in suicide-related PFDs, followed by antidepressants (30%). There was wide geographical variation in the writing of PFDs; coroners in Manchester wrote the most (18%). Coroners expressed 237 concerns; the most common were procedural inadequacies (14%, n=32), inadequate documentation and communication (10%, n=22), and inappropriate prescription access (9%, n=21). 203 recipients received these PFDs, most being sent to NHS trusts (31%), clinical commissioning groups (10%), and general practices (10%), of which only 58% responded to the coroner. We created the Preventable Deaths Tracker (https://preventabledeathstracker.net/) to increase dissemination and analysis of coroners' PFDs.

Conclusions: Concerns raised by coroners in suicide-related PFDs involving medicines highlight essential gaps in care that require actions from the Government, health services, and prescribers. To aid suicide prevention, PFDs should be disseminated nationally, and responses should be enforced so that actions are taken to prevent suicides.

Study protocol registration: https://doi.org/10.17605/OSF.IO/EXJK3

Keywords: Suicide; mental health; medicines; overdose; coroners; preventable deaths

Introduction

Suicide is a leading cause of preventable deaths, involving over 700,000 deaths every year globally ([1](#)). In England and Wales, 5691 suicides occurred in 2019, a 12% increase since 2000 ([2](#)). In 2021, there were 5583 registered suicides in England and Wales ([3](#)). Various factors are associated with the risk of suicide, including a previous attempt, self-harm, bereavement, mental ill-health, social isolation, and recent adverse life events ([1](#), [4](#), [5](#)).

Legislation to reduce pack sizes of over-the-counter (OTC) paracetamol has been effective in reducing deaths, with a 35% reduction in paracetamol poisonings between 1998 and 2009 ([6](#)). Co-proxamol was removed from UK markets in 2007 owing to its toxicity in suicide, with an estimated 500 lives saved between 2010 and 2015 ([7](#)). However, codeine is still available OTC in the UK and large volumes are being sold ([8](#)).

Coroners in England and Wales have a duty to communicate a death when they believe that actions should be taken to prevent future deaths ([9](#)). These reports, called Prevention of Future Deaths reports (PFD) ([10-12](#)), are sent to individuals or organisations with the power to act. Once received, recipients must respond to the coroner within 56 days, outlining actions taken or proposed. However, there is no mechanism to enforce responses or to analyse, monitor, or learn from PFDs ([13](#)).

Previous studies have assessed suicide-related PFDs ([13-15](#)) but these have focused on specific issues, including claims against the NHS. The Guardian reviewed 500 PFDs in 2018 to identify deaths related to mental illness ([16](#), [17](#)). A systematic case series of PFDs involving medicines was also conducted in 2013-2022, identifying opioids and antidepressants as the leading medicines involved in preventable deaths ([18](#)). The ONS has published an analysis of 164 suicide-related PFDs, covering 2021 and 2022 alone ([19](#), [20](#)). No research to date has investigated PFDs involving medicine-related suicides before the SARS-CoV-2 pandemic. We therefore aimed to assess all available PFDs detailing deaths by suicide in which a medicine caused or contributed to the death. Our primary objectives were to discover: 1) the medicines involved; 2) the concerns raised by coroners; 3) the recipients to whom PFDs were addressed; and 4) the response rates of addressees.

Methods

We designed a retrospective case series and pre-registered the study protocol on an open repository (https://doi.org/10.17605/OSF.IO/EXJK3).

Data sources

PFDs have been openly available on the Courts and Tribunals Judiciary website since 2013 ([21](#)). We used web scraping to automatically collect all published PFDs to create a database and build the Preventable Deaths Tracker (https://preventabledeathstracker.net/) ([22](#)). The code to create the web scraper is available on GitHub and has been previously described ([23](#), [24](#)). The database contains the case reference number; the date of the report; the name of the deceased; the coroner's name; the coroner's area; the category of death (defined by the Chief Coroner's Office); to whom the report was sent; and the url of the Judiciary website. Population data for suicides in England and Wales corresponding to the dates of PFDs included in the analysis (2013-2019) were obtained from the Office for National Statistics (ONS) ([2](#)).

Eligibility of PFDs

There were 3037 PFDs available at the time of the web scrape. PFDs were screened using a pre-defined algorithm (Supplementary Fig.1), in which suicide was defined as intentional self-harm that led to or contributed to death or death of undetermined intent by injury or poisoning. This definition was widened compared with the ONS definition of suicide, because the standard of proof for reaching a conclusion of suicide was changed from a criminal to a civil one in 2018 ([25](#)). The ONS relies on the conclusion supplied by the coroner, which is influenced by factors such as the age and sex of the deceased, the suicide method, marital status, and the individual likelihood that the coroner will supply a narrative verdict ([2](#), [26-31](#)). If there were discrepancies in eligibility, cases were discussed and agreed on.

Following the inclusion of all suicide-related PFDs, cases were reviewed for medicines. PFDs were included if a medicine was listed as a medical cause of death or was stated as contributing to the death by the coroner, or was the cause of toxicity resulting in death. A medicine was defined as any prescription-only or over-the-counter medication. Cases were excluded if the drug or drugs were obtained illicitly, or if the suicide involved non-medical products (e.g. alcohol), or if the medicine was mentioned in the report, but there was no indication that it directly or indirectly contributed to the death.

Data extraction

Following the screening for eligibility, one study author (GA) manually extracted data from the included PFDs into a predesigned Google Sheet. The data extracted, included:

1. the date of death,
2. the inquest start date and end date,
3. the age of the deceased (if reported),
4. the sex of the deceased,
5. the location of the death (if noted),

6. the medicine that caused or contributed to the death,

7. how the medicines were obtained,

8. health service contacts before the death,

9. relevant medical, mental health, and social history,

10. concern(s) raised by the coroner,

11. those who responded to the PFD, and

12. the date on which the response was received.

Data analysis

We descriptively analysed the extracted data using medians and interquartile ranges (IQR) for continuous variables and frequencies (percentages) for categorical variables. The types of medicines were categorised into drug classes using the British National Formulary (BNF). To examine geographical variation, we further grouped coroner areas into standard regions to create a choropleth map. We used content analysis with inductive logic to group concerns raised by coroners from included PFDs. Common themes and phrases were highlighted and drawn out, and frequencies were calculated using the total numbers of concerns as the denominators. We calculated a response rate to PFDs for each organisation as the proportion of reports to which a response was submitted over the total number received. In October 2020, responses were further classified as either early (delivered before 56 days), on-time (delivered on the 56th day), late (submitted after 56 days), or overdue (when no response was present on the Judiciary website).

Software and data sharing

We used Google Sheets to analyse the data, and Tableau to create figures. The data and study materials are openly available via the Open Science Framework, and the web scraper code is available on Github ([32](#)). Datawrapper was used to create a choropleth map of published PFDs by regions of England and Wales (available at https://www.datawrapper.de).

Results

734 PFDs (n=735 deaths) involved a suicide between July 2013 and December 2019 (Figure 1). Of these suicide-related PFDs, 14% (n=100) reported that a medicine caused or contributed to the death. Over the seven-year study period, there was a median of 118 (IQR: 102-119) suicide-related PFDs each year and a median of 17 (IQR: 10-19) medicine-related suicides (Figure 2; Supplementary Table S1).

Figure 1. Flow diagram of the screening of coroners' Prevention of Future Deaths reports (PFDs) assessed in this study.

Figure 2: Trends of coroners' Prevention of Future Deaths reports (PFDs) published between July 2013 and December 2019 involving suicides (light blue) and medicine-related suicides (green) compared with the total number of published PFDs (dark blue).

In the 100 PFDs involving medicines and suicides, 46 unique medicines were identified, involving a total of 152 medicines. Morphine (15%) was the most often reported, followed by paracetamol (12%) and zopiclone (10%) (Figure 3). Using BNF drug classes, opioids (n=40) caused or contributed to the most medicine-related suicides, followed by anti-depressants (n=30) (Supplement Table S2). In 16 cases the type of medicine was unspecified.

Figure 3 Types of medicines reported by coroners in suicide-related Prevention of Future Death reports (PFDs) in England and Wales between July 2013 and December 2019.

Across the 100 suicides involving medicines, there was an equal sex ratio of males (n=50) and females (n=50). The median age of the deceased was 42 years (IQR: 28-50 years; n=61). Depression (n=29) was the most common mental health problem reported by coroners, followed by anxiety disorders (n=12), emotionally unstable personality disorder/borderline personality disorder (n=11) and psychotic symptoms (n=10) (Supplement Figure S2).

Fifty-three coroners across 47 coroner areas wrote the 100 PFDs. Coroners in the Northwest (22%) and Southwest (16%) wrote the most, whereas coroners in the Northeast (3.0%) and Yorkshire and the Humber (3.0%) regions wrote the fewest (Figure 4). Manchester South issued the highest number of PFDs involving medicines and suicide (8.0%) (Supplement Table S3).

Figure 4: Choropleth map of medicine-related suicides reported in coroners' Prevention of Future Deaths reports (PFDs) published in England and Wales from July 2013 to December 2019.

237 concerns were raised by coroners in the 100 PFDs. These were categorised using content analysis, which identified 18 themes (Figure 5). The concerns most frequently reported included procedural inadequacies (14%, n=32), inadequate documentation and communication (9.3%, n=22), inappropriate prescription access (8.9%, n=21), and improper risk assessment (7.2%, n=17; Supplement Table S4).

Figure 5: Concerns reported by coroners in 100 suicide-related Prevention of Future Deaths report (PFDs) involving medicines in England and Wales published between July 2013 and December 2019.

Coroners sent 203 PFDs to 145 individuals or organisations (Table 1). NHS trusts (31%; n=63), clinical commission groups (CCGs; 9.9%; n=20), and general practices (9.9%; n=20) were sent the most PFDs. By law, recipients must respond to the coroner within 56 days, but only 42% (n=85) were received within that time. 13% (n=27) were late, and 42% (n=86) were overdue. Government bodies (63%) and NHS bodies (62%) had the highest response rates, while law enforcement (44%) had the lowest response rates.

Table 1. Recipients sent coroners' Prevention of Future Death reports (PFDs) involving medicine-related suicides in England and Wales published between July 2013 and December 2019. Responses were classified as early (delivered before 56 days), on-time (delivered on the 56th day), late (submitted after 56 days), or overdue (when no response was present on the Judiciary website).

Addressee	No. of reports received	No. of responses	Response rate (%)	% Early	%On time	%Late	%Overdue	%Unclear
NHS Body	**138**	**86**	**62**	**36**	**10**	**14**	**38**	**1**
CCGs	20	13	65	45	5	10	35	5
NHS addiction services	1	0	0	0	0	0	100	0
NHS England	9	6	67	22	0	44	33	0
NHS England south-west	1	0	0	0	0	0	100	0

NHS Trusts and University Health Boards	63	40	63	33	14	14	38	2
Ambulances	8	4	50	38	0	13	50	0
Hospitals	9	5	56	22	33	0	44	0
GP practices	20	11	55	35	0	20	45	0
Mental Health Trusts	6	6	100	100	0	0	0	0
NHS Pathways	1	1	100	0	100	0	0	0
Government	**24**	**15**	**63**	**33**	**0**	**25**	**38**	**4**
Local Authorities	6	4	67	50	0	17	33	0
DHSC	11	8	73	27	0	45	27	0
Home office	2	0	0	0	0	0	100	0
Home secretary	1	0	0	0	0	0	100	0
Secretary of State for Health	1	1	100	100	0	0	0	0
Welsh Minister for Health	2	2	100	50	0	0	0	50
Advisory Council on the misuse of drugs	1	0	0	0	0	0	100	0
Law enforcement	**18**	**8**	**44**	**22**	**11**	**11**	**56**	**0**
Police	5	3	60	60	0	0	40	0
MOJ	2	1	50	0	0	50	50	0
MOD	1	1	100	100	0	0	0	0
Mitie Care and Custody	1	0	0	0	0	0	100	0
Prisons	3	0	0	0	0	0	100	0
Director of public prosecutions	1	0	0	0	0	0	100	0
National offender management service	1	0	0	0	0	0	100	0
HM prison and probation service	2	2	100	0	50	50	0	0
National police chiefs' council	1	0	0	0	0	0	100	0
G4S	1	1	100	0	100	0	0	0
Professional bodies	**13**	**7**	**54**	**31**	**8**	**0**	**46**	**15**
CQC	1	1	100	0	0	0	0	100
PHE	1	1	100	100	0	0	0	0
MHRA	3	0	0	0	0	0	100	0
GMC	2	1	50	50	0	0	50	0
GPC	1	0	0	0	0	0	100	0
RCGP	2	2	100	100	0	0	0	0
Toxbase	1	1	100	0	0	0	0	100
University bodies	2	1	50	0	50	0	50	0
Other	**10**	**2**	**20**	**20**	**0**	**0**	**80**	**0**
Housing support/associations	4	2	50	50	0	0	50	0
Private healthcare companies	3	0	0	0	0	0	100	0
Blank	1	0	0	0	0	0	100	0
Livewell Southwest	1	0	0	0	0	0	100	0
Post-prison support charity	1	0	0	0	0	0	100	0

CCG: Crown Commissioning Group; CQC: Care Quality Commission; DHSC: Department Of Health and Social Care; G4S: Group 4 Securicor; GMC: General Medical Council: GPC: General Pharmaceutical Council; MHRA: Medicines and Healthcare Products Regulatory Agency; MOD: Ministry Of Defence; MOJ: Ministry of Justice; PHE: Public Health England; RCGP: Royal College of General Practitioners.

Discussion

Coroners reported 100 deaths in PFDs that involved suicides and medicines before the Covid-19 pandemic. Coroners often raised concerns regarding procedural inadequacies, inadequate documentation and communication, and inappropriate prescription access. Opioids, antidepressants, and paracetamol were the most common medicines involved in suicide-related PFDs. NHS organisations and Government bodies received the most PFDs, but only 58% responded to the coroner.

Comparison with previous research

A case series of all PFDs involving medicines similarly found that opioids and antidepressants were the most common medicines reported in PFDs (18). The ONS previously reported that opioids, antidepressants, and paracetamol were the most common drugs reported in self-poisoning suicides (33). The 2023 Healthcare Quality Improvement Partnership (HQIP) annual National Confidential Inquiry into Suicide and Safety in Mental Health also found that opioids were the most common type of drugs involved in suicides, most often obtained by prescription (34). An analysis of PFDs between January 2021 and October 2022 by the ONS found 164 PFDs categorised as suicide from the Judiciary website, which identified similar coroners' concerns to our study (20). NHS Resolution has also reported similar concerns highlighted in suicide-related PFDs between 2015 and 2017, identifying issues in communication, risk assessments and the observational processes (13).

Despite the statutory requirement that mandates a response to the coroners within 56 days (11), there is no mechanism to follow up responses to PFDs or the ability to assess and ensure that actions proposed in responses are implemented. We found that over 40% of responses were still outstanding, making it challenging to understand whether action has been taken to reduce future deaths. Others have similarly found low response rates to PFDs (18, 35, 36). Future research should further examine responses and assess the factors that affect the response rate, such as the nature of death or the types of recipients.

We identified significant variation in the writing of PFDs across England and Wales. Others have similarly reported this. For example, France et al. identified Manchester, Birmingham, and Inner London as the most common areas with medicines-related PFDs (14, 18, 35). Future research should

determine the factors driving this variation, including population statistics, mortality rates and coroners' attitudes towards writing PFDs.

Male suicides account for three-quarters of suicides in England and Wales, leading to increased initiatives to reduce male-related suicide ([2]). However, we found no sex differences for medicines-related suicides. Studies have further highlighted the fact that female deaths are less likely to be reported to the coroner, be taken to inquest, and reach a conclusion of suicide, with accident/ misadventure and narrative verdicts being favoured conclusions for females ([2], [30], [31], [37-39]). Future research should account for potential sex biases in the writing of PFDs and the reporting of such deaths in PFDs.

Strengths and limitations

This study is the first to have assessed medicines-related suicides reported in PFDs before the start of the Covid-19 pandemic. Our reproducible data collection methods allowed us to examine more than six years of data from over 3000 PFDs. Previous studies have investigated less than two years of data and used manual methods based on the Judiciary website categorisation ([14], [20], [35], [36]). Unlike the investigations conducted by HQIP and the Guardian ([16], [17], [40]), our study included all suicides, even if the individual had not been in contact with a mental healthcare professional in the past year and when an NHS trust or hospital was not addressed in the report. To be inclusive, we expanded previous definitions of suicide, as the Judiciary categorisation was available only from 2015 onwards, and the ONS definition ([27]) uses the coroners' conclusions of suicide, open, undetermined and narrative to assign a categorisation of suicide. We examined all PFDs and included cases where intentional self-harm led to or contributed to death, or death of undetermined intent by self-injury or poisoning. Our definition reduces the coronial variation of reaching a suicide conclusion, but it may have also resulted in over-estimation of suicide-related PFDs ([25], [31], [41]).

The 100 deaths included in this study do not represent all deaths deemed preventable involving suicides and medicines between July 2013 and December 2019, as the writing of PFDs depends on the coroner and the evidence presented during the death investigation. A qualitative interview of 32 senior coroners in 2009 showed that only 50% of respondents regarded public health as a key aspect of their duties ([42]). It may be possible that some PFDs and their responses may not have been published on the Judiciary website, resulting in publication and reporting biases. We also found that the data provided in PFDs varied. We encountered missing data for essential variables, including age (not reported in 39% of PFDs), types of medicines involved (not specified in 16%), where the medicine was obtained (not reported in 47%), and the location of the suicide (not reported in 19%).

Implications

Concerns raised by coroners in PFDs should aid in suicide prevention and patient care. Our study highlights the continued need for safer prescribing of opioids and antidepressants. In England, the prescribing of opioids and antidepressants has increased by 34% and 100% respectively between 2008 and 2018 (43, 44). The poor application of policies and procedures that were repeatedly identified by coroners should be addressed when updating NICE guidance on depression, which will be renewed in 2023 (45, 46). Medicines implicated in PFD suicides may be emerging evidence of risks associated with these medications; the MHRA are considering if further action is required following concerns raised regarding increased risk of suicide when using ciprofloxacin and quinolones (50, 51).

PFDs can have further policy impact, including the PFD of a 14-year old girl who died by suicide in 2017 following exposure to harmful online content, which has contributed to changes in the Online Safety bill (47, 48). In October 2023 the government will instate a new Health Services Safety Investigations Body, tasked with investigating mental health inpatient care settings, PFD reports could play a role in "how providers'...learn from tragic deaths that take place in their care" (49).

Our study examined suicide-related PFDs published before the Covid-19 pandemic (July 2013 to December 2019). However, there is substantial evidence that the Covid-19 pandemic negatively affected the mental health of many people, particularly the young, minorities, and women (52-56). The global burden of major depressive and anxiety disorders rose by 27% and 26% retrospectively during the Covid-19 pandemic (57). A analysis of PFDs during the Covid-19 pandemic (2021-2022) has since been conducted by the ONS, which found 164 suicide-related PFDs (20). In response to ONS's analysis of suicide-related PFDs, the Samaritans stated that *"It's hugely concerning that these common themes are consistently emerging and there's no evidence to suggest this will improve without urgent, deliberate action."*(58) Prescriptions for mental health conditions have been predicted to increase following the pandemic, and lockdowns may have increased the severity of mental illness (59, 60). Therefore, concerns raised by coroners in PFDs should be used by policy makers to mitigate the increased risk of suicide.

Wider dissemination of PFDs and their responses could contribute to national initiatives to reduce suicides, as detailed by the *NHS Mental Health Implementation Plan 2019/20 – 2023/24* (61), national charities, and NHS Resolution (13, 19, 58). Despite the legal requirement under Regulation 29 of The Coroners (Investigations) Regulations 2013 to respond to coroners within 56 days (11), 42% of responses remained overdue. NHS Resolution recommended in 2018 that neighbouring NHS trusts should share learning from their PFDs and their responses to them, highlighting good practice to be regular meetings with their local and area coroner, to identify common themes arising (13). The Justice Select Committee's report on the Coroner's Service in 2021/22 acknowledged that the absence of follow-up on coroners' PFDs is a missed opportunity and recommended that an Inspectorate be

established to follow up on actions proposed in PFDs.([62](#)) The Government responded, stating that they were "not in a position to accept the recommendation at this stage" and that the cost would be disproportionate to the benefit it may bring.([63](#)) We therefore created a method, database, and resource (https://preventabledeathstracker.net/) to disseminate lessons from PFDs.

Future research should investigate the barriers and facilitators of writing and responding to PFDs, the suitability of concerns reported in suicide-medicine PFDs, the suitability of recipients of such PFDs, and the actions taken and proposed in the 117 responses received for suicide-medicine PFDs. Others can use our data to analyse the 635 suicide-related PFDs that did not involve medicines between July 2013 and December 2019. The wide variation and missing data in PFDs also highlight areas for coronial training and the need to standardise data reporting for PFDs.

Conclusions

PFDs hold a wealth of information regarding gaps in suicide prevention measures and the risks of medicine-related suicides. However, it is unclear if these PFDs are being used to benefit patient care and prevent future deaths. To aid in suicide prevention, PFDs should be disseminated more widely, responses to PFDs should be audited, and regular analyses of the spatial and temporal changes in concerns should be conducted. This study provides a tool (https://preventabledeathstracker.net/) to increase lessons learnt from coroners' PFDs.

Funding: No funding was obtained to undertake this study.

Ethical approval: As the case reports are publicly available on the Judiciary website, approval from an ethics committee was not required.

Contributions: GA proposed the topic for research, wrote the study protocol and devised the algorithm for the eligibility of cases, screened the database for eligible case reports, extracted and analysed the data, and wrote the first draft of the manuscript. GCR designed the study, guided the writing of the study protocol, sourced the data, contributed to the web scraper, provided data management, dual screened the database for eligible cases, analysed the data, provided primary supervisory support and substantially contributed to subsequent manuscript drafts. JKA, RB, and CH reviewed the protocol and preliminary findings and provided supervisory support. All authors have full access to all the data in the study, read and approved the final manuscript, and accept responsibility to submit for publication.

Data availability: Study materials, protocol, data, and statistical code are openly available on the OSF (https://doi.org/10.17605/OSF.IO/AD4UP).

Declaration of interests: GA has no conflicts to declares. JKA has published articles and edited textbooks on adverse drug reactions and interactions and has often given medicolegal advice, including appearances as an expert witness in coroners' courts, dealing with the adverse effects of drugs. CH has received expenses and fees for his media work, received expenses from the World Health Organisation (WHO), Food and Drug Administration (FDA), and holds grant funding from the NIHR School for Primary Care Research (SPCR), and the NIHR SPCR Evidence Synthesis Working Group [Project 380], the NIHR BRC Oxford and the WHO. On occasion, CH receives expenses for teaching EBM and is also paid for his GP work in NHS out of hours (contract with Oxford Health NHS Foundation Trust). GCR was financially supported by the NIHR SPCR, the Naji Foundation, and the Rotary Foundation to study for a Doctor of Philosophy (2017-2020), is an Associate Editor of BMJ Evidence Based Medicine, and has a limited company that is independently contracted to conduct epidemiological research. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, or the Department of Health and Social Care.

References

1.   WHO. Suicide: World Health Organisation; 2021 [Available from: https://www.who.int/news-room/fact-sheets/detail/suicide.

2.   ONS. Suicides in England and Wales 1981-2021: Office for National Statistics; 2022 [Available from: https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/datasets/suicidesintheunitedkingdomreferencetables.

3.   ONS. Suicides in England and Wales: 2021 registrations 2022 [updated 6th September. Available from: https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/bulletins/suicidesintheunitedkingdom/2021registrations.

4.   Gunnell D, Appleby L, Arensman E, Hawton K, John A, Kapur N, et al. Suicide risk and prevention during the COVID-19 pandemic. The Lancet Psychiatry. 2020;7(6):468-71.

5.   Rodway C, Tham S-G, Ibrahim S, Turnbull P, Windfuhr K, Shaw J, et al. Suicide in children and young people in England: a consecutive case series. The Lancet Psychiatry. 2016;3(8):751-9.

6.   Hawton K, Bergen H, Simkin S, Dodd S, Pocock P, Bernal W, et al. Long term effect of reduced pack sizes of paracetamol on poisoning deaths and liver transplant activity in England and Wales: interrupted time series analyses. BMJ : British Medical Journal. 2013;346:f403.

7.   Hawton K, Bergen H, Simkin S, Wells C, Kapur N, Gunnell D. Six-year follow-up of impact of co-proxamol withdrawal in England and Wales on prescribing and deaths: time-series study. PLoS Med. 2012;9(5):e1001213.

8.   Richards GC, Aronson JK, MacKenna B, Goldacre B, Hobbs FDR, Heneghan C. Sales of Over-the-Counter Products Containing Codeine in 31 Countries, 2013–2019: A Retrospective Observational Study. Drug Safety. 2022;45(3):237-47.

9.   The Coroners Rules 1984 1984 [Available from: https://www.legislation.gov.uk/uksi/1984/552/article/43/made.

10.   Coroners and Justice Act 2009 2009 [Available from: https://www.legislation.gov.uk/ukpga/2009/25/pdfs/ukpga_20090025_en.pdf.

11.   The Coroners (Investigations) Regulations 2013 2013 [Available from: https://www.legislation.gov.uk/uksi/2013/1629/pdfs/uksi_20131629_en.pdf.

12.   Chief Coroner HHJP. Guidance No.5 Reports to Prevent Future Deaths 2016 [Available from: https://www.judiciary.uk/wp-content/uploads/2013/09/guidance-no-5-reports-to-prevent-future-deaths.pdf.

13.   Resolution NHS OA. Learning from Suicide-Related Claims: A thematic review of NHS Resolution data 2018 [Available from: https://resolution.nhs.uk/wp-content/uploads/2018/09/NHS-Resolution_learing_from_suicide_claims_148pp_ONLINE1.pdf.

14.   King R, Benbow EW. Are the recommendations in coronial prevention of future death (PFD) reports realistic and achievable? Med Leg J. 2022;90(1):27-31.

15.   Leary A, Bushe D, Oldman C, Lawler J, Punshon G. A thematic analysis of the prevention of future deaths reports in healthcare from HM coroners in England and Wales 2016–2019. Journal of Patient Safety and Risk Management. 2021;26(1):14-21.

16.   Duncan P CD, Bawden A. Hundreds of mental health patients died after NHS care failures 2018 [updated 5th March 2018. Available from: http://www.theguardian.com/society/2018/mar/05/hundreds-of-mental-health-patients-dying-after-nhs-care-failures.

17.   P D. How the Guardian investigated mental health patient deaths 2018 [updated 5th March 2018. Available from: http://www.theguardian.com/society/2018/mar/05/how-the-guardian-investigated-mental-health-patient-deaths.

18.   France HS, Aronson JK, Heneghan C, Ferner RE, Cox AR, Richards GC. Preventable Deaths Involving Medicines: A Systematic Case Series of Coroners' Reports 2013-22. Drug Saf. 2023;46(4):335-42.

19.   Papyrus. Analysis of suicides will help to save lives: Prevention of Young Suicide; 2023 [Available from: https://www.papyrus-uk.org/analysis-of-suicides-will-help-to-save-lives/.

20.   ONS. Prevention of Future Death Reports for Suicide submitted to coroners in England and Wales: January 2021 to October 2022 2023 [updated 29 March 2023. Available from: https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/mentalhealth/articles/preventionoffuturedeathreportsforsuicidesubmittedtocoronersinenglandandwales/january2021tooctober2022.

21.   Prevention of Future Deaths [Available from: https://www.judiciary.uk/prevention-of-future-death-reports/.

22.   Preventable Deaths Tracker 2021 [cited 2021 6th June]. Available from: https://preventabledeathstracker.net.

23.   Richards G. georgiarichards.github.io: GitHub; [cited 2021 6th June]. Available from: https://github.com/georgiarichards/georgiarichards.github.io.

24.   DeVito Nj RG, Inglesby P. How we learnt to stop worrying and love web scraping. Nature. 2020;585(7826):621-2.

25.   Maughan, R (On the Application Of) v Her Majesty's Senior Coroner for Oxfordshire [2019] EWCA Civ 809 (10 May 2019) 2019 [Available from: https://www.bailii.org/ew/cases/EWCA/Civ/2019/809.html.

26.   ONS. Suicides in the UK: 2013 Registrations- Narrative Verdicts in England and Wales 2015 [Available from: https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/bulletins/suicidesintheunitedkingdom/2015-02-19.

27.   ONS. Suicide rates in the UK QMI: Office for National Statistics; 2019 [cited 2020 30th October]. Available from: https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/methodologies/suicideratesintheukqmi.

28.   QC CCHJPT. Guidance no.17 Conclusions: Short-Form and Narrative 2016 Jan [Available from: https://www.judiciary.uk/wp-content/uploads/2013/09/guidance-no-17-conclusions.pdf.

29.   Gunnell D, Bennewith O, Simkin S, Cooper J, Klineberg E, Rodway C, et al. Time trends in coroners' use of different verdicts for possible suicides and their impact on officially reported incidence of suicide in England: 1990–2005. Psychological Medicine. 2013;43(7):1415-22.

30.   Carroll R, Hawton K, Kapur N, Bennewith O, Gunnell D. Impact of the growing use of narrative verdicts by coroners on geographic variations in suicide: analysis of coroners' inquest data. Journal of Public Health. 2011;34(3):447-53.

31.   Palmer BS, Bennewith O, Simkin S, Cooper J, Hawton K, Kapur N, et al. Factors influencing coroners' verdicts: an analysis of verdicts given in 12 coroners' districts to researcher-defined suicides in England in 2005. Journal of Public Health. 2014;37(1):157-65.

32.   Richards GC, Anthony, G. L., Aronson, J. K., Heneghan, C., Goldacre, B Preventable suicides: a systematic analysis of coroners' prevent future death reports in England and Wales.: OSF; 2020 [updated 28th October. Available from: https://doi.org/10.17605/OSF.IO/EXJK3.

33.   Paul Breen AB. Deaths related to drug poisoning by selected substances, England and Wales 2022 [Available from: https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/datasets/deathsrelatedtodrugpoisoningbyselectedsubstances.

34.   HQIP. National Confidential Inquiry into Suicide and Safety in Mental Health 2023 [Available from: https://www.hqip.org.uk/wp-content/uploads/2023/03/Ref-395-NCISH-2023-Annual-Report.pdf?vgo_ee=hm0jxNm4C2%2FaWjmR9d8idXUSKn30JrGyZ16N%2BJD6xewHE7BHlI0WgfM%3D%3ACcLCLm7Cgn5Euj%2FnMOEjqhoTo1eFYMKV

35.   Fox AW, Jacobson J. How well do Regulation 28 reports serve future public health and safety? Medicine, Science and the Law. 2021;61(3):186-92.

36.   Ferner RE, Easton C, Cox AR. Deaths from Medicines: A Systematic Analysis of Coroners' Reports to Prevent Future Deaths. Drug Safety. 2018;41(1):103-10.

37.   QC CCHJPT. Report of the Chief Coroner to the Lord Chancellor; Third annual report: 2015-2016 [2016 [Available from: https://www.judiciary.uk/wp-content/uploads/2016/09/chief_coroner_report_2016_web2.pdf.

38.   Roberts ISD, Gorodkin LM, Benbow EW. What is a natural cause of death? A survey of how coroners in England and Wales approach borderline cases. Journal of Clinical Pathology. 2000;53(5):367-73.

39.   Mclean M. Coroner consistency – The 10-jurisdiction, 10-year, postcode lottery? Medicine, Science and the Law. 2015;55(2):102-12.

40.   HQIP. The National Confidential Inquiry into Suicide and Safety in Mental Health. Annual Report: England, Northern Ireland, Scotland and Wales 2019 [Available from: https://www.hqip.org.uk/wp-content/uploads/2019/12/Mental-Health-CORP-Report-2019-FINAL.pdf.

41.   Mclean M. THE CORONER IN ENGLAND AND WALES; CORONIAL DECISION-MAKING AND LOCAL VARIATION IN CASE OUTCOMES [Doctoral]: University of Huddersfield; 2015.

42.   McGowen C. Frustration of purpose: public health and the future of death investigation in England & Wales, p130-133 [Dotoral]2012.

43.   BMJ. Vastly differing opioid prescribing patterns in England even in similarly deprived areas 2020 [updated 14th December. Available from: https://www.bmj.com/company/newsroom/vastly-differing-opioid-prescribing-patterns-in-england-even-in-similarly-deprived-areas/.

44.   Iacobucci G. NHS prescribed record number of antidepressants last year. BMJ. 2019;364:l1508.

45.   NICE. Depression in adults: treatment and management: National Institute for Health and Care Excellence; [Available from: https://www.nice.org.uk/guidance/ng222.

46.   NICE. Depression in Adults (update): National Institue for Health and Care Excellence; 2011 [Available from: https://www.nice.org.uk/guidance/qs8.

47.   Walker A. Molly Russell: Prevention of future deaths report 2022 [Available from: https://www.judiciary.uk/prevention-of-future-death-reports/molly-russell-prevention-of-future-deaths-report/.

48.   Commons Ho. Online Safety Bill 87 (Rev) 2023 [Available from: https://bills.parliament.uk/publications/49376/documents/2822.

49.   Department of Health and Social Care MCM, and The Rt Hon Steve Barclay MP. Government acts to improve patient safety in mental health care 2023 [Available from: https://www.gov.uk/government/news/government-acts-to-improve-patient-safety-in-mental-health-care.

50.   Wilkinson E. Rare ciprofloxacin suicide link to be reviewed by MHRA after coroner verdict: hospital pharmacy europe; 2023 [updated 22nd June 2023. Available from: https://hospitalpharmacyeurope.com/clinical-zones/mental-health/rare-ciprofloxacin-suicide-link-to-be-reviewed-by-mhra-after-coroner-verdict/.

51.   Fleming M. Robert Stevenson 2023 [Available from: https://www.judiciary.uk/prevention-of-future-death-reports/robert-stevenson-prevention-of-future-deaths-report/.

52.   Nomura S, Kawashima T, Yoneoka D, Tanoue Y, Eguchi A, Gilmour S, et al. Trends in suicide in Japan by gender during the COVID-19 pandemic, up to September 2020. Psychiatry Research. 2021;295:113622.

53.   Monteith LL, Holliday R, Hoffmire CA. Understanding women's risk for suicide during the COVID-19 pandemic: A call to action. Psychiatry Research. 2021;295:113621.

54.   Digital N. Mental Health of Children and Young People in England, 2020: Wave 1 follow up to the 2017 survey 2020 [Available from: https://digital.nhs.uk/data-and-information/publications/statistical/mental-health-of-children-and-young-people-in-england/2020-wave-1-follow-up.

55.   Iob E, Steptoe A, Fancourt D. Abuse, self-harm and suicidal ideation in the UK during the COVID-19 pandemic. The British Journal of Psychiatry. 2020;217(4):543-6.

56.   O'Connor RC, Wetherall K, Cleare S, McClelland H, Melson AJ, Niedzwiedz CL, et al. Mental health and well-being during the COVID-19 pandemic: longitudinal analyses of adults in the UK COVID-19 Mental Health & Wellbeing study. The British Journal of Psychiatry. 2021;218(6):326-33.

57.   Santomauro DF, Mantilla Herrera AM, Shadid J, Zheng P, Ashbaugh C, Pigott DM, et al. Global prevalence and burden of depressive and anxiety disorders in 204 countries and territories in 2020 due to the COVID-19 pandemic. The Lancet. 2021;398(10312):1700-12.

58.   Samaritans. Samaritans response to today's ONS report on Prevention of Future Deaths reports for suicide in England and Wales: Samaritans; 2023 [updated 29th March. Available from: https://www.samaritans.org/news/samaritans-response-to-todays-ons-report-on-prevention-of-future-deaths-reports-for-suicide-in-england-and-wales/.

59.   Carr MJ, Steeg S, Webb RT, Kapur N, Chew-Graham CA, Abel KM, et al. Effects of the COVID-19 pandemic on primary care-recorded mental illness and self-harm episodes in the UK: a population-based cohort study. The Lancet Public Health. 2021;6(2):e124-e35.

60.   Williams R, Jenkins DA, Ashcroft DM, Brown B, Campbell S, Carr MJ, et al. Diagnosis of physical and mental health conditions in primary care during the COVID-19 pandemic: a retrospective cohort study. The Lancet Public Health. 2020;5(10):e543-e50.

61.   NHS. NHS Mental Health Implementation Plan 2019/20 – 2023/24 2019 [Available from: https://www.longtermplan.nhs.uk/wp-content/uploads/2019/07/nhs-mental-health-implementation-plan-2019-20-2023-24.pdf.

62.   Committee HoCJ. Bereaved people are
not yet at the heart
of the coroner service: House of Commons Committees; 2021 [Available from: https://houseofcommons.shorthandstories.com/justice-coroner-service/index.html.

63.   The Coroner Service: Government Response to the Committee's First Report- Letter from the Rt Hon Robert Buckland QC MP, Lord Chancellor and Secretary of State for Justice 2021 [Available from:

https://committees.parliament.uk/publications/7221/documents/77640/default/#:~:text=On%2027%20
May%202021%20the,received%20on%2027%20July%202021.