Outbreak risks, cases, and costs of vaccination strategies against wild poliomyelitis in polio-free settings: a modelling study

Megan Auzenbergs[1,2], Kaja Abbas[1,2], Arie Voorman[3,4], Corey Peak[4], Mark Jit, Kathleen M O'Reilly[1,2]

[1] Centre for Mathematical Modelling of Infectious Diseases, London School of Hygiene and Tropical Medicine, Keppel Street, London, UK
[2] Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, Keppel Street, London, UK
[3] Institute for Disease Modelling, Seattle, WA, USA
[4] Bill and Melinda Gates Foundation, Seattle, WA, USA

Corresponding author:
Megan Auzenbergs
Megan.auzenbergs@lshtm.ac.uk

Abstract

Background

Polio eradication was threatened in 2022 by importation of wild poliovirus serotype 1 into Malawi with subsequent international spread, representing the first wild polio cases in Africa since 2014. Preventing importations and spread of wild poliovirus is critical, and dependent on population immunity provided through routine immunisation and supplementary immunisation activities (SIAs). However, the scale of preventative SIAs has reduced in recent years due to financial constraints.

Methods

We developed a mathematical model of polio transmission dynamics to evaluate the probability of an outbreak, expected number of poliomyelitis cases, and the costs associated with vaccination delivery through routine immunisation (RI), outbreak response SIAs (oSIAs) and preventative SIAs (pSIAs). Across varying levels of routine immunisation coverage, we explore three key strategies: RI+oSIAs, RI+oSIAs+annual pSIAs, and RI+oSIAs+biannual pSIAs.

Results

The annual pSIA strategy (RI+oSIAs+annual pSIAs) had higher costs but greater probability of no outbreaks: under our model assumptions, annual pSIAs result in 80% probability of no outbreaks when routine immunisation coverage ≥50%. The biannual pSIA strategy (RI+oSIAs+biannual pSIAs) costs less and averts more outbreaks than RI+oSIAs, but RI coverage ≥65% was required to achieve equivalent risk of no outbreaks. The strategy with no pSIAs (RI+oSIAs) had the lowest costs but required ≥75% RI coverage to achieve equivalent risk of no outbreaks.

Conclusion

Prioritisation of pSIAs must balance outbreak risk against implementation costs, ideally favouring the smallest manageable outbreak risk compatible with elimination. We infer that there are few short-term risks due to population immunity from RI, but without pSIAs, long-term risks accumulate and can result in outbreaks with potential for international spread. We do not consider the costs of further delaying the eradication timeline or societal implications of outbreaks, both of which emphasise the need for pSIAs.

Introduction

In 2019, the African Region was certified free from endogenous transmission of wild poliovirus (WPV), with the last case reported in Nigeria in July 2014 (1). However, in early 2022, Malawi and Mozambique reported WPV serotype 1 (WPV1) cases, linked to ongoing circulation in Pakistan (2, 3). The geographic distribution and genetic linkage of these WPV1 cases suggest missed transmission of an unknown geographic extent (2). These WPV1 cases highlight the importance of ensuring high and homogeneous levels of population immunity despite decreasing global incidence and elimination within the African continent (4). Historically, WPV1 outbreaks in Africa had a varied epidemiology—ranging in size from a few cases to explosive outbreaks that resulted in international spread and wide-scale outbreak responses. This variation can be attributed to population exposure to circulating virus and serotype-specific population immunity against poliovirus, among other factors (5).

Since 2016, the recommended routine immunisation schedule is with bivalent oral polio vaccine (bOPV) and at least 1 dose of the inactivated polio vaccine (IPV) to provide population immunity. However, variable routine immunisation coverage leads to differential population immunity in different countries. The joint assessment of immunization coverage by UNICEF and WHO (WUENIC) estimated in 2021 routine vaccine coverage in Malawi and Mozambique was 93% and 61%, respectively. Also, low levels of population immunity persist in other areas, such as Liberia, where a recent study demonstrated poliovirus antibody seroprevalence was only 60% against serotype 1 (6). Further, the number of circulating vaccine derived poliovirus serotype 1 (cVDPV1) outbreaks has increased since 2022, suggesting that population immunity against this serotype has declined in some areas despite a rise in IPV coverage (which is expected to reduce the incidence of paralysis but has limited impact on transmission) (7).

The Global Polio Eradication Initiative (GPEI) is responsible for the coordination of activities to support polio eradication. The activities include surveillance for acute flaccid paralysis (AFP) caused by polio infection, environmental surveillance (ES) for poliovirus, and providing polio vaccinations through both supplementary immunisation activities (SIAs) and routine immunisation (through the Expanded Programme for Immunisation (EPI)). The SIAs aim to vaccinate hard-to-reach children otherwise missed by RI (8). Despite an annual expenditure of hundreds of millions of USD, decision makers within polio eradication often must make difficult decisions in resource allocation. The global budget has remained relatively stable over the last decade with noted decreases in the last three years. Alongside, the frequency of preventative supplementary immunisation activities (pSIAs), which aim to strengthen population immunity and prevent the frequency and impact of outbreaks has decreased in almost all countries in Africa since 2017 (Figure 1) (9). However, the annual costs for vaccination and programmatic activities such as antiviral development and vaccine stockpiling carried out by GPEI have increased considerably (10, 11).

Figure 1. Historical pSIAs and RI coverage for 30 African countries (A) Average number of pSIAs per year from 2013-2017 and 2018-2022 and (B) year of last pSIA and WUENIC estimates of DTP3 coverage. Preventative SIAs were defined as either a national or subnational immunisation day (NID, SNID) with bOPV (or tOPV pre-2016) and did not occur within 365 days of a WPV1 or VDPV1 detection, to distinguish historic pSIAs from oSIAs. Any SIAs that occurred within 365 days of a WPV1 or VDPV1 outbreak were not included in the pSIA count. CAR= Central African Republic; DRC = Democratic Republic of the Congo

Since the World Health Assembly's 1988 resolution to eradicate polio by the year 2000, economic analyses have informed strategies to progress towards this goal. These studies inferred that polio eradication is cost-effective, assuming all vaccination stops following certification of global

eradication (12-14). Together, polio vaccination and achieving global polio eradication are estimated
to save approximately 40 million DALYs by 2050 (15). However, existing economic analyses do not
consider the cost-effectiveness of specific operational decisions for preventative SIAs (pSIAs) and
outbreak response SIAs (oSIAs) given a shrinking budget (16) or are outdated and apply to limited
geographies (17). The distinction between pSIAs and oSIAs is important because pSIAs and oSIAs
have different strategic goals and funding approaches, and programmatic decisions between
vaccination strategies are required imminently, not just in the future context of OPV cessation. We
aim to compare strategies of differing frequency of pSIAs to that of RI and oSIAs to identify at what
levels of RI the risks of outbreaks and polio cases may outweigh the associated costs of
implementing pSIAs. This is an evidence gap identified by stakeholders that is important to address if
and when we approach the final few years of WPV1 transmission and start to consider OPV
cessation.
109

Methods

The GPEI works with a fixed annual budget for their operations, based on donations from
stakeholders. This budget is used to support the GPEI's objectives, including the global interruption
of poliovirus transmission. Operationally, the planned budget is further divided into planned SIAs
and outbreak response, and additional budget lines (not considered further in this study). Outbreak
response includes oSIAs, and planned SIAs are either preventive in nature or used in endemic
settings (not consider further here); pSIAs lower the risk of outbreaks due to the increase in
population immunity. oSIAs include vaccination following a detected case (or at least two separate
environmental detections that were collected from two different ES sites with no overlap in
catchment area or collected from the same ES site at least two months apart (18)) and aim to raise
population immunity in the affected areas to stop transmission. Operationally, pSIAs and oSIAs differ
both in the target populations for vaccination as well as the funding and planning for activities –
oSIAs must be implemented quickly and require more resource for rapid mobilisation. In this study,
we consider only the risks of WPV1 outbreaks in a polio-free setting; the bOPV and IPV used in RI
provide immunity against serotype 1. SIAs using bOPV also provide protection against infection as
well as protection against paralysis. Outbreaks of vaccine-derived poliovirus serotype 2 require
alternative vaccines and assumptions and are therefore beyond the scope of this analysis.
127
We evaluated different pSIA vaccination strategies in addition to RI and oSIAs for a population of 8
million children under five years of age, reflecting an average population size of a LMIC in Sub-
Saharan Africa. We estimated costs alongside health benefits and risks for each vaccination strategy
over a five-year time-period to align with the current GPEI timeline for polio eradication. Model
outputs from each strategy include: probability of an outbreak, estimated cases of paralytic
poliomyelitis and VAPP, number of outbreaks and estimated disability adjusted life years (DALYs)
associated with these cases (Figure 2).
135
136   ***Figure 2. Health benefits, risks, and costs of polio vaccination.*** *Schematic*
137   *showing the relationships between vaccination, population immunity, paralytic*
138   *poliomyelitis cases, adverse events and subsequent costs.*
139

Vaccination strategies

We explored three vaccination strategies (Table 1). We assume that vaccination via RI occurs in all
strategies and follows a sequential immunisation schedule that includes 3 doses of bOPV given orally
via drops in the mouth and 1 dose IPV administered intramuscularly or subcutaneously. The first two
comparator strategies (RI+oSIAs+annual pSIAs and RI+oSIAs+biannual pSIAs) maintain the same

assumptions about baseline RI coverage, but additionally include one pSIA either every year or every
two years, respectively, across time.

Table 1. Polio vaccination strategies. *RI = Routine Immunisation, oSIA = outbreak response*
supplementary immunisation activity, pSIA = preventative supplementary immunisation activity. An
outbreak response was only conducted if a simulation had at least 1 case of paralytic polio.

Vaccination strategy	Specification
Baseline strategy	RI + oSIAs *No preventative SIAs*
Comparator strategy 1	RI + oSIAs + *annual* pSIAs
Comparator strategy 2	RI + oSIAs + *biannual* pSIAs

In all strategies, we define an outbreak as at least one case of paralytic polio (18). An oSIA is
conducted in all simulations for each strategy where at least one case was detected within any 90-
day interval, in line with the median interval from notification to response as per the standard
operating procedures (SOP) for polio outbreaks (18, 19). oSIAs continue until all cases are stopped
over the five-year time horizon. We do not account for ES detections in this analysis. Children who
were not vaccinated via RI, can receive additional doses of bOPV vaccine through either pSIAs or
oSIAs. We assume that all SIAs reach 25% of children missed by RI, as prior evidence suggests that
SIA coverage can be variable across locations and preliminary analysis of higher coverage for zero-
dose children resulted in unrealistically high population immunity when compared to empirical data
(20, 21). Appendix Table 4 includes a sensitivity analysis of different SIA assumptions.

Model structure

We developed a stochastic non-linear mathematical model to simulate polio transmission dynamics,
whereby infectious individuals develop either asymptomatic or symptomatic infection, both of
which are assumed to be infectious. We specify vaccine induced immunity based on OPV and IPV
doses (Appendix Figure 1). In the model, children under five years of age are either susceptible, fully
vaccinated and protected from poliovirus infection, or have received an incomplete vaccination
series (less than 3 bOPV doses + 1 IPV dose). Each subsequent dose of vaccine corresponds to
additional protection and an opportunity for a child to seroconvert and be considered fully
protected from poliovirus infection. Further explanation of model compartments and transitions is in
Appendix Table 1.

Time horizon and model assumptions

The modelled time horizon is five years, in line with the current GPEI strategic plan (22). The WPV1
case to infection ratio was assumed to be 1:200, which is consistent with estimates for poliovirus
serotype 1 (23). For polio, the basic reproductive number, R_0, can vary substantially between
locations depending on sanitation and hygiene conditions (24). Here we have used an R_0 of 3 for the
primary analysis, supported by recent work exploring the parameter space of variable R_0 values in a
non-endemic setting in Africa for children under five years of age (20, 25), and explore other R_0
assumptions in a sensitivity analysis (Appendix Table 5). The proportion of children receiving three
bOPV doses through RI at 6, 10, and 14 weeks is varied between 25%-100%, and for simplicity is
assumed to occur upon entering the model. Children who have not seroconverted following
vaccination or did not receive three or more doses of bOPV remain susceptible to infection. The
proportion of children vaccinated with one dose of IPV is assumed to be equal to the third dose of
bOPV RI coverage, which is in line with WUENIC data (26) (Appendix Figure 2). As of 2020, WHO
recommends an additional IPV dose as part of RI (27), however, the number of countries that have
introduced IPV2 as of 2023 are limited, therefore, we only consider IPV1 in our model.

We assume a fully mixed population of children under five years of age with no heterogeneity in the
probability of a child being vaccinated in an SIA or in transmission of poliovirus. Simulations are run
for 50 years before virus introduction, then one infection was introduced into the population at the
start of the simulation and further importations were assumed to occur randomly at a rate of two
importations per year. Different importation rates are reported in a sensitivity analysis in Appendix
Table 6. The model was repeated for 1,000 stochastic simulations and run for five years. All models
were run using the R package SimInf (28).

Probability of an outbreak occurring

For each vaccination strategy, the probability of an outbreak was calculated using the proportion of
stochastic simulations that resulted in at least one case of paralytic polio (i.e., a polio AFP case) over
the entire modelled time horizon. The probability of an outbreak is comparable to maintaining
elimination in an infection-free setting such as most African countries. This definition is not directly
comparable towards the WHO criteria for elimination status (29), but is useful for understanding
outbreak risk. We do not include the specific criteria for elimination status as modelling of the
processes that lead towards achieving eradication are beyond the scope of this analysis.

DALYs

DALYs were calculated for each strategy assuming that in LMICs, the average discounted lifetime
DALYs associated with one paralytic poliomyelitis case, with no age-weighting is 14 DALYs per
paralytic case of polio (30), assuming that one in 200 infections leads to irreversible paralysis and
among those paralysed, 5-10% die when respiratory muscles become paralysed (31) and long-term
mortality is approximately 20% higher in paralytic polio cases than the general population (32). The
proportional contribution of YLD (years lost to disability) and YLL (years of life lost due to premature
mortality) assuming 14 DALYs on average per case is 60% YLDs on average per case in addition to
40% YLLs on average per case (33). We assume no international transmission during outbreaks for
calculations of DALYs.

Health and economic outcomes

Incremental costs and incremental outcomes (in terms of DALYs and outbreaks averted) were
estimated under each pSIA vaccination strategy in comparison to the baseline comparator
(RI+oSIAs). We used a 3% discount rate for costs and 0% for health with no age-weighting, with
other discounting assumptions explored in Appendix Figure 5. The total cost for each comparator
strategy included baseline RI vaccine costs for both bOPV and IPV. We present both a health system
(excludes GPEI costs), and GPEI perspective (includes only GPEI costs) for specification of costs, but
do not include indirect costs of vaccination, such as opportunity costs of time spent for vaccination.

Vaccine costs

Vaccine costs per dose for bOPV and IPV in Gavi supported countries were obtained from the latest
UNICEF update in USD 2023, costing on average $0.18 and $2.00, respectively (34, 35) and
administrative costs associated with polio RI were obtained from costs used in previous research
(36), Appendix Table 3. All costs have been adjusted to 2023 USD. GPEI estimates OPV wastage
ranges from 5-15% in SIAs and wastage is lower in SIAs than in RI (37). Estimates of IPV wastage vary
between 5-20% in the literature (38) and country-specific analyses from Nigeria and The Gambia
suggest IPV wastage can even be higher in practice, exceeding 20% for IPV wastage in some
instances (39, 40). Accordingly, the main analysis assumes an average of 10% wastage for OPV in
SIAs, 15% for OPV in RI and 15% for IPV, but further wastage assumptions are explored in a
sensitivity analysis (Appendix Figure 6).

Data and analysis on SIAs in Africa

The Polio Information System (POLIS) (41) was used to obtain SIA data for 2013-2022 and further analysis was done to distinguish preventative campaigns from outbreak response campaigns. We assumed that pSIAs included National or Subnational Immunisation Days (NID, SNID) with bOPV (or trivalent OPV (tOPV) pre-2016) that did not occur within 365 days of a WPV1 or VDPV1 detection by AFP surveillance or ES. Any SIA that occurred within this interval was not included in the overall count of pSIAs. Country level historic pSIA number and year of last pSIA were presented alongside WUENIC estimates of Diphtheria Tetanus Toxoid and Pertussis vaccine third dose (DTP3) coverage (42), an indicator of vaccination via RI as the DTP vaccine is administered concurrently with OPV in the routine immunisation series.

SIA costs

The cost per child for both pSIAs and oSIAs was obtained from GPEI data (Appendix Table 7), for which an average across countries within the African Regional Office (AFRO) was used. Historic cost data for oSIAs has been less readily available, therefore, we assume oSIAs cost 2.0 times the cost of a pSIA, as per GPEI estimated costs for 2023. We also explore a range of proportional costs between pSIAs and oSIAs that align with previous assumptions in polio economic modelling (40) (Appendix Figures 7 and 8). Due to the stochasticity of epidemic simulations, there is considerable variability in outcomes for identical assumptions. The number of outbreaks, which affects total estimated costs, is variable and contributes to the variability in expected costs across all strategies (Appendix Figure 3).

Adverse events

The expected risk of adverse events, such as vaccine associated paralytic polio (VAPP) in countries using OPV is 1 case of VAPP per 0.9 million doses of bOPV administered and declines with subsequent doses (43). VAPP rate across settings with and without SIAs has been shown to vary (44), although this variation is not well understood or documented. OPV administration following IPV receipt is associated with a further risk reduction of VAPP (43) and is accounted for in model assumptions (Appendix Table 3). VAPP cases contribute to overall risk and costs for each strategy. A VAPP case was considered equivalent to a case of wild-acquired paralytic polio for calculation of the expected costs of VAPP.

Perspectives

Budget and incremental cost results are analysed from both the GPEI and health system perspectives. The GPEI costs include those associated with SIAs and IPV in RI. Treatment costs associated with paralytic polio (including VAPP) are paid for by the country and are included as health system costs. The probability of an outbreak under each vaccination strategy is analysed from both the global perspective for polio eradication and the country health system perspective.

Ethics

Ethical approval for this project was received from the London School of Hygiene and Tropical Medicine, project ID 15873.

Results

On average across all simulations, the expected number of WPV1 cases over five years is greatest in the baseline strategy (RI+oSIAs) and least in the annual pSIA strategy (RI+oSIAs+annual pSIAs) (Figure 3A). Under the base case assumptions (including R_0 and proportion of zero dose children reached by SIAs) annual pSIAs achieve and maintain >80% probability of no outbreaks when baseline RI coverage is 50% (Figure 3B). The biannual pSIA strategy (RI+oSIAs+biannual pSIAs) achieves >80% probability of no outbreaks when RI is above 65% and the baseline strategy requires ≥75% RI coverage to achieve >80% probability of no outbreaks. The above results are from simulations assuming two importations per year during the five-year period of simulations. The frequency of importations was varied in a sensitivity analysis (Appendix Table 6); when RI was 50% or lower, the increased frequency of importation had a large increase on the probability of outbreaks, but for simulations with an RI ≥75%, the probability of no outbreaks remained above 80%.

Figure 3. Estimated number of paralytic polio cases and probability that no outbreaks occur over the five-year modelled time horizon, (A) number of expected paralytic polio cases (presenting as a polio AFP case) and (B) probability of no outbreaks occurring across all vaccination strategies. Outbreak probability was based on 1,000 simulations per vaccination strategy. The red dashed line corresponds to 80% probability that no outbreaks occur.

Expected VAPP cases over five years are greatest in the annual pSIA strategy, corresponding to the strategy with the greatest number of vaccine doses administered, but fewest expected WPV cases over five-years. Estimated costs from both the health system and the GPEI perspectives are shown in Figure 4. From a health system perspective, estimated costs are lowest in the annual pSIA strategy due to fewer polio cases. From the GPEI perspective, estimated costs of the annual pSIA strategy always exceed the cost of the baseline strategy due to the number of preventative campaigns.

Figure 4. Health system and GPEI estimated total costs over five years. The size of the circle is proportional to the mean number of expected paralytic polio cases across all model simulations. The solid circles correspond to >80% probability the strategy had no outbreaks over a five-year period.

Incremental costs per DALY averted by both pSIA strategies in comparison to the baseline strategy are shown in Figure 5 with further explanation of the quadrants in a cost-effectiveness plane (Appendix Figure 9). For non-GPEI costs (Figure 5A), when RI coverage exceeds 50%, points for both pSIA strategies are densely clustered in the bottom-right quadrant highlighting approaches with relatively lower costs and more DALYs averted. Assuming herd immunity is calculated by $(1 − 1/R_0)$, when RI coverage exceeds 65%, the point when herd immunity is achieved in this simple homogenously mixed model, the annual pSIA strategy becomes less efficient in averting DALYS. For GPEI costs (Figure 5B), which do not account for treatment or RI bOPV costs, at all levels of RI coverage, the annual pSIA strategy incurs relatively higher costs in comparison to biannual pSIAs, with densely clustered points in top-right quadrant or along the positive y-axis. The cost effectiveness planes are presented in Appendix Figure 5 when a discount rate of 3% has been considered for both health and costs.

Figure 5. Incremental costs per DALY averted. Incremental costs and DALYs averted under the annual pSIA (RI+oSIAs+annual pSIAs) and biannual pSIA

Incremental costs per outbreak averted capture the global perspective of polio eradication as a
single outbreak under any vaccination strategy has implications for global polio eradication. This
analysis has shown a greater number of outbreaks, albeit substantially smaller size, in the annual
pSIA strategy in comparison to the baseline strategy when RI coverage is below 45% (under base-
case assumptions), but the health system incremental costs are kept down by the small number of
polio cases (Figure 6A). From the health system perspective, when RI coverage ranges from 45-70%,
the annual pSIA strategy averts many outbreaks in comparison to the baseline strategy, but above
70% RI coverage, the annual pSIA strategy does not avert many outbreaks and is a costly approach.
Both pSIA strategies avert few to no outbreaks when RI coverage exceeds 75%. Under the GPEI
perspective, the annual and biannual pSIA strategies avert a similar number of outbreaks above 60%
RI coverage, but annual pSIAs have higher incremental costs per outbreak averted, corresponding to
the greater number of SIAs conducted (Figure 6B).

Figure 6. Incremental costs per outbreak averted. *Incremental costs and*
outbreak averted under the annual pSIA (RI+oSIAs+annual pSIAs) and biannual
pSIA (RI+oSIAs+biannual pSIAs) strategies in comparison to the baseline strategy
(RI+oSIAs) assuming a 3% discount rate for costs and 0% discount rate for health.
The points correspond to 1,000 model simulations. Here, it is assumed that the
cost per child in an oSIA is two times the cost per child in a pSIA. An outbreak is
defined as at least one case of paralytic polio. It is important to note that this
figure does not suggest that the RI+oSIAs strategy is better performing at low RI
coverage levels. Instead, negative (or no) outbreaks averted by both pSIA
strategies at low levels of RI coverage correspond to two phenomena: (i) the
RI+oSIA strategy has larger explosive outbreaks when RI coverage is low, so the
susceptible population is depleted quicker, whilst the pSIA strategies have much
smaller, albeit more frequent, outbreaks when RI coverage is low.

The implications for decision making are outlined in Table 2. Focussing on estimated risk if historic
pSIAs have been regularly conducted, when RI coverage falls below 40%, the annual pSIA strategy
averts many cases, so removal of pSIAs entirely would create substantial risk. However, pSIAs do not
avert many outbreaks. It is important to note that this finding does not suggest that oSIAs are
necessarily better performing at these coverage levels. Instead, the baseline strategy instead leads
to quicker depletion of susceptibles or, results in more frequent vaccination via outbreak response
following *at least one* paralytic case, whereas the pSIA strategies only vaccinate the target
population annually or biannually. Countries with 50-90% RI coverage and historic pSIAs have a
higher probability of no outbreaks occurring, with the outbreak risk reducing with higher RI.
However, the risk of an outbreak is not removed entirely until the probability of no outbreaks
reaches 100% (when WPV1 transmission is interrupted globally). All strategies require >95%
coverage for 100% probability that no outbreaks occur, however, above 80% RI coverage, outbreak
probability is low and, if an outbreak does occur, the expected outbreak size is very small. Figure 1
further summarises country-level historic pSIAs alongside the date since the last pSIA and current
estimates of RI coverage. In summary, countries with <60% RI coverage have conducted more recent
and frequent pSIAs and countries with >90% RI coverage have, on average, conducted fewer historic
pSIAs per year. However, a few countries that fall below 80% RI coverage have not conducted a pSIA
in over a decade.

Table 2. Policy implications of polio vaccination strategies. *Implications for adopting each*
vaccination strategy based on a country's RI coverage and historic SIA schedule according to GPEI SIA
data (column 2). Average values across all model simulations are references for DALYs averted and
outbreaks averted while expected paralytic polio cases are conditional averages amongst simulations
that resulted in at least one case. The probability of no outbreaks occurring is obtained from the
proportion of model simulations (out of 1,000 simulations) that resulted in zero paralytic cases and
correspond to important implications for polio eradication.

RI coverage	Did country conduct pSIAs in the past?	DALYs averted*?	Outbreaks averted*?	Estimated risk with annual pSIAs	Estimated risk with biannual pSIAs	Estimated risk relying on oSIAs only†	Implications for decision making
				Average number of polio cases if an outbreak occurs‡ (Probability <u>no</u> outbreaks occur)			
35%	∟	∟	X	5 (35%)	894 (3%)	15,753 (0%)	pSIA removal would have high risks and consequences
50%	∟	∟	∟	1 (80%)	4 (33%)	3,549 (1%)	Removal of pSIAs altogether could lead to a high risk of outbreaks in subsequent years
55%	∟	∟	∟	1 (89%)	2 (60%)	617 (3%)	
60%	∟	∟	∟	1 (92%)	1 (76%)	34 (11%)	
65%	∟	∟	∟	1 (94%)	1 (85%)	3 (37%)	
70%	∟	X	∟	1 (96%)	1 (91%)	1 (72%)	
80%	∟	X	∟	1 (97%)	1 (96%)	1 (92%)	Reducing the frequency of pSIAs could still maintain a low risk of large outbreaks
90%	∟	X	∟	1 (99%)	1 (99%)	1 (98%)	
100%	∟	X	X	0 (100%)	0 (100%)	0 (100%)	Even if pSIAs are removed, there is low to no risk of outbreaks

385 * Estimated DALYS and outbreaks averted are obtained from the annual pSIA strategy
386 † Estimated risk is obtained from the oSIA strategy assuming no pSIAs occur
387 ‡ Expected paralytic polio cases were calculated by taking the average number of paralytic polio cases across all stochastic model
388 simulations that had <u>at least one</u> case, so this is the average number of polio cases if an outbreak occurs

Discussion

The key messages of our study are: (i) with higher RI coverage of three doses of bOPV and 1 dose IPV, the probability of outbreaks reduces considerably – under our model assumptions, outbreak size and risk are minimal when RI is above 65%. At lower values of RI, specific vaccination strategies should be considered; (ii) in scenarios where RI is below 50%, annual pSIAs limit outbreak risk and result in high cost, an annual pSIA strategy when RI is 50-80% results in a low probability of outbreaks and continues to be costly, (iii) biannual pSIAs require higher baseline RI coverage (65%) to achieve an equivalent probability of no outbreaks, but the reduction in cost has few negative outcomes; (iv) a strategy with only RI and oSIAs implicitly accepts some level of outbreak risk, but if RI is above 70-80% the risk of outbreaks is considerably less than other settings where RI is below this value.

At low levels (<50%) of RI coverage, the probability of outbreaks is high (99%) following an importation if pSIAs do not occur at any frequency. Further, the incremental costs show a clear benefit of a pSIA strategy, outweighing the increased number of expected VAPP cases and cost of vaccination. Countries such as Madagascar or South Sudan where 55% and 49% of the population under five years of age, respectively, are vaccinated with three or more DTP doses (used to approximate polio RI), have many subpopulations that could benefit from regular pSIAs. Indeed, between 2018-2022 South Sudan has implemented annual pSIAs, and Madagascar has implemented oSIAs in response to cVDPV1. Some countries in Africa have many subpopulations that fall towards the middle or upper range of analysed RI coverage estimates. In both Burkina Faso and Sierra Leone, for example, future SIAs would not seem necessary as routine immunisation coverage of three doses of bOPV and 1 dose IPV exceeds 90% without reliance on historic pSIAs, unless there are subpopulations with substantially lower coverage. The proportion of the population in Malawi that has received three doses of bOPV peaked in 2011, but has been unstable since, falling to 83% in 2016 (26), with no historic reliance on pSIAs. WPV1 was reported in Malawi and local transmission and international spread has since been reported. In our analysis, RI exceeding 80% would be unlikely to result in an outbreak, but we have not included heterogeneity in immunity in the simulations, likely over-estimating the probability of no outbreaks. It may also be that the frequency of WPV1 importation was under-estimated for the Malawi setting, and close monitoring of migration routes from endemic countries remains an important component of preventing polio outbreaks. Whilst WHO now recommends two doses of IPV (IPV2) as part of RI, many countries in Africa have not yet implemented this strategy. As the number of countries adopting this policy decision increases, future studies will be needed to analyse the impact of IPV2.

Our study has health policy implications for decision making on global polio eradication. Despite the high costs, the ability of pSIAs to decrease the probability of an outbreak is an important consideration for polio eradication. Should the GPEI adopt an annual pSIA strategy irrespective of estimated RI and importations? Annual pSIAs that include all children under five years of age in LMICs in Sub-Saharan Africa would consume most of the GPEI annual budget for activities and would be an inefficient use of funds, potentially reducing funds for other activities (surveillance, vaccination against other serotypes). However, prioritising SIAs in countries with low RI and perceived risk of introductions is a necessary compromise to which GPEI already adheres, and here we provide a framework to support decision making. When considering the vaccination strategies based on perceived outbreak risk (as described here) and the fixed budget for these activities, we must balance making the best use of limited resources while limiting WPV1 outbreak risk. The GPEI is reliant on a combination of country donations and philanthropic contributions to support activities, and budget has remained the same or reduced since 2010 and reduced further since the COVID-19 pandemic (45, 46). From the perspective of donors committing to polio eradication, reducing commitment prior to WPV1 interruption is a false saving: for every outbreak experienced, the timeline to WPV1 interruption become longer and the costs of eradication grows (22). Renewed

commitment by donors was requested in 2022 (47) in light of the 2022-2026 Strategic Plan and
these commitments remain essential to resource the activities needed to meet the objectives of
polio eradication, including interrupting WPV transmission.

Our study has limitations. The main results assume a R_0 of 3 in a homogenous population (both with
respect to transmission and population immunity), a frequency of poliovirus introduction of two
infections per year and SIAs reaching 25% of children missed by routine vaccination. These
assumptions are a necessary simplification of real-world complexity. Several assumptions for R_0,
poliovirus introductions, and SIA reach were considered in a univariate sensitivity analysis. Should
SIAs reach up to 50% of zero-dose children, the impact of SIAs on reducing outbreak risk is further
reduced, consequently for the same costs a better outcome is achieved. Assuming a higher R_0 and
increasing the frequency of importations would also increase the outbreak risk. One of the most
uncertain inputs of the analysis is importation rate: as poliovirus infection is typically asymptomatic
this is not directly observable (although ES has frequently illustrated introductions in the absence of
AFP cases (48)), and due to the changing epidemiology of polio globally, the importation rate will
vary in time. We have not considered heterogeneity in the population. If there are pockets of the
target population with higher rates of transmission and/or lower vaccination coverage, then the
probability of an outbreak occurring would be higher. Additional discussion on the model framework
is presented in Appendix Section 1, page 5.

Further, the model structure assumes that children are vaccinated when they are born into the
population, not accounting for infection prior to vaccination. Because the average age of infection
with poliomyelitis occurs after the vaccination schedule of bOPV given at 6, 10 and 14 weeks of age
alongside the effect of maternal antibodies, this limitation is unlikely to impact the expected number
of AFP cases. However, because the size of the birth cohort vaccinated may be overestimated in the
absence of infant mortality assumptions, the costs of RI may be over-estimated here. Additionally,
we do not include indirect costs of vaccination, such as opportunity costs of time spent for
vaccination, which may result in an underestimation of costs across all strategies. The indirect
benefits of these additional interventions (such as productivity increases from averting polio cases)
are also not quantified in the costs and benefits of SIAs described here, therefore the deaths and
burden averted are polio specific and exclude indirect economic effects, in contrast to some other
analyses. Since 2020, pSIAs have been reduced or stopped altogether across most LMICs in Sub-
Saharan Africa, meaning 2023 represents the fourth consecutive year of diminished immunity. This
analysis does not directly assess the consequences of this diminished immunity but presents
baseline research that can be used to inform these future research questions. We do not consider
the costs of further delaying the eradication timeline through outbreaks, or the societal implications
of outbreaks on polio eradication, both of which may further emphasise the need to implement
pSIAs even when the outbreak risks are small. By limiting our analysis to a five-year time horizon, we
underestimate the benefits of SIAs (particularly pSIAs) as they will increase the likelihood of
eradication, meaning that control efforts after eradication can be scaled back (though not stopped).
Further, the pSIA health system costs only consider the geographical remit stated in the model and
ignores the potential for further international spread. International spread would be far more likely
with larger outbreaks, consequently the health system costs are under-estimated.

In settings that have not recently experienced a WPV1 outbreak, it is challenging to decide between
responding to short-term versus long-term risks, especially with a fixed budget. From the global
perspective striving to achieve polio eradication, investing in pSIAs results in a greater probability of
polio elimination, but still require justification in a pragmatic environment of finite resources.
Therefore, future funding provisions and advocacy for the importance of pSIAs rely on research that
communicates risk and burden in their absence. Here, we communicate the utility of different
vaccination strategies and critically assess the risks associated with adopting different strategies

given baseline RI coverage. Decisions made solely based on fixed budget, cost-effectiveness or burden reduction may not fully capture all consequences or benefits associated with adopting a particular vaccination strategy. Urgently, as importations of WPV remain a threat to the AFRO region, this analysis serves as an invaluable tool to estimate risk and plan vaccination activities across a range of settings.

Authors' contributions
MA and KMOR conceptualised the study. MA curated the data, conducted the modelling analysis, prepared visualisations of results and wrote the original draft. MA, KMOR and KA independently validated the data and results. All authors contributed to reviewing results and editing of the manuscript and have approved the final version. The authors alone are responsible for the views expressed in this article and they do not necessarily present the decisions, policy or views of their affiliated organisations.

Acknowledgements

This project was funded by the Bill and Melinda Gates Foundation (BMGF), OPP1191821. We would like to thank Rachel M. Burke (BMGF), Grace Macklin (WHO), John Edmunds (LSHTM), Anna Vassall (LSHTM), Stephan Widgren (SimInf), and others in the GPEI Subgroup for Analysis and Modelling for their time and feedback over the course of this project.

Declaration of interests
The authors have no conflicts of interest to declare.

Data sharing
Simulation code used in the analysis and sample model outputs can be accessed at: https://github.com/mauzenbergs/polio_econ.

Acronyms
Supplementary Immunisation Activities (SIAs), outbreak response SIAs (oSIAs), planned preventative SIAs (pSIAs), Routine Immunisation (RI), Global Polio Eradication Initiative (GPEI), bOPV (bivalent oral polio vaccine), IPV (inactivated polio vaccine), circulating vaccine derived poliovirus (cVDPV), vaccine associated paralytic polio (VAPP), disability adjusted life years (DALYs), wild poliovirus serotype 1 (WPV1), WHO (World Health Organization), The United Nations Children's Fund (UNICEF), WUENIC (WHO and UNICEF estimates of national immunization coverage), acute flaccid paralysis (AFP), environmental surveillance (ES), African Regional Office (AFRO), Bill and Melinda Gates Foundation (BMGF), low- or middle-income country (LMIC), London School of Hygiene and Tropical Medicine (LSHTM), standard operating procedures (SOP).

References

1.   Etsano A, Gunnala R, Shuaib F, Damisa E, Mkanda P, Ticha JM, et al. Progress toward
poliomyelitis eradication—Nigeria, January 2014–July 2015. Morbidity and Mortality Weekly Report.
2015;64(32):878.

2.   Malawi Ministry of Health, Global Polio Eradication Initiative, Davlantes E. Notes from the Field:
Initial Outbreak Response Activity Following Wild Poliovirus Type 1 Detection—Malawi, February 2022.
Morbidity and Mortality Weekly Report. 2022;71(23):776-7.

3.   Rachlin A, Patel JC, Burns CC, Jorba J, Tallis G, O'Leary A, et al. Progress Toward Polio Eradication
- Worldwide, January 2020-April 2022. MMWR Morb Mortal Wkly Rep. 2022;71(19):650-5.

4.   McCarthy K, Howard W, Yousif M, Moonsamy S, Suchard M. The show is not over–wild-type
polio in Malawi is a wake up call and an opportunity for elimination efforts. International Journal of
Infectious Diseases. 2022.

5.   O'Reilly KM, Lamoureux C, Molodecky NA, Lyons H, Grassly NC, Tallis G. An assessment of the
geographical risks of wild and vaccine-derived poliomyelitis outbreaks in Africa and Asia. BMC Infect Dis.
2017;17(1):367.

6.   Stephen B Kennedy GRM, Gloria Mason Ross. Poliovirus antibodies following two rounds of
campaigns with a type 2 novel oral poliovirus vaccine in Liberia: a clustered, population-based
seroprevalence survey. Lancet Global Health. 2023.

7.   Franco-Paredes C, Rodriguez-Morales AJ, Henao-Martínez AF, Carrasco P, Tuells J. The growing
threat of wild poliovirus 1 and vaccine-derived cases in the COVID-19 era. The Lancet Infectious Diseases.
2022;22(10):1412-4.

8.   Global Polio Eradication Initiative. Polio endgame strategy 2019-2023: eradication, integration,
certification and containment. World Health Organization; 2019.

9.   Voorman A, Lyons H, Bennette C, Kovacs S, Makam JK, F Vertefeuille J, et al. Analysis of
population immunity to poliovirus following cessation of trivalent oral polio vaccine. Vaccine. 2022.

10.   Global Polio Eradication Initiative. GPEI Multi-year Budget 2019-2023: Enabling GPEI to
Interrupt Poliovirus Transmission & Achieve Certification.

11.   Thompson KM, Kalkowska DA. An updated economic analysis of the Global Polio Eradication
Initiative. Risk Analysis. 2021;41(2):393-406.

12.   Thompson KM, Tebbens RJD. Eradication versus control for poliomyelitis: an economic analysis.
The Lancet. 2007;369(9570):1363-71.

13.   Zimmermann M, Hagedorn B, Lyons H. Projection of Costs of Polio Eradication Compared to
Permanent Control. The Journal of Infectious Diseases. 2019;221(4):561-5.

14.   Thompson KM, Kalkowska DA. Review of poliovirus modeling performed from 2000 to 2019 to
support global polio eradication. Expert Rev Vaccines. 2020;19(7):661-86.

15.   Khan MM, Ehreth J. Costs and benefits of polio eradication: a long-run global perspective.
Vaccine. 2003;21(7):702-5.

16.   Tebbens RJD, Hampton LM, Wassilak SGF, Pallansch MA, Cochi SL, Thompson KM. Maintenance
and Intensification of Bivalent Oral Poliovirus Vaccine Use Prior to its Coordinated Global Cessation. J
Vaccines Vaccin. 2016;7(5).

17.   Kalkowska DA, Duintjer Tebbens RJ, Thompson KM. Modeling Strategies to Increase Population
Immunity and Prevent Poliovirus Transmission in 2 High-Risk Areas in Northern India. The Journal of
Infectious Diseases. 2014;210(suppl_1):S398-S411.

18.   Global Polio Eradication Initiative. Standard operating procedures: responding to a poliovirus
event or outbreak 2022 [Version 4:[Available from: https://polioeradication.org/wp-
content/uploads/2022/04/Standard-Operating-Procedures-For-Responding-to-a-Poliovirus-Event-Or-
Outbreak-20220417_OBR_SOP_final_pre_pub_website.pdf.

19.   Blake IM, Chenoweth P, Okayasu H, Donnelly CA, Aylward RB, Grassly NC. Faster Detection of
Poliomyelitis Outbreaks to Support Polio Eradication. Emerg Infect Dis. 2016;22(3):449-56.

20.   Blake IM, Martin R, Goel A, Khetsuriani N, Everts J, Wolff C, et al. The role of older children and
adults in wild poliovirus transmission. Proceedings of the National Academy of Sciences.
2014;111(29):10604-9.

21.   Voorman A, Lyons HM. Measuring polio immunity to plan immunization activities. Vaccine.
2016;34(48):5946-52.
22.   World Health Organization, Global Polio Eradication Initiative. Polio eradication strategy 2022–
2026: delivering on a promise. 2021.
23.   Nathanson N, Kew OM. From emergence to eradication: the epidemiology of poliomyelitis
deconstructed. American journal of epidemiology. 2010;172(11):1213-29.
24.   Fine PEM, Carneiro IAM. Transmissibility and Persistence of Oral Polio Vaccine Viruses:
Implications for the Global Poliomyelitis Eradication Initiative. American Journal of Epidemiology.
1999;150(10):1001-21.
25.   McCarthy KA, Chabot-Couture G, Shuaib F. A spatial model of Wild Poliovirus Type 1 in Kano
State, Nigeria: calibration and assessment of elimination probability. BMC Infectious Diseases.
2016;16(1):521.
26.   World Health Organization. Poliomyelitis vaccination coverage.
27.   World Health Organization. Weekly Epidemiological Record, 2020, vol. 95, 22 [full issue].
2020;95(22):241-56.
28.   Widgren S, Bauer P, Eriksson R, Engblom S. SimInf: An R package for data-driven stochastic
disease spread simulations. arXiv preprint arXiv:160501421. 2016.
29.   Dowdle WR. The principles of disease elimination and eradication. Bulletin of the World Health
Organization. 1998;76(Suppl 2):22.
30.   Duintjer Tebbens RJ, Pallansch MA, Cochi SL, Wassilak SGF, Linkins J, Sutter RW, et al. Economic
analysis of the global polio eradication initiative. Vaccine. 2010;29(2):334-43.
31.   World Health Organization. Poliomyelitis   [Available from: https://www.who.int/news-
room/fact-sheets/detail/poliomyelitis.
32.   Nielsen NM, Rostgaard K, Juel K, Askgaard D, Aaby P. Long-term mortality after poliomyelitis.
Epidemiology. 2003:355-60.
33.   World Health Organization. The global burden of disease: 2004 update: World Health
Organization; 2008.
34.   UNICEF.   Inactivated   Polio   Vaccine   (IPV)   price   data   [Available   from:
https://www.unicef.org/supply/media/16266/file/IPV-vaccine-prices-14032023.pdf.
35.   UNICEF.   oral   polio   vaccine   price   data   2022   [Available   from:
https://www.unicef.org/supply/media/12671/file/oral-polio-vaccine-price-data-08062022.pdf.
36.   Kalkowska DA, Thompson KM. Health and economic outcomes associated with polio vaccine
policy options: 2019–2029. Risk Analysis. 2021;41(2):364-75.
37.   UNICEF. Guidance Note on Cold Chain Logistics and Vaccine Management During Polio
Supplementary Immunization Activities. UNICEF, New York, NY; 2015.
38.   Mvundura M, Hsu J-S, Frivold C, Kristensen D, Boyle S, Zehrung D, et al. Evaluating the cost per
child vaccinated with full versus fractional-dose inactivated poliovirus vaccine. Vaccine: X. 2019;2:100032.
39.   Wallace AS, Willis F, Nwaze E, Dieng B, Sipilanyambe N, Daniels D, et al. Vaccine wastage in
Nigeria: An assessment of wastage rates and related vaccinator knowledge, attitudes and practices.
Vaccine. 2017;35(48 Pt B):6751-8.
40.   Thompson KM, Kalkowska DA. Potential future use, costs, and value of poliovirus vaccines. Risk
Analysis. 2021;41(2):349-63.
41.   Global Polio Eradication Initiative. Polio Information System. 2023.
42.   World Health Organization. Diphtheria tetanus toxoid and pertussis (DTP) vaccination coverage.
43.   Ni L, Seward JF, Santibanez TA, Pallansch MA, Kew OM, Prevots DR, et al. Vaccine policy changes
and epidemiology of poliomyelitis in the United States. Jama. 2004;292(14):1696-701.
44.   Tebbens RJD, Pallansch MA, Kew OM, Cáceres VM, Jafari H, Cochi SL, et al. Risks of paralytic
disease due to wild or vaccine-derived poliovirus after eradication. Risk Analysis. 2006;26(6):1471-505.
45.   OECD.   The   0.7%   ODA/GNI   target   -   a   history   [Available   from:
https://www.oecd.org/dac/financing-sustainable-development/development-finance-
standards/the07odagnitarget-ahistory.htm.
46.   Philip Loft PB. The 0.7% aid target. 2022.

47.   Global Polio Eradication Initiative. GPEI calls for renewed commitments to achieve promise of a
polio-free world 2022 [Available from: https://polioeradication.org/news-post/global-polio-eradication-
initiative-calls-for-renewed-commitments-to-achieve-promise-of-a-polio-free-world/.
48.   Asghar H, Diop OM, Weldegebriel G, Malik F, Shetty S, El Bassioni L, et al. Environmental
surveillance for polioviruses in the Global Polio Eradication Initiative. The Journal of infectious diseases.
2014;210(suppl_1):S294-S303.

Country	DTP3 coverage 2021	Year of last pSIA
RI coverage <50%		
CAR	42%	2021
Somalia	42%	2021
Guinea	47%	2021
South Sudan	49%	2021
>50% RI coverage <80%		
Madagascar	55%	2019
Nigeria	56%	2021
Chad	58%	2021
Mozambique	61%	2011
DRC	65%	2021
Ethiopia	65%	2021
Cameroon	69%	2021
Côte d'Ivoire	76%	2018
Benin	76%	2018
Mali	77%	2019
>80% RI coverage <90%		
Gambia	82%	2014
Niger	82%	2019
Togo	83%	2014
Sudan	84%	2019
Senegal	85%	2016
South Africa	86%	2010
Zimbabwe	86%	2007
>90% RI coverage		
Burkina Faso	91%	2016
Kenya	91%	2019
Uganda	91%	2019
Zambia	91%	2009
Sierra Leone	92%	2019
Malawi	93%	2010
Burundi	94%	2011
Botswana	95%	2004
Ghana	98%	2015

A **Health system perspective**

B **GPEI perspective**

● RI + oSIAs + annual pSIAs ● RI + oSIAs + biannual pSIAs

Baseline comparator strategy is RI + oSIAs
3% discounting for costs, 0% discounting for health

A Health system perspective

B GPEI perspective

25% RI coverage · 30% RI coverage · 35% RI coverage · 40% RI coverage
45% RI coverage · 50% RI coverage · 55% RI coverage · 60% RI coverage
65% RI coverage · 70% RI coverage · 75% RI coverage · 80% RI coverage

Incremental costs (USD$2023)

Outbreaks averted over 5 years

● RI + oSIAs + annual pSIAs   ● RI + oSIAs + biannual pSIAs

Baseline comparator strategy is RI + oSIAs
3% discounting for costs, 0% discounting for health