Prevalence and Clinical Significance of Commonly Diagnosed Genetic Disorders in Preterm Infants

Selin S. Everett[a], Miles Bomback[b], Rakesh Sahni, MD[c], Ronald J. Wapner, MD[a], Veeral N. Tolia, MD[d,e,f], Reese H. Clark, MD[d], Alex Lyford, PhD[g], Thomas Hays, MD, PhD[c]

Affiliations: [a] Division of Maternal Fetal Medicine, Department of Obstetrics and Gynecology, Columbia University Irving Medical Center, New York City
[b] Feinberg School of Medicine, Northwestern University, Chicago [3]The MEDNAX Center for Research, Education, Quality and Safety, Sunrise, Florida
[c] Division of Neonatology, Department of Pediatrics, Columbia University Irving Medical Center, New York City
[d] The Pediatrix Center for Research, Education, Quality and Safety, Sunrise, Florida
[e] Division of Neonatology, Department of Pediatrics, Baylor University Medical Center, Dallas, Texas
[f] Pediatrix Medical Group, Dallas, Texas
[g] Department of Statistics, Middlebury College, Middlebury, Vermont

Address correspondence to: Thomas Hays, Division of Neonatology, Department of Pediatrics, Columbia University Irving Medical Center, 622 West 168th S, Suite PH 17W-303, New York, New York, 10032, th2712@cumc.columbia.edu

Short title: Genetic Disorders in Preterm Infants

Conflict of Interest Disclosures (includes financial disclosures): The authors have no example conflicts of interest to disclose.

Funding/Support: No funding was secured for this study. TH is supported by a *Thrasher Research Fund Early Career Award*.

Role of Funder/Sponsor (if any): The Thrasher Research Fund had no role in the design and conduct of the study.

Abbreviations: AKI (acute kidney injury), BPD (bronchopulmonary dysplasia), ICH (intracerebral hemorrhage), OR (odds ratio) NEC (necrotizing enterocolitis), ROP (retinopathy of prematurity)

Article Summary

Genetic disorders were found in 1.3% of preterm infants and at a higher rate (2.0%) in infants who died or developed severe morbidity.

What's Known on This Subject
Previous research described the prevalence and associated short-term morbidity and mortality of trisomy 13, 18, and 21 in preterm infants. The prevalence of other commonly diagnosed genetic disorders and associated short-term morbidity and mortality in preterm infants is unknown.

What This Study Adds
In a multicenter, retrospective cohort of 320,582 preterm (<34 weeks' gestation) infants, we found that 1.3% had genetic disorders diagnosed through standard care. Multiple disorders were associated with increased adjusted odds of morbidities or mortality prior to hospital discharge.

Contributors Statement Page

Selin S. Everett conceptualized and designed the study, conducted analyses, drafted the initial manuscript, and critically reviewed and revised the manuscript.

Dr. Thomas Hays conceptualized and designed the study, drafted the initial manuscript, and critically reviewed and revised the manuscript.

Miles Bomback conceptualized and designed the study and critically reviewed and revised the manuscript.

Drs. Veeral N. Tolia and Reese H. Clark coordinated and supervised data collection and critically reviewed and revised the manuscript.

Dr. Rakesh Sahni conceptualized and designed the study and critically reviewed and revised the manuscript.

Dr. Alex Lyford conducted analyses and critically reviewed and revised the manuscript.

Dr. Ronald J. Wapner reviewed and critically revised the manuscript.

All authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

Abstract

Background and Objectives: Preterm infants (<34 weeks' gestation) experience high rates of
morbidity and mortality before hospital discharge. Genetic disorders substantially contribute to
morbidity and mortality in related populations. The prevalence and clinical impact of genetic
disorders is unknown in this population. We sought to determine the prevalence of commonly
diagnosed genetic disorders in preterm infants, and to determine the association of disorders with
morbidity and mortality.
132
133   **Methods:** This was a retrospective multicenter cohort study of infants born from 23 to 33 weeks'
134   gestation between 2000 and 2020. Genetic disorders were abstracted from diagnoses present in
135   electronic health records. We excluded infants transferred from or to other health care facilities
136   prior to discharge or death when analyzing clinical outcomes. We determined the adjusted odds
137   of pre-discharge morbidity or mortality after adjusting for known risk factors.
138
139   **Results:** Of 320,582 infants, 4196 (1.3%) had genetic disorders. Infants with trisomy 13, 18, 21,
140   or cystic fibrosis had greater adjusted odds of severe morbidity or mortality. Of the 17,427
141   infants who died, 566 (3.2%) had genetic disorders. Of the 65,968 infants with a severe
142   morbidity, 1319 (2.0%) had genetic disorders.
143
Conclusions: Genetic disorders are prevalent in preterm infants, especially those with life-
threatening morbidities. Clinicians should consider genetic testing for preterm infants with
severe morbidity and maintain a higher index of suspicion for life-threatening morbidities in
preterm infants with genetic disorders. Prospective genomic research is needed to clarify the
prevalence of genetic disorders in this population, and the contribution of genetic disorders to
preterm birth and subsequent morbidity and mortality.
150

Introduction

Preterm infants, particularly those born before 34 weeks' gestation, experience high rates of

severe morbidity and mortality prior to discharge.[1-3] Multiple factors have been shown to

contribute to these outcomes including gestational age, birth weight, and social determinants and

health.[1-3] Genetic disorders have been shown to strongly contribute to morbidity and mortality in

related populations including full-term infants and children with critical illness.[4-7] Genetic

disorders are particularly prevalent in stillbirth and fetal disorders.[8-10] Despite this, there are

limited descriptions of the prevalence of genetic disorders in preterm infants. Boghossian et al.

described the prevalence, morbidity, and mortality of very low birthweight infants with common

aneuploidies.[11,12] However, no studies to our knowledge have provided a comprehensive

description of genetic disorders diagnosed in this population. We therefore sought to achieve the

following objectives: 1) determine the prevalence of commonly diagnosed genetic disorders in

preterm infants, and 2) determine the association of these genetic disorders with severe morbidity

and mortality in preterm infants.

We recently utilized the Pediatrix Clinical Data Warehouse (CDW) in related studies, to

investigate the association of congenital renal anomalies with genetic disorders in preterm

infants.[13] CDW contains detailed records of 20% of preterm infants born in the United States.

Given this utility and broad representation, we sought to employ a similar strategy in this study.

Methods

This study was approved by the institutional review board of Columbia University. This report

followed the guidelines for cohort studies as outlined by Strengthening the Reporting of

Observational Studies in Epidemiology (STROBE). This was a retrospective study using the

CDW multicenter cohort of infants born from 23 to 34 weeks' gestation.

Prevalence of commonly diagnosed genetic disorders in preterm infants.

Multiple genetic disorders, including common aneuploidies, are associated with fetal growth

restriction. Therefore, we used a gestational age-based cutoff rather than weight-based. We

extracted records from 409,704 infants discharged from 2000 to 2020. We excluded infants with

missing covariate data, as well as those transferred after birth (often because of complex illness)

and infants transferred prior to discharge (in which the clinical outcome is unknown) to limit

ascertainment biases.[15] After these exclusions, 320,582 preterm infants born < 34 weeks'

gestation remained. Data from NICU admission to discharge were collected, including clinician-

noted diagnosis of genetic disorders. This cohort did not undergo prospective, genome-wide

genetic testing. Genetic diagnoses were limited to those made during standard care.

Association of commonly diagnosed genetic disorders with severe morbidity and mortality in

preterm infants.

We collected infant characteristics (genetic diagnoses, gestational age, normalized birth weight

and categorization by Fenton criteria at 10th and 90th percentiles for gestational age and sex,[14]

maternal self-identified race, sex, mode of delivery, exposure to antenatal betamethasone,

intubation in the first 72 hours of life, year of discharge, and the presence of congenital

anomalies) and data regarding clinical outcomes (diagnoses of severe co-morbidities, mortality,

and disposition). Severe morbidities were defined as the following: acute kidney injury (AKI)

defined by the diagnosis of oliguria or anuria (ICD-9 788.5 or ICD-10: R34); severe intracranial

hemorrhage (ICH) defined as grade III or IV intraventricular hemorrhage by Papile criteria or

cystic periventricular leukomalacia;6 medically or surgically treated necrotizing enterocolitis

(NEC); severe bronchopulmonary dysplasia (BPD) defined as invasive mechanical ventilation at

36 weeks' postmenstrual age; severe retinopathy of prematurity (ROP) defined as need for any

medical or surgical intervention; culture-positive sepsis defined by any positive blood or urine

culture; or shock defined as administration of any vasopressor or inotrope.

We used logistic regression to model the relationship between genetic disorders and severe

morbidity or mortality. The following were covariates: year of discharge, gestational age, birth

weight (here as a Z-score normalized to gestational age and sex to account for continuous rather

than categorical effects), sex, maternal race, mode of delivery, intubation in the first 72 hours of

life, presence of a congenital anomaly, and exposure to antenatal betamethasone. We determined

crude and adjusted odds ratios (OR) after controlling for these covariates. OR were reported with

99.97% confidence intervals (determined as $1 - \frac{\alpha}{n}$) to correct for n comparisons of 9 clinical

outcomes in 22 categories of genetic disorders and accepting statistical significance at a

Bonferroni corrected, two-tailed α of 0.05. Analyses were made using RStudio (version

2022.02.0).[15]

Results

Prevalence of commonly diagnosed genetic disorders in preterm infants.

Genetic disorders were identified in 4,196 (1.31 %) of the 320,582 infants in this cohort that had

data for each of our covariates (**Table 1**). Of the 4,196 infants with a genetic diagnosis, 1927

(45.92%) were females and 2269 (54.08%) were males. The disorders included in the dataset

were comprised of 19 specific diagnoses, as well as unspecified aneuploidy, unspecified copy

number variant, and infants found to have multiple genetic disorders. Aneuploidies (trisomy 13,

18, 21, and 22, Klinefelter syndrome, Turner syndrome) were present in 1,863 infants, with

trisomy 21 accounting for more than half (973) of these cases. Copy number variants (13q

deletion, Cri-du-chat syndrome, DiGeorge syndrome) were present in 87 infants. Unspecified

aneuploidy and copy number variants were noted in 548 and 35 infants respectively. Eleven

distinct monogenic disorders were identified in 2,246 infants. Most (1,986) of these had

hematologic disorders. Thalassemia accounted for 1,174 cases. Sickle cell disease was identified

in 194 infants and cystic fibrosis in 165. Thirteen infants with multiple genetic disorders were

identified. We did not detect differences in the prevalence of classes of genetic disorders

diagnosed over time in this cohort (**Figure 1**).

Association of commonly diagnosed genetic disorders with severe morbidity and mortality in

preterm infants.

Following exclusions, 320,582 preterm infants remained for analysis of clinical outcomes.

Genetic disorders were prevalent in 4,196 (1.3%) of these individuals. The rate of mortality in

preterm infants without genetic disorders in this cohort, and following exclusions was 5.3%.

Genetic disorders were present in 566 (3.2%) of preterm infants who died, in 1,319 (2.0%) of

preterm infants who experienced any severe morbidity or mortality, in 3,630 (1.2%) of preterm

infants who survived to discharge, and in 2,664 (1.1%) of preterm infants who survived to

discharge free of severe morbidity. The mortality rate after exclusions for each genetic disorder

is provided in Table 1. The highest crude rates of mortality were found in cases of aneuploidy.

84.5 and 83.9% of preterm infants with trisomy 13 and 18 died before discharge. 14.7% of

preterm infants with trisomy 21 died before discharge. Mortality rates for preterm infants with

copy number variants varied from 12.5% in Cri-du-chat syndrome to 20.0% in cases of

unspecified copy number variants. Preterm infants with hematologic disorders had lower

observed rates of mortality, with the highest rates found in cases of sickle cell disease (2.6%) and

thalassemia (2.2%). Varying rates of mortality were noted in preterm infants with miscellaneous

monogenic disorders, including 9.7% mortality for infants with cystic fibrosis.

We then determined the crude and adjusted odds of severe morbidity and mortality for preterm

infants with diagnosed genetic disorders as compared to preterm infants without known genetic

disorders. The baseline characteristics for covariates used in these analyses are provided for

preterm infants with aneuploidies (**Supplemental Table 1**), copy number variants

(**Supplemental Table 2**), hematologic disorders (**Supplemental Table 3**), and miscellaneous

monogenic disorders (**Supplemental Table 4**). The absolute numbers of preterm infants

experiencing each morbidity and mortality for each diagnosed genetic disorder were also

determined (**Supplemental Table 5).**

Preterm infants with aneuploidies had significantly greater adjusted odds of several outcomes

(**Table 2**). Preterm infants with trisomy 13, 18, and 21 had greater adjusted odds of death.

Preterm infants with trisomy 13 also had greater adjusted odds of AKI. Preterm infants with

trisomy 21 had greater adjusted odds of AKI, BPD, sepsis, shock, and death. Preterm infants

with copy number variants and hematologic disorders were not found to have significantly

greater adjusted odds of morbidity or mortality (**Tables 3 and 4**). The only monogenic disorder

diagnosis associated with significantly increased adjusted odds of morbidity or mortality was

cystic fibrosis **(Table 5)**. Preterm infants with cystic fibrosis had significantly increased adjusted

odds of NEC, BPD, and ROP. Elevated risk of morbidities in preterm infants with

hypophosphatasia did not persist after adjustment for known risk factors.

Discussion

This study provides the most comprehensive determination to our knowledge of the prevalence

of genetic disorders in preterm infants and short-term risk of morbidity and mortality conferred

by genetic disorders in this population. We found that genetic disorders, consisting of

aneuploidies, copy number variants, and monogenic disorders, were diagnosed in 1.3% of

preterm infants. Trisomy 13, 18, and 21, and cystic fibrosis were associated with increased risk

of morbidity or mortality after adjustment for known risk factors. Rates of comorbidities for

preterm infants with trisomy 13 and 18 may be underestimated given mortality as a competing

outcome, particularly following redirection to palliative goals of care.

The overall prevalence of genetic disorders may reflect a contribution of genetic disease to

shorter gestation. Several of the most prevalent genetic disorders we analyzed are associated with

preterm birth including hemoglobinopathies,[16] cystic fibrosis,[17] and common aneuploidies.[18] This

results from a complex interplay of factors including disrupted placentation,[18] increased risk of

fetal growth restriction and stillbirth, closer monitoring of pregnancy, greater maternal age

contributing to aneuploidy and risk of preterm birth,[19] as well as idiopathic factors.

We found prevalence rates of trisomy 13 and 18 lower than previously described in preterm

infants,[12] however our analysis used a gestational age-based cutoff rather than birth weight.

Trisomy 13 and 18 cause fetal growth restriction. Given the same study population, a weight-

based cutoff will select for infants with gestational age > 33 weeks with these genetic disorders

and estimate a higher prevalence rate. Given these selection differences, it is difficult to compare

morbidity and mortality between our cohorts and previous descriptions.[11]

Interestingly, we found increased adjusted odds of NEC, BPD, and ROP in preterm infants with

cystic fibrosis. Infants with cystic fibrosis have been found to be born at earlier gestational age

and with lower birth weight.[17,20] However, after adjustment for these factors, we found an

increased risk of morbidity. Intestinal disease caused by meconium peritonitis and meconium

ileus may mimic or predispose infants to NEC.[21,22] BPD and ROP may be attributable to

disrupted lung disease, and exposure to oxygen.[23,24] Maternal cystic fibrosis also contributes to

neonatal morbidity,[25] and may account for a portion of observed outcomes. However, maternal

diagnoses were not available in our cohort.

This retrospective study has significant strengths and important weaknesses. It encompasses a

large number of infants, cared for in multiple institutions, and is highly generalizable. The depth

of data, including patient characteristics, diagnoses of genetic disorders, and co-morbidities

should enable clinicians to better counsel families and inform clinical decision-making in rare

presentations. It encompasses a broad period, during which genetic testing has undergone

considerable change. Interestingly, we did not detect differences in the broad classes of genetic

disorders diagnosed in preterm infants over time (**Figure 1**). However, our dataset lacked

information regarding genetic testing modalities, and negative results. There may be underlying,

dynamic changes that contributed to heterogeneity in the cohort with respect to year of birth. As

genome-wide testing becomes more common, rare and uncommon genetic disorders may be

diagnosed more frequently in this population.

This dataset lacks information regarding social determinants of health and maternal access to

care (including genetic testing). Multiple factors could have contributed to over- or

underestimation of the prevalence of genetic disorders. Preterm infants experience high rates of

morbidity and mortality, which may prompt testing for genetic disease in some settings. Genetic

disorders often present non-specifically in preterm infants,[4,26] compared to more complete and

well-known phenotypes found in term infants or pediatric or adult populations. Therefore,

genetic disorders may have gone undetected in this cohort. Unfortunately, we do not have data

regarding the extent of pre- and postnatal genetic testing, including negative results, in this

cohort. At this time, broad, prospective screening for neonatal genetic disorders is not routine.

Given the impact of genetic disorders on preterm morbidity and mortality in our study, there is a

crucial need for broad, prospective studies to establish this. At present, a number of studies are

underway to evaluate the potential of sequencing based newborn screening. In addition to

determining the technical advantage of obtaining a molecular diagnosis of a limited number of

rare biochemical disorders, this work may offer an opportunity to investigate the impact of a

more diverse group of genetic disorders on neonatal care.

In summary, using a multicenter retrospective cohort, we found that genetic disorders are present

in 1.3% of infants born before 34 weeks' gestation, and that trisomy 13, 18, 21, and cystic

fibrosis are associated with greater adjusted odds of death or severe morbidity. We determined

the prevalence of 19 distinct genetic disorders, and the crude and adjusted risk of mortality and

severe morbidities. These data may help clinicians provide more informed guidance to families

and lead to better clinical decision-making in rare presentations.[27] These data indicate the need

for a greater index of suspicion for multiple critical illnesses, such as sepsis in the setting of

trisomy 21 and NEC in the setting of cystic fibrosis. Given the trend towards greater adjusted

odds of multiple morbid outcomes, clinicians should also consider genetic evaluation when

caring for preterm infants with critical illnesses. Prospective, genome-wide screening is needed

to determine the true prevalence of genetic disorders in this high-risk population, the contribution

of genetic disorders to preterm birth, and the portion of disease in preterm infants attributable to

genetic disorders.

Acknowledgments
The authors wish to gratefully acknowledge Dr. Ali G. Gharavi (Columbia University Irving Medical Center) for their mentorship and thoughtful review of this project.

Table 1. Prevalence of genetic disorders in preterm infants and associated rates of mortality.

Genetic Disorder	n (Prevalence per 100,000 Preterm Infants)	n (Mortality Rate %)
Aneuploidy	1863 (581.1)	480/1863 (25.8)
Trisomy 13	110 (34.3)	93/110 (84.5)
Trisomy 18	155 (48.3)	130/155 (83.9)
Trisomy 21	973 (303.5)	143/973 (14.7)
Trisomy 22	1 (0.3)	1/1 (100)
Klinefelter syndrome	35 (10.9)	2/35 (5.7)
Turner syndrome	41 (12.8)	5/41 (12.2)
Unspecified Aneuploidy	548 (170.9)	106/548 (19.3)
Copy number variation disorder	87 (27.1)	14/87 (16.1)
13q deletion syndrome	7 (2.2)	1/7 (14.3)
Cri-du-chat syndrome	16 (5)	2/16 (12.5)
DiGeorge syndrome	29 (9)	4/29 (13.8)
Unspecified CNV	35 (10.9)	7/35 (20)
Monogenic disorder	2233 (696.5)	71/2233 (3.2)
Hematologic disorder	1973 (615.4)	39/1973 (2)
Thalassemia	1174 (366.2)	26/1174 (2.2)
Hemoglobin bart's	384 (119.8)	4/384 (1)
Sickle cell disease	194 (60.5)	5/194 (2.6)
G6PD deficiency	206 (64.3)	2/206 (1)
Hemophilia A	10 (3.1)	2/10 (20)
Hemophilia B	5 (1.6)	0/5 (0)
Miscellaneous	260 (81.1)	32/260 (12.3)
Cystic fibrosis	165 (51.5)	16/165 (9.7)
Fragile X syndrome	7 (2.2)	0/7 (0)
Hypophosphatasia	86 (26.8)	15/86 (17.4)
Fanconi Anemia	2 (0.6)	1/2 (50)
Multiple Disorders	13 (4.1)	1/13 (7.7)

Table 2. Odds of severe morbidity or mortality in preterm infants with aneuploidies[1]

Outcome	Trisomy 13 (n=110)		Trisomy 18 (n=155)		Trisomy 21 (n=973)		Klinefelter syndrome (n=35)		Turner syndrome (n=41)	
	Crude OR	Adjusted OR	Crude OR	Adjusted OR	Crude OR	Adjusted OR	Crude OR	Adjusted OR	Crude OR	Adjusted OR
Mortality or any severe morbidity	53.42 (15.61, 356.05)*	61.87 (16.15,437.72)*	29.47 (12.34, 90.49)	28.09 (10.56,93.09)*	2.04 (1.58, 2.62)	2.39 (1.75,3.26)*	1.66 (0.35, 6.18)	0.98 (0.16,4.75)	1.88 (0.48, 6.27)	2.43 (0.47,10.51)
Mortality	97.18 (39.3, 303.91)*	265.24 (93.12,916.46)*	92.37 (43.1, 231.52)	196.84 (81.82,539.66)*	3.06 (2.14, 4.26)	5.93 (3.83,8.95)*	1.08 (0.01, 8.2)	0.81 (0.01,8.14)	2.47 (0.22, 11.21)	4.17 (0.3,29.12)
AKI	6.67 (2.19, 16.27)*	4.17 (1.24,11.54)*	3.12 (0.84, 8.17)	1.91 (0.5,5.34)	3.39 (2.15, 5.07)	4.02 (2.43,6.35)*	2.06 (0.02, 15.82)	1.47 (0.01,13.72)	4.73 (0.43, 21.61)	6.91 (0.55,40.06)
ICH	—	—	0.32 (0, 2.17)	0.28 (0,1.99)	1.09 (0.56, 1.9)	1.29 (0.64,2.36)	1.5 (0.01, 11.48)	1.12 (0.01,10.61)	1.95 (0.06, 11.49)	2.2 (0.07,17.02)
NEC	1.33 (0.12, 5.36)	0.77 (0.07,3.31)	0.93 (0.09, 3.68)	0.58 (0.05,2.36)	1.55 (0.85, 2.57)	1.44 (0.78,2.45)	—	—	1.43 (0.01, 10.69)	1.55 (0.01,12.75)

BPD	5.37 (0.8, 18.46)	1.84 (0.26,7.04)	5.45 (1.19, 15.76)	1.78 (0.37,5.5)	4.1 (2.21, 6.91)	3.32 (1.71,5.94)*	—	—	—	—
ROP	—	—	—	—	0.34 (0.02, 1.44)	0.47 (0.03,2.24)	—	—	—	—
Sepsis	0.89 (0.17, 2.73)	0.49 (0.09,1.65)	0.76 (0.18, 2.11)	0.44 (0.1,1.28)	1.49 (1.02, 2.11)	1.56 (1.04,2.29)*	2.95 (0.47, 11.77)	2.31 (0.34,10.54)	2.8 (0.51, 10.24)	3.73 (0.6,16.07)
Shock	2.11 (0.74, 5)	1.08 (0.34,2.89)	2.38 (1.04, 4.85)	1.44 (0.58,3.25)	1.8 (1.28, 2.48)	2.49 (1.64,3.7)*	1.23 (0.08, 6.41)	0.89 (0.05,6.22)	1.63 (0.19, 6.86)	2.03 (0.2,11.97)

Abbreviations: AKI (acute kidney injury), BPD (bronchopulmonary dysplasia), ICH (intracerebral hemorrhage), OR (odds ratio) NEC (necrotizing enterocolitis), ROP (retinopathy of prematurity)

[1]odd ratio (99.97% confidence interval), * indicates two-tailed, Bonferroni-corrected p-value < 0.05

Table 3. Odds of severe morbidity or mortality in preterm infants with copy number variants[1]

Outcome	DiGeorge syndrome (n=29)		Cri-du-chat syndrome (n=16)	
	Crude OR	**Adjusted OR**	**Crude OR**	**Adjusted OR**
Mortality or any severe morbidity	2.56 (0.54, 10.68)	2.02 (0.29,12.01)	1.65 (0.13, 11.54)	1.64 (0.09,17.42)
Mortality	2.84 (0.17, 15.59)	2.59 (0.12,23.83)	2.54 (0.02, 24.38)	4.6 (0.04,73.4)
AKI	—	—	4.87 (0.05, 47.1)	5.67 (0.05,79.46)
ICH	—	—	—	—
NEC	—	—	3.99 (0.04, 38.63)	3.93 (0.04,44.41)
BPD	5.85 (0.06, 46.55)	2.62 (0.02,32.83)	—	—
ROP	—	—	—	—
Sepsis	2.6 (0.3, 12.16)	2.22 (0.22,12.98)	3.32 (0.19, 24.16)	3.91 (0.19,35.63)
Shock	1.52 (0.09, 8.35)	0.98 (0.05,8.17)		

Abbreviations: AKI (acute kidney injury), BPD (bronchopulmonary dysplasia), ICH (intracerebral hemorrhage), OR (odds ratio) NEC (necrotizing enterocolitis), ROP (retinopathy of prematurity)

[1]odd ratio (99.97% confidence interval), * indicates two-tailed, Bonferroni-corrected p-value < 0.05

Table 4. Odds of severe morbidity or mortality in preterm infants with hematologic disorders[1]

Outcome	Thalassemia (n=1174)		Hemoglobin Bart's (n=384)		Sickle cell disease (n=194)		G6PD deficiency (n=206)	
	Crude OR	Adjusted OR	Crude OR	Adjusted OR	Crude OR	Adjusted OR	Crude OR	Adjusted OR
Mortality or severe morbidity	1.28 (0.99, 1.64)	0.83 (0.6,1.14)	0.75 (0.44, 1.23)	0.86 (0.45,1.57)	0.91 (0.44, 1.73)	0.64 (0.26,1.45)	0.72 (0.33, 1.39)	0.78 (0.31,1.77)
Mortality	0.4 (0.17, 0.79)*	0.24 (0.1,0.49)*	0.19 (0.01, 0.8)*	0.21 (0.01,1)	0.47 (0.04, 1.83)	0.32 (0.03,1.47)	—	—
AKI	1.42 (0.76, 2.4)	1.04 (0.54,1.82)	0.36 (0.02, 1.55)	0.4 (0.03,1.86)	1.66 (0.33, 4.94)	1.4 (0.26,4.64)	0.85 (0.08, 3.31)	1.05 (0.1,4.5)
ICH	1.36 (0.79, 2.16)	1.07 (0.61,1.77)	0.53 (0.09, 1.61)	0.63 (0.11,2.09)	0.79 (0.1, 2.82)	0.67 (0.08,2.64)	0.87 (0.13, 2.89)	1.09 (0.16,4.04)
NEC	1.37 (0.78, 2.24)	0.98 (0.54,1.62)	0.67 (0.13, 1.95)	0.71 (0.14,2.13)	0.29 (0, 1.94)	0.23 (0,1.59)	1.43 (0.31, 4.06)	1.54 (0.33,4.58)
BPD	1.37 (0.51, 2.92)	0.99 (0.36,2.19)	1.04 (0.1, 4)	1.26 (0.12,5.23)	2.09 (0.2, 8.18)	1.82 (0.17,7.81)	1.96 (0.19, 7.67)	2.58 (0.24,10.96)
ROP	1.77 (0.74, 3.53)	1.42 (0.56,3.02)	—	—	0.85 (0.01, 5.65)	1.04 (0.01,8.08)	—	—
Sepsis	1.53 (1.09, 2.09)	1.14 (0.78,1.61)	1 (0.48, 1.85)	1.18 (0.53,2.31)	1.21 (0.45, 2.66)	1.02 (0.36,2.46)	0.78 (0.24, 1.91)	0.84 (0.24,2.2)

| Shock | 1.13 (0.78, 1.6) | 0.75 (0.49,1.11) | 0.66 (0.27, 1.34) | 0.86 (0.32,1.96) | 0.63 (0.16, 1.66) | 0.44 (0.1,1.34) | 0.43 (0.09, 1.29) | 0.54 (0.1,1.87) |

Abbreviations: AKI (acute kidney injury), BPD (bronchopulmonary dysplasia), ICH (intracerebral hemorrhage), OR (odds ratio) NEC (necrotizing enterocolitis), ROP (retinopathy of prematurity)

[1]odd ratio (99.97% confidence interval), * indicates two-tailed, Bonferroni-corrected p-value < 0.05

Table 5. Odds of severe morbidity or mortality in preterm infants with miscellaneous genetic disorders[1]

Outcome	Cystic Fibrosis (n=165)		Hypophosphatasia (n=86)	
	Crude OR	Adjusted OR	Crude OR	Adjusted OR
Mortality or severe morbidity	5.18 (2.86, 9.57)*	2.97 (1.38, 6.47)*	7.14 (3.09, 17.88)*	2.08 (0.7, 6.74)
Mortality	1.91 (0.6, 4.62)	0.69 (0.2, 1.96)	3.75 (1.09, 9.97)*	0.9 (0.24, 2.7)
AKI	4.17 (1.4, 9.77)*	1.92 (0.6, 5.03)	7.79 (2.35, 20.38)*	2.43 (0.69, 7)
ICH	2.29 (0.66, 5.83)	1.07 (0.29, 3)	3.25 (0.69, 9.91)	1.21 (0.25, 4.02)
NEC	7.52 (3.44, 14.88)*	4.03 (1.73, 8.59)*	4.53 (1.12, 12.95)*	1.62 (0.39, 4.92)
BPD	9.67 (3.25, 22.68)*	4.63 (1.45, 12.03)*	7 (1.03, 24.62)*	1.92 (0.27, 7.29)
ROP	9.36 (3.04, 22.35)*	3.95 (1.13, 11.45)*	11.95 (2.76, 35.24)*	3.16 (0.67, 10.7)
Sepsis	3.73 (1.85, 7.05)*	1.91 (0.87, 4.01)	6.2 (2.57, 14.14)*	2.05 (0.78, 5.21)
Shock	4.26 (2.18, 7.89)*	1.93 (0.87, 4.17)	6.22 (2.6, 14.16)*	1.82 (0.67, 4.89)

Abbreviations: AKI (acute kidney injury), BPD (bronchopulmonary dysplasia), ICH (intracerebral hemorrhage), OR (odds ratio) NEC (necrotizing enterocolitis), ROP (retinopathy of prematurity)

[1] odd ratio (99.97% confidence interval), * indicates two-tailed, Bonferroni-corrected p-value < 0.05

Figure 1. Prevalence of diagnosed genetic disorders over time. We determined the prevalence of the three broad classes (single gene disorder, aneuploidy, and copy number variants) of disorders within preterm infants in this cohort. We did not detect differences in these prevalence rates over time.

Supplemental Content

Supplemental Table 1. Baseline characteristics of preterm infants diagnosed with aneuploidies.
Supplemental Table 2. Baseline characteristics of preterm infants diagnosed with copy number variants.
Supplemental Table 3. Baseline characteristics of preterm infants diagnosed with hematologic disorders.
Supplemental Table 4. Baseline characteristics of preterm infants diagnosed with miscellaneous genetic disorders.
Supplemental Table 5. Instances of severe morbidity and mortality in preterm infants with genetic disorders.

References

1.   Ray JG, Park AL, Fell DB. Mortality in Infants Affected by Preterm Birth and Severe Small-for-Gestational Age Birth Weight. *Pediatrics*. Dec 2017;140(6)doi:10.1542/peds.2017-1881

2.   Horbar JD, Carpenter JH, Badger GJ, et al. Mortality and neonatal morbidity among infants 501 to 1500 grams from 2000 to 2009. *Pediatrics*. Jun 2012;129(6):1019-26. doi:10.1542/peds.2011-3028

3.   Baer RJ, Rogers EE, Partridge JC, et al. Population-based risks of mortality and preterm morbidity by gestational age and birth weight. *J Perinatol*. Nov 2016;36(11):1008-1013. doi:10.1038/jp.2016.118

4.   Hays T, Wapner RJ. Genetic testing for unexplained perinatal disorders. *Curr Opin Pediatr*. Apr 1 2021;33(2):195-202. doi:10.1097/MOP.0000000000000999

5.   Kingsmore SF, Henderson A, Owen MJ, et al. Measurement of genetic diseases as a cause of mortality in infants receiving whole genome sequencing. *NPJ Genom Med*. 2020;5(1):49. doi:10.1038/s41525-020-00155-8

6.   Owen MJ, Wright MS, Batalov S, et al. Reclassification of the Etiology of Infant Mortality With Whole-Genome Sequencing. *JAMA Network Open*. 2023;6(2):e2254069-e2254069. doi:10.1001/jamanetworkopen.2022.54069

7.   Clark MM, Hildreth A, Batalov S, et al. Diagnosis of genetic diseases in seriously ill children by rapid whole-genome sequencing and automated phenotyping and interpretation. *Sci Transl Med*. Apr 24 2019;11(489):eaat6177-eaat6177. doi:10.1126/scitranslmed.aat6177

8.   Stanley KE, Giordano J, Thorsten V, et al. Causal Genetic Variants in Stillbirth. *N Engl J Med*. Sep 17 2020;383(12):1107-1116. doi:10.1056/NEJMoa1908753

9.   Petrovski S, Aggarwal V, Giordano JL, et al. Whole-exome sequencing in the evaluation of fetal structural anomalies: a prospective cohort study. *Lancet*. Feb 23 2019;393(10173):758-767. doi:10.1016/S0140-6736(18)32042-7

10.   Lord J, McMullan DJ, Eberhardt RY, et al. Prenatal exome sequencing analysis in fetal structural anomalies detected by ultrasonography (PAGE): a cohort study. *Lancet*. Feb 23 2019;393(10173):747-757. doi:10.1016/S0140-6736(18)31940-8

11.   Boghossian NS, Hansen NI, Bell EF, et al. Mortality and morbidity of VLBW infants with trisomy 13 or trisomy 18. *Pediatrics*. Feb 2014;133(2):226-35. doi:10.1542/peds.2013-1702

12.   Boghossian NS, Horbar JD, Carpenter JH, Murray JC, Bell EF. Major chromosomal anomalies among very low birth weight infants in the Vermont Oxford Network. *J Pediatr*. May 2012;160(5):774-780.e11. doi:10.1016/j.jpeds.2011.11.005

13.   Hays T, Thompson MV, Bateman DA, et al. The Prevalence and Clinical Significance of Congenital Anomalies of the Kidney and Urinary Tract in Preterm Infants. *JAMA Netw Open*. Sep 1 2022;5(9):e2231626. doi:10.1001/jamanetworkopen.2022.31626

14.   Fenton TR, Kim JH. A systematic review and meta-analysis to revise the Fenton growth chart for preterm infants. *BMC Pediatr*. Apr 20 2013;13:59. doi:10.1186/1471-2431-13-59

15.   *R: A language and environment for statistical computing.* R Foundation for Statistical Computing; 2022. https://www.R-project.org/

16.   ACOG Practice Bulletin No. 78: hemoglobinopathies in pregnancy. *Obstet Gynecol*. Jan 2007;109(1):229-37. doi:10.1097/00006250-200701000-00055

17.   Festini F, Taccetti G, Repetto T, et al. Gestational and neonatal characteristics of children with cystic fibrosis: a cohort study. *J Pediatr*. Sep 2005;147(3):316-20. doi:10.1016/j.jpeds.2005.04.031

18.   Bianco K, Gormley M, Farrell J, et al. Placental transcriptomes in the common aneuploidies reveal critical regions on the trisomic chromosomes and genome-wide effects. *Prenat Diagn*. Sep 2016;36(9):812-22. doi:10.1002/pd.4862

19.   Savva GM, Walker K, Morris JK. The maternal age-specific live birth prevalence of trisomies 13 and 18 compared to trisomy 21 (Down syndrome). *Prenat Diagn*. Jan 2010;30(1):57-64. doi:10.1002/pd.2403

20.   Schlüter DK, Griffiths R, Adam A, et al. Impact of cystic fibrosis on birthweight: a population based study of children in Denmark and Wales. *Thorax*. 2019;74(5):447-454. doi:10.1136/thoraxjnl-2018-211706

21.   Neu J. Necrotizing Enterocolitis: The Future. *Neonatology*. 2020;117(2):240-244. doi:10.1159/000506866

22.   Sathe M, Houwen R. Meconium ileus in Cystic Fibrosis. *J Cyst Fibros*. Nov 2017;16 Suppl 2:S32-s39. doi:10.1016/j.jcf.2017.06.007

23.   Sly PD, Brennan S, Gangell C, et al. Lung disease at diagnosis in infants with cystic fibrosis detected by newborn screening. *Am J Respir Crit Care Med*. Jul 15 2009;180(2):146-52. doi:10.1164/rccm.200901-0069OC

24.   Meyerholz DK, Stoltz DA, Gansemer ND, et al. Lack of cystic fibrosis transmembrane conductance regulator disrupts fetal airway development in pigs. *Lab Invest*. Jun 2018;98(6):825-838. doi:10.1038/s41374-018-0026-7

25.   Jelin AC, Sharshiner R, Caughey AB. Maternal co-morbidities and neonatal outcomes associated with cystic fibrosis(). *J Matern Fetal Neonatal Med*. Jan 2017;30(1):4-7. doi:10.3109/14767058.2016.1161747

26.   Petrikin JE, Willig LK, Smith LD, Kingsmore SF. Rapid whole genome sequencing and precision neonatology. *Semin Perinatol*. Dec 2015;39(8):623-31. doi:10.1053/j.semperi.2015.09.009

27.   Hays T, Hernan R, Disco M, et al. Implementation of Rapid Genome Sequencing for Critically Ill Infants With Complex Congenital Heart Disease. *Circ Genom Precis Med*. Jul 7 2023:e004050. doi:10.1161/circgen.122.004050

Supplemental Table 1. Baseline characteristics of preterm infants diagnosed with aneuploidies.

	Trisomy 13 (n=110)	Trisomy 18 (n=155)	Trisomy 21 (n=973)	Trisomy 22 (n=1)	Klinefelter syndrome (n=35)	Turner syndrome (n=41)
Gestational age (weeks)						
Mean (SD)	30.9 (2.30)	31.0 (2.16)	31.0 (2.19)	32.0 (NA)	29.6 (2.57)	30.4 (2.63)
Median [Min, Max]	32.0 [24.0, 33.0]	32.0 [24.0, 33.0]	32.0 [23.0, 33.0]	32.0 [32.0, 32.0]	30.0 [23.0, 33.0]	31.0 [24.0, 33.0]
Birth weight						
SGA	27 (24.5%)	101 (65.2%)	190 (19.5%)	0 (0%)	9 (25.7%)	9 (22.0%)
AGA	77 (70.0%)	50 (32.3%)	666 (68.4%)	1 (100%)	26 (74.3%)	25 (61.0%)
LGA	6 (5.5%)	4 (2.6%)	117 (12.0%)	0 (0%)	0 (0%)	7 (17.1%)
Race						
Asian	1 (0.9%)	2 (1.3%)	30 (3.1%)	0 (0%)	2 (5.7%)	2 (4.9%)
Black	27 (24.5%)	29 (18.7%)	134 (13.8%)	0 (0%)	7 (20.0%)	2 (4.9%)
Hispanic	31 (28.2%)	54 (34.8%)	238 (24.5%)	1 (100%)	9 (25.7%)	7 (17.1%)
Other	8 (7.3%)	12 (7.7%)	81 (8.3%)	0 (0%)	1 (2.9%)	2 (4.9%)
White	43 (39.1%)	58 (37.4%)	490 (50.4%)	0 (0%)	16 (45.7%)	28 (68.3%)
Sex						
Female	47 (42.7%)	96 (61.9%)	425 (43.7%)	1 (100%)	8 (22.9%)	41 (100%)
Male	63 (57.3%)	59 (38.1%)	548 (56.3%)	0 (0%)	27 (77.1%)	0 (0%)
Delivery mode						
Cesarean Section	81 (73.6%)	119 (76.8%)	766 (78.7%)	1 (100%)	28 (80.0%)	32 (78.0%)
Vaginal Delivery	29 (26.4%)	36 (23.2%)	207 (21.3%)	0 (0%)	7 (20.0%)	9 (22.0%)
Antenatal betamethasone						
Absent	40 (36.4%)	72 (46.5%)	214 (22.0%)	0 (0%)	7 (20.0%)	9 (22.0%)
Present	70 (63.6%)	83 (53.5%)	759 (78.0%)	1 (100%)	28 (80.0%)	32 (78.0%)

Intubation within 72 hours						
Absent	31 (28.2%)	54 (34.8%)	628 (64.5%)	0 (0%)	18 (51.4%)	24 (58.5%)
Present	79 (71.8%)	101 (65.2%)	345 (35.5%)	1 (100%)	17 (48.6%)	17 (41.5%)
Discharge Year						
2000 to 2004	27 (24.5%)	31 (20.0%)	144 (14.8%)	0 (0%)	0 (0%)	10 (24.4%)
2005 to 2009	23 (20.9%)	48 (31.0%)	257 (26.4%)	0 (0%)	7 (20.0%)	11 (26.8%)
2010 to 2014	31 (28.2%)	35 (22.6%)	258 (26.5%)	0 (0%)	13 (37.1%)	7 (17.1%)
2015 to 2020	29 (26.4%)	41 (26.5%)	314 (32.3%)	1 (100%)	15 (42.9%)	13 (31.7%)
Anomaly						
Absent	38 (34.5%)	64 (41.3%)	579 (59.5%)	0 (0%)	28 (80.0%)	32 (78.0%)
Present	72 (65.5%)	91 (58.7%)	394 (40.5%)	1 (100%)	7 (20.0%)	9 (22.0%)
Morbidity						
Absent	7 (6.4%)	17 (11.0%)	623 (64.0%)	0 (0%)	24 (68.6%)	27 (65.9%)
Present	103 (93.6%)	138 (89.0%)	350 (36.0%)	1 (100%)	11 (31.4%)	14 (34.1%)
Disposition						
Alive	17 (15.5%)	25 (16.1%)	830 (85.3%)	0 (0%)	33 (94.3%)	36 (87.8%)
Died	93 (84.5%)	130 (83.9%)	143 (14.7%)	1 (100%)	2 (5.7%)	5 (12.2%)

Abbreviations: SGA (small for gestational age), AGA (appropriate for gestational age), LGA

(large for gestational age)

Supplemental Table 2. Baseline characteristics of preterm infants diagnosed with copy number variants.

	DiGeorge syndrome (n=29)	Cri-du-chat syndrome (n=16)	13q-deletion syndrome (n=7)
Gestational age (weeks)			
Mean (SD)	30.6 (2.40)	31.3 (2.65)	31.1 (1.35)
Median [Min, Max]	31.0 [25.0, 33.0]	32.0 [24.0, 33.0]	31.0 [29.0, 33.0]
Birth weight			
SGA	3 (10.3%)	2 (12.5%)	3 (42.9%)
AGA	23 (79.3%)	14 (87.5%)	4 (57.1%)
LGA	3 (10.3%)	0 (0%)	0 (0%)
Race			
Black	9 (31.0%)	6 (37.5%)	4 (57.1%)
Hispanic	9 (31.0%)	4 (25.0%)	3 (42.9%)
Other	2 (6.9%)	1 (6.3%)	0 (0%)
White	8 (27.6%)	5 (31.3%)	0 (0%)
Asian	1 (3.4%)	0 (0%)	0 (0%)
Sex			
Female	17 (58.6%)	12 (75.0%)	3 (42.9%)
Male	12 (41.4%)	4 (25.0%)	4 (57.1%)
Delivery mode			
Cesarean Section	24 (82.8%)	12 (75.0%)	5 (71.4%)
Vaginal Delivery	5 (17.2%)	4 (25.0%)	2 (28.6%)
Antenatal betamethasone			
Absent	9 (31.0%)	1 (6.3%)	2 (28.6%)
Present	20 (69.0%)	15 (93.8%)	5 (71.4%)
Intubation within 72 hours			
Absent	14 (48.3%)	8 (50.0%)	4 (57.1%)
Present	15 (51.7%)	8 (50.0%)	3 (42.9%)
Discharge Year			
2000 to 2004	1 (3.4%)	2 (12.5%)	0 (0%)
2005 to 2009	8 (27.6%)	5 (31.3%)	0 (0%)
2010 to 2014	6 (20.7%)	2 (12.5%)	1 (14.3%)
2015 to 2020	14 (48.3%)	7 (43.8%)	6 (85.7%)
Anomaly			
Absent	13 (44.8%)	8 (50.0%)	5 (71.4%)

Present	16 (55.2%)	8 (50.0%)	2 (28.6%)
Morbidity			
Absent	17 (58.6%)	11 (68.8%)	5 (71.4%)
Present	12 (41.4%)	5 (31.3%)	2 (28.6%)
Disposition			
Alive	25 (86.2%)	14 (87.5%)	6 (85.7%)
Died	4 (13.8%)	2 (12.5%)	1 (14.3%)

Abbreviations: SGA (small for gestational age), AGA (appropriate for gestational age), LGA

(large for gestational age)

Supplemental Table 3. Baseline characteristics of preterm infants diagnosed with hematologic disorders.

	Thalassemia (n=1174)	Hemoglobin Bart's (n=384)	Sickle cell anemia (n=194)	G6PD deficiency (n=206)	Hemophilia A (n=10)	Hemophilia B (n=5)
Gestational age (weeks)						
Mean (SD)	29.5 (2.73)	30.1 (2.43)	29.7 (2.81)	30.3 (2.48)	29.6 (3.06)	29.4 (3.36)
Median [Min, Max]	30.0 [23.0, 33.0]	31.0 [23.0, 33.0]	30.0 [23.0, 33.0]	31.0 [23.0, 33.0]	31.0 [25.0, 33.0]	28.0 [26.0, 33.0]
Birth weight						
SGA	137 (11.7%)	32 (8.3%)	35 (18.0%)	25 (12.1%)	1 (10.0%)	0 (0%)
AGA	996 (84.8%)	332 (86.5%)	152 (78.4%)	173 (84.0%)	9 (90.0%)	5 (100%)
LGA	41 (3.5%)	20 (5.2%)	7 (3.6%)	8 (3.9%)	0 (0%)	0 (0%)
Race						
Black	692 (58.9%)	296 (77.1%)	180 (92.8%)	169 (82.0%)	2 (20.0%)	0 (0%)
Asian	93 (7.9%)	22 (5.7%)	2 (1.0%)	9 (4.4%)	0 (0%)	0 (0%)
Hispanic	88 (7.5%)	15 (3.9%)	6 (3.1%)	9 (4.4%)	1 (10.0%)	1 (20.0%)
Other	81 (6.9%)	22 (5.7%)	1 (0.5%)	13 (6.3%)	0 (0%)	0 (0%)
White	220 (18.7%)	29 (7.6%)	5 (2.6%)	6 (2.9%)	7 (70.0%)	4 (80.0%)
Sex						
Female	569 (48.5%)	208 (54.2%)	104 (53.6%)	18 (8.7%)	0 (0%)	0 (0%)
Male	605 (51.5%)	176 (45.8%)	90 (46.4%)	188 (91.3%)	10 (100%)	5 (100%)
Delivery mode						
Cesarean Section	795 (67.7%)	280 (72.9%)	144 (74.2%)	138 (67.0%)	8 (80.0%)	3 (60.0%)
Vaginal Delivery	379 (32.3%)	104 (27.1%)	50 (25.8%)	68 (33.0%)	2 (20.0%)	2 (40.0%)
Antenatal betamethasone						
Present	970 (82.6%)	325 (84.6%)	152 (78.4%)	178 (86.4%)	8 (80.0%)	4 (80.0%)
Absent	204 (17.4%)	59 (15.4%)	42 (21.6%)	28 (13.6%)	2 (20.0%)	1 (20.0%)
Intubation within 72 hours						

Absent	659 (56.1%)	274 (71.4%)	114 (58.8%)	152 (73.8%)	6 (60.0%)	2 (40.0%)
Present	515 (43.9%)	110 (28.6%)	80 (41.2%)	54 (26.2%)	4 (40.0%)	3 (60.0%)
Discharge Year						
2000 to 2004	290 (24.7%)	4 (1.0%)	23 (11.9%)	1 (0.5%)	2 (20.0%)	0 (0%)
2005 to 2009	337 (28.7%)	117 (30.5%)	49 (25.3%)	51 (24.8%)	1 (10.0%)	4 (80.0%)
2010 to 2014	229 (19.5%)	107 (27.9%)	58 (29.9%)	93 (45.1%)	2 (20.0%)	0 (0%)
2015 to 2020	318 (27.1%)	156 (40.6%)	64 (33.0%)	61 (29.6%)	5 (50.0%)	1 (20.0%)
Anomaly						
Present	86 (7.3%)	28 (7.3%)	10 (5.2%)	12 (5.8%)	1 (10.0%)	0 (0%)
Absent	1088 (92.7%)	356 (92.7%)	184 (94.8%)	194 (94.2%)	9 (90.0%)	5 (100%)
Morbidity						
Absent	868 (73.9%)	318 (82.8%)	155 (79.9%)	172 (83.5%)	6 (60.0%)	2 (40.0%)
Present	306 (26.1%)	66 (17.2%)	39 (20.1%)	34 (16.5%)	4 (40.0%)	3 (60.0%)
Disposition						
Alive	1148 (97.8%)	380 (99.0%)	189 (97.4%)	204 (99.0%)	8 (80.0%)	5 (100%)
Died	26 (2.2%)	4 (1.0%)	5 (2.6%)	2 (1.0%)	2 (20.0%)	0 (0%)

Abbreviations: SGA (small for gestational age), AGA (appropriate for gestational age), LGA (large for gestational age)

Supplemental Table 4. Baseline characteristics of preterm infants diagnosed with miscellaneous genetic disorders.

	Cystic fibrosis (n=165)	Hypophosphatasia (n=86)	Fragile X syndrome (n=7)	Fanconi anemia (n=2)
Gestational age (weeks)				
Mean (SD)	29.0 (3.12)	27.6 (2.84)	30.1 (2.34)	29.5 (4.95)
Median [Min, Max]	29.0 [23.0, 33.0]	27.0 [23.0, 33.0]	31.0 [26.0, 32.0]	29.5 [26.0, 33.0]
Birth weight				
SGA	26 (15.8%)	32 (37.2%)	0 (0%)	2 (100%)
AGA	126 (76.4%)	52 (60.5%)	5 (71.4%)	0 (0%)
LGA	13 (7.9%)	2 (2.3%)	2 (28.6%)	0 (0%)
Race				
Asian	1 (0.6%)	4 (4.7%)	0 (0%)	2 (100%)
Black	17 (10.3%)	21 (24.4%)	1 (14.3%)	0 (0%)
Hispanic	14 (8.5%)	22 (25.6%)	0 (0%)	0 (0%)
Other	6 (3.6%)	6 (7.0%)	0 (0%)	0 (0%)
White	127 (77.0%)	33 (38.4%)	6 (85.7%)	0 (0%)
Sex				
Female	71 (43.0%)	40 (46.5%)	2 (28.6%)	2 (100%)
Male	94 (57.0%)	46 (53.5%)	5 (71.4%)	0 (0%)
Delivery mode				
Cesarean Section	115 (69.7%)	69 (80.2%)	5 (71.4%)	2 (100%)
Vaginal Delivery	50 (30.3%)	17 (19.8%)	2 (28.6%)	0 (0%)
Antenatal betamethasone				
Absent	33 (20.0%)	18 (20.9%)	2 (28.6%)	2 (100%)
Present	132 (80.0%)	68 (79.1%)	5 (71.4%)	0 (0%)
Intubation within 72 hours				

Absent	52 (31.5%)	21 (24.4%)	3 (42.9%)	1 (50.0%)
Present	113 (68.5%)	65 (75.6%)	4 (57.1%)	1 (50.0%)
Discharge Year				
2000 to 2004	68 (41.2%)	27 (31.4%)	2 (28.6%)	0 (0%)
2005 to 2009	56 (33.9%)	26 (30.2%)	0 (0%)	0 (0%)
2010 to 2014	24 (14.5%)	19 (22.1%)	1 (14.3%)	1 (50.0%)
2015 to 2020	17 (10.3%)	14 (16.3%)	4 (57.1%)	1 (50.0%)
Anomaly				
Absent	126 (76.4%)	78 (90.7%)	5 (71.4%)	2 (100%)
Present	39 (23.6%)	8 (9.3%)	2 (28.6%)	0 (0%)
Morbidity				
Absent	68 (41.2%)	29 (33.7%)	6 (85.7%)	1 (50.0%)
Present	97 (58.8%)	57 (66.3%)	1 (14.3%)	1 (50.0%)
Disposition				
Alive	149 (90.3%)	71 (82.6%)	7 (100%)	1 (50.0%)
Died	16 (9.7%)	15 (17.4%)	0 (0%)	1 (50.0%)

SGA: small for gestational age; AGA: appropriate for gestational age; LGA: large for gestational age.

Supplemental Table 5. Instances of severe morbidity and mortality in preterm infants with genetic disorders.

Genetic Disorder	Mortality or Any Severe Morbidity	Mortality	AKI	ICH	NEC	BPD	ROP	Sepsis	Shock
No genetic diagnosis	*68326*	*16861*	*9021*	*12289*	*10933*	*3953*	*3836*	*28874*	*30087*
Any genetic diagnosis	*1526*	*565*	*279*	*204*	*232*	*172*	*72*	*585*	*615*
Aneuploidy	883	480	171	93	101	105	16	258	341
Trisomy 13	103	93	18	0	5	7	1	9	20
Trisomy 18	138	130	13	2	5	10	0	11	31
Trisomy 21	350	143	88	41	51	48	4	127	155
Trisomy 22	1	1	1	0	0	0	0	0	0
Klinefelter syndrome	11	2	2	2	0	1	0	8	4
Turner syndrome	14	5	5	3	2	1	0	9	6
Unspecified Aneuploidy	266	106	44	45	38	38	11	94	125
Copy number variation disorder	35	14	7	1	8	5	0	16	13
13q deletion syndrome	2	1	0	1	0	0	0	0	0
Cri-du-chat syndrome	5	2	2	0	2	1	0	4	2
DiGeorge syndrome	12	4	1	0	1	2	0	6	4
Unspecified CNV	16	7	4	0	5	2	0	6	7
Monogenic disorder	608	71	101	110	123	62	56	311	261
Hematologic disorder	452	39	66	85	76	36	28	232	174
Thalassemia	306	26	47	61	55	20	25	156	125
Hemoglobin bart's	66	4	4	8	9	5	0	35	25
Sickle cell disease	39	5	9	6	2	5	2	21	12
G6PD deficiency	34	2	5	7	10	5	1	15	9
Hemophilia A	4	2	1	2	0	1	0	3	2
Hemophilia B	3	0	0	1	0	0	0	2	1
Miscellaneous	156	32	35	25	47	26	28	79	87
Cystic fibrosis	97	16	18	14	35	18	17	45	51
Fragile X syndrome	1	0	0	0	0	0	0	1	1
Hypophosphatasia	57	15	16	10	12	7	11	33	34
Fanconi Anemia	1	1	1	1	0	1	0	0	1
Multiple Disorders	6	1	2	1	3	2	1	5	2

Abbreviations: AKI (acute kidney injury), BPD (bronchopulmonary dysplasia), ICH (intracerebral hemorrhage), OR (odds ratio) NEC (necrotizing enterocolitis), ROP (retinopathy of prematurity)