Title: Admissions for eating disorders and other mental health diagnoses during the COVID-19 pandemic

Authors: Sydney C. Jones, BS[1]*, Megan Jacobs, MD[2]*, Emile Latour, MS[3], Rebecca Marshall MD, MPH[4], Michelle Noelck, MD[2], Byron A. Foster, MD, MPH[2,5]
* Indicates co first-authorship

Affiliations:
[1] School of Medicine, Oregon Health & Science University
2730 SW Moody Avenue Portland, OR 97201 U.S.
jonessy@ohsu.edu
[2] Department of Pediatrics, Oregon Health & Science University
707 SW Gaines St CDRC Portland, OR 97239 U.S.
jacobsme@ohsu.edu, noelck@ohsu.edu, fosterb@ohsu.edu
[3] Biostatistics Shared Resource, Knight Cancer Institute, Oregon Health & Science University
2720 SW Moody Ave, Portland, OR 97201 U.S.
latour@ohsu.edu
[4] Department of Psychiatry, Oregon Health & Science University
3181 SW Sam Jackson Park Rd Portland, OR 97239 Multnomah Pavilion, Room 2316 U. S.
marshare@ohsu.edu
[5] OHSU-PSU School of Public Health
 3181 SW Sam Jackson Park Road Portland, OR 97239 U.S.

Address correspondence to:
Megan Jacobs, MD
Department of Pediatrics, Oregon Health & Science University
707 SW Gaines St
CDRC
Portland, OR 97239
Jacobsme@ohsu.edu
503-418-5700

Source of Funding: This work was supported by Oregon Health & Science University institutional funding.

Acknowledgment: Jose Rodriguez extracted multiple iterations of the data from the medical records.

Conflict of Interest: Dr. Jacobs, Dr. Foster, Dr. Marshall, Dr. Noelck, Mr. Latour, and Ms. Jones have no conflicts of interest to disclose.

Key Words:
COVID-19, Pediatrics, Adolescent, Eating Disorder, Mental Health

Abstract:

Objective: To examine how hospital admissions for mental health and eating disorders changed

at the beginning of the COVID-19 pandemic and with the return to fully in-person school with

increased vaccine availability.

Methods: Data from a tertiary care children's hospital were examined for admissions to the

hospital from March 2018 through March 2022, including children 6-20 years old admitted with

ICD-10 codes for mental health and eating disorders. Interrupted time series (ITS) analyses were

used to examine for changes at specific time points.

Results: In the time between March 2018 through March 2022, 851 admissions met inclusion

criteria for eating disorders and 1,505 admissions for other mental health diagnoses. In the first

year of the pandemic, the ITS analysis showed a significant increase in admissions per month for

eating disorders with a slope of 1.2 (95% CI: 0.2, 2.2) and for other mental health diagnoses, a

slope of 1.9 (95% CI: 1.1, 2.7). In a longer-term ITS analysis, return to fully in-person school

was associated with a subsequent decrease in admissions over time at -1.0 per month (95% CI: -

2.1, 0.1). For other mental health diagnoses, return to school was associated with an initial drop

and then an increase in admissions over time, slope of 2.2 (95% CI: -0.5, 3.8).

Conclusion: The COVID-19 pandemic had an initial impact on admissions for eating disorders

and other mental health that attenuated over time. We note differences in the association of

return to school on eating disorders compared with other mental health diagnoses.

Introduction:

The COVID-19 pandemic has drastically impacted the pediatric population.[1] When schools physically closed and students moved to primarily virtual learning, many children experienced social isolation from their peers and lost access to school-based mental health resources.[2] Preliminary research suggests that mental health problems have increased among children and young adults since the onset of the COVID pandemic, with potential contributors including social isolation, difficulty accessing mental health resources, the economic recession, and the stress surrounding a public health crisis.[2,3,4,5] There is emerging evidence from across the country that school closures had a detrimental impact on the health and wellbeing of adolescents. Two medical centers from the East Coast and Midwest found an increase in eating disorder hospital admissions in the months following their states' stay at home orders.[6,7] Other studies reported a general increase in eating disorder behavior since the start of the pandemic. [8, 9, 10, 11, 12,13] Several hospitals reported increased emergency department mental health visits for pediatric patients during the later parts of the pandemic compared to before the pandemic.[14,15,16]

While several reports have described the initial increase in mental health-associated visits and admissions, few studies have described the impact of the overall trajectory of the COVID-19 pandemic,[1] nor examined the association of changing pandemic school restrictions and vaccine availability on pediatric mental health hospital admissions. This study was conducted in Oregon where around 1500 to 1600 public, private, and charter schools tracked their virtual, hybrid, and in-person learning statuses. Following statewide school closures in March of 2020, by October 2020, approximately 83% of schools were using a virtual learning model. By the end of the 2020-2021 school year, only around 2% of the schools were virtual, with 61% of schools reporting hybrid instruction, and 34% reporting in person only.[17] It was not until the start of the

2021-2022 school year, that all of Oregon schools returned to a mostly in-person full academic

year.[18] During this same time period, vaccine availability for the 12+ age group was approved for

emergency use authorization (EUA) on May 12th 2021.[19] As of August 30, 2021, 52.1% of 12-17

year olds in Oregon had received one COVID-19 vaccine.[20] The aim of this study was to

examine how hospital admissions for mental health conditions and eating disorders changed at

both the beginning of the pandemic and the start of the 2021-2020 school year, representing the

first fully in-person school year and a greater return to pre-pandemic normalcy. We also

examined whether there were differences in sex, gender, and racial and ethnic categories among

children admitted for eating disorders and other mental health diagnoses over the course of the

pandemic.

Methods

Study design and sample selection

A retrospective chart review of the electronic medical record was conducted at a 150-bed

academic tertiary-care children's hospital in the Pacific Northwest. We examined all admission

encounters from March 1, 2018, two years before the onset of the COVID-19 pandemic, through

March 31, 2022, for patients 6-20 years of age. Patients could be included more than once in the

study period. Demographic data were abstracted from the medical record. We examined

patients' self-identified race and ethnicity as well as gender variables. Over the course of the

study, the method used to categorize gender changed in the medical record, leading to a

substantial number of patients with unspecified gender. We also examined biologic sex. This

study was approved by our institutional IRB.

Inclusion and exclusion criteria

Patients meeting the following criteria were included in this study: age 6-20 years and

having an International Classification of Diseases, 10th revision (ICD-10) code in their electronic

health record for either the primary diagnosis or encounter diagnosis of an eating disorder or

another mental health disorder including major depressive disorder, suicidal ideation, and

intentional self-harm. Please see appendix for full list of ICD-10 codes.

We a priori defined children as having an eating disorder admission if they had an eating

disorder ICD-10 code associated with their admission or an ICD-10 code of malnutrition that

was not one of the excluded codes (e.g. Kwashiorkor). Similarly, we defined children as having a

mental health admission if they had any mental health related ICD-10 codes associated with their

admission encounter in the medical record (see appendix).

Statistical analysis

Overall, we stratified the analysis by mental health diagnoses and eating disorder

diagnoses. Descriptive statistics (means and standard deviations, medians and interquartile

ranges, frequencies and percentages) were used to summarize patient characteristics. Group

comparisons for continuous variables were made using one-way ANOVA, and comparisons for

categorical variables using Fisher's exact test. We conducted two interrupted time series (ITS)

analyses: (1) examining the interruption of pandemic related school closures, and (2) examining

the start of the first fully in person school year with vaccines available to adolescents.

The first ITS analysis examines the interruption of pandemic related school closures. The

pandemic was assumed to start on April 1, 2020, just after the date of local pandemic-related

school closures. The pre-pandemic period was considered as those hospital admissions between

March 1, 2018 through March 31, 2020. The post-pandemic period includes the first year of

pandemic admissions, April 1, 2020 through March 31, 2021.

133   The second ITS analysis examines whether the return to the fully in person academic year

134 was associated with a change in hospital admissions. It examined two interruptions and three

135 time periods: pre-pandemic period (March 1, 2018 to March 31, 2020); the pandemic school

136 closure interruption (April 1, 2020); the pandemic year with school closure, hybrid school

137 models, and some return to in-person learning (April 1, 2020 to August 31, 2021); the return to

138 the fully in-person school year interruption (September 1, 2021); and the full school year return

139 to in-person learning (September 1, 2021 to March 31, 2022). All analyses used R: A Language

140 and Environment for Statistical Computing.[21] A *p*-value < 0.05 was defined as statistically

141 significant.

142 **Results**

143 *All hospital admissions*

144   All admitted patients from March 1, 2018 through March 31, 2022 can be seen in Figure

145 1. Before the pandemic, there were an average of 261.7 admissions per month (95% CI: 248.6,

146 274.8) with a stable slope over time of -0.5 admissions per month (95% CI: -1.7, 0.7; p=0.44).

147 Starting April 1, 2020, there was a decrease in all admissions by -60.3 admissions (95% CI: -

148 27.2, -93.4; p<0.001). Subsequently, the rate of admissions per month increased during the

149 pandemic by 4.5 admissions per month (95% CI: 0.4, 8.5; p<0.001). On September 1, 2021, with

150 the return to in-person learning, the number of admissions did not substantially change (p=0.60),

151 and there was no change in the slope of admissions per month after that period (p=0.24) (Figure

152 1b). In summary, for all hospitalizations, we observed a decrease in admission numbers at the

153 start of the pandemic, and then a slow rise in admission rate throughout the course of the

154 pandemic lockdown and virtual to hybrid school period with no change in admission rate

155 associated with the return to in person learning.

Patient Characteristics:

Between March 1, 2018 and March 31, 2022, there were a total of 851 patients who were

admitted with a primary or secondary ICD-10 code for an eating disorder. During this same time

period, there were 1,506 patients admitted with a mental health primary or secondary ICD-10

code. Patient characteristics can be found in Table 1. There was a relative increase in youth

admitted with female sex for eating disorders after the onset of the pandemic (Table 1). An

increase in the proportion of admissions for patients with a transgender or other gender was

noted in the first part of the pandemic, increasing from 2.6% to 3.3% from April 1, 2020 through

August 31, 2021, and then back down to 2.4% in the final period after return to school. For other

mental health diagnoses, we saw a similar increase from 2.6% pre-pandemic to 4.2% during the

April 1, 2020 through August 31, 2021 time frame, and a subsequent decrease to 1.6% post

September 1, 2021. The proportion of those reporting Hispanic ethnicity decreased in both

groups over time, as missing ethnicity data increased.

Statistical Model:

In the ITS analysis examining the effect of the pandemic interruption on April 1, 2020

and the subsequent year of admissions data, we modeled admissions separately for eating

disorders (Figure 2) and other mental health diagnoses (Figure 3). The figures show the observed

trend from the ITS models and the predicted trend had the pre-pandemic trends continued.

Eating disorder admissions:

For eating disorders, prior to the COVID-19 pandemic (prior to April 1, 2020), the

average number of admissions per month was estimated to be 20.6 (95% CI: 15.9, 25.4). The rate

of admissions per month was stable over time with a slope of 0.1 admissions per month (95% CI:

-0.3, 0.5; p=0.588) (Figure 2).

On April 1, 2020, the number of eating disorder admissions showed a non-significant

decrease of -7.0 (95% CI: -16.3, 2.3; p=0.142) admissions. The number of eating disorder

admissions per month then increased over time during the first year of the pandemic with a slope

of 1.2 (95% CI: 0.2, 2.2; p=0.020) admissions per month (Figure 2a).

The second set of ITS models (Figure 2b) included admissions over a longer time period

and a second interruption on September 1, 2021, symbolizing time of increased socialization due

to opening of more in-person school and increasing immunization rates. In examining the longer

time period in this model, the number of admissions per month stayed the same during the

pandemic with a slope of 0.3 admissions per month (95% CI: -0.4, 1.1; p=0.376). At the second

interruption (September 1, 2021), the number of admissions remained stable at an estimated 0.3

admissions per month (95% CI: -14.2, 13.6; p=0.968). From September 1, 2021 through March

31, 2022, the number of admissions per month decreased slightly over time with a slope of -1.0

admissions per month (95% CI: -2.1, 0.1; p=0.072) without statistical significance (Figure 2b).

Other mental health admissions:

For other mental health diagnoses (Figure 3), prior to the COVID-19 pandemic, the

average number of admissions per month was 35.4 (95% CI: 31.3, 39.5). The rate of admissions

per month was stable over time with a slope of 0.2 admissions per month (95% CI: -0.1, 0.5;

p=0.13).

At the first interruption on April 1, 2020, the number of mental health admissions

decreased by 15.8 admissions (95% CI: 9.7, 21.9; p<0.001) (Figure 3a). Subsequently, the

number of mental health admissions per month increased over time during the first year of the

pandemic with a slope of 1.9 more admissions per month (95% CI: 1.1, 2.7; p<0.001).

Looking at the longer period of time of 18 months in the second ITS model (Figure 3b),

the number of mental health admissions per month stayed the same over time during the

pandemic with a slope of 0.4 admissions per month (95% CI: -0.5, 1.4; p=0.384). Using

September 1, 2021 as the second interruption, the number of admissions decreased by -16.9

(95% CI: -26.5, -7.3; p<0.001). Subsequently, the number of admissions per month showed an

increase with a slope of 2.2 admissions per month (95% CI: -0.5, 3.8; p=0.012).

Discussion

The COVID-19 pandemic has had a significant impact on youth mental health.[1] This

study sought to determine impact at the level of hospitalizations at an urban tertiary care

children's hospital with a relationship to mental health etiologies including suicidality,

depression, anxiety disorders, and medical stabilization for disordered eating. In the single

interrupted time series model, both eating disorder and other mental health admission rates

increased post-April 1, 2020 intervention with statistical significance. However, although the

April 1, 2020 date seemed to be a significant intervention date for mental health admissions, this

was not necessarily the case for eating disorder admissions. This lack of significance in eating

disorder admissions on intervention dates with the second intervention model was also true with

September 1, 2021. Admissions for eating disorder medical stabilizations at the children's

hospital has a protocolized length of stay of an average of a week, and there is a limitation on the

number of rooms capable of meeting the needs of these types of admissions. These limitations

may have created a cap on admissions of this type.

After the initial drop in admissions associated with the start of the pandemic, both eating

disorder and other mental health admissions increased over the course of the first year of the

pandemic, consistent with other reports,[6,7] though overall admissions over the longer course of

the pandemic were stable compared to prior. Following the return to mostly in-person school

with vaccines widely available in Oregon, we observed an initial decrease in mental health

admissions suggesting that improved social interactions and possible perceived community

safety related to increased immunization rates in youth may have acutely improved youth mental

health. The significance in increased mental health admissions in the 2021 Fall and Winter

months may reflect annual trends in depression, seasonal changes, or Omicron COVID-19

variant emergence in November of 2021.[22] While these ecologic associations between major

societal shifts and changes in admissions to hospitals have inherent epidemiological weaknesses

in terms of causality, they can provide some insights into the effects associated with such

changes.

The relative short-term increase in hospital admissions for disordered eating in the first

year of the pandemic may be attributed to many factors, including the toll of pandemic stress and

the changes in schooling on pediatric eating disorder health.[23,24] Times of change or transition

are associated with increases in and onset of eating disorders in adolescents.[25] These findings are

consistent with other studies demonstrating an increase in hospital admissions for eating

disorders,[6,7,13] increase in eating disorder behaviors,[8,9,10,11] and increase in eating disorder disease

severity[12] during the pandemic worldwide.

After lockdown, we observed a relative and short-term increase in admissions for both

patients with eating disorders and other mental health diagnoses compared to the predicted

hospitalization rate before the lockdown, similar to other interrupted time series on eating

disorders hospital admissions before and during the pandemic.[6,7] One possible explanation for

this temporary increase in admissions includes the loss of school resources and structure for the

pediatric patient population. It has been reported that of adolescents who use mental health

services in a year, 57% used a school based mental health service and 35% accessed their mental

health care exclusively from a school.[2] Disruption of routines and peer connections, exposure to

increased stress or abuse at home, and increased social media usage could also be contributors to

increasing rates of admissions.[3]

Of interest is our finding that the return to school may have been associated with

differential effects between eating disorders and other mental health admissions. While there are

likely varied and multifactorial explanations for this finding, we hypothesize that the social

isolation of school closure had variable impact on the two different patient populations. While

there was lack of access to mental health resources through schools, it is possible that increased

time spent with family units was in fact a protective factor for patients at risk for suicidality, in

the midst of all of the societal stress of the pandemic.[26,27] For adolescents with disordered eating,

we hypothesize that more time spent in the home and with parent(s) could have led to higher

awareness of the adolescents' disordered eating behaviors that would have otherwise been easier

to hide while at school. One other factor to consider with adolescents no longer being around

peers in the school setting is the possibility that they experienced less bullying- which might

have been a protective feature for all students.[27] In addition, Oregon experienced one of the

lowest COVID-19 infection rates per 100,000 citizens in the United States, and it is possible the

stress of the pandemic did not affect Oregon's pediatric psychiatric and eating disorder patient

population as greatly as other states.[28,29] There is a studied seasonality of pediatric mental health

with increasing numbers of emergency department mental health presentations during the fall

and spring and decreasing numbers during the summer.[30,31] The hospital admissions greatly

decreased in July with a slow increase towards September, consistent with the observed

seasonality pattern of pediatric mental health. The level change also could be that after a

complete lack of in-person school, returning to the classroom briefly improved pediatric mental

health before the stresses of school contributed to increased hospital admissions.

We observed a slight increase in transgender patients admitted for both mental health and

eating disorders in the pandemic period which may reflect increases in transgender identity

among youth, which has been described recently using CDC data.[32] However, this population

also experiences discrimination and stress at baseline and may have been disproportionately

impacted by the stressors of the pandemic. Given that only one third of transgender youth live in

home with family acceptance,[33,34] the extended school closures in Oregon may have been a

particular challenge for these youth, in addition to increased barriers to transgender care.

Anecdotally, the severity of pediatric mental health and eating disorder admissions

increased since April 2020. Need for complex care plans, psychiatry, psychology, and 1:1

observation safety needs were higher than normal. Changes to medical unit security and safety

plans were reassessed frequently and dedicated nursing staff were called on to develop de-

escalation plans to keep patients and staff protected. Those with disordered eating diagnoses

were noted to have more significant bradycardia than had been experienced in the years prior in

many cases, and patients reported more severe symptoms temporal to the onset of pandemic

isolation. (personal communication, medical director of acute care unit).

Limitations:

This study has some inherent limitations including examining ecological level changes

and examining changes in admissions associated with those changes. Additionally, chart review

of ICD-10 codes tied to a patient admission is complex. Codes are assigned based on many

system and billing provider structures. For example, a patient being admitted is assigned a code,

each consulting and primary billing provider assigns diagnostic codes to their visit each day, and

there is a discharge diagnosis finally assigned. We were inclusive in examining the number of

patients with an eating disorder, for example, by including all malnutrition diagnostic codes. This

methodology may have included some patients with malnutrition without eating disorders.

Conclusion:

This study adds new information on the impact of the COVID-19 pandemic on children's

mental health using a period of two years of hospital admissions data post-pandemic start,

highlighting both the initial increases observed in other studies but also importantly the

association with return to in-person school. The COVID-19 pandemic highlighted many pre-

existing challenges within our healthcare system— caring for patients with mental health crises

and eating disorder diagnoses within a medical model of care is one of them.

References

1.   General O of the S. U.S. Surgeon General Issues Advisory on Youth Mental Health

Crisis Further Exposed by COVID-19 Pandemic. HHS.gov. Published December 7, 2021.

Accessed March 2, 2022. https://www.hhs.gov/about/news/2021/12/07/us-surgeon-general-

issues-advisory-on-youth-mental-health-crisis-further-exposed-by-covid-19-pandemic.html

2.   Golberstein E, Wen H, Miller BF. Coronavirus Disease 2019 (COVID-19) and Mental

Health for Children and Adolescents. *JAMA Pediatrics*. 2020;174(9):819-820.

doi:10.1001/jamapediatrics.2020.1456

3.   Marques de Miranda D, da Silva Athanasio B, Sena Oliveira AC, Simoes-e-Silva AC.

How is COVID-19 pandemic impacting mental health of children and adolescents? *International*

Journal of Disaster Risk Reduction. 2020;51:101845. doi:10.1016/j.ijdrr.2020.101845

4.   Meherali S, Punjani N, Louie-Poon S, et al. Mental Health of Children and Adolescents

Amidst COVID-19 and Past Pandemics: A Rapid Systematic Review. *Int J Environ Res Public*

Health. 2021;18(7):3432. doi:10.3390/ijerph18073432

5.   Mayne SL, Hannan C, Davis M, et al. COVID-19 and Adolescent Depression and

Suicide Risk Screening Outcomes. *Pediatrics*. Published online June 17, 2021:e2021051507.

doi:10.1542/peds.2021-051507

6.   Otto AK, Jary JM, Sturza J, et al. Medical Admissions Among Adolescents With Eating

Disorders During the COVID-19 Pandemic. *Pediatrics*. Published online July 8,

2021:e2021052201. doi:10.1542/peds.2021-052201

7.   Lin JA, Hartman-Munick SM, Kells MR, et al. The Impact of the COVID-19 Pandemic

on the Number of Adolescents/Young Adults Seeking Eating Disorder-Related Care. *Journal of*

Adolescent Health. Published online July 2021:S1054139X21002792.

doi:10.1016/j.jadohealth.2021.05.019

8.   Phillipou A, Meyer D, Neill E, et al. Eating and exercise behaviors in eating disorders

and the general population during the COVID-19 pandemic in Australia: Initial results from the

COLLATE project. *International Journal of Eating Disorders*. 2020;53(7):1158-1165.

doi:https://doi.org/10.1002/eat.23317

9.   Castellini G, Cassioli E, Rossi E, et al. The impact of COVID-19 epidemic on eating

disorders: A longitudinal observation of pre versus post psychopathological features in a sample

of patients with eating disorders and a group of healthy controls. *Int J Eat Disord*.

2020;53(11):1855-1862. doi:10.1002/eat.23368

10.   Vitagliano JA, Jhe G, Milliren CE, et al. COVID-19 and eating disorder and mental

health concerns in patients with eating disorders. *J Eat Disord*. 2021;9(1):80.

doi:10.1186/s40337-021-00437-1

11.   Termorshuizen JD, Watson HJ, Thornton LM, et al. Early impact of COVID-19 on

individuals with self-reported eating disorders: A survey of ~1,000 individuals in the United

States and the Netherlands. *Int J Eat Disord*. 2020;53(11):1780-1790. doi:10.1002/eat.23353

12.   Springall G, Cheung M, Sawyer SM, Yeo M. Impact of the coronavirus pandemic on

anorexia nervosa and atypical anorexia nervosa presentations to an Australian tertiary paediatric

hospital. *J Paediatr Child Health*. Published online September 27, 2021. doi:10.1111/jpc.15755

13.   Agostino H, Burstein B, Moubayed D, et al. Trends in the Incidence of New-Onset

Anorexia Nervosa and Atypical Anorexia Nervosa Among Youth During the COVID-19

Pandemic in Canada. *JAMA Netw Open*. 2021;4(12):e2137395.

doi:10.1001/jamanetworkopen.2021.37395

14.   Haripersad YV, Kannegiesser-Bailey M, Morton K, et al. Outbreak of anorexia nervosa

admissions during the COVID-19 pandemic. *Arch Dis Child*. 2021;106(3):e15-e15.

doi:10.1136/archdischild-2020-319868

15.   Radhakrishnan L. Pediatric Emergency Department Visits Associated with Mental Health

Conditions Before and During the COVID-19 Pandemic — United States, January 2019–January

2022. *MMWR Morb Mortal Wkly Rep*. 2022;71. doi:10.15585/mmwr.mm7108e2

16.   Hill RM, Rufino K, Kurian S, Saxena J, Saxena K, Williams L. Suicide Ideation and

Attempts in a Pediatric Emergency Department Before and During COVID-19. *Pediatrics*.

2021;147(3):e2020029280. doi:10.1542/peds.2020-029280

17.   Oregon Department of Education : 2020-21 School Status : Health, Safety & Wellness :

State of Oregon. Accessed March 18, 2023. https://www.oregon.gov/ode/students-and-

family/healthsafety/Pages/2020-21-School-Status.aspx

18.   Oregon Department of Education : Health, Safety & Wellness : State of Oregon.

Accessed May 21, 2023. https://www.oregon.gov/ode/students-and-

family/healthsafety/pages/planning-for-the-2020-21-school-year.aspx

19.   Children and COVID-19 Vaccinations Trends: AAP Analysis of Data Posted by the

Centers for Disease Control and Prevention. Presented at: American Academy of Pediatrics;

January 4, 2023.

https://downloads.aap.org/AAP/PDF/Child%20Vaccinations%20Report%20US%20Cumulative

%20and%20Weekly%201.4.2023%20edits.pdf?_ga=2.261804557.1497222321.1676950142-

1546576798.1676950142

20.   Oregon COVID-19 Vaccine Effort Metrics. Tableau Software. Accessed March 18, 2023.

https://public.tableau.com/views/OregonCOVID-

19VaccineEffortMetrics/RaceandEthnicityData?%3Adisplay_static_image=y&%3AbootstrapWh

enNotified=true&%3Aembed=true&%3Alanguage=en-

US&:embed=y&:showVizHome=n&:apiID=host0#navType=0&navSrc=Parse

21.   R: The R Project for Statistical Computing. Accessed August 6, 2022. https://www.r-

project.org/

22.   Coronavirus Disease 2019 (COVID-19). Centers for Disease Control and Prevention.

Published February 11, 2020. Accessed May 21, 2023. https://www.cdc.gov/coronavirus/2019-

ncov/variants/index.html

23.   Sanzari CM, Liu RT. Temporal Trends in Treatment Utilization for Disordered Eating in

U.S. Adolescents From 2004 Through 2017: A Nationally Representative Study. *J Adolesc*

Health. 2019;65(4):564-566. doi:10.1016/j.jadohealth.2019.05.005

24.   Kessler RC, Avenevoli S, Costello EJ, et al. Prevalence, persistence, and

sociodemographic correlates of DSM-IV disorders in the National Comorbidity Survey

Replication Adolescent Supplement. *Arch Gen Psychiatry*. 2012;69(4):372-380.

doi:10.1001/archgenpsychiatry.2011.160

25.   Berge JM, Loth K, Hanson C, Croll-Lampert J, Neumark-Sztainer D. Family life cycle

transitions and the onset of eating disorders: a retrospective grounded theory approach. *J Clin*

Nurs. 2012;21(9-10):1355-1363. doi:10.1111/j.1365-2702.2011.03762.x

26.   Janiri D, Doucet GE, Pompili M, et al. Risk and protective factors for childhood

suicidality: a US population-based study. *Lancet Psychiatry*. 2020;7(4):317-326.

doi:10.1016/S2215-0366(20)30049-3

27.   Ati NAL, Paraswati MD, Windarwati HD. What are the risk factors and protective factors

of suicidal behavior in adolescents? A systematic review. *J Child Adolesc Psychiatr Nurs*.

2021;34(1):7-18. doi:10.1111/jcap.12295

28.   CDC. COVID Data Tracker. Centers for Disease Control and Prevention. Published

March 28, 2020. Accessed May 21, 2023. https://covid.cdc.gov/covid-data-tracker

29.   Covid in the U.S.: Latest Maps, Case and Death Counts - The New York Times.

Accessed May 21, 2023. https://www.nytimes.com/interactive/2021/us/covid-cases.html

30.   Copeland JN, Babyak M, Inscoe AB, Maslow GR. Seasonality of Pediatric Mental Health

Emergency Department Visits, School, and COVID-19. *Pediatric Emergency Care*. Published

online March 15, 2022. doi:10.1097/PEC.0000000000002671

31.   Marshall R, Ribbers A, Sheridan D, Johnson KP. Mental Health Diagnoses and Seasonal

Trends at a Pediatric Emergency Department and Hospital, 2015–2019. *Hospital Pediatrics*.

2021;11(3):199-206. doi:10.1542/hpeds.2020-000653

32.   Herman JL, Flores AR, O'Neill KK. *How Many Adults and Youth Identify as*

Transgender in the United States? Williams Institute School of Law UCLA; 2022:25.

https://williamsinstitute.law.ucla.edu/wp-content/uploads/Trans-Pop-Update-Jun-2022.pdf

33.   Bleckmann C, Leyendecker B, Busch J. Sexual and Gender Minorities Facing the

Coronavirus Pandemic: A Systematic Review of the Distinctive Psychosocial and Health-related

Impact. *J Homosex*. Published online May 16, 2022:1-22. doi:10.1080/00918369.2022.2074335

34.   Grante JM, Mottet LA, Tanis J, Harrison J, Herman JL, Keisling M. *Injustice at Every*

Turn: A Report of the National Transgender Discrimination Survey. Washington: National

Center for Transgender Equality and National Gay and Lesbian Task Force; 2011:220.

https://transequality.org/sites/default/files/docs/resources/NTDS_Report.pdf

Table 1. Characteristics of patients with admissions in the three time periods examined

Patients with an eating disorder diagnosis					
	Overall (n=851)	Before April 2020 (n=425)	April 2020 through August 2021 (n=302)	September 2021 through April 2022 (n=124)	p-value
Admission age, years (mean (SD))	14.57 (2.48)	14.48 (2.57)	14.63 (2.37)	14.72 (2.43)	0.566
Sex (% female)	579 (68.0)	272 (64.0)	215 (71.2)	92 (74.2)	0.043
Gender identity (%)					0.423
Female	279 (32.8)	156 (36.7)	109 (36.1)	14 (11.3)	
Male	108 (12.7)	64 (15.1)	40 (13.2)	4 (3.2)	
Transgender and other	24 (2.8)	11 (2.6)	10 (3.3)	3 (2.4)	
Unspecified	440 (51.7)	194 (45.6)	143 (47.4)	103 (83.1)	
Ethnicity, Race (%)					0.062
Hispanic	105 (12.3)	55 (12.9)	38 (12.6)	12 (9.7)	
Non-Hispanic, White	589 (69.2)	312 (73.4)	194 (64.2)	83 (66.9)	
Non-Hispanic, Black	27 (3.2)	10 (2.4)	12 (4.0)	5 (4.0)	
Non-Hispanic, Asian	29 (3.4)	14 (3.3)	12 (4.0)	3 (2.4)	
Non-Hispanic, Other	19 (2.2)	4 (0.9)	8 (2.6)	7 (5.6)	
Missing	82 (9.6)	30 (7.1)	38 (12.6)	14 (11.3)	
Length of stay, days (median [IQR])	6.00 [4.00, 9.00]	6.00 [4.00, 9.00]	6.00 [4.00, 9.00]	6.00 [4.00, 8.00]	0.961
Insurance (%)					0.088
Medicaid / Medicare	353 (41.5)	192 (45.3)	114 (37.7)	47 (37.9)	
Private	497 (58.5)	232 (54.7)	188 (62.3)	77 (62.1)	
Patients with a mental health diagnosis					
	Overall (n=1506)	Before April 2020 (n=815)	April 2020 through August 2021 (n=502)	September 2021 through April 2022 (n=189)	p-value
Admission age, years (mean (SD))	13.72 (3.25)	13.67 (3.34)	13.69 (3.26)	14.07 (2.83)	0.292
Sex (% female)	953 (63.3)	500 (61.3)	323 (64.3)	130 (68.8)	0.139
Gender identity (%)					0.156
Female	512 (34.0)	311 (38.2)	177 (35.3)	24 (12.7)	
Male	256 (17.0)	163 (20.0)	86 (17.1)	7 (3.7)	
Transgender and other	45 (3.0)	21 (2.6)	21 (4.2)	3 (1.6)	
Unspecified	693 (46.0)	320 (39.3)	218 (43.4)	155 (82.0)	
Ethnicity, Race (%)					0.373
Hispanic	248 (16.5)	145 (17.8)	80 (15.9)	23 (12.2)	
Non-Hispanic, White	971 (64.5)	540 (66.3)	319 (63.5)	112 (59.3)	
Non-Hispanic, Black	51 (3.4)	27 (3.3)	17 (3.4)	7 (3.7)	
Non-Hispanic, Asian	38 (2.5)	27 (3.3)	5 (1.0)	6 (3.2)	
Non-Hispanic, Other	41 (2.7)	23 (2.8)	12 (2.4)	6 (3.2)	
Missing	157 (10.4)	53 (6.5)	69 (13.8)	35 (18.6)	
Length of stay, days (median [IQR])	3.00 [2.00, 6.00]	3.00 [2.00, 6.00]	4.00 [2.00, 7.00]	3.00 [2.00, 6.00]	0.188
Insurance (%)					0.209
Medicaid / Medicare	750 (49.9)	422 (52.0)	241 (48.1)	87 (46.0)	
Private	752 (50.1)	390 (48.0)	260 (51.9)	102 (54.0)	

IQR = interquartile range; SD = standard deviation; Note that p-values for Fisher's exact tests do not reflect the missing categories

Figure 1. a) Graphical representation looking at hospital admissions by eating disorder, other mental health and non-mental health encounters over time and b) Interrupted time series analysis of the same data with onset of the pandemic and return to in-person school examined as interruptions

a)

b)

Figure 2. Interrupted time series analysis of admissions to the hospital with a diagnosis of eating disorder in a) the first year post-COVID and b) over the longer term (two years)

a) eating disorders in the first year of the pandemic

b) eating disorders over the first two years of the pandemic

Figure 3. Interrupted time series analysis of admissions to the hospital with a diagnosis of other mental health (non-eating disorder) in a) the first year post-COVID and b) over the longer term (two years)

a) Other mental health disorder admissions in the first year of the pandemic

b) Other mental health admissions over the first two years of the pandemic