Validity, reliability, and readability of single-item and short physical activity

questionnaires for use in surveillance: a systematic review

Authors:

Antonina Tcymbal[1*], Sven Messing[1], Rachel Mait[1], Roberto Galindo Perez[1], Taiyeba

Akter[1], Ivo Rakovac[2], Peter Gelius[1], Karim Abu-Omar[1]

1 Department of Sport Science and Sport, Friedrich-Alexander-Universität Erlangen-

Nürnberg, Gebbertstr. 123b, 91058 Erlangen, Germany

2 World Health Organization Regional Office for Europe, Marmorvej 51, DK-2100

Copenhagen, Denmark

* Corresponding author:

Dr. Antonina Tcymbal

antonina.tcymbal@fau.de

Abstract

Background: Accurate and fast measurement of physical activity is important for surveillance. Even though many physical activity questionnaires (PAQ) are currently used in research, it is unclear which of them is the most reliable, valid, and easy to use. This systematic review aimed to identify existing brief PAQs, describe and compare their measurement properties, and assess their level of readability.

Methods: We performed a systematic review based on the PRISMA statement. Literature searches were conducted in six scientific databases in March 2022. Articles were included if they evaluated validity and/or reliability of brief (i.e., with a maximum of three questions) physical activity or exercise questionnaires intended for healthy adults. Due to the heterogeneity of studies, data were summarized narratively. The level of readability was calculated according to the Flesch-Kincaid formula.

Results: In total, 34 articles published in English or Spanish were included, evaluating 31 distinct brief PAQs. The studies indicated moderate to good levels of reliability for the PAQs. However, the majority of results showed weak validity when validated against objective measurements and demonstrated weak to moderate validity when validated against other PAQs. Most of the assessed PAQs met the criterion of being "short," allowing respondents to complete them in less than one minute either by themselves or with an interviewer. However, only 17 questionnaires had a readability level that indicates that the PAQ is easy to understand for the majority of the population.

Conclusions: This review identified a variety of brief PAQs, but most of them were evaluated in only a single study. The methods used to assess measurement properties varied widely across studies, limiting the comparability between different

PAQs and making it challenging to identify a single tool as the most suitable.

Furthermore, PAQs employed different concepts for measuring physical activity,

necessitating consideration of measurement properties and assessment goals when

selecting a specific tool. None of the evaluated brief PAQs allowed for the

measurement of whether a person fulfills the main WHO physical activity

recommendations. Future development or adaptation of PAQs should prioritize

readability as an important factor to enhance their usability.

Keywords: physical activity, questionnaire, measurement properties, validity,

reliability, readability, surveillance

Background

It has been demonstrated that regular physical activity (PA) can help improve

physical and mental functions as well as reverse some effects of chronic diseases

[1]. Regularly engaging in 150 minutes of PA per week [2], alongside following a

healthy diet and abstaining from smoking and alcohol consumption, is seen as being

key to prevent non-communicable diseases.

However, accurate measurement of PA levels is important to determine the amount

of activity needed to improve health and identify links with other health outcomes and

behaviors [3]. To this day, self-report questionnaires are the most common measures

to collect PA data, often as part of surveillance systems such as the Behavioral Risk

Factor Surveillance System [4], the WHO STEPwise approach to noncommunicable

disease risk factor surveillance [5], and the European Health Interview Survey [6].

While such self-report questionnaires have seen widespread use because of their

efficiency, they have also been shown to have limitations related to bias and data accuracy [7].

Many different self-report questionnaires have been developed over the years with the Global Physical Activity Questionnaire (GPAQ), the International Physical Activity Questionnaire (IPAQ; also available as a short form, IPAQ-SF), and the European Health Interview Survey Physical Activity Questionnaire (EHIS-PAQ) being the most widely utilized in global PA surveillance [8]. All three questionnaires assess PA across different domains, asking respondents to report their PA in a typical week (GPAQ, EHIS-PAQ) or during the last 7 days (IPAQ-SF). All three are comparatively complex and range from seven items (IPAQ-SF) to 16 (GPAQ). Nevertheless, the measurement properties of these questionnaires are modest. For GPAQ [9], IPAQ-SF [10, 11], and the EHIS-PAQ [12], different studies have demonstrated reasonable reliability but comparatively low validity. In a recent review [8] of IPAQ-SF, GPAQ, and EHIS-PAQ, the questionnaires showed low-to-moderate validity against objective measures of PA such as accelerometers, and moderate-to-high validity against subjectively measured PA such as other questionnaires.

It is well known that questionnaires with many items can increase the response burden on respondents [13, 14], which has resulted in the development of several short self-report instruments. In contrast to the detailed questionnaires mentioned above, the purpose of short PA questionnaires is to simplify and speed up the procedure for assessing PA levels.

In addition, language-related difficulties are currently an important topic in public health. Patient education materials can increase patient compliance, but only if they are written in a language that is easy for the patient to understand [15]. Regarding PA questionnaires, Altschuler et al. [16] found significant gaps between respondents'

interpretations of some PA questions and researchers' original assumptions about

what those questions were intended to measure. One of the characteristics of

language difficulty is the level of readability, which indicates how easily readers can

understand the text. Research of texts used in healthcare consistently shows that

materials intended for patients often require a high level of education and are too

complicated for the average person [15, 17]. In relation to physical activity

questionnaires (PAQs), readability can influence the amount of time a person needs

to understand the question and, if the text is too complicated, may potentially

decrease the response rate and correctness of the answer.

A number of existing reviews have investigated the measurement properties of PA

questionnaires. Van Poppel et al. [18] reviewed the validity and reliability

methodology of 85 versions of PAQs with no clear consensus regarding the best

questionnaire for PA measurement. Helmerhost et al. [19] studied reliability and

objective criterion-related validity of 34 newly-developed and 96 existing PAQs.

However, to our knowledge, a dedicated review of short and brief PA questionnaires

measurement properties has not yet been performed. Therefore, the aim of this

systematic review is to identify existing brief PAQs and to comparatively describe

their measurement properties and level of readability.

Methods

This review follows the guidelines of the PRISMA (Preferred Reporting Items for

Systematic reviews and Meta-Analyses) statement [20] and the Consensus-based

Standards for the Selection of Health Measurement Instruments (COSMIN) guideline

for systematic reviews of patient-reported outcome measures [21].

Information sources and search strategy

A systematic search was performed in six databases (PubMed, Web of Science,

Scopus, CINAHL, SportDiscus, PsycInfo) in March 2022. Additionally, reference lists

from previous reviews of PA questionnaires and other relevant publications were

screened to identify additional studies. A comprehensive search strategy was

developed with a combination of keywords in the categories of measured construct

(physical activity) and type of instruments (brief/short questionnaire).

The resulting search string was as follows:

("physical activit*" OR "physical inactivit*")

AND

(questionnaire OR measure OR evaluat* OR assess* OR surveillance OR monitor*

OR screening)

AND

("single item" OR single-item OR "single question" OR "one item" OR one-item OR

"one question" OR "brief questionnaire " OR "short questionnaire " OR "brief

assessment" OR "short assessment" OR "two-item*" OR "two-item*" OR "two

questions" OR "brief physical activity assessment" OR "Single response" OR "Single-

response")

No restrictions were made regarding language or publication date.

Eligibility criteria

Articles were included in the review if they fulfilled the following eligibility criteria:

1. The article described a self-report PA or exercise questionnaire intended for

healthy adults.

2. The evaluated questionnaire was brief and included a maximum of three

questions.

3. The article investigated one or more measurement properties of the

questionnaire.

4. The article was published in a peer-reviewed journal.

Articles were excluded based on the following exclusion criteria:

1. The article focused on a questionnaire that measured only physical inactivity,

screen time, or sedentary behavior.

2. The evaluated questionnaire was intended solely for children and adolescents or

people with specific conditions.

3. The article did not investigate the measurement properties of the questionnaire.

Study selection

Two reviewers independently screened and selected the relevant articles. First, all

articles were screened based on titles and abstracts. If the title and/or abstract

indicated that the study fulfilled the inclusion criteria, both reviewers screened the full

text for eligibility. When necessary, supplementary files were also reviewed for

additional information. Disagreements between the reviewers were discussed within

the research team until a consensus was reached.

Records were managed using the Covidence systematic review software (Veritas

Health Innovation, Melbourne, Australia; www.covidence.org) and EndNote X9

(Clarivate Analytics, Philadelphia, PA, USA).

Data extraction

Data of included studies were extracted and summarized by one reviewer and

verified by a second reviewer to reduce bias and error. Discrepancies were

discussed between reviewers to achieve consensus. Extracted information included:

publication details (first author, year of publication, country), sample characteristics

(number of participants, age category, special health conditions), the measurement

tool(s) explored, who assessed PA levels of participants, assessed measurement

properties, other measurement tool(s) used as a comparison, reliability test-retest

interval, and the results of the study.

Risk of bias assessment

The methodological quality of the individual studies included in this review was

assessed with 18 questions based on the Appraisal tool for Cross-Sectional Studies

(AXIS) [22]. In order to specifically assess the quality of the tools' validity and

reliability measurement properties, some questions were modified based on the

COSMIN risk of bias checklist [21] and on suggestions from previous systematic

reviews examining PA assessment measures [23]. This quality assessment

evaluated articles based on study design, sample size, participant selection process,

appropriate blinding, examiner experience, method of measurement, adequate data

reporting, internal consistency, and six other categories (see S1 file). Risk of bias

assessment was conducted independently by two reviewers, with any discrepancies

resolved through discussion in the research team. Studies were scored 1 if they

satisfied the quality element, and 0 if they did not. The summary score (range: 0–18)

indicates the risk of bias, with a higher score indicating higher quality and therefore a

lower risk of bias.

Data synthesis and analysis of measurement properties

The primary objective of this systematic review was to identify existing short

questionnaires suitable for assessing PA levels in surveillance and primary care

settings, and to compare their measurement properties. To accomplish this goal,

relevant information from the included studies was summarized separately for each

questionnaire. In order to evaluate the reliability and validity of the identified

questionnaires, a range of tests were employed in the included studies. Some

studies reported results for a total questionnaire summary score, while others

assessed reliability and validity for specific aspects, intensities, or domains of the

questionnaire. Additionally, certain studies examined these measurement properties

within subgroups of the test population. Due to the heterogeneity in the methods

used and the lack of standardized reporting across studies, a quantitative meta-

analysis was not feasible. Consequently, the information from the included studies

was summarized narratively, highlighting the key findings for each questionnaire.

This narrative synthesis allows for a comprehensive overview of the reliability and

validity findings, highlighting strengths and limitations of each questionnaire.

Analysis of length and readability of questionnaires

To determine how quickly questionnaires could be answered and how easy it is to

understand the questions, identified PAQs were analyzed for word count and

readability level. The expected time the tool would take for self-administration (silent

reading speed) and interviewer administration (spoken word speed) was calculated

based on the respective questionnaire's word count and English reading speeds

established by Brysbaert [24]. The level of readability was calculated according to the

Flesch-Kincaid formula, which was chosen because it is the most commonly used

tool to calculate the readability level of written health information [25]. The formula

has two forms: the Flesch Reading-Ease-Score, and the Flesch–Kincaid Grade

Level. The Flesch Reading-Ease-Score test produces a score from 0 to 100, and

higher scores indicate material that is easier to read; lower scores mark passages

that are more difficult to read. The Flesch–Kincaid Grade Level formula matches the

text to the grade level achievement (number of years of education) required to

understand the text. We applied both formulas for each questionnaire using an online

calculator [26].

Results

Study selection process

The search across six databases resulted in 2,422 publications. After removing 1,255

duplicates, 1,167 articles were screened based on title and abstract. 65 studies were

found eligible for full text assessment. One recently published article [27] and one

article identified while screening the reference lists of relevant publications [28]

satisfied all eligibility criteria and were therefore included in the analysis.

Subsequently, 33 full text articles were excluded due to the length of the

questionnaire (n=14), the publication type (n=14), a lack of self-reporting on PA and

exercise (n=2), and a study design that did not consider validity and reliability (n=3).

In total, 34 studies were included for data extraction and methodological quality

assessment. A summary of the search results is presented in Figure 1.

Figure 1. PRISMA flowchart of study eligibility (PA=physical activity)

Characteristics of included studies

A summary of the characteristics of the 34 included studies is presented in S1 Table.

The included studies were conducted in Western European countries (n=14), the

USA (n=10), Australia (n=5), Canada (n= 2), New Zealand (n=2), and Japan (n=1).

A total of 114,143 adults were assessed using the 31 unique brief PAQs (with sample

sizes ranging from 9 to 39,379). In 25 studies, the sample consisted only of healthy

adults, and nine studies also included specific populations such as older adults [29]

[30] [31], overweight/obese adults [32, 33], as well as patients with rheumatoid

arthritis [34], coronary heart disease [35] [36], and chronic obstructive pulmonary

disease [37].

Thirty-one studies documented who completed the PAQ. In 16 studies, respondents

filled out the questionnaire by themselves, and in 15 studies the PA level was

collected from an interviewer reading a questionnaire.

Questionnaires asked for the amount of PA (n=6), the number of days per week with

a sufficient amount of PA (n=4), general exercise participation (n= 3), self-reported

activity compared with peers (n=2), or for respondents to choose a categorical

descriptor of PA levels ranging from "inactive" to "very active" (n=16).

Seven of the included studies investigated the measurement properties of the Single

Item Physical Activity Measure (SIPAM) [27, 30, 38-42], five were related to the Brief

Physical Activity Assessment Tool (BPAAT) [37, 43-46], and two to the Stanford

Leisure-Time Activity Categorical Item (L-Cat) [32, 33]. The other 28 PAQs were

assessed in a single study.

Risk of bias assessment

The average quality score of included studies was 11.75, with a range between 7 and

17 (out of 18). Thirteen studies received 14 or more points, which can be considered

a high methodological quality. In most of the studies, the aims and objectives, the

process of the measurement properties investigation, the statistical analysis and the

results were sufficiently described, and the study design was appropriate. However,

the included papers reported poorly on whether the examiners had enough

experience with the tool and whether they were blinded to participant characteristics,

previous findings, or other observed findings. Only four studies justified their choice

of sample size. Additionally, many studies failed to consistently report reasons for

dropout and characteristics of non-responders. Quality scores of the individual

studies are reported in S1 Table.

Questionnaires' measurement properties

A summary of the reliability, validity, and diagnostic test accuracy data found in the

identified studies is presented in S2 Table. Studies commonly use a number of

different statistical analyses to define absolute (agreement between the two

measurement tools) or relative (the degree to which the two measurement tools rank

individuals in the same order) validity and reliability. These types of statistical

analysis include correlations (Pearson's; Spearman's; interclass), regression, kappa

statistics, and area under the receiver operating curve (AUC-ROC).

Eleven studies measured the reliability of brief PAQs. All of them used a test-retest

procedure to measure the consistency of the PAQs. Statistical methods and test-

retest intervals varied widely between studies. Overall, studies showed moderate to

good reliability levels of the PAQs.

The validity of PAQs was assessed in 33 studies. As a "gold standard" for validation,

13 studies used other PAQs, 15 studies validated brief PAQs against accelerometers

or pedometers, and 11 studies compared results of the brief PAQs to other objective

measurements, such as BMI [28, 32, 47, 48], VO_2 max [47, 49], or doubly labeled

water [34]. Validity coefficients were in general considerably lower than reliability

coefficients. The majority of results showed weak validity of brief PAQs against

objective measurements and weak to moderate validity against other PAQs.

Length and complexity of the questionnaires

The texts of all questionnaires and information about their length and readability level

are presented in S3 Table. It should be noted that, while some PAQs were created

and conducted in other languages, length and readability were calculated for the

English versions.

Most of the assessed PAQs (n=26) can be read by in silence by the respondent or

aloud by the interviewer in less than one minute. However, some questionnaires

presented as "brief" by the authors of the identified studies contain a large amount of

text and require more than one minute for reading: the Nordic Physical Activity

Questionnaire (NPAQ) short, the Total Activity Measure (TAM), the Self-report Scale

to Assess Habitual Physical Activity, the Stanford Leisure-Time Activity Categorical

Item, and the Stanford Brief Activity Survey.

The calculation of the readability levels with the Flesch-Kincaid formula showed that

only 17 out of 31 brief PAQs had readability levels of "easy to read" (n=9) or "plain

English," (n=8) and can be easily understood by the majority of the population (see

Figure 2). Other questionnaires have long sentences and/or many complicated words

(three syllables and more), making them difficult to read. Seven questionnaires

scored as "difficult to read" and require college-level education; it is likely that people

with lower education levels will find them hard to understand.

Figure 2. Readability levels for brief PA questionnaires (English language versions).

Comparison of brief PAQs

Data on both validity and reliability were only available for nine of the 31 assessed

PAQs. These brief PAQs were chosen for an in-depth comparison. Their summarized

measurement properties and linguistic characteristics are presented in Table 1.

Table 1. Comparison of the brief physical activity questionnaires by reliability, validity,

length and readability.

Tool	Reliability	Validity		Word count	Readability
		Against objective measurement	Against other PAQ		
Single Item Physical Activity Measure (SIPAM)	Good (0,63-0,82)	Poor to Good (0,22-0,81)	Poor to Good (0,33-0,81)	63 – original 111 – parents version	Fairly difficult to read Parents version - Very difficult to read

Brief Physical Activity Assessment Tool (BPAAT)	Moderate to good (0,53-0,72)	Poor to moderate (0,18-0,43)	Moderate to good (0,45-0,64)	106	Difficult to read
Three-Question Assessment	Moderate to good (0,56-0,63)	Poor (0,24-0,31)	Moderate (0,39-0,43)	140	Fairly difficult to read
Nordic physical activity questionnaire (NPAQ) - short (open ended) *	Good (0,82)	Poor (0,33)	NM	237	Difficult to read
Total Activity Measure (Version 2)	Good (0,82)	Poor (0,36-0,38)	NM	241	Fairly difficult to read
Japan Collaborative Cohort (JACC) Questionnaire*	Moderate (0.39 - 0.56)	NM	Moderate (0.43-0.60)	90	Plain English
Relative PA Question (Compared to peers)	Moderate (0,56)	Poor to moderate (0,28-0,57)	NM	25	Easy to read
The Seven-level Single-Question Scale *	Good (0,7)	Poor (0,33)	NM	144	Fairly difficult to read
Absolute PA Question	Good (0,7)	Poor (0,1-0,33)	NM	34	Fairly difficult to read
A single-item 5-point rating of usual PA (Usual PA Scale)	Good (0,68-0,88)	NM	Good (0,66)	147	Plain English
The Stanford Leisure-Time Activity Categorical Item (L-CAT)	Good (0,64-0,8)	Poor (0,36-0,38)	NM	331	Plain English

* = the study used not the English version of the questionnaire, PAQ=physical activity questionnaire

<u>The Single Item Physical Activity Measure (SIPAM)</u> uses a single question to assess

the number of days per week on which 30 minutes or more of PA are performed

(excluding housework and work-related PA). For this PAQ, the largest number of

studies on validity and reliability were found. The SIPAM showed good reliability

levels in all studies, however, results on validity varied considerably between studies

– from poor to good – against both objective measurements and other subjective

PAQs. Zwolinsky et al. (2015) also measured SIPAM's ability to identify people that

meet or do not meet WHO PA recommendations. The tool showed a low diagnostic

capacity compared to the IPAQ. The agreement between the SIPAM and the IPAQ

was low for identification of participants who meet WHO PA recommendations

(kappa= 0.13, 95% CI 0.12 to 0.14) and moderate for the classification of inactive

participants (kappa= 0.45, 95% CI 0.43 to 0.47) [42].

The Brief Physical Activity Assessment Tool (BPAAT) consists of two questions, one

regarding the frequency and duration of vigorous PA and the other regarding

moderate PA and walking performed in an individual's usual week. By combining the

results of both questions (scores can range from 0 to 8), the subject can be classified

as insufficiently (0–3 score) or sufficiently active (≥4 score). Results of reviewed

studies showed that the BPAAT has moderate to good levels of reliability and validity

in comparison with other PAQs, however comparison with accelerometers identified

only poor to moderate validity.

Smith et al. (2005) evaluated the Three-Question Assessment variant of the BPAAT

which has separate questions about moderate PA and walking. The results of the

study did not find a considerable difference in validity and reliability between the two-

and three-question versions. One study also reported that more physicians preferred

the two-question version (BPAAT) as it is shorter and therefore easier to use (Smith,

2005).

The Nordic Physical Activity Questionnaire-short (NPAQ-short) includes one question

about moderate to vigorous PA and a second question about vigorous PA. Danquah

et al. (2018) compared open-ended and closed-ended questions with the open-

ended version achieving better results. The open-ended version showed good

reliability but performed similarly to other questionnaires, showing poor validity when

compared against objective PA measures. The analyses showed that the

questionnaire was one of the longest and was rated "difficult to read". The agreement

with accelerometer data in identification of people that meet or do not meet the WHO

PA recommendations was low (kappa=0.42).

The Total Activity Measures (TAM) includes three open-ended questions about strenuous, moderate, and mild PA. The revised second version, TAM2, asks about the total time spent at each activity level over a 7-day period. The TAM2 showed good reliability when validated against objectively-measured PA. The word count was high at 241, and readability was rated as "fairly difficult".

The Japan Collaborative Cohort (JACC) Questionnaire has three items. Two questions focus on leisure-time PA, i.e., time per week engaging in sport or PA (with options ranging from "little" to "at least 5 hours"), and frequency of engagement in sport over the past year (options from "seldom" to "at least twice a week"). The third question asks about daily walking patterns (options from "little" to "more than 1 hour"). The questionnaire showed moderate reliability after one year and moderate validity when compared against a more in-depth interview by a trained researcher. The questions are 90 words in total and were rated as "plain English."

The Relative PA Question is the shortest of the compared PAQs and has an easy readability level. It allows respondents to compare their level of PA with other people of the same age and to choose from five categories ranging from "much more active" to "much less active." The questionnaire has a moderate reliability level and showed poor validity against objective measures.

The Absolute PA Question asked participants to choose what best describes their activity level from three options: (1) vigorously active for at least 30 minutes, three times per week; (2) moderately active at least three times per week; or (3) seldom active, preferring sedentary activities. The questionnaire is relatively short, but as most of the other brief PAQs, it showed high reliability and poor validity against objective measures.

The <u>Seven-Level Single-Question Scale (SR-PA L7)</u> requires respondents to assign

themselves to a level ranging from "I do not move more than is necessary in my daily

routines/chores" and "I participate in competitive sports and maintain my fitness

through regular training." These seven items are supposed to categorize respondents

as maintaining low, medium, or high PA. The SR-PA L7 showed poor validity when

compared to objectively measured PA but good reliability. The scale is moderately

long at 144 words and was rated as "fairly difficult to read."

<u>A single-item 5-point rating of usual PA (Usual PA Scale)</u> includes descriptions of

three PA levels: highly active, moderately active, and inactive. Respondents need to

read a description of each category and identify their usual PA level from the list. The

questionnaire's readability level was ranked as "plain English," and it showed good

reliability and validity levels. However, validation was done only against another PAQ

which is not well-known and requires further research.

<u>The Stanford Leisure-Time Activity Categorical Item (L-CAT)</u> is a single-item

questionnaire that consists of six descriptive PA categories ranging from inactive to

very active. Each category consists of one or two statements describing common

activity patterns over the past month, differing in frequency, intensity, duration, and

types of activity. The categories are described in "plain English," but this made the

questionnaire longer than the other questionnaires. The L-CAT showed high

reliability but poor validity against pedometer and accelerometer validation.

Discussion

This review assessed the validity, reliability, and readability of brief PAQs for adults.

To our knowledge, it is the first review focused specifically on brief PAQs and also

the first to assess their readability. The sheer number of brief PAQs we identified

(n=31) indicates a high research interest in such instruments. However, it also

indicates a lack of harmonization when it comes to PA assessment using brief

questionnaires.

Overall, most PAQs were reported to have poor or moderate validity. They showed

higher validity levels against other self-reported tools than against objective PA

measures. Although reliability is an important measurement property, it was

assessed only in 11 studies. About half of all PAQs showed a moderate to good level

of reliability. These results are in line with other reviews of PAQs [18, 19, 50, 51].

A significant difficulty in conducting this review was that the studies used different

methods for validation, varying time-intervals between repeated measurements, and

different statistical methods to analyze data. Complete data on validity and reliability

were only available for nine PAQs, and only those could be compared in greater

detail. All of these factors prevented a more detailed comparison of results across

PAQs. In general, the methodological quality of the included studies was modest.

The most common flaws were comparably small sample sizes, a lack of sample size

justification, and a poor description of the validity and reliability assessment process.

Additionally, most studies utilized convenience samples, making it impossible to

assess if measurement properties would differ between adults with different levels of

socioeconomic status and/or educational attainment.

Also, the included PAQs used different concepts of measuring PA, further

complicating a direct comparison between instruments. For example, the SIPAM

measures on how many days per week respondents perform 30 minutes or more of

PA; the Absolute PAQ requires respondents to choose from several descriptions of

different PA levels; and the Relative PA question asks them to compare their level of

PA with peers. Some PAQs, such as the TAM, aim to assess the total volume of

moderate-to-vigorous PA. Others focus on particular PA domains. And yet others

assess only leisure-time PA. This variety can be partly explained by efforts to keep

PAQs short at the price of excluding some PA dimensions (type, duration, intensity,

volume). However, it could also be interpreted as a lack of common understanding

about which dimensions of PA should be assessed with brief PAQs. It should also be

taken into account that none of the reviewed brief PAQs allowed for the

measurement of whether a person fulfills the WHO physical activity

recommendations [52].

All of the included studies were conducted in highly developed nations, and 23 of

them took place in English-speaking countries. This most certainly biased the results,

since terminology related to PA differs between languages, as do levels of literacy

[53]. Ultimately, to assess reliability and validity of PAQs, many more studies should

be conducted in developing nations to obtain a more accurate assessment of their

suitability for international surveillance systems. Certainly, in order to stimulate such

research, funding opportunities need to be made available to research teams from

such nations.

The results of this study point out that many PAQs have low readability levels. This is

particularly disturbing when considering that most of these PAQs were tested in

developed nations with comparatively high levels of literacy. Potentially, poor

readability is related to the modest measurement properties that PAQs commonly

come with. Questionnaires that are more readable or might even have been co-

developed [54] with the intended population groups are likely to yield better

measurement properties. This would also strengthen the case of integrating them

into PA surveillance systems. However, at this point in time, there is a dearth of

research relating PAQs and their measurement properties to readability. Advancing

knowledge in this field would also benefit longer PA questionnaires (such as IPAQ

and GPAQ) that are widely utilized in surveillance and potentially score low on

readability as well.

This review comes with certain limitations. The search was limited to scientific

databases, and only studies published in peer reviewed scientific journals were

included. This can lead to a publication bias, as all other types of publications and

gray literature were excluded. The varying measurement methods and conditions

complicated the comparison of findings from different studies and limited data

analysis to a narrative description of differences between tools. This led to more

subjective results and greater difficulty in identifying the tools with the best

measurement properties. It should be also taken into account that the Flesch-Kincaid

readability formula was created for longer texts and not adapted for short

questionnaires. Consequently, the readability levels of PAQs presented here should

be interpreted with caution.

Conclusion

This systematic review sheds light on the validity, reliability, and readability of short

PAQs. Results indicate that additional research is needed, notably regarding their

reliability and validity, their readability, and their applicability to non-English speaking

and/or developing countries.

Recent years have seen a shift from self-report questionnaires towards objective

measures in PA surveillance. This shift has been partially motivated by the

persistently modest measurement properties of PAQs, as well as technological

advancements in the quality and affordability of accelerometers and other PA

measurement tools. However, objective PA assessment comes with its own set of

limitations [55], and self-report and objective measures have been described as

measuring entirely different parameters. Right now, it seems unclear what the future

for PA surveillance might hold and if brief PAQs will continue to play a role.

In this regard, the key merit of short PAQs is that they require substantially less time

to complete than the established PA surveillance questionnaires (GPAQ, IPAQ, and

EHIS-PAQ). This makes them highly appealing for use in surveillance systems. In

order for them to be included in existing systems, more studies would need to

investigate how they compare to longer PAQs. Potentially, if their validity and

reliability are found to be at a comparable level, we might see surveillance systems

gradually shift from longer towards briefer PAQs.

List of abbreviations:

BPAAT: Brief Physical Activity Assessment Tool

EHIS-PAQ: European Health Interview Survey Physical Activity Questionnaire

GPAQ: Global Physical Activity Questionnaire

IPAQ: International Physical Activity Questionnaire

JACC: Japan Collaborative Cohort Questionnaire

L-CAT: The Stanford Leisure-Time Activity Categorical Item

NPAQ-short: The Nordic Physical Activity Questionnaire-short

PA: physical activity

PAQ: physical activity questionnaire

SIPAM: Single Item Physical Activity Measure

SR-PA L7: The Seven-level Single-Question Scale for Self-Reported Leisure Time
Physical Activity

TAM: Total Activity Measure

Declarations

Ethics approval and consent to participate: Not applicable.

Availability of data and materials: Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.

Competing interests: The authors declare that they have no competing interests.

Funding: This research was conducted as part of projects funded by the German Federal Ministry of Health (ZMI5-2522WHO001, ZMI5-2523WHO001). The ministry was neither involved in writing this manuscript nor in the decision to submit the article for publication. Furthermore, we acknowledge financial support by Deutsche Forschungsgemeinschaft and Friedrich-Alexander-Universität Erlangen-Nürnberg within the funding programme "Open Access Publication Funding".

Authors' contributions: AT, KA-O and IR conceptualized the study. AT conducted the literature search. AT, RA, RGP, and TA performed abstract/title screening, full-text selection, data extraction, and quality assessment. AT with support from KA-O, SM and PG prepared a draft of the manuscript. All authors critically reviewed the draft and approved the final manuscript.

References:

1.   McPhee JS, French DP, Jackson D, Nazroo J, Pendleton N, Degens H. Physical activity in older age: perspectives for healthy ageing and frailty. Biogerontology. 2016;17(3):567-80. Epub 20160302. doi: 10.1007/s10522-016-9641-0. PubMed PMID: 26936444; PubMed Central PMCID: PMCPMC4889622.

2.   World Health Organization. Guidelines on physical activity and sedentary behaviour. Geneva: World Health Organization; 2020.

3.   Welk G. Physical activity assessments for health-related research: Human

Kinetics; 2002.

4.   Centers for Disease Control and Prevention. Behavioral Risk Factor

Surveillance System Survey Questionnaire. Atlanta, Georgia: U.S. Department of

Health and Human Services, Centers for Disease Control and Prevention; 2021.

5.   World Health Organization. WHO STEPS Surveillance Manual: The WHO

STEPwise approach to noncommunicable disease risk factor surveillance. 2020.

6.   Finger JD, Tafforeau J, Gisle L, Oja L, Ziese T, Thelen J, et al. Development

of the European Health Interview Survey - Physical Activity Questionnaire (EHIS-

PAQ) to monitor physical activity in the European Union. Arch Public Health.

2015;73:59. Epub 20151202. doi: 10.1186/s13690-015-0110-z. PubMed PMID:

26634120; PubMed Central PMCID: PMCPMC4667448.

7.   Sallis JF, Saelens BE. Assessment of physical activity by self-report: status,

limitations, and future directions. Research quarterly for exercise and sport.

2000;71:1-14. doi: https://doi.org/10.1080/02701367.2000.11082780.

8.   Sember V, Meh K, Sorić M, Starc G, Rocha P, Jurak G. Validity and Reliability

of International Physical Activity Questionnaires for Adults across EU Countries:

Systematic Review and Meta Analysis. International journal of environmental

research and public health. 2020;17(19):7161. Epub 20200930. doi:

https://doi.org/10.3390/ijerph17197161. PubMed PMID: 33007880; PubMed Central

PMCID: PMCPMC7579664.

9.   Bull FC, Maslin TS, Armstrong T. Global physical activity questionnaire

(GPAQ): nine country reliability and validity study. J Phys Act Health. 2009;6(6):790-

804. doi: 10.1123/jpah.6.6.790. PubMed PMID: 20101923.

10.   Craig CL, Marshall AL, Sjöström M, Bauman AE, Booth ML, Ainsworth BE, et

al. International physical activity questionnaire: 12-country reliability and validity.

Medicine and science in sports and exercise. 2003;35(8):1381–95. doi: 10.1249/01.MSS.0000078924.61453.FB.

11.   Lee PH, J. MD, Lam TH, Stewart SM. Validity of the International Physical Activity Questionnaire Short Form (IPAQ-SF): a systematic review. The international journal of behavioral nutrition and physical activity. 2011;8:115. doi: https://doi.org/10.1186/1479-5868-8-115.

12.   Baumeister SE, Ricci C, Kohler S, Fischer B, Topfer C, Finger JD, et al. Physical activity surveillance in the European Union: reliability and validity of the European Health Interview Survey-Physical Activity Questionnaire (EHIS-PAQ). The international journal of behavioral nutrition and physical activity. 2016;13:61. Epub 20160523. doi: 10.1186/s12966-016-0386-6. PubMed PMID: 27215626; PubMed Central PMCID: PMCPMC4877949.

13.   Ware J, Jr., M. K, Keller SD. A 12-Item Short-Form Health Survey: construction of scales and preliminary tests of reliability and validity. Medical Care. 1996;34(3):220-33. doi: https://doi.org/10.1097/00005650-199603000-00003.

14.   Rolstad S, Adler J, Rydén A. Response burden and questionnaire length: is shorter better? A review and meta-analysis. Value in health : the journal of the International Society for Pharmacoeconomics and Outcomes Research. 2011;14(8):1101-8. doi: https://doi.org/10.1016/j.jval.2011.06.003.

15.   Grabeel KL, Russomanno J, Oelschlegel S, Tester E, Heidel RE. Computerized versus hand-scored health literacy tools: a comparison of Simple Measure of Gobbledygook (SMOG) and Flesch-Kincaid in printed patient education materials. J Med Libr Assoc. 2018;106(1):38-45. Epub 2018/01/18. doi: 10.5195/jmla.2018.262. PubMed PMID: 29339932; PubMed Central PMCID: PMCPMC5764592.

16.   Altschuler A, Picchi T, Nelson M, Rogers JD, Hart J, Sternfeld B. Physical

activity questionnaire comprehension: lessons from cognitive interviews. Med Sci

Sports Exerc. 2009;41(2):336-43. doi: 10.1249/MSS.0b013e318186b1b1. PubMed

PMID: 19127192; PubMed Central PMCID: PMCPMC2694667.

17.   Wallace LS, Lennon ES. American Academy of Family Physicians patient

education materials: Can patients read them? Family Medicine. 2004;36(8):571-4.

Epub 2004/09/03. PubMed PMID: 15343418.

18.   van Poppel MN, Chinapaw MJ, Mokkink LB, van Mechelen W, Terwee CB.

Physical activity questionnaires for adults: a systematic review of measurement

properties. Sports Medicine (Auckland, N Z). 2010;40(7):565-600. doi:

https://doi.org/10.2165/11531930-000000000-00000.

19.   Helmerhorst HJH, S. B, Warren J, Besson H, Ekelund U. A systematic review

of reliability and objective criterion-related validity of physical activity questionnaires.

The international journal of behavioral nutrition and physical activity. 2012;9:103. doi:

10.1186/1479-5868-9-103.

20.   Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et

al. The PRISMA 2020 statement: an updated guideline for reporting systematic

reviews. BMJ. 2021;372:n71. doi: 10.1136/bmj.n71.

21.   Prinsen CAC, Mokkink LB, Bouter LM, Alonso J, Patrick DL, de Vet HCW, et

al. COSMIN guideline for systematic reviews of patient-reported outcome measures.

Quality of Life Research : An International Journal of Quality of Life Aspects of

Treatment, Care, and Rehabilitation. 2018;27(5):1147-57. Epub 2018/02/13. doi:

https://doi.org/10.1007/s11136-018-1798-3. PubMed PMID: 29435801; PubMed

Central PMCID: PMCPMC5891568.

22.   Downes MJ, Brennan ML, Williams HC, Dean RS. Development of a critical

appraisal tool to assess the quality of cross-sectional studies (AXIS). BMJ Open.

2016;6(12):e011458. doi: 10.1136/bmjopen-2016-011458.

23.   Williams MA, McCarthy CJ, Chorti A, Cooke MW, Gates S. A systematic

review of reliability and validity studies of methods for measuring active and passive

cervical range of motion. Journal of Manipulative and Physiological Therapeutics.

2010;33(2):138-55. Epub 2010/02/23. doi: 10.1016/j.jmpt.2009.12.009. PubMed

PMID: 20170780.

24.   Brysbaert M. How many words do we read per minute? A review and meta-

analysis of reading rate. Journal of Memory and Language. 2019;109. doi:

https://doi.org/10.1016/j.jml.2019.104047.

25.   Wang LW, Miller MJ, Schmitt MR, Wen FK. Assessing readability formula

differences with written health information materials: application, results, and

recommendations. Res Social Adm Pharm. 2013;9(5):503-16. Epub 2012/07/28. doi:

10.1016/j.sapharm.2012.05.009. PubMed PMID: 22835706.

26.   charactercalculator.com. Flesch Reading Ease 2023. Available from:

https://charactercalculator.com/flesch-reading-ease/.

27.   Bauman AE, Richards JA. Understanding of the Single-Item Physical Activity

Question for Population Surveillance. Journal of Physical Activity and Health.

2022;19(10):681-6. doi: 10.1123/jpah.2022-0369.

28.   Schechtman KB, Barzilai B, Rost K, B. FJE. Measuring physical activity with a

single question. American journal of public health. 1991;81(6):771-3.

29.   Gill DP, Jones GR, Zou G, Speechley M. Using a single question to assess

physical activity in older adults: A reliability and validity study. BMC Medical

Research Methodology. 2012;12(20). doi: 10.1186/1471-2288-12-2022373159.

30.   Macdonald HM, Nettlefold L, Bauman A, Sims-Gould J, McKay HA. Pragmatic

Evaluation of Older Adults' Physical Activity in Scale-Up Studies: Is the Single-Item

Measure a Reasonable Option? Journal of Aging and Physical Activity.

2022;30(1):25-32. doi: 10.1123/japa.2020-0412.

31.   Portegijs E, Sipilä S, Viljanen A, Rantakokko M, Rantanen T. Validity of a

single question to assess habitual physical activity of community-dwelling older

people. Scandinavian Journal of Medicine & Science in Sports. 2017;27(11):1423-30.

doi: 10.1111/sms.12782.

32.   Kiernan M, Schoffman DE, Lee K, Brown SD, Fair JM, Perri MG, et al. The

stanford leisure-time activity categorical item (L-Cat): A single categorical item

sensitive to physical activity changes in overweight/obese women. International

Journal of Obesity. 2013;37(12):1597-602. doi: 10.1038/ijo.2013.36.

33.   Ross KM, Leahey TM, Kiernan M. Validation of the Stanford Leisure-Time

Activity Categorical Item (L-Cat) using armband activity monitor data. *Obesity*

science & practice. 2018;4(3):276-82. doi: 10.1002/osp4.155.

34.   Johansson G, Westerterp KR. Assessment of the physical activity level with

two questions: Validation with doubly labeled water. International Journal of Obesity.

2008;32(6):1031-3. doi: 10.1038/ijo.2008.42.

35.   Blomqvist A, Bäck M, Klompstra L, Strömberg A, Jaarsma T. Utility of single-

item questions to assess physical inactivity in patients with chronic heart failure. ESC

Heart Fail. 2020;7(4):1467-76. doi: 10.1002/ehf2.12709.

36.   Orrell A, Doherty P, Miles J, Lewin R. Development and validation of a very

brief questionnaire measure of physical activity in adults with coronary heart disease.

European Journal of Cardiovascular Prevention and Rehabilitation. 2007;14(5):615-

23. doi: 10.1097/HJR.0b013e3280ecfd56.

37.   Cruz J, Jácome C, Oliveira A, Paixão C, Rebelo P, Flora S, et al. Construct

validity of the brief physical activity assessment tool for clinical use in COPD. The

Clinical Respiratory Journal. 2021;15(5):530-9. doi: 10.1111/crj.13333.

38.   Hamilton K, White KM, Cuddihy T. Using a single-item physical activity

measure to describe and validate parents' physical activity patterns. Research

Quarterly for Exercise and Sport. 2012;83(2):340-5. doi:

10.1080/02701367.2012.10599865.

39.   Milton K, Bull FC, Bauman A. Reliability and validity testing of a single-item

physical activity measure. British Journal of Sports Medicine. 2011;45(3):203-8. doi:

10.1136/bjsm.2009.068395.

40.   Milton K, Clemes S, Bull F. Can a single question provide an accurate

measure of physical activity? British Journal of Sports Medicine. 2013;47(1):44-8.

doi: 10.1136/bjsports-2011-090899.

41.   O'Halloran P, Kingsley M, Nicholson M, Staley K, Randle E, Wright A, et al.

Responsiveness of the single item measure to detect change in physical activity.

PLoS ONE. 2020;15(6). doi: 10.1371/journal.pone.0234420.

42.   Zwolinsky S, McKenna J, Pringle A, Widdop P, Griffiths C. Physical activity

assessment for public health: Efficacious use of the single-item measure. Public

Health. 2015;129(12):1630-6. doi: 10.1016/j.puhe.2015.07.015.

43.   Marshall AL, Smith BJ, Bauman AE, Kaur S. Reliability and validity of a brief

physical activity assessment for use by family doctors. British Journal of Sports

Medicine. 2005;39(5):294-7. doi: 10.1136/bjsm.2004.013771.

44.   Puig Ribera A, Peña Chimenis Ò, Romaguera Bosch M, Duran Bellido E,

Heras Tebar A, Solà Gonfaus M, et al. How to identify physical inactivity in Primary

Care: Validation of the Catalan and Spanish versions of 2 short questionnaires.

Atencion Primaria. 2012;44(8):485-93. doi: 10.1016/j.aprim.2012.01.005.

45.   Puig-Ribera A, Martín-Cantera C, Puigdomenech E, Real J, Romaguera M,

Magdalena-Belio JF, et al. Screening physical activity in family practice: Validity of

the Spanish version of a brief physical activity questionnaire. PLoS ONE. 2015;10(9).

doi: 10.1371/journal.pone.0136870.

46.   Smith BJ, Marshall AL, Huang N. Screening for physical activity in family

practice: Evaluation of two brief assessment tools. American journal of preventive

medicine. 2005;29(4):256-64. doi: 10.1016/j.amepre.2005.07.005.

47.   Gionet NJ, Godin G. Self-Reported Exercise Behavior of Employees - A

Validity Study. Journal of Occupational and Environmental Medicine.

1989;31(12):969-73. doi: 10.1097/00043764-198912000-00007.

48.   Li S, Carlson E, Holm K. Validation of a single-item measure of usual physical

activity. Perceptual and Motor Skills. 2000;91(2):593-602. doi:

10.2466/pms.2000.91.2.593.

49.   Graff-Iversen S, Anderssen SA, Holme IM, Jenum AK, Raastad T. Two short

questionnaires on leisure-time physical activity compared with serum lipids,

anthropometric measurements and aerobic power in a suburban population from

Oslo, Norway. European journal of epidemiology. 2008;23(3):167-74. doi:

10.1007/s10654-007-9214-2.

50.   Forsén L, Loland NW, Vuillemin A, Chinapaw MJ, van Poppel MN, Mokkink

LB, et al. Self-administered physical activity questionnaires for the elderly: a

systematic review of measurement properties. Sports Medicine. 2010;40(7):601-23.

doi: 10.2165/11531350-000000000-00000. PubMed PMID: 20545382.

51.   Smith TO, McKenna MC, Salter C, Hardeman W, Richardson K, Hillsdon M, et

al. A systematic review of the physical activity assessment tools used in primary

care. Family Practice. 2017;34(4):384-91. doi:

https://doi.org/10.1093/fampra/cmx011.

52.   Organization WH. Guidelines on physical activity and sedentary behaviour.

Geneva: World Health Organization; 2020.

53.   Levin-Zamir D, Leung AYM, Dodson S, Rowlands G. Health Literacy in

Selected Populations: Individuals, Families, and Communities from the International

and Cultural Perspective. Studies in Health Technology and Informatics.

2017;240:392-414. PubMed PMID: 28972530.

54.   Marsilio M, Fusco F, Gheduzzi E, Guglielmetti C. Co-Production Performance

Evaluation in Healthcare. A Systematic Review of Methods, Tools and Metrics.

International Journal of Environmental Research and Public Health. 2021;18(7).

Epub 20210324. doi: 10.3390/ijerph18073336. PubMed PMID: 33804862; PubMed

Central PMCID: PMCPMC8037812.

55.   Hankinson A. Association of activity and chronic disease risk factors: utility

and limitations of objectively measured physical activity data. Journal of the American

Dietetic Association. 2008;108(6):945-7. doi: 10.1016/j.jada.2008.03.018. PubMed

PMID: 18502224.

56.   Ball TJ, Joy EA, Goh TL, Hannon JC, Gren LH, Shaw JM. Validity of two brief

primary care physical activity questionnaires with accelerometry in clinic staff.

Primary Health Care Research & Development. 2015;16(1):100-8. doi: DOI:

10.1017/S1463423613000479.

57.   Cruz J, Jácome C, Oliveira A, Paixão C, Rebelo P, Flora S, et al. Construct

validity of the brief physical activity assessment tool for clinical use in COPD. Clin

Respir J. 2021;15(5):530-9. doi: 10.1111/crj.13333.

58.   Danquah IH, Petersen CB, Skov SS, Tolstrup JS. Validation of the NPAQ-

short - A brief questionnaire to monitor physical activity and compliance with the

WHO recommendations. BMC Public Health. 2018;18(1). doi: 10.1186/s12889-018-

5538-y.

59. Hyvärinen M, Sipilä S, Kulmala J, Hakonen H, Tammelin TH, Kujala UM, et al.

Validity and reliability of a single question for leisure-time physical activity

assessment in middle-aged women. Journal of Aging and Physical Activity.

2020;28(2):231-41. doi: 10.1123/JAPA.2019-0093.

60. Iwai N, Hisamichi S, Hayakawa N, Inaba Y, Nagaoka T, Sugimori H, et al.

Validity and reliability of single-item questions about physical activity. Journal of

Epidemiology. 2001;11(5):211-8. doi: 10.2188/jea.11.211.

61. Jackson AW, Morrow JR, Jr., Bowles HR, Fitzgerald SJ, Blair SN. Construct

validity evidence for single-response items to estimate physical activity levels in large

sample studies. Research Quarterly for Exercise and Sport. 2007;78(2):24-31. doi:

10.1080/02701367.2007.10599400.

62. Moreno-Llamas A, García-Mayor J, De la Cruz-Sánchez E. Concurrent and

convergent validity of a single, brief question for physical activity assessment.

International Journal of Environmental Research and Public Health. 2020;17(6):1989.

doi: 10.3390/ijerph17061989.

63. Rose SB, Elley CR, Lawton BA, Dowell AC. A single question reliably

identifies physically inactive women in primary care. The New Zealand Medical

Journal. 2008;121(1268):U2897.

64. Taylor-Piliae RE, Norton LC, Haskell WL, Mahbouda MH, Fair JM, Iribarren C,

et al. Validation of a new brief physical activity survey among men and women aged

60-69 years. American Journal of Epidemiology. 2006;164(6):598-606. doi:

10.1093/aje/kwj248.

65. Webster S, Khan A, Nitz JC. A brief questionnaire is able to measure

population physical activity levels accurately: A comparative validation study. Journal

of Clinical Gerontology and Geriatrics. 2011;2(3):83-7. doi:

10.1016/j.jcgg.2011.06.003.

66.   Weiss TW, Slater CH, Green LW, Kennedy VC, Albright DL, Wun CC. The

validity of single-item, self-assessment questions as measures of adult physical

activity. Journal of Clinical Epidemiology. 1990;43(11):1123-9. doi: 10.1016/0895-

4356(90)90013-F.

67.   Yore MM, Bowles HR, Ainsworth BE, Macera CA, Kohl HW, III. Single versus

multiple item questions on occupational physical activity. Journal of Physical Activity

and Health. 2006;3(1):102-11. doi: 10.1123/jpah.3.1.102.

68.   Bauman AE, Richards JA. Understanding of the Single-Item Physical Activity

Question for Population Surveillance. Journal of Physical Activity and Health. 2022.

doi: 10.1123/jpah.2022-0369.

Supporting information

S1 Table. Studies overview.

S2 Table. Summary of measurement properties.

S3 Table. Length and readability levels of short physical activity questionnaires.

S1 File. Blank for studies quality assessment.

Figure 1

Figure 2