Association between PM$_{10}$ exposure and risk of myocardial infarction in adults: a

systematic review and meta-analysis

Kleiton Strobl[1*], Syed Asad Irfan[1], Hassan Masood[1], Noor Latif[2], Om Kurmi[1,3]

Affiliation

[1]Faculty of Health Sciences, McMaster University, Hamilton, Canada

[2]Faculty of Science, McMaster University, Hamilton, Canada

[3]Faculty Centre for Intelligent Healthcare, Coventry University, Coventry, UK

9
10
11

12   **Correspondence***
13
14   Kleiton Strobl M.Sc.

15   McMaster University

16   1280 Main Street West, Hamilton, ON L8S4L8, Canada

17   (289) 684-3075

18   kstrobl1@gmail.com

Abstract

Background: Air pollution has several negative health effects. Particulate matter (PM) is a

pollutant that is often linked to health adversities. $PM_{2.5}$ (PM with an aerodynamic diameter of

≤2.5μm) exposure has been associated with negative cardiovascular outcomes. However, the

impact of PM_{10} (PM with an aerodynamic diameter of ≤10μm) exposure is often overlooked due

to its limited ability to pass the alveolar barrier. This study aims to assess the association

between PM_{10} exposure and risk of myocardial infarction (MI) amongst adults (≥18 years of age)

as this has been poorly studied.

Methods: The study protocol was published on PROSPERO (CRD42023409796) on March 31,

2023. Literature searches were conducted on 4 databases (OVID Medline, Embase, CINAHL,

and Web of Science) for studies looking at associations between PM and MI. English studies

from all time periods were assessed. Studies selected for review were time-series, case-

crossover, and cohort studies which investigated the risk of MI as an outcome upon PM_{10}

exposure. The quality of evidence was assessed using Cochrane's GRADE approach. Data for

different risk outcomes (risk ratio (RR), odds ratio (OR), hazard ratio (HR)) and 3 lags was meta-

analyzed using an inverse variance statistical analysis using a random effects model. The pooled

effect sizes and the 95% confidence intervals (CIs) were reported in forest plots.

Results: Among the 1,099 studies identified, 41 were included for review and 23 were deemed

eligible for meta-analysis. Our analysis revealed that there is an increased risk (OR=1.01; 95%

CI:1.00 - 1.02) of MI with a 10 $\mu g/m^3$ increase in PM_{10} after a lag 0 and lag 1 delay.

Conclusions: Our findings indicate that PM_{10} exposure is associated with an increased risk of

MI. This can aid in informing environmental policy-making, personal-level preventative

measures, and global public health action.

1. Introduction

Air pollution is a complex combination of gaseous and particle constituents, which are hazardous to human health [1]. In air pollution, particulate matter (PM) comprises carbonaceous particles containing adsorbed organic compounds and reactive metals [1]. Nitrates, sulfates, polycyclic aromatic hydrocarbons, endotoxin, and metals such as iron, copper, nickel, zinc, and vanadium are all common constituents of PM [1]. PM can be further classified relative to the particle size into coarse (PM_{10}: diameter <10μm), fine ($PM_{2.5}$: diameter <2.5μm), and ultrafine ($PM_{0.1}$: diameter <0.1μm) [1]. Earlier scientific research has reported that $PM_{2.5}$ is associated with adverse health outcomes, including poor cardiovascular health outcomes. PM_{10} is present in dust from roads, farms, construction sites, and mines [2]. However, the chemical composition and size distribution of PM_{10} varies widely depending on where it originates and how it forms in the environment [3]. Although PM_{10} is also associated with adverse impact health as it is an irritant for the nose, throat, and eyes; however, in general, this is not a major focus airborne pollutant of study [2].

The short-term impacts of PM_{10} exposure on respiratory pathologies, such as chronic obstructive pulmonary disease and asthma, are well known and studied [3]. However, in cardiovascular health research, the effects of $PM_{2.5}$ are more often researched as $PM_{2.5}$ has been shown to pass the alveolar barrier and cause an inflammatory response in blood vessels [4]. This exacerbates atherosclerosis and increases the risk of myocardial infarction (MI), ischemic heart disease, and thrombotic stroke [4]. The risk of MI was chosen as it is one of the largest causes of death worldwide, with approximately three million deaths annually [4]. Different populations are also exposed to varying levels of PM, and understanding its health adversities is important for understanding subsequent health disparities. Interestingly, the impact of PM_{10} on the risk of MI

is unknown and poorly studied, despite instances being reported where exposure to PM_{10} has led

to a surge in cardiovascular-related hospital admissions [3]. Although there is a higher

probability of $PM_{2.5}$ passing the alveolar barrier, it is still possible for PM_{10} to enter the

bloodstream in smaller proportions [5]. When searching this topic on the PROSPERO database,

only one matching systematic review and meta-analysis was found, and it solely assessed the

impact of $PM_{2.5}$ and not PM_{10} [6]. The primary aim of this system review and meta-analysis is to

assess the effect of PM_{10} on the risk of MI to better understand its burden of disease.

2. Methods

2.1. Search strategy

Studies were obtained from a systematic search of the following databases: OVID

Medline, Embase, CINAHL, and Web of Science. The literature searches were conducted on Jan

17, 2023, and included studies from all time periods. See the specific search terms utilized (S1

Table). The study protocol was published on PROSPERO (CRD42023409796) on March 31,

2023, after completing the literature search.

2.2. Inclusion/exclusion criteria

The inclusion criteria for this study were: (1) study design had to be time-series, case-

crossover, or cohort; (2) PM_{10} had to be an exposure; (3) risk of myocardial infarction had to be

an outcome (risk ratio/relative risk (RR), odds ratio (OR), hazard ratio (HR)); (4) population ≥18

years of age.

The exclusion criteria for this study were: (1) Other primary study designs (RCTs,

clinical trials, cross-sectional designs, protocols, pilot studies, etc.); (2) secondary studies

(narrative reviews, systematic reviews, meta-analyses, scoping reviews); (3) Risk outcome(s)

only reported in graph(s) (4) studies that only look at $PM_{2.5}$ or coarse particulate matter ($PM_{2.5-}$

$_{10}$); (5) population <18 years of age.

2.3. Lag periods

PM_{10} exposure lags were based on the following definitions: lag 0 = same day (0-24

hours); lag 1 = 1-3 day delay (24-96 hours); lag 2 = 3 day delay or more (>96 hours). If there

were multiple data points, the value closest to the midpoint of the range was selected for meta-

analysis for lag 0 and 1 (lag 0 = 12 hours; lag 1 = 60 hours). For lag 2, the value closest to >96

hours was selected for meta-analysis.

2.4. Study selection

Covidence was used to manage the screening phase of this study [7]. For the abstract

screening, two authors (HM & NL) independently screened abstracts based on the

inclusion/exclusion criteria, and another two (KS & SI) resolved conflicts. For full-text

screening, two authors independently (KS & SI) screened the full manuscripts, and the same

authors (KS & SI) discussed any conflicts and reached a consensus. The study drafting process

was recorded using "Preferred Reporting Items for Systematic Reviews and Meta-Analyses"

(PRISMA) guidelines [8].

2.5. Data extraction

Two authors independently extracted data from one half of the included studies (KS &

SI), while another two (HM & NL) independently extracted data from the other half of the

included studies. Any discrepancies in data values were corrected cohesively amongst the two

authors for their extracted data half, respectively.

Data extracted from studies included: study name, design, country, sample size,

male/female ratio, participant characteristics, exposure increments for risk measure, risk scale,

lag intervals, and effect size (risk outcomes).

For studies with multiple models, outcome data values adjusting for the greatest number

of confounding variables were meta-analyzed to maintain data validity. Industrial values were

used for studies reporting industrial and non-industrial location outcome values because they

were more likely to report reliable PM_{10} exposure MI risk outcomes. For studies that reported

data on multiple regions of a country, data was selected from the most central geographical

region.

2.6. Methodological risk of bias assessment

Risk of bias was assessed by two authors (KS & SI) using the NHLBI Quality

Assessment Tool for Observational Cohort and Cross-Sectional Studies [9] which assesses

methodological risk of bias through 14 criteria: (1) clear research question/objective; (2) clearly

defined study population; (3) 50% participation rate of eligible persons; (4) all participants

selected from identical/similar populations in the same time period using pre-specified inclusion

and exclusion criteria; (5) provision of sample size justification, power description, or variance

and effect estimates; (6) exposure measured before outcome in analysis; (7) sufficient time frame

to expect association; (8) examination of differing levels of exposure as related to the outcome;

(9) clearly defined, valid, reliable, widely-applied exposure measures; (10) >1 exposure

assessments; (11) clearly defined, valid, reliable, widely-applied outcome measures; (12)

blinding of outcome assessors to participant exposure status; (13) <20% loss to follow-up after

baseline; (14) measurement and adjustment of potential confounders affecting impact between

exposure and outcome.

Each criterion was deemed to be satisfied or unsatisfied based on the author's rating.

Overall quality ratings included: 'good', 'fair', and 'poor'. The overall quality of each study

began at 'good', and got demoted by one level per unsatisfied criteria. Conflicts in the

methodological risk of bias assessment were resolved by the same two authors (KS & SI)

through consensus.

2.7. Meta-analysis

Meta-analysis was conducted using Review Manager 5.4.1.

2.7.1. *MI risk outcome measures*

The primary outcome was MI risk. For the meta-analysis, data was included for studies

reporting risk based on PM_{10} concentration increments of $10\mu g/m^3$. MI risk outcome measures

included RR, OR, and HR with 95% confidence intervals (CIs). A RR, OR, or HR >1 indicates a

higher PM_{10}-associated MI risk.

MI risk was stratified by MI risk outcome measure (RR, OR, HR) and lag (0, 1, 2) to

preserve data validity.

2.7.2. *Data synthesis*

The pooled risk outcome was considered if the following criteria were met: two or more

studies reported the same MI risk outcome measure and lag, the I^2 statistic for measuring

statistical heterogeneity ≤60%.

An I^2 >60% was considered heterogeneous on a statistically significant level. An inverse

variance statistical analysis was conducted with a random-effects model when creating the forest

plots, as there was heterogeneity, differences in study design, setting, and adjustment models. A

P-value <0.05 was considered statistically significant.

Publication bias was assessed through analysis of funnel plot symmetricity. Publication

bias was not assessed for quantitative analyses with <5 studies due to a lack of statistical power.

2.7.3. *Quality of evidence evaluation*

Quality of evidence was evaluated by two authors (KS & SI) independently using

Cochrane's GRADE approach (Schünemann et al., 2013), which assesses: (1) risk of bias; (2)

inconsistency; (3) indirectness; (4) imprecision; (5) publication bias; (6) large magnitude of

effect; (7) dose-response gradient; (8) residual confounding. Overall quality ratings include:

'high', 'moderate', 'low' and 'very low'. All resulting outcomes began with a rating of 'high'

and were demoted one level for each unsatisfied criteria 1-5; upgrading could occur for

satisfying criteria 6-8. Any evaluation conflicts were resolved by the same two authors (KS &

SI) through consensus.

3. Results

3.1. Study selection

A breakdown of study identification and screening can be found in the PRISMA

flowchart (Figure 1). A systematic literature search from databases/registers identified 1,099

potential studies viable for inclusion. After 396 duplicates were removed, 703 total studies were

available for abstract screening. Five hundred sixty-nine studies were excluded following

abstract screening, and 12 studies were not retrieved due to a lack of full-text availability. One

hundred twenty-two studies remained for full-text screening, and 81 studies were excluded after

the full-text screening: 64 for having the wrong outcome measure, 11 for having different study

design (not meeting the inclusion criteria), 3 for analyzing $PM_{2.5}$ exclusively and not PM_{10}, 2 for

having different patient population, and 1 study for not being in English. Forty-one studies were

included for review, with 23 of 41 deemed eligible for meta-analysis.

196
197 **Fig 1. PRISMA flow diagram**.

198 **3.2. Study Characteristics**

199   Table 1 provides the characteristics of studies included in the meta-analysis [10-31], adjusted confounding variables can be

200 found for these respective studies (S2 Table). See study characteristics for other studies included in the review (S3 Table) (reference).

202 **Table 1. Study characteristics for studies included in the meta-analysis.**

#	Author/ publication year	Study Design	Country	Sample size (n)	# Male (%)	Risk Measure	Lag Intervals
1	Argacha 2016 [10]	Case-crossover	Belgium	11,428	8,607 (75.3)	OR	Lag 0: 24hrs post-exposure
2	Bard 2014 [11]	Case-crossover	France	2,134	1,642 (76.9)	OR	Lag 0: Same day Lag 1: 1-day delay Lag 0-1: Average of same day and 1 day previous
3	Bhaskaran 2011 [12]	Case-crossover	United Kingdom	79,288	50,988 (64.8)	RR	Lag 0: 1-6hrs Lag 1: 7-12hrs Lag 2: 13-18hrs Lag 3: 19-24hrs Lag 4: 25-72hrs
4	Buszman 2020 [13]	Case-crossover	Poland	1,957	NR	OR	Lag 0: Same day Lag 1: 1-day delay
5	Cheng 2021 [14]	Case-crossover	Australia	3,307	2,162 (65.4)	RR	Lag 0: 1hr Lag 1: 2-6hrs Lag 2: 7-12hrs Lag 3: 13-24hrs
6	Claeys 2015 [15]	Time-series	Belgium	15,963	11,995 (75.1)	RR	5-day delay
7	Collart 2017 [16]	Time-series	Belgium	21,491	14,377 (66.9)	RR	Single-day lags: Lag 0, 1, 2, 3, 4, 5, 6

8	Davoodabadi 2019 [17]	Case-crossover	Iran	319	238 (74.6)	OR	Lag 0: 24h post-exposure Lag 1: 48h post-exposure
9	Downward 2018 [18]	Cohort	Netherlands	33,381	7,846 (23.5)	HR	1-year delay
10	Huss 2010 [19]	Cohort	Switzerland	4,580,311	NR	HR	Average exposure in the year 2000
11	Kim 2020 [20]	Time-series	South Korea	196,167	104,949 (53.5)	HR	5-year delay
12	Konduracka 2019 [21]	Time-series	Poland	3,545	1,602 (45.2)	OR	Single-day lags: Lag 0, 1, 2, 3, 4, 5, 6
13	Kuzma 2021 [22]	Time-series	Poland	9,046	5,692 (62.9)	OR	Single-day lags: Lag 0, 1, 2, 3, 4, 5, 6
14	Lipsett 2011 [23]	Cohort	USA	124,614	0 (0)	HR	Median of 8.3-year delay
15	Nuvolone 2011 [24]	Case-crossover	Italy	11,450	6,985 (61.0)	OR	Single-day lags: Lag 0, 1, 2, 3, 4, 5 Multiple-day lags: Lag 0–2, 0–5, 3–5
16	Puett 2008 [25]	Cohort	USA	66,250	0 (0)	HR	Lag 1: 1-month post-exposure average Lag 2: 3-month post-exposure average Lag 3: 12-month post-exposure average Lag 4: 48-month post-exposure average
17	Roye 2019 [26]	Time-series	Spain	9,871	7,008 (71.0)	RR	14-day delay
18	Soleimani 2019 [27]	Cohort	Iran	6,425	3,652 (56.8)	RR	Single-day lags: Lag 0, 1, 2, 3, 4, 5, 6, 7, 8, 9
19	Vidale 2017 [28]	Cohort	Italy	4,110	2,672 (65.0)	HR	Single-day lags: Lag 0, 1, 2, 3, 4
20	Yang 2022 [29]	Case-crossover	China	25,299	17,100 (67.6)	RR	Lag 1: 1-day delay
21	Yen 2022 [30]	Case-crossover	China	979,979	458,195 (46.8)	OR	Lag 0: Same day Lag 1: 1-day delay Lag 0-1: Average of same day and 1 day previous

| 22 | Zhang 2016 [31] | Case-crossover | China | 2,749 | 1,612 (58.6) | OR | Single-day lags: Lag 0, 1, 2, 3, 4, 5 |
| 23 | Zhu 2019 [32] | Cohort | China | 147,422 | NR | RR | Single-day lags: Lag 0, 1, 2, 3, 4, 5, 6 |

RR = Risk ratio; OR = Odds ratio; HR = Hazard ratio; NR = Not reported

3.2.1. *Study design characteristics*

Amongst the 41 studies included in the systematic review, 22 studies (53.7%) had a case-

crossover design, 12 studies (29.3%) had an observational cohort design, and 7 studies (17.1%)

had a time-series design.

For the meta-analysis, ten studies (43.5%) had a case-crossover design, seven studies

(30.4%) had an observational cohort design, and six studies (26.1%) had a time-series design.

3.2.2. *Participant characteristics*

For qualitative synthesis, data was calculated from a total of 6,663,870 participants.

For the meta-analysis, data was calculated from a total of 6,336,506 participants. All but three

studies reported a male/female participant ratio [13, 19, 32]. Of 1,606,816 participants from the

remaining studies with a reported male/female participant ratio, there were 707,322 male

participants (44.0%) and 899,494 (56.0%) female participants.

3.2.3. *MI risk outcome measures*

Amongst the 41 studies included in the review, 18 studies (43.9%) used OR as their

effect size for MI risk outcome measure, 12 studies (29.3%) used RR, and 11 studies (26.8%)

used HR.

For the meta-analysis, studies most often utilized OR (n=9; 39.1%), followed by RR

(n=8; 34.8%) and HR (n=6; 26.1%) for MI risk outcome measures.

3.3. Methodological risk of bias analysis for included studies

After the quality assessment was conducted (S4 Table), a large majority of studies were

rated at a quality score of good (38/41), with the remainder of the studies rated at a quality of fair

(3/41). No studies were rated at a poor quality score. All three studies demoted to a fair quality

score were done so for the same reason: failure to adjust for confounders [13, 38, 27].

3.4. Primary MI risk outcomes – meta-analysis

See the meta-analysis summary of primary MI risk outcomes (Table 2). For lag 0 and lag

1 HR outcomes, meta-analysis could not be conducted as there was only one associated study

for each respective outcome [28] (S1, S2 Fig). Evidence assessment could also not be conducted

due to one study (S11, S12 Table).

Table 2. Summary of primary MI risk outcomes

	Lag 0 Effect Size (95% CI)	Lag 1 Effect Size (95% CI)	Lag 2 Effect Size (95% CI)
Risk Ratio	1.02 (0.97,1.07)	1.01 (0.99,1.03)	1.01 (1.00,1.03)
Odds Ratio	1.01 (1.00,1.02)	1.01 (1.00,1.02)	1.01 (0.99,1.02)
Hazard Ratio	1.00 (1.00,1.01)	1.00 (1.00,1.01)	1.00 (0.99,1.01)

3.4.1. *Risk Ratio – Lag 0 (Same day [0-24hrs])*

Three studies with three outcomes (n = 89,020) were reported for RR. Soleimani et al.

[27] and Cheng et al. [14] reported an increased risk of MI with PM_{10} exposure, while Bhaskaran

et al. [12] reported a decreased risk. Statistical pooling was appropriate due to statistical

homogeneity (I^2 = 60%) (Figure 2). Although there was an overall increased risk of MI (RR =

1.02, 95% CI: 0.97-1.07) it was not statistically significant (P-value = 0.43). Evidence quality

was rated high on the GRADE scale (S5 Table).

Fig 2. Lag 0 RR for MI after 10 μg/m^3 increase in PM$_{10}$ exposure.

3.4.2. *Risk Ratio – Lag 1 (1-3 days [24-96hrs])*

Four studies with four outcomes (n = 132,503) were reported for RR. Statistical pooling

was inappropriate for the lag 1 RR values because of statistical heterogeneity (I^2 = 72%) (Figure

3). Evidence quality was rated moderate on the GRADE scale due to inconsistency (S6 Table).

Fig 3. Lag 1 RR for MI after 10 μg/m^3 increase in PM$_{10}$ exposure.

3.4.3. *Risk Ratio – Lag 2 (3+ days [96hrs+])*

Four studies with four outcomes (n = 179,681) were reported for RR. Roye et al. [26] and

Zhu et al. [32] reported an increased risk of MI with PM$_{10}$ exposure, while Soleimani et al. [27]

reported a decreased risk. Claeys et al. [15] reported no change in risk. Statistical pooling was

appropriate due to statistical homogeneity (I^2 = 60%) (Figure 3). Although there was an overall

increased risk of MI (RR = 1.01, 95% CI: 1.00-1.03), it was not statistically significant (P-value

= 0.10). Evidence quality was rated high on the GRADE scale (S7 Table).

Fig 4. Lag 2 RR for MI after 10 μg/m^3 increase in PM$_{10}$ exposure.

3.4.4. *Odds Ratio – Lag 0 (Same day [0-24hrs])*

Eight studies with eleven outcomes (n = 1,019,062) were reported for OR. Kuzma et al.
[22] reported a reduced risk of STEMI with PM$_{10}$ exposure, while all other studies reported an
increased risk within their respective groups/subgroups (MI, STEMI, NSTEMI). Statistical
pooling was appropriate due to statistical homogeneity (I^2 = 60%) (Figure 4). There was an
overall increased risk of MI (RR = 1.01, 95% CI: 1.00-1.02), and it was statistically significant
(P-value = 0.01). Evidence quality was rated high on the GRADE scale (S8 Table).

Fig 5. Lag 0 OR for MI after 10 μg/m³ increase in PM₁₀ exposure.

3.4.5. *Odds Ratio – Lag 1 (1-3 days [24-96hrs])*

Seven studies with ten outcomes (n = 1,007,634) were reported for OR. Kuzma et al. [22]

reported a reduced risk of STEMI with PM₁₀ exposure, while all other studies reported an

increased risk within their respective groups/subgroups (MI, STEMI, NSTEMI). Statistical

pooling was appropriate due to statistical homogeneity (I^2 = 50%) (Figure 5). There was an

overall increased risk of MI (RR = 1.01, 95% CI: 1.00-1.02), and it was statistically significant

(P-value = 0.02). Evidence quality was rated high on the GRADE scale (S9 Table).

Fig 6. Lag 1 OR for MI after 10 μg/m³ increase in PM₁₀ exposure.

3.4.6. *Odds Ratio – Lag 2 (3+ days [96hrs+])*

Five studies with seven outcomes (n = 27,109) were reported for OR. Zhang et al. [31]

reported a reduced risk of MI with PM₁₀ exposure, while all other studies reported an increased

risk within their respective group/subgroup (MI, STEMI, NSTEMI). Statistical pooling was

appropriate due to statistical homogeneity ($I^2 = 41\%$) (Figure 6). There was an overall increased

risk of MI (RR = 1.01, 95% CI: 0.99-1.02), but it was not statistically significant (P-value =

0.28). Evidence quality was rated high on the GRADE scale (S10 Table).

Fig 7. Lag 2 OR for MI after 10 μg/m³ increase in PM₁₀ exposure.

3.4.7. *Hazard Ratio – Lag 2 (3+ days [96hrs+])*

Six studies with six outcomes (n = 4,913,615) were reported for HR. Nuvolone et al. [24] and Konduracka et al. [21] reported an increased risk of MI with PM₁₀ exposure, while Zhang et al. [31] reported a reduced risk. Statistical pooling was appropriate due to statistical homogeneity ($I^2 = 6\%$) (Figure 8). There was no impact on risk of MI (RR = 1.00, 95% CI: 0.99-1.01), and the result was not statistically significant (P-value = 0.69). Evidence quality was rated high on the GRADE scale (S13 Table).

Fig 8. Lag 2 HR for MI after 10 μg/m^3 increase in PM$_{10}$ exposure.

3.4.8. *Sensitivity Analyses*

  Two sensitivity analyses were performed. The first sensitivity analysis was the exclusion

of risk outcome values which had more than a 7-day post-exposure delay. This was done in order

to eliminate values which had relatively long post-exposure delays lasting multiple weeks,

months, or years. This sensitivity was conducted to all lag 2 forest plots and consisted of six

excluded values: 1 for RR [26] and 5 for HR [18, 19, 20, 23, 25]. However, no significant

differences were found in this sensitivity analysis.

  The second sensitivity analysis was the exclusion of Buszman et al. [13] from the OR

forest plots for lag 0 and lag 1. This was done in order to evaluate whether this study was a

detrimental outlier and played a significant role in the overall risk outcome. However, no

significant differences were found in this sensitivity analysis.

4. Discussion

  This study aimed to evaluate the risk of MI after exposure to PM$_{10}$. This was important

considering that cardiovascular disease is the leading cause of death in the world [51]. Assessing

potential exacerbators is crucial to reducing the burden of disease. Notably, our analysis showed

an increased risk (OR=1.01; 95% CI:1.00 - 1.02) of myocardial infarction with a 10 μg/m^3

increase in PM_{10} after a lag 0 and lag 1 delay. Considering the high prevalence of MI, the

statistically significant risk increase of 1% indicates the absolute risk of PM10 is high.

Farhadi et al. [6]investigated short-term exposure to $PM_{2.5}$ and its effects on the risk of

MI. Our study and their study both looked at RR and OR as a risk outcome, and assessed studies

with a case-crossover and time-series design. However, our study additionally looked at HR as a

risk outcome, and assessed studies with a cohort design. Both studies also looked at the effects of

lags, but lag definitions were different. Our study looked at lags with short delays (lag 0 = 0-24

hours, lag 1 = 24-96 hours, lag 2 = >96 hours), whereas Farhadi et al. [6] looked at long delays

(short follow-up period = <4 years, long follow-up period = >4 years). The results of their study

are comparable to our findings as the RR and OR in our meta-analysis were similar. Their meta-

analysis reported a RR of 1.02 (95% CI: 1.01–1.03; P-value $\leq$ 0.0001) [6]. However, our results

had more homogeneity, likely due to stratification of studies by risk outcome measures and

smaller lag ranges. Our statistical analysis yielded ORs of 1.01 (95% CI 1.00–1.02; P-value

<0.05) for lag 0 and lag 1. These results indicate that while the risk of MI from exposure to PM_{10}

is 1% less in comparison to $PM_{2.5}$ exposure, there is a potential impact of $PM_{2.5-10}$ pollutants on

MI risk. This indicates that the risk of $PM_{2.5-10}$ entering the bloodstream through the alveolar

barrier and causing cardiovascular adversities, is similar to that of $PM_{2.5}$.

When assessing cardiovascular health concerns related to particulate matter, PM_{10} should

not be discredited – especially in communities of low socioeconomic status, as these populations

are most vulnerable to the health effects of air pollution [52].

4.1. *Strengths and Limitations*

This study had some limitations. The first one was the varying adjustment models utilized

by included studies. There are multiple participant confounders (hypertension, diet, sleep, etc.) to

consider since PM_{10} association to MI risk is observational, which could explain potential

differences in MI risk outcomes. The retrospective nature of case-crossover and time-series

designs is also a potential drawback considering the ambiguity of lag definitions. Another factor

to consider is the heterogeneity of the studies. All studies varied in study design, population and

age characteristics, and the exposure assessment methods. This review's studies included a

variety of population sizes from various places, which can limit how the results can be applied to

specific people or environments. The strength and direction of the relationship between PM_{10}

exposure and the risk of MI may be influenced by variables in air pollution sources, climate, and

demographic factors.

4.2. *Next Steps*

To enhance the precision and applicability of the findings, certain measures can be taken

in the future. The compatibility and consistency of studies assessing MI risk can be increased by

standardizing a specific risk outcome measure. While conversion algorithms exist for RR and

OR, they depend on an estimated incidence variable (P0) which can constrain data synthesis

validity. Future research should seek to study the distinct risk of STEMI and NSTEMI.

Understanding these MI subtypes' diverse relationships with PM_{10} exposure can offer more

specialized insights for prevention. Comprehensive adjustment models should also be more

widely employed to ensure confounding variables are taken into account for all associative PM_{10}

studies. This will allow for more robust and reliable risk estimates. By taking these additional

actions, researchers can improve the accuracy and application of the findings, and clarify the link

between exposure to PM_{10} and the risk of MI.

5. Conclusions

The results of this meta-analysis showed that there is an increased risk of MI with a

$10\mu g/m^3$ increase in PM_{10} exposure (OR=1.01; 95% CI:1.00 - 1.02). Risk of MI is also marginal

between PM_{10} and $PM_{2.5}$ exposure. It is important to recognize and assess the cardiovascular

impacts of PM_{10} alongside other pollutants. This can help guide environmental policy,

individual-level preventive measures, and global public health initiatives focused on lessening

the disease burden of MI caused by PM_{10} exposure.

6. Acknowledgments

We have no funding source or competing interests to report. All data used within this
review can be found in the manuscript and supporting information.

7. References

1. Hamanaka RB, Mutlu GM. Particulate matter air pollution: effects on the cardiovascular system. Frontiers in endocrinology. 2018 Nov 16;9:680.

2. Particle Pollution | Air | CDC [Internet]. Available from: https://www.cdc.gov/air/particulate_matter.html

3. Tornevi A, Olstrup H, Forsberg B. Short-term associations between PM10 and respiratory health effects in Visby, Sweden. Toxics. 2022 Jun 17;10(6):333.

4. Thangavel P, Park D, Lee YC. Recent insights into particulate matter (PM2. 5)-mediated toxicity in humans: an overview. International journal of environmental research and public health. 2022 Jun 19;19(12):7511.

5. Nemmar A, Hoet PM, Vanquickenborne B, Dinsdale D, Thomeer M, Hoylaerts MF, Vanbilloen H, Mortelmans L, Nemery B. Passage of inhaled particles into the blood circulation in humans. Circulation. 2002 Jan 29;105(4):411-4.

6. Farhadi Z, Abulghasem Gorgi H, Shabaninejad H, Aghajani Delavar M, Torani S. Association between PM 2.5 and risk of hospitalization for myocardial infarction: a systematic review and a meta-analysis. BMC Public Health. 2020 Dec;20:1-2.

7. Veritas Health Innovation. Covidence systematic review software [Internet]. Covidence. [cited 2023 Jul 03]. Available from: www.covidence.org

8. Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart LA. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. Systematic reviews. 2015 Dec;4(1):1-9.

9. Study quality assessment tools. National Heart, Lung, and Blood Institute. NHLBI [Internet]. https://www.nhlbi.nih.gov/health-topics/study-qualityassessment-tools

10. Argacha JF, Collart P, Wauters A, Kayaert P, Lochy S, Schoors D, Sonck J, de Vos T,

Forton M, Brasseur O, Beauloye C. Air pollution and ST-elevation myocardial infarction:

a case-crossover study of the Belgian STEMI registry 2009–2013. International journal of

cardiology. 2016 Nov 15;223:300-5.

11. Bard D, Kihal W, Schillinger C, Fermanian C, Segala C, Glorion S, Arveiler D, Weber C.

Traffic-related air pollution and the onset of myocardial infarction: disclosing benzene as

a trigger? A small-area case-crossover study. PloS one. 2014 Jun 16;9(6):e100307.

12. Bhaskaran K, Hajat S, Armstrong B, Haines A, Herrett E, Wilkinson P, Smeeth L. The

effects of hourly differences in air pollution on the risk of myocardial infarction: case

crossover analysis of the MINAP database. Bmj. 2011 Sep 20;343.

13. Buszman PE, Derbisz K, Kwasiborski P, Chrząszcz P, Mularska M, Baron D, Sobieszek

A, Mendyk A, Skoczylas P, Cisowski M, Buszman PP. Impact of air pollution on

hospital patients admitted with ST-and non-ST-segment elevation myocardial infarction

in heavily polluted cities within the European Union. Cardiology Journal.

2020;27(5):541-7.

14. Cheng J, Tong S, Su H, Xu Z. Hourly air pollution exposure and emergency department

visit for acute myocardial infarction: vulnerable populations and susceptible time

window. Environmental Pollution. 2021 Nov 1;288:117806.

15. Claeys MJ, Coenen S, Colpaert C, Bilcke J, Beutels P, Wouters K, Legrand V, Van

Damme P, Vrints C. Environmental triggers of acute myocardial infarction: results of a

nationwide multiple-factorial population study. Acta Cardiologica. 2015 Dec

1;70(6):693-701.

435 16. Collart P, Dramaix M, Levêque A, Coppieters Y. Short-term effects of air pollution on

436  hospitalization for acute myocardial infarction: age effect on lag pattern. International

437  journal of environmental health research. 2017 Jan 2;27(1):68-81.

438 17. Davoodabadi Z, Soleimani A, Pourmoghaddas A, Hosseini SM, Jafari-Koshki T, Rahimi

439  M, Shishehforoush M, Lahijanzadeh A, Sadeghian B, Moazam E, Mohebi MB.

440  Correlation between air pollution and hospitalization due to myocardial infarction.

441  ARYA atherosclerosis. 2019 Jul;15(4):161.

442 18. Downward GS, van Nunen EJ, Kerckhoffs J, Vineis P, Brunekreef B, Boer JM, Messier

443  KP, Roy A, Verschuren WM, van der Schouw YT, Sluijs I. Long-term exposure to

444  ultrafine particles and incidence of cardiovascular and cerebrovascular disease in a

445  prospective study of a Dutch cohort. Environmental health perspectives. 2018 Dec

446  19;126(12):127007.

447 19. Huss A, Spoerri A, Egger M, Röösli M. Aircraft noise, air pollution, and mortality from

448  myocardial infarction. Epidemiology. 2010 Nov 1:829-36.

449 20. Kim OJ, Lee SH, Kang SH, Kim SY. Incident cardiovascular disease and particulate

450  matter air pollution in South Korea using a population-based and nationwide cohort of

451  0.2 million adults. Environmental Health. 2020 Dec;19(1):1-2.

452 21. Konduracka E, Niewiara Ł, Guzik B, Kotynia M, Szolc P, Gajos G, Nessler J, Podolec P,

453  Żmudka K. Effect of short-term fluctuations in outdoor air pollution on the number of

454  hospital admissions due to acute myocardial infarction among inhabitants of Kraków,

455  Poland. Polskie Archiwum Medycyny Wewnętrznej= Polish Archives of Internal

456  Medicine. 2019;129(2).

22. Kuźma Ł, Wańha W, Kralisz P, Kazmierski M, Bachórzewska-Gajewska H, Wojakowski

W, Dobrzycki S. Impact of short-term air pollution exposure on acute coronary syndrome

in two cohorts of industrial and non-industrial areas: A time series regression with

6,000,000 person-years of follow-up (ACS-Air Pollution Study). Environmental

Research. 2021 Jun 1;197:111154.

23. Lipsett MJ, Ostro BD, Reynolds P, Goldberg D, Hertz A, Jerrett M, Smith DF, Garcia C,

Chang ET, Bernstein L. Long-term exposure to air pollution and cardiorespiratory

disease in the California teachers study cohort. American journal of respiratory and

critical care medicine. 2011 Oct 1;184(7):828-35.

24. Nuvolone D, Balzi D, Chini M, Scala D, Giovannini F, Barchielli A. Short-term

association between ambient air pollution and risk of hospitalization for acute myocardial

infarction: results of the cardiovascular risk and air pollution in Tuscany (RISCAT)

study. American journal of epidemiology. 2011 Jul 1;174(1):63-71.

25. Puett RC, Schwartz J, Hart JE, Yanosky JD, Speizer FE, Suh H, Paciorek CJ, Neas LM,

Laden F. Chronic particulate exposure, mortality, and coronary heart disease in the

nurses' health study. American journal of epidemiology. 2008 Nov 15;168(10):1161-8.

26. Royé D, Zarrabeitia MT, Fdez-Arroyabe P, Gutiérrez AÁ, Santurtún A. Role of apparent

temperature and air pollutants in hospital admissions for acute myocardial infarction in

the North of Spain. Revista Española de Cardiología (English Edition). 2019 Aug

1;72(8):634-40.

27. Soleimani Z, Darvishi Boloorani A, Khalifeh R, Griffin DW, Mesdaghinia A. Short-term

effects of ambient air pollution and cardiovascular events in Shiraz, Iran, 2009 to 2015.

Environmental Science and Pollution Research. 2019 Mar 8;26:6359-67.

28. Vidale S, Arnaboldi M, Bosio V, Corrado G, Guidotti M, Sterzi R, Campana C. Short-

term air pollution exposure and cardiovascular events: a 10-year study in the urban area

of Como, Italy. International journal of cardiology. 2017 Dec 1;248:389-93.

29. Yang J, Liu M, Cheng Q, Yang L, Sun X, Kan H, Liu Y, Bell ML, Dasan R, Gao H, Yao

X. Investigating the impact of air pollution on AMI and COPD hospital admissions in the

coastal city of Qingdao, China. Frontiers of Environmental Science & Engineering. 2022

May;16:1-3.

30. Yen CC, Chen PL. Effect of short-term exposure to fine particulate matter and particulate

matter pollutants on triggering acute myocardial infarction and acute heart failure. The

American Journal of Cardiology. 2022 Jul 15;175:158-63.

31. Zhang Q, Qi W, Yao W, Wang M, Chen Y, Zhou Y. Ambient particulate matter (PM2.

5/PM10) exposure and emergency department visits for acute myocardial infarction in

Chaoyang District, Beijing, China during 2014: a case-crossover study. Journal of

Epidemiology. 2016 Oct 5;26(10):538-45.

32. Zhu X, Qiu H, Wang L, Duan Z, Yu H, Deng R, Zhang Y, Zhou L. Risks of hospital

admissions from a spectrum of causes associated with particulate matter pollution.

Science of The Total Environment. 2019 Mar 15;656:90-100.

33. Akbarzadeh MA, Khaheshi I, Sharifi A, Yousefi N, Naderian M, Namazi MH, Safi M,

Vakili H, Saadat H, Parsa SA, Nickdoost N. The association between exposure to air

pollutants including PM10, PM2. 5, ozone, carbon monoxide, sulfur dioxide, and

nitrogen dioxide concentration and the relative risk of developing STEMI: A case-

crossover design. Environmental research. 2018 Feb 1;161:299-303.

34. Atkinson RW, Carey IM, Kent AJ, van Staa TP, Anderson HR, Cook DG. Long-term exposure to outdoor air pollution and incidence of cardiovascular diseases. Epidemiology. 2013 Jan 1:44-53.

35. Cheng MF, Tsai SS, Yang CY. Air pollution and hospital admissions for myocardial infarction in a tropical city: Kaohsiung, Taiwan. Journal of Toxicology and Environmental Health, Part A. 2009 Aug 31;72(19):1135-40.

36. Cramer J, Jørgensen JT, Hoffmann B, Loft S, Bräuner EV, Prescott E, Ketzel M, Hertel O, Brandt J, Jensen SS, Backalarz C. Long-term exposure to air pollution and incidence of myocardial infarction: a Danish nurse cohort study. Environmental health perspectives. 2020 May 6;128(5):057003.

37. Kim H, Kim J, Kim S, Kang SH, Kim HJ, Kim H, Heo J, Yi SM, Kim K, Youn TJ, Chae IH. Cardiovascular effects of long□term exposure to air pollution: a population□based study with 900 845 person□years of follow□up. Journal of the American Heart Association. 2017 Nov 8;6(11):e007170.

38. Kuźma Ł, Pogorzelski S, Struniawski K, Bachórzewska-Gajewska H, Dobrzycki S. Exposure to air pollution—a trigger for myocardial infarction? A nine-year study in Bialystok—the capital of the Green Lungs of Poland (BIA-ACS registry). International Journal of Hygiene and Environmental Health. 2020 Aug 1;229:113578.

39. Lee MW, Choi BG, Kim SW, Rha SW, Shim MS, Kim DJ, Seo HS, Oh DJ, Jeong MH, Korea Acute Myocardial Infarction Registry (KAMIR) Investigators, Kim YH. Air pollution and short□term clinical outcomes of patients with acute myocardial infarction. Clinical and Experimental Pharmacology and Physiology. 2017 Jun;44(6):631-8.

40. Liu X, Bertazzon S, Villeneuve PJ, Johnson M, Stieb D, Coward S, Tanyingoh D,

Windsor JW, Underwood F, Hill MD, Rabi D. Temporal and spatial effect of air

pollution on hospital admissions for myocardial infarction: a case-crossover study.

Canadian Medical Association Open Access Journal. 2020 Oct 1;8(4):E619-26.

41. Pan HY, Cheung SM, Chen FC, Wu KH, Cheng SY, Chuang PC, Cheng FJ. Short-term

effects of ambient air pollution on ST-elevation myocardial infarction events: are there

potentially susceptible groups?. International journal of environmental research and

public health. 2019 Oct;16(19):3760.

42. Rasche M, Walther M, Schiffner R, Kroegel N, Rupprecht S, Schlattmann P, Schulze PC,

Franzke P, Witte OW, Schwab M, Rakers F. Rapid increases in nitrogen oxides are

associated with acute myocardial infarction: A case-crossover study. European journal of

preventive cardiology. 2018 Nov 1;25(16):1707-16.

43. Rodins V, Lucht S, Ohlwein S, Hennig F, Soppa V, Erbel R, Joeckel KH, Weimar C,

Hermann DM, Schramm S, Moebus S. Long-term exposure to ambient source-specific

particulate matter and its components and incidence of cardiovascular events–the Heinz

Nixdorf Recall study. Environment international. 2020 Sep 1;142:105854.

44. Sahlén A, Ljungman P, Erlinge D, Chan MY, Yap J, Hausenloy DJ, Yeo KK, Jernberg T.

Air pollution in relation to very short-term risk of ST-segment elevation myocardial

infarction: case-crossover analysis of SWEDEHEART. International Journal of

Cardiology. 2019 Jan 15;275:26-30.

45. Sen T, Astarcioglu MA, Asarcikli LD, Kilit C, Kafes H, Parspur A, Yaymaci M, Pinar M,

Tüfekcioglu O, Amasyali B. The effects of air pollution and weather conditions on the

incidence of acute myocardial infarction. The American Journal of Emergency Medicine.

2016 Mar 1;34(3):449-54.

46. Wang XD, Zhang XM, Zhuang SW, Luo Y, Kang S, Liu YL. Short-term effects of air

pollution on acute myocardial infarctions in Shanghai, China, 2013–2014. Journal of

geriatric cardiology: JGC. 2016 Feb;13(2):132.

47. Wichmann J, Rosengren A, Sjöberg K, Barregard L, Sallsten G. Association between

ambient temperature and acute myocardial infarction hospitalisations in Gothenburg,

Sweden: 1985–2010. PloS one. 2013 Apr 30;8(4):e62059.

48. Wichmann J, Sjöberg K, Tang L, Haeger-Eugensson M, Rosengren A, Andersson EM,

Barregard L, Sallsten G. The effect of secondary inorganic aerosols, soot and the

geographical origin of air mass on acute myocardial infarction hospitalisations in

Gothenburg, Sweden during 1985–2010: a case-crossover study. Environmental Health.

2014 Dec;13(1):1-6.

49. Wolf K, Schneider A, Breitner S, Meisinger C, Heier M, Cyrys J, Kuch B, von Scheidt

W, Peters A, KORA Study Group. Associations between short-term exposure to

particulate matter and ultrafine particles and myocardial infarction in Augsburg,

Germany. International journal of hygiene and environmental health. 2015 Aug

1;218(6):535-42.

50. Yu Y, Yao S, Dong H, Ji M, Chen Z, Li G, Yao X, Wang SL, Zhang Z. Short-term

effects of ambient air pollutants and myocardial infarction in Changzhou, China.

Environmental Science and Pollution Research. 2018 Aug;25:22285-93.

51. Mc Namara K, Alzubaidi H, Jackson JK. Cardiovascular disease as a leading cause of

death: how are pharmacists getting involved?. Integrated pharmacy research and practice.

2019 Feb 4:1-1.

52. Hajat A, Hsia C, O'Neill MS. Socioeconomic disparities and air pollution exposure: a

global review. Current environmental health reports. 2015 Dec;2:440-50.

8. Supporting Information Captions

S1 Table. Search strategy

S2 Table. Adjusted confounding variables for studies included in the meta-analysis

S3 Table. Study characteristics of remaining studies included in the review

S4 Table. Methodological risk of bias assessment

S5 Table. GRADE assessment for RR – Lag 1 (1-3 days [24-96hrs]). O = Criteria Satisfied, -1
= Criteria Unsatisfied

S6 Table. GRADE assessment for RR – Lag 1 (1-3 days [24-96hrs]). O = Criteria Satisfied, -1
= Criteria Unsatisfied

S7 Table. GRADE assessment for RR – Lag 2 (3+ days [96hrs+]). O = Criteria Satisfied, -1 =
Criteria Unsatisfied

S8 Table. GRADE assessment for OR – Lag 0 (Same day [0-24hrs]). O = Criteria Satisfied, -
1 = Criteria Unsatisfied

S9 Table. GRADE assessment for OR – Lag 1 (1-3 days [24-96hrs]). O = Criteria Satisfied, -1
= Criteria Unsatisfied

S10 Table. GRADE assessment for OR – Lag 2 (3+ days [96hrs+]). O = Criteria Satisfied, -1
= Criteria Unsatisfied

S11 Table. GRADE assessment for HR – Lag 0 (Same day [0-24hrs]). O = Criteria Satisfied,
-1 = Criteria Unsatisfied

S12 Table. GRADE assessment for HR – Lag 1 (1-3 days [24-96hrs]). O = Criteria Satisfied, -
1 = Criteria Unsatisfied

S13 Table. GRADE assessment for HR – Lag 2 (3+ days [96hrs+]). O = Criteria Satisfied, -1
= Criteria Unsatisfied

S1 Fig. Lag 0 HR for MI after 10 $\mu g/m^3$ increase in PM_{10} exposure.

S2 Fig. Lag 1 HR for MI after 10 $\mu g/m^3$ increase in PM_{10} exposure.

S3 Fig. Protocol

S4 Fig. PRISMA Checklist