Multivariate Associations Between Structural Brain Networks, Genetics, Environments, and Cognitive-Psychopathological Traits in Children

Jungwoo Seo[1], Eunji Lee[2], Bo-Gyeom Kim[2], Gakyung Kim[1], Yoonjung Yoonie Joo[2,3,4], Jiook Cha[1,2,5,6]

1. Department of Brain and Cognitive Sciences, College of Natural Sciences, Seoul National University, Seoul, 08826, South Korea
2. Department of Psychology, College of Social Sciences, Seoul National University, Seoul, 08826, South Korea
3. Department of Digital Health, Samsung Advanced Institute for Health Sciences & Technology (SAIHST), Sungkyunkwan University, Seoul, 06355, South Korea
4. Research Institute for Future Medicine, Samsung Medical Center, Seoul, 06351, South Korea
5. Institute of Psychological Science, Seoul National University, Seoul, 08826, South Korea
6. Graduate School of Artificial Intelligence, Seoul National University, Seoul, 08826, South Korea

Correspondence to: Jiook Cha, PhD

Gwanak-ro 1, Building 16, Suite M512, Gwanakgu, Seoul, 08826, South Korea, connectome@snu.ac.kr

Short title: Children's brain network, gene, environment, and cognitive-behavior

Keywords: structural brain network, genetic factor, environmental factor, cognitive-behavior, preadolescent, multivariate analysis

Abstract

Brain development in childhood is shaped by complex interactions between genetic predispositions, environmental influences, and neural connectivity, yet how these factors jointly contribute to cognitive and mental health outcomes remains unclear. Structural brain networks, quantified through graph-theoretic measures, have been linked to cognition and psychiatric risk, but the extent to which genetic architecture and environmental exposures shape these networks, and whether brain networks mediate these influences, is not well understood. Here we show that genetic predispositions related to cognitive ability and socioeconomic status (SES) exhibit the strongest covariation with structural brain network topology in children. Using sparse canonical correlation analysis (SCCA) on ABCD Study data (N = 10,343), we identified robust associations between brain network properties, polygenic scores for cognition, SES indicators, and cognitive-psychopathological phenotypes. Mediation analysis further demonstrated that structural brain networks partially mediate the influence of genetic and environmental factors on cognitive performance and mental health outcomes, suggesting that neurodevelopmental trajectories may be shaped by both genetic liability and modifiable environmental conditions. These findings provide empirical support for a multivariate, systems-level perspective on brain development and cognitive-psychopathological variation in youth. By elucidating shared neural substrates underlying genetic and environmental influences, this work advances our understanding of brain network development and highlights potential pathways for individualized interventions and predictive modeling in developmental psychiatry and neuroscience.

Introduction

Childhood and adolescence are critical periods for brain development (Bethlehem et al., 2022). Proper brain development during this time is vital for cognitive and behavioral maturation (Bunge & Wright, 2007; Luna et al., 2010) and mental health (Fornito et al., 2015; Paus et al., 2008). This development is shaped by a complex interplay between genetic predispositions, environmental influences, and brain network dynamics, yet the precise mechanisms underlying these relationships remain poorly understood. Therefore, understanding the connections between the brain, cognitive-behavioral traits in children, and the impact of genetics and the environment on brain development is crucial in developmental and clinical neuroscience.

The Adolescent Brain and Cognitive Development (ABCD) study provides an unprecedented opportunity to examine these interrelations in a large, representative sample (Jernigan et al., 2018). Leveraging multimodal neuroimaging, genome-wide polygenic scores (PGSs), and extensive environmental and behavioral assessments, the ABCD dataset enables a more comprehensive modeling of developmental trajectories than previously possible.

The brain is best understood as a complex, interconnected network, rather than a collection of isolated regions. Graph-theoretic approaches to brain connectivity provide powerful tools to quantify structural network properties such as integration (e.g., global efficiency), segregation (e.g., modularity), and centrality (e.g., degree centrality) (Rubinov & Sporns, 2010). Diffusion MRI and tractography allow for the reconstruction of white matter networks, enabling researchers to examine how these

structural connections support cognition (Jeurissen et al., 2019; Sotiropoulos & Zalesky, 2019). Notably, individual differences in brain network organization have been linked to cognitive abilities (Bathelt, Gathercole, Butterfield, et al., 2018; Kim et al., 2016; Koenis et al., 2015; Ma et al., 2017; Suprano et al., 2020), psychiatric risk (Alexander-Bloch et al., 2010; Collin et al., 2017; Rudie et al., 2012), and environmental influences such as socioeconomic status (Kim et al., 2019; Tooley et al., 2020). However, how genetic and environmental factors jointly shape structural brain networks, and whether these networks mediate their influence on cognitive and psychiatric outcomes, remains poorly understood.

Twin and heritability studies suggest that brain network properties are, at least in part, genetically influenced, with estimates of heritability for measures like global efficiency ranging from 25% to 70% (Koenis et al., 2015; van den Heuvel et al., 2013). Yet, the specific genetic variants shaping these structural networks—particularly in the context of cognitive and psychiatric outcomes—remain poorly characterized. Investigating this gene-brain-behavior interplay is essential for understanding how biological predispositions manifest in cognitive and mental health outcomes during development.

Despite valuable insights from previous studies, most prior research has employed univariate approaches, limiting the ability to capture the multivariate complexity of the gene/environment-brain-behavior relationship. To address this limitation, canonical correlation analysis (CCA) is useful for investigating holistic relationships underlying a set of variables. CCA models multivariate associations linking sets of variables from two or more domains by maximizing the canonical correlation between them (Hotelling, 1936; Wang et al., 2020). Indeed, CCA has been

widely applied in studying links between brain connectivity, cognitive function, genetics, and environmental factors (Alnaes et al., 2020; Fernandez-Cabello et al., 2022; Modabbernia et al., 2021; Smith et al., 2015; Wang et al., 2020). However, traditional CCA methods may struggle with high-dimensional datasets, necessitating more advanced approaches. To further address these limitations, we apply sparse canonical correlation analysis (SCCA) (Witten et al., 2009), a multivariate method that identifies maximally covarying patterns between genetic factors, brain network properties, and behavioral traits while reducing dimensionality and enhancing interpretability.

Utilizing the ABCD dataset's expansive genetic-environmental-neuroimaging-behavioral data, this study aims to answer two fundamental questions. Firstly, "How do genetic predispositions/environmental influences shape structural brain network properties in children?"; Secondly, "Do brain network properties mediate the influence of genetic and environmental factors on cognitive or psychiatric phenotypes?"

By integrating genetics, environmental, neuroimaging, and behavioral assessments in a large, developmentally significant cohort, this study seeks to elucidate the complex gene-environmental-brain-behavior pathways that shape cognitive and mental health outcomes. This research contributes to a more comprehensive understanding of the biological and environmental foundations of brain development and lays the groundwork for future interventions aimed at optimizing cognitive and psychological well-being in youth.

Methods and Materials

ABCD participants

We used genetic, neuroimaging, environmental, and phenotypic data from the Adolescent Brain Cognitive Development (ABCD) study (http://abcdstudy.org), specifically from release 2.0 for genetic and neuroimaging data, release 3.0 for genetic ancestry information, and release 5.1 for environmental and phenotypic data. The ABCD study, the largest longitudinal investigation of brain development and child health in the United States, recruited multiethnic children (N=11,875) aged 9-10 years from 21 research sites. The self-reported ethnicities of participants included 52.3% White, 20.3% Hispanic, 14.7% Black, and 12.5% Asian and others. All participants provided informed assent, and their parents or legal guardians provided informed consent before participating in the study.

Genotype Data and Polygenic Scores

The genotype data used in this study were obtained from the ABCD cohort, with DNA samples genotyped at Rutgers University Cell and DNA Repository (RUCDR) using the Affymetrix NIDA Smoke Screen Array. Standard quality control (QC) procedures were applied to remove variants with low genotype call rate and minor allele frequency (MAF). Genotype imputation was performed using the Michigan Imputation Server (Das et al., 2016) with the 1000 Genomes reference panel (Genomes Project et al., 2015). We further filtered out imputed variants that did not meet our QC criteria. To account for population structure and relatedness, principal component analysis (PCA) and kinship-based filtering were used to exclude close relatives (Conomos et al., 2015; Conomos et al., 2016).

Polygenic scores (PGSs) for 30 cognitive, psychiatric, and behavioral traits were calculated based on genome-wide association study (GWAS) summary statistics. These included PGSs for cognitive performance, educational attainment, schizophrenia, bipolar disorder, depression, insomnia, body mass index, and automobile speed propensity, among others. The PGSs were derived using PRS-CS (Ge et al., 2019), a Bayesian regression approach, with optimal hyperparameter

selection.

The genotype data and polygenic scores used in this study are identical to

those reported in (Joo et al., 2024). A complete list of the 30 polygenic scores, along

with further methodological details on genotype data processing, GWAS sources,

ancestry-based adjustments, and validation procedures can be found in the

Supplementary Materials or (Joo et al., 2024).

Environmental Factors

To investigate the relationship between children's brain network properties

and their environment, we analyzed 56 variables representing various environmental

aspects. These variables included indicators of family and neighborhood

socioeconomic status, such as family income, parental education level, marital

status, and neighborhood deprivation based on the Area Deprivation Index (ADI).

Additionally, variables related to prenatal substance exposure (e.g., tobacco, alcohol,

cocaine, and marijuana), as well as parental and family factors such as parental

acceptance and family conflict, were included.

To maximize statistical power, we imputed missing data to include as many

samples as possible in the analysis. To account for uncertainty in imputing missing

values and to improve the accuracy and reliability of the imputed values, we

employed the Multiple Imputation by Chained Equations (MICE) method (Van

Buuren & Groothuis-Oudshoorn, 2011). We set the number of imputation iterations to

40 to ensure stable convergence. All imputation procedures were conducted using

the `statsmodels` package in Python. To ensure that our analysis was robust to

imputation, we also conducted analyses using only the complete data without any

missing values and included these results as supplementary material.

Phenotype Data

To explore the relationship between children's brain network properties and

their cognitive ability, mental well-being, and physical health, we examined 86

phenotype variables. To evaluate cognitive ability, we examined the NIH Toolbox

measurements, which included fluid, crystallized, and overall cognition scores, along

with domain-specific task scores for episodic memory, executive function, language,

and processing speed (Weintraub et al., 2013). For assessing the children's mental

health, our analysis encompassed a broad range of emotional and behavioral

measurements, including instruments such as the Child Behavioral Checklist

(CBCL), the Kiddie-Structured Assessment for Affective Disorders and Schizophrenia

for DSM-5 (KSADS-5), and the Prodromal Psychosis Scale (PPS). We also included

measures of behavioral tendencies, such as behavioral inhibition and activation

(BIS/BAS), which assess avoidance behaviors, reward sensitivity, and behavioral

control; impulsivity traits (UPPS), which measure dimensions such as urgency,

perseverance, and sensation seeking; and sleep-related problems, which provide

insights into sleep patterns, quality, and related disorders. For the children's physical

health, we included data such as physical activity. To maximize statistical power,

missing values in the phenotype data were imputed using the Multiple Imputation by

Chained Equations method (Van Buuren & Groothuis-Oudshoorn, 2011).

Structural Brain Network Construction

Detailed procedures for acquiring and preprocessing MRI data are described

in (Kim et al., 2022). In brief, we used structural and diffusion MRI data acquired by

the ABCD study (Casey et al., 2018) from data release 2.0. The preprocessing steps,

as detailed in (Kim et al., 2022), included eddy current and head motion correction,

diffusion gradient adjustment, and various distortion corrections. Quality control (QC)

was performed using freesurfer QC metric (fsqc_qc) and raw dMRI QC metric

(iqc_dmri_1_qc_score). To estimate brain structural networks from neuroimaging,

individual connectome data was generated. This was achieved by applying MRtrix3

(Tournier et al., 2019) to the preprocessed dMRI data to estimate whole-brain white

matter tracts and generate individualized connectomes. Probabilistic tractography

was performed using constrained-spherical deconvolution (CSD) (Calamante et al.,

2010; Tournier et al., 2007) with random seeding across the brain and target

streamline counts of 20 million. Initial tractograms were filtered using spherical-

deconvolution informed filtering (2:1 ratio)(Smith et al., 2013), resulting in a final

streamline count of 10 million. An 84x84 whole-brain connectome matrix was

generated for each participant using the T1-based parcellation and segmentation

from FreeSurfer with Desikan-Killiany atlas (Desikan et al., 2006)(68 nodes for the

cortical regions and 16 nodes for the subcortical regions). This approach ensured

that individual participants' connectomes were based on their neuroanatomy. The

computation was conducted on supercomputers at Argonne Leadership Computing

Facility Theta and Texas Advanced Computing Center Stampede2.

Brain Network Measures (BNMs)

We used the connectome matrix to construct an undirect weighted graph

representing the structural brain network. Nodes and edges in the graph represent

parcellated gray matter regions and connections between them, respectively.

Connection strength was quantified by the streamline counts. To account for the

potential false positive connections generated by probabilistic tractography and their

impact on network topology, we eliminated extremely weak connections (streamline

counts less than 3). After thresholding, we excluded individuals with at least one

isolated node, assuming all brain regions are communicable via at least one path.

We calculated 13 different types of brain network measurements (BNMs)

representing different aspects of brain network's property (Rubinov & Sporns, 2010;

van den Heuvel & Sporns, 2011). We calculated eight global graph metrics (including

network density, modularity, normalized modularity, normalized average clustering

coefficient, normalized characteristic path length, global efficiency, normalized global

efficiency, small worldness) and five nodal graph metrics (including degree, strength,

clustering coefficient, betweenness centrality, nodal efficiency) to represent brain

network's global and regional properties. All graph measures were calculated using

the package Brain Connectivity Toolbox (https://sites.google.com/site/bctnet/).

Statistical analysis

236   *Sparse Canonical Correlation Analysis*

Brain Network Measures (BNMs)

We used the connectome matrix to construct an undirect weighted graph

representing the structural brain network. Nodes and edges in the graph represent

parcellated gray matter regions and connections between them, respectively.

Connection strength was quantified by the streamline counts. To account for the

potential false positive connections generated by probabilistic tractography and their

impact on network topology, we eliminated extremely weak connections (streamline

counts less than 3). After thresholding, we excluded individuals with at least one

isolated node, assuming all brain regions are communicable via at least one path.

We calculated 13 different types of brain network measurements (BNMs)

representing different aspects of brain network's property (Rubinov & Sporns, 2010;

van den Heuvel & Sporns, 2011). We calculated eight global graph metrics (including

network density, modularity, normalized modularity, normalized average clustering

coefficient, normalized characteristic path length, global efficiency, normalized global

efficiency, small worldness) and five nodal graph metrics (including degree, strength,

clustering coefficient, betweenness centrality, nodal efficiency) to represent brain

network's global and regional properties. All graph measures were calculated using

the package Brain Connectivity Toolbox (https://sites.google.com/site/bctnet/).

Statistical analysis

Sparse Canonical Correlation Analysis

To examine a latent mode of covariation between structural brain network

properties and various polygenic scores, environmental factors, and phenotypic

outcomes, we used sparse canonical correlation analysis (Witten et al., 2009)

between brain network measures and three types of non-imaging data (i.e., PGSs,

environmental variables, phenotype variables) separately. While traditional CCA can

be effective, it often suffers from overfitting and interpretability issues in high-

dimensional datasets. SCCA, with its L1 regularization, addresses these issues by

producing sparse solutions that enhance interpretability and reduce overfitting.

Although our dataset is not high-dimensional enough to make traditional CCA

infeasible, we chose SCCA to identify interpretable patterns and ensure robust

results.

The most popular algorithm for sparse canonical correlation analysis is

penalized matrix decomposition (PMD)(Witten et al., 2009), which solves

optimization problem of below equation for given two sets of data matrix $X_{n \times p}$, $Y_{n \times q}$.

(n: sample size; p, q: the number of variables of domain X and Y respectively; u, v:

canonical weights of domain X and Y respectively; c1, c2: regularization parameter)

$$\max cov(Xu, \ Yv)$$

$$s.t. \ \| u \|_2 = \| v \|_2 = 1, \ \| u \|_1 \leq c_1, \ \| v \|_1 \leq c_2$$

To interpret Witten's sparse canonical correlation analysis as correlation

maximization, we need to assume covariance matrices $X^T X$, $Y^T Y$ are identity

matrices (Witten et al., 2009). But in our study with the high dimensional brain

datasets, the assumption is hardly satisfied. For this reason, we interpreted Witten's

sparse canonical correlation analysis as a maximizing covariance algorithm between

two sets of variables rather than maximizing correlation.

To test the generalizability of the sparse canonical correlation analysis

results, we split the dataset into a training and test set. To reduce ABCD site-

sensitive bias, we performed a stratified train (80%) - test (20%) split based on site

ID. For the sparse canonical correlation analysis with PGS and brain network

measures, we only used participants classified as genetically European ancestry to

control genetic confounding effects as the main analysis. To ensure that these

analyses generalize to a multiethnic dataset, we conducted additional analyses on

the multiethnic dataset. In the multiethnic analysis, genetic ancestry was included as

an additional covariate. **Table 1 and Supplementary Table 1** summarize the

demographic information of the samples included in main and supplementary

analysis, respectively.

We attempted to control for potential confounding effects from age, sex, self-

reported race/ethnicity, ABCD study site, and handedness. Similar to (Modabbernia

et al., 2021), we controlled these potential confounding effects by regressing out the

variance explained by age, sex, age * sex, age^2, age^2 * sex, self-reported

race/ethnicity, ABCD study site, and handedness from all variables prior to

performing SCCA analysis. For binary variables such as KSADS diagnosis, we used

logistic regression to regress out these effects. The residualized data was then used

as input for sparse canonical correlation analysis (SCCA). We used `SCCA_PMD`

function from the Python 'cca-zoo' package. To ensure that the main findings are

robust and not driven by the selection of specific covariates, we performed additional

analyses using a reduced set of covariates (age, sex, self-reported race/ethnicity,

ABCD study site, and handedness).

We selected optimal L1 regularization parameters from 5-fold cross validation

searching from 0.1 to 1 with a step size of 0.05 for both X and Y variables

respectively. The optimal L1 parameter combination was selected to maximize the

covariance of validation set between canonical variates of the first component.

For each sparse canonical correlation analysis, we extracted five modes of

covariance. To examine the statistical significance of each mode, we used a

permutation test. By randomly shuffling the rows of one dataset and remaining the

other, we generated 5,000 permutation sets. The p-value of each component was

calculated based on the number of permutation sets having greater covariance than

that obtained from the original dataset, and FDR-correction was done within each

CCA.

295
$$p_{uncorrected} = \frac{N_{null\ cov > cov}}{N_{null}}$$

Selected variables and their loading depend on the input sample. To find

variables reliably related to each mode, we used bootstrap resampling. We randomly

resampled 5,000 times with replacement and assessed the 95% confidence interval

of each variable's loading and how consistently it was selected. We interpreted the

significant modes based on loading patterns of variables whose 95% confidence

interval of loading does not cross zero (Xia et al., 2018) and selected more

frequently than expected by chance (i.e., more frequently selected than expected by

binomial distribution). Because sparse canonical correlation analysis with bootstrap

sample may change the order of components (axis rotation) and signs (reflection)

(Misic et al., 2016; Xia et al., 2018), the re-alignment procedure is needed to

estimate confidence interval of loading properly. We matched the components and

signs based on cosine similarity of weight vectors obtained from original dataset and

bootstrap sample. To assess the reproducibility of the findings, we applied the model

to the held-out test set and estimated significance of each mode through the

permutation test.

Mediation Analysis

After performing sparse canonical correlation analysis (SCCA), we tested

whether brain network properties mediate the influence of genetic and environmental

factors on cognitive and psychiatric phenotypes. Since SCCA identifies latent modes

of covariation between brain network properties and multiple domains, the SCCA-

derived variates serve as optimal summary representations that capture the

dominant axes of covariation among brain network properties, genetic

predispositions, environmental influences, and phenotypic traits. We used these

summary scores in mediation analysis to test the hypothesized gene/environment–

brain network–phenotype pathway while reducing high-dimensional data into

interpretable components.

We used the same covariates as in the SCCA, which include age, sex, age *

sex, age^2, age^2 * sex, self-reported race/ethnicity, ABCD study site, and

handedness. The two-sided p-values for each path were estimated from 500

bootstrap samples using the `mediation_analysis` function from the Python 'pingouin'

package.

Results

328   Using sparse canonical correlation analysis (SCCA), we investigated

329 relationships between structural brain network properties and three domains:

330 genome-wide polygenic scores (PGSs), environmental factors, and phenotypic

331 outcomes. **Figure 1** presents only variables whose 95% confidence intervals for

332 loadings do not cross zero, ensuring reliable positive or negative associations.

Covariation between polygenic scores and structural brain network properties

334   Our analysis identified a statistically significant mode of covariation between 30

335 genome-wide polygenic scores and brain network measures (PGS-BNM mode 1, $p =$

336 0.001 , $cov = 0.895$, $r = 0.143$, FDR corrected).

337   In this mode of PGSs **(Figure 1A)**, polygenic scores reflecting cognitive ability

338 (e.g., IQ and educational attainment) showed the strongest positive loadings, while the

339 polygenic score for BMI showed a negative loading. Additionally, polygenic scores for

340 automobile speed propensity and cannabis use had moderate positive loadings,

341 whereas those for insomnia showed moderate negative loadings. Given these

342 dominant loading patterns, this mode represents variance along the 'Cognitive-

343 Obesity Genetic Axis,' primarily reflecting genetic traits related to cognitive ability.

344 Loadings for polygenic scores not shown in **(Figure 1A)** are provided in

345 **(Supplementary Figure 1)**.

346   Higher values in this mode were linked to increased nodal efficiency in the

347 temporal gyri, betweenness centrality in the supramarginal and post-central gyri, and

348 higher clustering coefficients in the temporal, parietal, central, and inferior frontal gyri

349 **(Figure 1B)**. Conversely, higher values in this mode were associated with lower

connectivity in the cingulate cortex, insula, and subcortical regions, along with reduced

nodal efficiency in the insula and subcortical regions. Regarding global brain network

measures, density (-0.613) and global efficiency (-0.421) showed negative loadings,

while the normalized average clustering coefficient (0.542) showed a positive loading.

Results from the multi-ethnic analysis were consistent with those observed in

the European-only analysis **(Supplementary Figure 2)**. When we controlled for

confounding effects using a simplified set of covariates, the results remained largely

unchanged. However, when the SCCA model trained on the training set was applied

to the test set, the p-value was marginally significant ($p = 0.0534$).

**Covariation between environmental factors and structural brain network
properties**

Among the five modes analyzed, only the first mode was statistically

significant and generalized to the hold-out test set (ENV-BNM mode 1, $p < 0.001$,

cov = 1.251, cor = 0.146, FDR corrected). The environmental component of this

mode predominantly captured variance related to socioeconomic status (SES)

(Figure 1C).

Notably, neighborhood-level SES variables, such as ADI median family

income and ADI education level, showed the strongest positive loadings. Similarly,

family-level SES variables, including household income, also showed positive

associations, though their loadings were slightly lower than those of neighborhood-

level SES measures. Variables reflecting neighborhood-level socioeconomic

deprivation, such as ADI poverty indices, showed strong negative loadings.

Additionally, prenatal substance exposures—such as tobacco exposure during

pregnancy—showed moderate negative associations with this component. These

results suggest that this component reflects a spectrum of socioeconomic status

from socioeconomic advantage to disadvantage.

On the brain network side **(Figure 1D)**, connection strength and nodal

efficiency in the temporal gyrus showed positive loadings, and connection strength,

betweenness centrality, and clustering coefficient in regions such as the cingulate

cortex and precuneus showed negative loadings. Regarding global brain network

measures, normalized global efficiency (0.360) showed positive loading.

These results were robust across different sets of covariates. When missing

values were handled by dropping incomplete cases rather than through imputation,

the environmental component remained consistent. The general loading patterns of

brain network properties were also similar; but, the overall magnitude of the loadings

was slightly lower than in the main analysis, possibly due to reduced statistical power

(Supplementary Figure 4).

Covariation between phenotypes and structural brain network properties

Among the five modes of covariation identified by SCCA, the first two modes

were statistically significant and generalized to the hold-out test sets (Pheno-BNM

mode 1: $p < 0.001$, cov = 1.421, r = 0.143; Pheno-BNM mode 2: $p < 0.001$, cov = 0.923,

r = 0.123, all p-values were FDR corrected).

The first mode captured covariation between brain network properties and an

integrated measure of cognitive ability and psychopathology in children **(Figure 1E)**.

Cognitive ability-related scores (e.g., NIH Toolbox scores) showed positive loadings,

while psychopathology-related scores (e.g., CBCL scores) showed negative

loadings. These loading patterns suggest that the first phenotype mode reflects

variation along the cognitive ability-psychopathology axis.

Brain network properties associated with this mode showed positive loadings

for nodal efficiency and betweenness centrality in the temporal and parietal cortices.

In contrast, connection strength in the insula, cingulate, precuneus, and subcortical

regions showed negative loadings. Additionally, nodal efficiency in the insula and

subcortical regions was negatively associated with this mode **(Figure 1F)**. No

significant global brain network measures were identified for this mode.

In phenotype mode 2 **(Figure 2)**, positive loadings were observed for

standing height, being born prematurely, having hearing or vision issues, and

experiencing obstetric complications. Although the uncertainty in loading estimation

is considerable, abnormal behavior (CBCL scores) also exhibited the highest positive

loading in this mode **(Supplementary Figure 7)**. On the brain network properties

side, lower connection strength and nodal efficiency were observed in the precentral

and postcentral gyri, superior frontal gyrus, and thalamus. Degree in the middle

temporal gyrus, superior frontal gyrus, and postcentral gyrus also tended to be lower.

Among global brain network metrics, normalized modularity (-0.195), global

efficiency (-0.343), and normalized global efficiency (-0.487) showed negative

loadings, while modularity (0.534) showed positive loading.

The findings remained stable across different covariate sets. When missing

values were handled by excluding incomplete cases instead of using imputation, the

mode 1 results remained consistent **(Supplementary Figure 6)**. However, in mode

2, unlike the imputed case, only height, weight, and BMI exhibited reliable positive

loadings, while CBCL, despite its high estimation uncertainty, showed a strong

negative loading **(Supplementary Figure 8).**

Shared Covariation Patterns in Brain Network Properties

Our SCCA identified certain brain network properties that showed similar

covariation patterns across genetic factors related to cognitive ability, socioeconomic

status, and phenotypes of cognitive ability-psychopathology. **(Figure 1G)** presents

the brain network properties that shared common loading patterns with the primary

SCCA modes for polygenic scores (PGS mode 1), environmental factors (ENV mode

1), and phenotypes (Pheno mode 1). Across these domains, brain network measures

such as nodal efficiency in the temporal gyrus consistently showed positive loadings,

while measures like connection strength in the posterior cingulate and subcortical

regions consistently showed negative loadings.

Mediation Analysis

The observed covariation patterns in brain network properties across genetic,

environmental, and phenotypic domains led us to hypothesize that structural brain

networks mediate two key relationships: (1) between genetic factors related to

cognitive ability and the cognitive ability-psychopathology phenotype, and (2)

between socioeconomic status and the cognitive ability-psychopathology phenotype.

To test these hypotheses, we conducted mediation analyses using SCCA-derived

summary scores from each domain.

Our analysis revealed significant mediation effects in both cases. Structural

brain network properties partially mediated the relationship between polygenic

scores for cognitive ability and the cognitive ability-psychopathology phenotype

(indirect effect = 0.023, p < 0.001; **Figure 3A**), as well as between socioeconomic

status and the cognitive ability-psychopathology phenotype (indirect effect = 0.015, p

< 0.001; **Figure 3B)**.

Discussion

This study investigated how structural brain network properties covary with

genetic, environmental, and phenotypic factors in 9–10-year-old children and

whether these properties mediate genetic and environmental effects on cognitive-

behavioral outcomes. By integrating polygenic scores (PGS), environmental

variables, and cognitive-behavioral traits using sparse canonical correlation analysis

(SCCA), we identified shared covariation patterns in brain network properties across

these domains. Previous studies have demonstrated that structural brain network

properties are heritable (Koenis et al., 2015; van den Heuvel et al., 2013), but the

specific genetic contributions to brain network organization in children remain

underexplored. Our findings address this gap by providing evidence that cognitive

ability-related genetic factors, socioeconomic status, and cognitive-

psychopathological phenotypes are key determinants of structural brain network

variations in preadolescents. Moreover, mediation analyses reveal that structural

brain network properties serve as intermediaries between genetic/environmental

influences and cognitive-psychopathological outcomes. This suggests that variations

in brain network organization may provide a mechanistic pathway through which

early-life genetic and environmental factors contribute to individual differences in

cognitive and mental health outcomes.

Our results reveal that structural brain network properties exhibit distinct yet

overlapping covariation patterns with genetic, environmental, and phenotypic factors.

Specifically, brain network measures such as nodal efficiency in the temporal and

parietal cortices were consistently associated with cognitive ability-related genetic

factors, higher socioeconomic status, and better cognitive performance. Conversely,

weaker connectivity in the posterior cingulate, insula, and subcortical regions was

commonly linked to genetic risk for lower cognitive ability, socioeconomic

disadvantage, and increased psychopathology. Consistent with prior studies (Alnaes

et al., 2020; Fernandez-Cabello et al., 2022; Modabbernia et al., 2021; Smith et al.,

2015), these findings suggest that children's brain network properties also covary

along spectrums of 'positive-negative' factors across genetic, environmental, and

phenotypic dimensions. Unlike previous studies that explored covariation between

brain features and only some of these domains, our study examined all three factors

simultaneously, identifying a common set of brain network properties that covary

across genetic, environmental, and phenotypic factors. These findings highlight the

potential of these brain networks as key substrates underlying cognitive and mental

health disparities, shaped by both genetic predisposition and environmental

influences during childhood.

Our findings align with established neurodevelopmental models, which

suggest that childhood and adolescence are characterized by a shift from

subcortical-driven to cortico-cortical-dominated network organization (Baker et al.,

2015; Langen et al., 2018; Menon, 2013; Sato et al., 2015). Specifically, favorable

genetic and phenotypic traits related to cognitive ability and SES were associated

with greater nodal efficiency and betweenness centrality in the temporal and parietal

cortices and reduced connectivity in the posterior cingulate, insula, and subcortical

regions. These patterns are consistent with established neurodevelopmental

trajectories, which involve a progressive strengthening of long-range cortico-cortical

connectivity (Hwang et al., 2013; Oldham & Fornito, 2019) and reduction in

subcortical connectivity as higher-order networks become more specialized (Baker et

al., 2015; Langen et al., 2018; Sato et al., 2015). Interestingly, the associations

between genetic, environmental, and phenotypic factors and brain network

organization mirror expected patterns of cortical-subcortical reorganization, wherein

cortico-cortical integration strengthens while subcortical connectivity decreases.

These findings suggest that genetic and environmental influences may shape the

timing or pace of this cortical-subcortical transition, potentially accelerating or

delaying neurodevelopmental trajectories and influencing individual differences in

cognitive and mental health outcomes (Heller et al., 2016).

Children with higher cognitive performance tend to have greater nodal

efficiency and betweenness centrality in the temporal, parietal, and superior frontal

regions—areas crucial for higher-order cognitive functions such as language,

semantic processing, abstract reasoning, and working memory (Binder et al., 2009;

Culham & Kanwisher, 2001; du Boisgueheneuc et al., 2006; Price, 2012; Visser et

al., 2012). This suggests that enhanced network integrity in these regions supports

cognitive ability by efficient communication across the brain. Consistent with this, our

results support the parieto-frontal integration theory (P-FIT) of intelligence (Basten et

al., 2015; Jung & Haier, 2007), which posits that effective information integration

across distributed networks, particularly involving the frontal, parietal, and temporal

regions, underlies intelligence. Notably, while previous studies have linked global

efficiency with cognitive performance (Bathelt, Gathercole, Butterfield, et al., 2018;

Kim et al., 2016; Ma et al., 2017), our findings—derived from a largest to date

sample—suggest that regional network efficiency, rather than global efficiency, is

more strongly associated with cognitive ability. This highlights the importance of

region-specific network topologies in understanding neurodevelopmental differences.

Although neighborhood-level socioeconomic status (SES) and family-level

SES are both associated with children's brain development, neighborhood-level SES

has a unique relationship with brain structure and functional networks, distinct from

that of family-level SES (Rakesh et al., 2022; Tooley et al., 2020). In our results, the

link between neighborhood-level SES and structural brain network properties was

more pronounced than associations with family-level SES.

Prenatal substance exposure, particularly to tobacco and marijuana,

exhibited moderate negative loadings within the SES-related brain network mode,

reinforcing prior findings that lower socioeconomic status is associated with

increased prenatal exposure to neurotoxic substances (Gu et al., 2024; Metz et al.,

2018; Mravcik et al., 2020). This association likely reflects a complex interplay

between socioeconomic adversity and prenatal environmental stressors, both of

which have been implicated in shaping neurodevelopmental trajectories (El Marroun

et al., 2016; Ross et al., 2015; Thompson et al., 2009). Given that SES-related

disparities in brain connectivity may stem from a combination of prenatal exposures,

postnatal environments, and genetic predispositions, the observed negative loading

may capture broader socioeconomic influences rather than a direct teratogenic effect

of prenatal substance exposure. However, disentangling these effects is challenging,

as the precise mechanisms through which prenatal exposures contribute to structural

brain network alterations remain unclear. Future studies leveraging genome-

environment interaction analyses, longitudinal neuroimaging, and causal inference

approaches (e.g., Mendelian randomization) are essential to elucidate how prenatal

risk factors, SES, and genetic predispositions collectively shape neurodevelopmental

outcomes.

Phenotype mode 2 presents a complex pattern, linking standing height,

perinatal risk factors, and sensorimotor brain networks. The lower network efficiency

in the precentral and postcentral gyri, superior frontal gyrus, and thalamus suggests

a potential connection to early neurodevelopmental processes, particularly those

involved in motor and sensory integration. Given the variability in loading estimates,

further research is needed to determine whether these associations reflect specific

neurodevelopmental mechanisms or statistical artifacts.

Several limitations should be acknowledged. First, the cross-sectional design

precludes causal inferences, necessitating longitudinal research to validate the

mediating role of brain networks in shaping developmental trajectories of cognitive

and mental health. Second, gene-by-environment (G×E) interactions were not

explicitly modeled, limiting our ability to assess whether socioeconomic status (SES)

moderates genetic influences on brain connectivity. Future research should integrate

G×E interaction analyses to better understand how genetic predispositions interact

with environmental contexts in shaping neurodevelopment. Finally, although sparse

canonical correlation analysis (SCCA) provided a powerful multivariate approach, it

assumes linear relationships between brain networks and genetic/environmental

factors, which may oversimplify complex neurodevelopmental processes. Future

studies should consider non-linear modeling approaches and more sophisticated

causal inference methods (e.g., Mendelian randomization, structural equation

modeling) to capture the intricate interplay of genes, environments, and brain

development.

Addressing these gaps will enhance the robustness of future research,

ultimately contributing to a more comprehensive understanding of how genetic and

environmental factors shape brain development. These insights may inform early

intervention strategies aimed at mitigating neurodevelopmental disparities, such as

targeted cognitive training, socioeconomic policy reforms, or school-based

enrichment programs that support children from disadvantaged backgrounds.

Ethics Statement

This study analyzed data from the Adolescent Brain Cognitive Development

(ABCD) study. All participants provided informed assent, and their parents or legal

guardians provided informed consent before participating in the ABCD study.

Declaration of Competing Interest

The authors declare no competing financial interests.

Acknowledgments

This work was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MSIT) (No. 2021R1C1C1006503, RS-2023-00266787, RS-2023-00265406, RS-2024-00421268), by Creative-Pioneering Researchers Program through Seoul National University(No. 200-20240057), by Semi-Supervised Learning Research Grant by SAMSUNG(No.A0426-20220118), by Identify the network of brain preparation steps for concentration Research Grant by LooxidLabs(No.339-20230001), by Institute of Information & communications Technology Planning & Evaluation (IITP) grant funded by the Korea government(MSIT) [NO.RS-2021-II211343, Artificial Intelligence Graduate School Program (Seoul National University)] by the MSIT(Ministry of Science, ICT), Korea, under the Global Research Support Program in the Digital Field program(RS-2024-00421268) supervised by the IITP(Institute for Information & Communications Technology Planning & Evaluation), by the National Supercomputing Center with supercomputing resources including technical support(KSC-2023-CRE-0568) and by the Ministry of Education of the Republic of Korea and the National Research Foundation of Korea (NRF-2021S1A3A2A02090597), and by Artificial intelligence industrial convergence cluster development project funded by the Ministry of Science and ICT(MSIT, Korea) & Gwangju Metropolitan City.

Data Availability

All original data are publicly available from the NDA
(https://nda.nih.gov/abcd/). For rapid replication, we provide synthetic data, whose
distributions were matched to the original data, generated using conditional GAN for
tabular data (Xu et al., 2019) ([https://github.com/Transconnectome/ABCD-brain-](https://github.com/Transconnectome/ABCD-brain-network-SCCA)
[network-SCCA](https://github.com/Transconnectome/ABCD-brain-network-SCCA)).

Code Availability

Code is available from here: [https://github.com/Transconnectome/ABCD-](https://github.com/Transconnectome/ABCD-brain-network-SCCA)
[brain-network-SCCA](https://github.com/Transconnectome/ABCD-brain-network-SCCA).

Author contributions

- Jungwoo Seo, Conceptualization, Data curation, Formal analysis, Investigation,
  Methodology, Project administration, Software, Validation, Visualization, Writing –
  original draft, Writing – review and editing
- Eunji Lee, Data curation, Writing – original draft, Writing – review and editing
- Bo-Gyeom Kim, Data curation, Writing – review and editing
- Gakyung Kim, Data curation, Writing – original draft, Writing – review and editing
- Yoonjung Yoonie Joo, Data curation, Writing – review and editing
- Jiook Cha, Conceptualization, Supervision, Project administration, Funding
  acquisition, Resources, Data curation, Writing – original draft, Writing – review and
  editing.

Tables

Table 1. Demographic information of the main analysis participants.

		PGS – BNM (n=5,784)		ENV – BNM (n=10,343)		Pheno – BNM (n=10,343)	
		train	test	train	test	train	test
N	total	4,626	1,157	8,274	2,069	8,274	2,069
Sex	Male	2,453	627	4,343	1,065	4,343	1,065
	Female	2,172	529	3,929	1,003	3,929	1,003
	Other	1	1	2	1	2	1
Race	White	3,225	802	4,350	1,128	4,350	1,128
	Black	29	7	1,194	295	1,194	295
	Hispanic	902	223	1,686	408	1,686	408
	Asian	8	0	173	37	173	37
	Other	462	125	871	201	871	201

Figure 1 Loading patterns of principal mode of sparse canonical correlation analysis. The

SCCA results for the first modes of PGS-BNM, ENV-BNM, and Pheno-BNM. Only variables with 95%

confidence intervals, estimated from 5,000 bootstrap samples, that do not cross zero are shown.

Results for other variables can be found in supplementary figures. (A) The loadings of significant PGS

variables in the PGS-BNM mode 1. The error bars represent the 95% confidence interval of the

loading, estimated from the 5,000 bootstrap samples. The color of each bar represents the category

to which the variable belongs. (B) The loadings of significant nodal brain network measures in the

PGS-BNM mode 1. The loading patterns were visualized with R-package '*ggseg*' (Mowinckel & Vidal-

Pineiro, 2020). (C) The loadings of significant environmental variables in the ENV-BNM mode 1. (D)

The loadings of significant nodal brain network measures in the ENV-BNM mode 1. (E) The loadings

of significant phenotype variables in the Pheno-BNM mode 1. (F) The loadings of significant nodal

brain network measures in the Pheno-BNM mode 1. (G) Brain network measures that their loadings

are commonly significant in PGS-BNM mode 1, ENV-BNM mode1, and Pheno-BNM mode1.

Figure 2 Loading patterns of second mode of Pheno-BNM sparse canonical correlation analysis. (A) The loadings of significant phenotype variables in the Pheno-BNM SCCA mode 2. (B) The loadings of significant nodal brain network measures in the Pheno-BNM SCCA mode 2.

Figure 3 Results of mediation analysis investigating the gene-brain network-phenotype and

environment-brain network-phenotype pathways. Mediation analysis was conducted using

canonical variates derived from SCCA for polygenic scores, brain network measures, phenotypes,

and environmental factors. (A) Brain network properties mediating the relationship between polygenic

scores (reflecting cognitive ability) and phenotype scores (reflecting cognitive ability-

psychopathological traits). (B) Brain network properties mediating the relationship between

environmental factors (reflecting socioeconomic status) and phenotype scores (cognitive ability-

psychopathological traits).

References

Akiyama, M., Okada, Y., Kanai, M., Takahashi, A., Momozawa, Y., Ikeda, M., Iwata, N., Ikegawa, S., Hirata, M., Matsuda, K., Iwasaki, M., Yamaji, T., Sawada, N., Hachiya, T., Tanno, K., Shimizu, A., Hozawa, A., Minegishi, N., Tsugane, S., . . . Kamatani, Y. (2017). Genome-wide association study identifies 112 new loci for body mass index in the Japanese population. *Nat Genet*, *49*(10), 1458-1467. https://doi.org/10.1038/ng.3951

Alexander-Bloch, A. F., Gogtay, N., Meunier, D., Birn, R., Clasen, L., Lalonde, F., Lenroot, R., Giedd, J., & Bullmore, E. T. (2010). Disrupted modularity and local connectivity of brain functional networks in childhood-onset schizophrenia. *Front Syst Neurosci*, *4*, 147. https://doi.org/10.3389/fnsys.2010.00147

Alnaes, D., Kaufmann, T., Marquand, A. F., Smith, S. M., & Westlye, L. T. (2020). Patterns of sociocognitive stratification and perinatal risk in the child brain. *Proc Natl Acad Sci U S A*, *117*(22), 12419-12427. https://doi.org/10.1073/pnas.2001517117

Bai, S., Liu, W., & Guan, Y. (2021). The Visuospatial and Sensorimotor Functions of Posterior Parietal Cortex in Drawing Tasks: A Review. *Front Aging Neurosci*, *13*, 717002. https://doi.org/10.3389/fnagi.2021.717002

Baker, S. T., Lubman, D. I., Yucel, M., Allen, N. B., Whittle, S., Fulcher, B. D., Zalesky, A., & Fornito, A. (2015). Developmental Changes in Brain Network Hub Connectivity in Late Adolescence. *J Neurosci*, *35*(24), 9078-9087. https://doi.org/10.1523/JNEUROSCI.5043-14.2015

Basten, U., Hilger, K., & Fiebach, C. J. (2015). Where smart brains are different: A quantitative meta-analysis of functional and structural brain imaging studies on intelligence. *Intelligence*, *51*, 10-27.

Bathelt, J., Gathercole, S. E., Butterfield, S., team, C., & Astle, D. E. (2018). Children's academic attainment is linked to the global organization of the white matter connectome. *Dev Sci*, *21*(5), e12662. https://doi.org/10.1111/desc.12662

Bathelt, J., Gathercole, S. E., Johnson, A., & Astle, D. E. (2018). Differences in brain morphology and working memory capacity across childhood. *Dev Sci*, *21*(3), e12579. https://doi.org/10.1111/desc.12579

Bethlehem, R. A. I., Seidlitz, J., White, S. R., Vogel, J. W., Anderson, K. M., Adamson, C., Adler, S., Alexopoulos, G. S., Anagnostou, E., Areces-Gonzalez, A., Astle, D. E., Auyeung, B., Ayub, M., Bae, J., Ball, G., Baron-Cohen, S., Beare, R., Bedford, S. A., Benegal, V., . . . Alexander-Bloch, A. F. (2022). Brain charts for the human lifespan. *Nature*, *604*(7906), 525-533. https://doi.org/10.1038/s41586-022-04554-y

Binder, J. R., Desai, R. H., Graves, W. W., & Conant, L. L. (2009). Where is the semantic system? A critical review and meta-analysis of 120 functional neuroimaging studies. *Cereb Cortex*, *19*(12), 2767-2796. https://doi.org/10.1093/cercor/bhp055

Bipolar, D., Schizophrenia Working Group of the Psychiatric Genomics Consortium. Electronic address, d. r. v. e., Bipolar, D., & Schizophrenia Working Group of the Psychiatric Genomics, C. (2018). Genomic Dissection of Bipolar Disorder and Schizophrenia, Including 28 Subphenotypes. *Cell*, *173*(7), 1705-1715 e1716. https://doi.org/10.1016/j.cell.2018.05.046

Buckner, R. L., Andrews-Hanna, J. R., & Schacter, D. L. (2008). The brain's default network: anatomy, function, and relevance to disease. *Ann N Y Acad Sci*, *1124*, 1-38. https://doi.org/10.1196/annals.1440.011

Bunge, S. A., & Wright, S. B. (2007). Neurodevelopmental changes in working memory and cognitive control. *Curr Opin Neurobiol*, *17*(2), 243-250. https://doi.org/10.1016/j.conb.2007.02.005

Calamante, F., Tournier, J. D., Jackson, G. D., & Connelly, A. (2010). Track-density imaging (TDI): super-resolution white matter imaging using whole-brain track-density

mapping. *Neuroimage*, *53*(4), 1233-1243.
https://doi.org/10.1016/j.neuroimage.2010.07.024
Casey, B. J., Cannonier, T., Conley, M. I., Cohen, A. O., Barch, D. M., Heitzeg, M. M.,
Soules, M. E., Teslovich, T., Dellarco, D. V., Garavan, H., Orr, C. A., Wager, T. D.,
Banich, M. T., Speer, N. K., Sutherland, M. T., Riedel, M. C., Dick, A. S., Bjork, J. M.,
Thomas, K. M., . . . Workgroup, A. I. A. (2018). The Adolescent Brain Cognitive
Development (ABCD) study: Imaging acquisition across 21 sites. *Dev Cogn*
Neurosci, *32*, 43-54. https://doi.org/10.1016/j.dcn.2018.03.001
Collin, G., Scholtens, L. H., Kahn, R. S., Hillegers, M. H. J., & van den Heuvel, M. P. (2017).
Affected Anatomical Rich Club and Structural-Functional Coupling in Young Offspring
of Schizophrenia and Bipolar Disorder Patients. *Biol Psychiatry*, *82*(10), 746-755.
https://doi.org/10.1016/j.biopsych.2017.06.013
Conomos, M. P., Miller, M. B., & Thornton, T. A. (2015). Robust inference of population
structure for ancestry prediction and correction of stratification in the presence of
relatedness. *Genet Epidemiol*, *39*(4), 276-293. https://doi.org/10.1002/gepi.21896
Conomos, M. P., Reiner, A. P., Weir, B. S., & Thornton, T. A. (2016). Model-free Estimation of
Recent Genetic Relatedness. *Am J Hum Genet*, *98*(1), 127-148.
https://doi.org/10.1016/j.ajhg.2015.11.022
Cross-Disorder Group of the Psychiatric Genomics, C. (2013). Identification of risk loci with
shared effects on five major psychiatric disorders: a genome-wide analysis. *Lancet*,
381(9875), 1371-1379. https://doi.org/10.1016/S0140-6736(12)62129-1
Culham, J. C., & Kanwisher, N. G. (2001). Neuroimaging of cognitive functions in human
parietal cortex. *Curr Opin Neurobiol*, *11*(2), 157-163. https://doi.org/10.1016/s0959-
4388(00)00191-4
Das, S., Forer, L., Schonherr, S., Sidore, C., Locke, A. E., Kwong, A., Vrieze, S. I., Chew, E.
Y., Levy, S., McGue, M., Schlessinger, D., Stambolian, D., Loh, P. R., Iacono, W. G.,
Swaroop, A., Scott, L. J., Cucca, F., Kronenberg, F., Boehnke, M., . . . Fuchsberger,
C. (2016). Next-generation genotype imputation service and methods. *Nat Genet*,
48(10), 1284-1287. https://doi.org/10.1038/ng.3656
Demontis, D., Walters, R. K., Martin, J., Mattheisen, M., Als, T. D., Agerbo, E., Baldursson,
G., Belliveau, R., Bybjerg-Grauholm, J., Baekvad-Hansen, M., Cerrato, F., Chambert,
K., Churchhouse, C., Dumont, A., Eriksson, N., Gandal, M., Goldstein, J. I., Grasby,
K. L., Grove, J., . . . Neale, B. M. (2019). Discovery of the first genome-wide
significant risk loci for attention deficit/hyperactivity disorder. *Nat Genet*, *51*(1), 63-75.
https://doi.org/10.1038/s41588-018-0269-7
Deschamps, I., Baum, S. R., & Gracco, V. L. (2014). On the role of the supramarginal gyrus
in phonological processing and verbal working memory: evidence from rTMS studies.
Neuropsychologia, *53*, 39-46. https://doi.org/10.1016/j.neuropsychologia.2013.10.015
Desikan, R. S., Segonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., Buckner,
R. L., Dale, A. M., Maguire, R. P., Hyman, B. T., Albert, M. S., & Killiany, R. J. (2006).
An automated labeling system for subdividing the human cerebral cortex on MRI
scans into gyral based regions of interest. *Neuroimage*, *31*(3), 968-980.
https://doi.org/10.1016/j.neuroimage.2006.01.021
du Boisgueheneuc, F., Levy, R., Volle, E., Seassau, M., Duffau, H., Kinkingnehun, S.,
Samson, Y., Zhang, S., & Dubois, B. (2006). Functions of the left superior frontal
gyrus in humans: a lesion study. *Brain*, *129*(Pt 12), 3315-3328.
https://doi.org/10.1093/brain/awl244
El Marroun, H., Tiemeier, H., Franken, I. H., Jaddoe, V. W., van der Lugt, A., Verhulst, F. C.,
Lahey, B. B., & White, T. (2016). Prenatal Cannabis and Tobacco Exposure in
Relation to Brain Morphology: A Prospective Neuroimaging Study in Young Children.
Biol Psychiatry, *79*(12), 971-979. https://doi.org/10.1016/j.biopsych.2015.08.024
Fernandez-Cabello, S., Alnaes, D., van der Meer, D., Dahl, A., Holm, M., Kjelkenes, R.,
Maximov, II, Norbom, L. B., Pedersen, M. L., Voldsbekk, I., Andreassen, O. A., &
Westlye, L. T. (2022). Associations between brain imaging and polygenic scores of

mental health and educational attainment in children aged 9-11. *Neuroimage*, *263*, 119611. https://doi.org/10.1016/j.neuroimage.2022.119611

Fornito, A., Zalesky, A., & Breakspear, M. (2015). The connectomics of brain disorders. *Nat Rev Neurosci*, *16*(3), 159-172. https://doi.org/10.1038/nrn3901

Ge, T., Chen, C. Y., Ni, Y., Feng, Y. A., & Smoller, J. W. (2019). Polygenic prediction via Bayesian regression and continuous shrinkage priors. *Nat Commun*, *10*(1), 1776. https://doi.org/10.1038/s41467-019-09718-5

Genomes Project, C., Auton, A., Brooks, L. D., Durbin, R. M., Garrison, E. P., Kang, H. M., Korbel, J. O., Marchini, J. L., McCarthy, S., McVean, G. A., & Abecasis, G. R. (2015). A global reference for human genetic variation. *Nature*, *526*(7571), 68-74. https://doi.org/10.1038/nature15393

Grove, J., Ripke, S., Als, T. D., Mattheisen, M., Walters, R. K., Won, H., Pallesen, J., Agerbo, E., Andreassen, O. A., Anney, R., Awashti, S., Belliveau, R., Bettella, F., Buxbaum, J. D., Bybjerg-Grauholm, J., Baekvad-Hansen, M., Cerrato, F., Chambert, K., Christensen, J. H., . . . Borglum, A. D. (2019). Identification of common genetic risk variants for autism spectrum disorder. *Nat Genet*, *51*(3), 431-444. https://doi.org/10.1038/s41588-019-0344-8

Gu, Z., Barch, D. M., & Luo, Q. (2024). Prenatal substance exposure and child health: Understanding the role of environmental factors, genetics, and brain development. *PNAS Nexus*, *3*(1), pgae003. https://doi.org/10.1093/pnasnexus/pgae003

Heller, A. S., Cohen, A. O., Dreyfuss, M. F., & Casey, B. J. (2016). Changes in cortico-subcortical and subcortico-subcortical connectivity impact cognitive control to emotional cues across development. *Soc Cogn Affect Neurosci*, *11*(12), 1910-1918. https://doi.org/10.1093/scan/nsw097

Hotelling, H. (1936). Relations between two sets of variates. *Biometrika*, *28*, 321-377. https://doi.org/10.1093/biomet/28.3-4.321

Howard, D. M., Adams, M. J., Clarke, T. K., Hafferty, J. D., Gibson, J., Shirali, M., Coleman, J. R. I., Hagenaars, S. P., Ward, J., Wigmore, E. M., Alloza, C., Shen, X., Barbu, M. C., Xu, E. Y., Whalley, H. C., Marioni, R. E., Porteous, D. J., Davies, G., Deary, I. J., . . . McIntosh, A. M. (2019). Genome-wide meta-analysis of depression identifies 102 independent variants and highlights the importance of the prefrontal brain regions. *Nat Neurosci*, *22*(3), 343-352. https://doi.org/10.1038/s41593-018-0326-7

Hwang, K., Hallquist, M. N., & Luna, B. (2013). The development of hub architecture in the human functional brain network. *Cereb Cortex*, *23*(10), 2380-2393. https://doi.org/10.1093/cercor/bhs227

International Obsessive Compulsive Disorder Foundation Genetics, C., & Studies, O. C. D. C. G. A. (2018). Revealing the complex genetic architecture of obsessive-compulsive disorder using meta-analysis. *Mol Psychiatry*, *23*(5), 1181-1188. https://doi.org/10.1038/mp.2017.154

Jansen, P. R., Watanabe, K., Stringer, S., Skene, N., Bryois, J., Hammerschlag, A. R., de Leeuw, C. A., Benjamins, J. S., Munoz-Manchado, A. B., Nagel, M., Savage, J. E., Tiemeier, H., White, T., andMe Research, T., Tung, J. Y., Hinds, D. A., Vacic, V., Wang, X., Sullivan, P. F., . . . Posthuma, D. (2019). Genome-wide analysis of insomnia in 1,331,010 individuals identifies new risk loci and functional pathways. *Nat Genet*, *51*(3), 394-403. https://doi.org/10.1038/s41588-018-0333-3

Jernigan, T. L., Brown, S. A., & Dowling, G. J. (2018). The Adolescent Brain Cognitive Development Study. *J Res Adolesc*, *28*(1), 154-156. https://doi.org/10.1111/jora.12374

Jeurissen, B., Descoteaux, M., Mori, S., & Leemans, A. (2019). Diffusion MRI fiber tractography of the brain. *NMR Biomed*, *32*(4), e3785. https://doi.org/10.1002/nbm.3785

Joo, Y. Y., Lee, E., Kim, B. G., Kim, G., Seo, J., & Cha, J. (2024). Polygenic architecture of brain structure and function, behaviors, and psychopathologies in children. *bioRxiv*. https://doi.org/10.1101/2024.05.22.595444

Jung, R. E., & Haier, R. J. (2007). The Parieto-Frontal Integration Theory (P-FIT) of
intelligence: converging neuroimaging evidence. *Behav Brain Sci*, *30*(2), 135-154;
discussion 154-187. https://doi.org/10.1017/S0140525X07001185
Karlsson Linner, R., Biroli, P., Kong, E., Meddens, S. F. W., Wedow, R., Fontana, M. A.,
Lebreton, M., Tino, S. P., Abdellaoui, A., Hammerschlag, A. R., Nivard, M. G., Okbay,
A., Rietveld, C. A., Timshel, P. N., Trzaskowski, M., Vlaming, R., Zund, C. L., Bao, Y.,
Buzdugan, L., . . . Beauchamp, J. P. (2019). Genome-wide association analyses of
risk tolerance and risky behaviors in over 1 million individuals identify hundreds of
loci and shared genetic influences. *Nat Genet*, *51*(2), 245-257.
https://doi.org/10.1038/s41588-018-0309-3
Kim, D. J., Davis, E. P., Sandman, C. A., Glynn, L., Sporns, O., O'Donnell, B. F., & Hetrick,
W. P. (2019). Childhood poverty and the organization of structural brain connectome.
Neuroimage, *184*, 409-416. https://doi.org/10.1016/j.neuroimage.2018.09.041
Kim, D. J., Davis, E. P., Sandman, C. A., Sporns, O., O'Donnell, B. F., Buss, C., & Hetrick, W.
P. (2016). Children's intellectual ability is associated with structural network integrity.
Neuroimage, *124*(Pt A), 550-556. https://doi.org/10.1016/j.neuroimage.2015.09.012
Kim, K., Joo, Y. Y., Ahn, G., Wang, H. H., Moon, S. Y., Kim, H., Ahn, W. Y., & Cha, J. (2022).
The sexual brain, genes, and cognition: A machine-predicted brain sex score
explains individual differences in cognitive intelligence and genetic influence in young
children. *Hum Brain Mapp*, *43*(12), 3857-3872. https://doi.org/10.1002/hbm.25888
Koenis, M. M., Brouwer, R. M., van den Heuvel, M. P., Mandl, R. C., van Soelen, I. L., Kahn,
R. S., Boomsma, D. I., & Hulshoff Pol, H. E. (2015). Development of the brain's
structural network efficiency in early adolescence: A longitudinal DTI twin study. *Hum
Brain Mapp*, *36*(12), 4938-4953. https://doi.org/10.1002/hbm.22988
Lam, M., Chen, C. Y., Li, Z., Martin, A. R., Bryois, J., Ma, X., Gaspar, H., Ikeda, M.,
Benyamin, B., Brown, B. C., Liu, R., Zhou, W., Guan, L., Kamatani, Y., Kim, S. W.,
Kubo, M., Kusumawardhani, A., Liu, C. M., Ma, H., . . . Huang, H. (2019).
Comparative genetic architectures of schizophrenia in East Asian and European
populations. *Nat Genet*, *51*(12), 1670-1678. https://doi.org/10.1038/s41588-019-
0512-x
Langen, C. D., Muetzel, R., Blanken, L., van der Lugt, A., Tiemeier, H., Verhulst, F., Niessen,
W. J., & White, T. (2018). Differential patterns of age-related cortical and subcortical
functional connectivity in 6-to-10 year old children: A connectome-wide association
study. *Brain Behav*, *8*(8), e01031. https://doi.org/10.1002/brb3.1031
Lee, J. J., Wedow, R., Okbay, A., Kong, E., Maghzian, O., Zacher, M., Nguyen-Viet, T. A.,
Bowers, P., Sidorenko, J., Karlsson Linner, R., Fontana, M. A., Kundu, T., Lee, C., Li,
H., Li, R., Royer, R., Timshel, P. N., Walters, R. K., Willoughby, E. A., . . . Cesarini, D.
(2018). Gene discovery and polygenic prediction from a genome-wide association
study of educational attainment in 1.1 million individuals. *Nat Genet*, *50*(8), 1112-
1121. https://doi.org/10.1038/s41588-018-0147-3
Lehericy, S., Bardinet, E., Tremblay, L., Van de Moortele, P. F., Pochon, J. B., Dormont, D.,
Kim, D. S., Yelnik, J., & Ugurbil, K. (2006). Motor control in basal ganglia circuits
using fMRI and brain atlas approaches. *Cereb Cortex*, *16*(2), 149-161.
https://doi.org/10.1093/cercor/bhi089
Locke, A. E., Kahali, B., Berndt, S. I., Justice, A. E., Pers, T. H., Day, F. R., Powell, C.,
Vedantam, S., Buchkovich, M. L., Yang, J., Croteau-Chonka, D. C., Esko, T., Fall, T.,
Ferreira, T., Gustafsson, S., Kutalik, Z., Luan, J., Magi, R., Randall, J. C., . . .
Speliotes, E. K. (2015). Genetic studies of body mass index yield new insights for
obesity biology. *Nature*, *518*(7538), 197-206. https://doi.org/10.1038/nature14177
Loh, P. R., Danecek, P., Palamara, P. F., Fuchsberger, C., Y, A. R., H, K. F., Schoenherr, S.,
Forer, L., McCarthy, S., Abecasis, G. R., Durbin, R., & A, L. P. (2016). Reference-
based phasing using the Haplotype Reference Consortium panel. *Nat Genet*, *48*(11),
1443-1448. https://doi.org/10.1038/ng.3679
Luna, B., Padmanabhan, A., & O'Hearn, K. (2010). What has fMRI told us about the

development of cognitive control through adolescence? *Brain Cogn*, *72*(1), 101-113. https://doi.org/10.1016/j.bandc.2009.08.005

Ma, J., Kang, H. J., Kim, J. Y., Jeong, H. S., Im, J. J., Namgung, E., Kim, M. J., Lee, S., Kim, T. D., Oh, J. K., Chung, Y. A., Lyoo, I. K., Lim, S. M., & Yoon, S. (2017). Network attributes underlying intellectual giftedness in the developing brain. *Sci Rep*, *7*(1), 11321. https://doi.org/10.1038/s41598-017-11593-3

Menon, V. (2013). Developmental pathways to functional brain networks: emerging principles. *Trends Cogn Sci*, *17*(12), 627-640. https://doi.org/10.1016/j.tics.2013.09.015

Metz, V. E., Brown, Q. L., Martins, S. S., & Palamar, J. J. (2018). Characteristics of drug use among pregnant women in the United States: Opioid and non-opioid illegal drug use. *Drug Alcohol Depend*, *183*, 261-266. https://doi.org/10.1016/j.drugalcdep.2017.11.010

Misic, B., Betzel, R. F., de Reus, M. A., van den Heuvel, M. P., Berman, M. G., McIntosh, A. R., & Sporns, O. (2016). Network-Level Structure-Function Relationships in Human Neocortex. *Cereb Cortex*, *26*(7), 3285-3296. https://doi.org/10.1093/cercor/bhw089

Modabbernia, A., Janiri, D., Doucet, G. E., Reichenberg, A., & Frangou, S. (2021). Multivariate Patterns of Brain-Behavior-Environment Associations in the Adolescent Brain and Cognitive Development Study. *Biol Psychiatry*, *89*(5), 510-520. https://doi.org/10.1016/j.biopsych.2020.08.014

Mowinckel, A. M., & Vidal-Pineiro, D. (2020). Visualization of Brain Statistics With R Packages ggseg and ggseg3d. *Advances in Methods and Practices in Psychological Science*, *3*(4), 466-483. https://doi.org/10.1177/2515245920928009

Mravcik, V., Nechanska, B., Gabrhelik, R., Handal, M., Mahic, M., & Skurtveit, S. (2020). Socioeconomic characteristics of women with substance use disorder during pregnancy and neonatal outcomes in their newborns: A national registry study from the Czech Republic. *Drug Alcohol Depend*, *209*, 107933. https://doi.org/10.1016/j.drugalcdep.2020.107933

Nagel, M., Jansen, P. R., Stringer, S., Watanabe, K., de Leeuw, C. A., Bryois, J., Savage, J. E., Hammerschlag, A. R., Skene, N. G., Munoz-Manchado, A. B., andMe Research, T., White, T., Tiemeier, H., Linnarsson, S., Hjerling-Leffler, J., Polderman, T. J. C., Sullivan, P. F., van der Sluis, S., & Posthuma, D. (2018). Meta-analysis of genome-wide association studies for neuroticism in 449,484 individuals identifies novel genetic loci and pathways. *Nat Genet*, *50*(7), 920-927. https://doi.org/10.1038/s41588-018-0151-7

Namkung, H., Kim, S. H., & Sawa, A. (2017). The Insula: An Underestimated Brain Area in Clinical Neuroscience, Psychiatry, and Neurology. *Trends Neurosci*, *40*(4), 200-207. https://doi.org/10.1016/j.tins.2017.02.002

Nievergelt, C. M., Maihofer, A. X., Klengel, T., Atkinson, E. G., Chen, C. Y., Choi, K. W., Coleman, J. R. I., Dalvie, S., Duncan, L. E., Gelernter, J., Levey, D. F., Logue, M. W., Polimanti, R., Provost, A. C., Ratanatharathorn, A., Stein, M. B., Torres, K., Aiello, A. E., Almli, L. M., . . . Koenen, K. C. (2019). International meta-analysis of PTSD genome-wide association studies identifies sex- and ancestry-specific genetic risk loci. *Nat Commun*, *10*(1), 4558. https://doi.org/10.1038/s41467-019-12576-w

Okbay, A., Baselmans, B. M., De Neve, J. E., Turley, P., Nivard, M. G., Fontana, M. A., Meddens, S. F., Linner, R. K., Rietveld, C. A., Derringer, J., Gratten, J., Lee, J. J., Liu, J. Z., de Vlaming, R., Ahluwalia, T. S., Buchwald, J., Cavadino, A., Frazier-Wood, A. C., Furlotte, N. A., . . . Cesarini, D. (2016). Genetic variants associated with subjective well-being, depressive symptoms, and neuroticism identified through genome-wide analyses. *Nat Genet*, *48*(6), 624-633. https://doi.org/10.1038/ng.3552

Oldham, S., & Fornito, A. (2019). The development of brain network hubs. *Dev Cogn Neurosci*, *36*, 100607. https://doi.org/10.1016/j.dcn.2018.12.005

Otowa, T., Hek, K., Lee, M., Byrne, E. M., Mirza, S. S., Nivard, M. G., Bigdeli, T., Aggen, S. H., Adkins, D., Wolen, A., Fanous, A., Keller, M. C., Castelao, E., Kutalik, Z., der

Auwera, S. V., Homuth, G., Nauck, M., Teumer, A., Milaneschi, Y., . . . Hettema, J. M. (2016). Meta-analysis of genome-wide association studies of anxiety disorders. *Mol Psychiatry*, *21*(10), 1485. https://doi.org/10.1038/mp.2016.11

Pasman, J. A., Verweij, K. J. H., Gerring, Z., Stringer, S., Sanchez-Roige, S., Treur, J. L., Abdellaoui, A., Nivard, M. G., Baselmans, B. M. L., Ong, J. S., Ip, H. F., van der Zee, M. D., Bartels, M., Day, F. R., Fontanillas, P., Elson, S. L., andMe Research, T., de Wit, H., Davis, L. K., . . . Vink, J. M. (2019). Author Correction: GWAS of lifetime cannabis use reveals new risk loci, genetic overlap with psychiatric traits, and a causal effect of schizophrenia liability. *Nat Neurosci*, *22*(7), 1196. https://doi.org/10.1038/s41593-019-0402-7

Paus, T., Keshavan, M., & Giedd, J. N. (2008). Why do many psychiatric disorders emerge during adolescence? *Nat Rev Neurosci*, *9*(12), 947-957. https://doi.org/10.1038/nrn2513

Price, C. J. (2012). A review and synthesis of the first 20 years of PET and fMRI studies of heard speech, spoken language and reading. *Neuroimage*, *62*(2), 816-847. https://doi.org/10.1016/j.neuroimage.2012.04.062

Rakesh, D., Zalesky, A., & Whittle, S. (2022). Assessment of Parent Income and Education, Neighborhood Disadvantage, and Child Brain Structure. *JAMA Netw Open*, *5*(8), e2226208. https://doi.org/10.1001/jamanetworkopen.2022.26208

Ross, E. J., Graham, D. L., Money, K. M., & Stanwood, G. D. (2015). Developmental consequences of fetal exposure to drugs: what we know and what we still must learn. *Neuropsychopharmacology*, *40*(1), 61-87. https://doi.org/10.1038/npp.2014.147

Rubinov, M., & Sporns, O. (2010). Complex network measures of brain connectivity: uses and interpretations. *Neuroimage*, *52*(3), 1059-1069. https://doi.org/10.1016/j.neuroimage.2009.10.003

Rudie, J. D., Brown, J. A., Beck-Pancer, D., Hernandez, L. M., Dennis, E. L., Thompson, P. M., Bookheimer, S. Y., & Dapretto, M. (2012). Altered functional and structural brain network organization in autism. *Neuroimage Clin*, *2*, 79-94. https://doi.org/10.1016/j.nicl.2012.11.006

Sato, J. R., Salum, G. A., Gadelha, A., Vieira, G., Zugman, A., Picon, F. A., Pan, P. M., Hoexter, M. Q., Anes, M., Moura, L. M., Del'Aquilla, M. A., Crossley, N., Amaro Junior, E., McGuire, P., Lacerda, A. L., Rohde, L. A., Miguel, E. C., Jackowski, A. P., & Bressan, R. A. (2015). Decreased centrality of subcortical regions during the transition to adolescence: a functional connectivity study. *Neuroimage*, *104*, 44-51. https://doi.org/10.1016/j.neuroimage.2014.09.063

Savage, J. E., Jansen, P. R., Stringer, S., Watanabe, K., Bryois, J., de Leeuw, C. A., Nagel, M., Awasthi, S., Barr, P. B., Coleman, J. R. I., Grasby, K. L., Hammerschlag, A. R., Kaminski, J. A., Karlsson, R., Krapohl, E., Lam, M., Nygaard, M., Reynolds, C. A., Trampush, J. W., . . . Posthuma, D. (2018). Genome-wide association meta-analysis in 269,867 individuals identifies new genetic and functional links to intelligence. *Nat Genet*, *50*(7), 912-919. https://doi.org/10.1038/s41588-018-0152-6

Shen, H., Gelaye, B., Huang, H., Rondon, M. B., Sanchez, S., & Duncan, L. E. (2020). Polygenic prediction and GWAS of depression, PTSD, and suicidal ideation/self-harm in a Peruvian cohort. *Neuropsychopharmacology*, *45*(10), 1595-1602. https://doi.org/10.1038/s41386-020-0603-5

Sherman, S. M., & Guillery, R. W. (2002). The role of the thalamus in the flow of information to the cortex. *Philos Trans R Soc Lond B Biol Sci*, *357*(1428), 1695-1708. https://doi.org/10.1098/rstb.2002.1161

Smith, R. E., Tournier, J. D., Calamante, F., & Connelly, A. (2013). SIFT: Spherical-deconvolution informed filtering of tractograms. *Neuroimage*, *67*, 298-312. https://doi.org/10.1016/j.neuroimage.2012.11.049

Smith, S. M., Nichols, T. E., Vidaurre, D., Winkler, A. M., Behrens, T. E., Glasser, M. F., Ugurbil, K., Barch, D. M., Van Essen, D. C., & Miller, K. L. (2015). A positive-negative mode of population covariation links brain connectivity, demographics and behavior.

Nat Neurosci, 18(11), 1565-1567. https://doi.org/10.1038/nn.4125

Sotiropoulos, S. N., & Zalesky, A. (2019). Building connectomes using diffusion MRI: why,
how and but. *NMR Biomed, 32*(4), e3752. https://doi.org/10.1002/nbm.3752

Stahl, E. A., Breen, G., Forstner, A. J., McQuillin, A., Ripke, S., Trubetskoy, V., Mattheisen,
M., Wang, Y., Coleman, J. R. I., Gaspar, H. A., de Leeuw, C. A., Steinberg, S.,
Pavlides, J. M. W., Trzaskowski, M., Byrne, E. M., Pers, T. H., Holmans, P. A.,
Richards, A. L., Abbott, L., . . . Bipolar Disorder Working Group of the Psychiatric
Genomics, C. (2019). Genome-wide association study identifies 30 loci associated
with bipolar disorder. *Nat Genet, 51*(5), 793-803. https://doi.org/10.1038/s41588-019-
0397-8

Suprano, I., Kocevar, G., Stamile, C., Hannoun, S., Fourneret, P., Revol, O., Nusbaum, F., &
Sappey-Marinier, D. (2020). White matter microarchitecture and structural network
integrity correlate with children intelligence quotient. *Sci Rep, 10*(1), 20722.
https://doi.org/10.1038/s41598-020-76528-x

Thompson, B. L., Levitt, P., & Stanwood, G. D. (2009). Prenatal exposure to drugs: effects
on brain development and implications for policy and education. *Nat Rev Neurosci,*
10(4), 303-312. https://doi.org/10.1038/nrn2598

Tooley, U. A., Mackey, A. P., Ciric, R., Ruparel, K., Moore, T. M., Gur, R. C., Gur, R. E.,
Satterthwaite, T. D., & Bassett, D. S. (2020). Associations between Neighborhood
SES and Functional Brain Network Development. *Cereb Cortex, 30*(1), 1-19.
https://doi.org/10.1093/cercor/bhz066

Tournier, J. D., Calamante, F., & Connelly, A. (2007). Robust determination of the fibre
orientation distribution in diffusion MRI: non-negativity constrained super-resolved
spherical deconvolution. *Neuroimage, 35*(4), 1459-1472.
https://doi.org/10.1016/j.neuroimage.2007.02.016

Tournier, J. D., Smith, R., Raffelt, D., Tabbara, R., Dhollander, T., Pietsch, M., Christiaens,
D., Jeurissen, B., Yeh, C. H., & Connelly, A. (2019). MRtrix3: A fast, flexible and open
software framework for medical image processing and visualisation. *Neuroimage,*
202, 116137. https://doi.org/10.1016/j.neuroimage.2019.116137

Van Buuren, S., & Groothuis-Oudshoorn, K. (2011). mice: Multivariate imputation by chained
equations in R. *Journal of statistical software, 45*, 1-67.

van den Heuvel, M. P., & Sporns, O. (2011). Rich-club organization of the human
connectome. *J Neurosci, 31*(44), 15775-15786.
https://doi.org/10.1523/JNEUROSCI.3539-11.2011

van den Heuvel, M. P., van Soelen, I. L., Stam, C. J., Kahn, R. S., Boomsma, D. I., &
Hulshoff Pol, H. E. (2013). Genetic control of functional brain network efficiency in
children. *Eur Neuropsychopharmacol, 23*(1), 19-23.
https://doi.org/10.1016/j.euroneuro.2012.06.007

Visser, M., Jefferies, E., Embleton, K. V., & Lambon Ralph, M. A. (2012). Both the middle
temporal gyrus and the ventral anterior temporal area are crucial for multimodal
semantic processing: distortion-corrected fMRI evidence for a double gradient of
information convergence in the temporal lobes. *J Cogn Neurosci, 24*(8), 1766-1778.
https://doi.org/10.1162/jocn_a_00244

Walters, R. K., Polimanti, R., Johnson, E. C., McClintick, J. N., Adams, M. J., Adkins, A. E.,
Aliev, F., Bacanu, S. A., Batzler, A., Bertelsen, S., Biernacka, J. M., Bigdeli, T. B.,
Chen, L. S., Clarke, T. K., Chou, Y. L., Degenhardt, F., Docherty, A. R., Edwards, A.
C., Fontanillas, P., . . . Agrawal, A. (2018). Transancestral GWAS of alcohol
dependence reveals common genetic underpinnings with psychiatric disorders. *Nat*
Neurosci, 21(12), 1656-1669. https://doi.org/10.1038/s41593-018-0275-1

Wang, H. T., Smallwood, J., Mourao-Miranda, J., Xia, C. H., Satterthwaite, T. D., Bassett, D.
S., & Bzdok, D. (2020). Finding the needle in a high-dimensional haystack: Canonical
correlation analysis for neuroscientists. *Neuroimage, 216*, 116745.
https://doi.org/10.1016/j.neuroimage.2020.116745

Watson, H. J., Yilmaz, Z., Thornton, L. M., Hubel, C., Coleman, J. R. I., Gaspar, H. A.,

Bryois, J., Hinney, A., Leppa, V. M., Mattheisen, M., Medland, S. E., Ripke, S., Yao,
S., Giusti-Rodriguez, P., Anorexia Nervosa Genetics, I., Hanscombe, K. B., Purves,
K. L., Eating Disorders Working Group of the Psychiatric Genomics, C., Adan, R. A.
H., . . . Bulik, C. M. (2019). Genome-wide association study identifies eight risk loci
and implicates metabo-psychiatric origins for anorexia nervosa. *Nat Genet*, *51*(8),
1207-1214. https://doi.org/10.1038/s41588-019-0439-2
Weintraub, S., Dikmen, S. S., Heaton, R. K., Tulsky, D. S., Zelazo, P. D., Bauer, P. J.,
Carlozzi, N. E., Slotkin, J., Blitz, D., Wallner-Allen, K., Fox, N. A., Beaumont, J. L.,
Mungas, D., Nowinski, C. J., Richler, J., Deocampo, J. A., Anderson, J. E., Manly, J.
J., Borosh, B., . . . Gershon, R. C. (2013). Cognition assessment using the NIH
Toolbox. *Neurology*, *80*(11 Suppl 3), S54-64.
https://doi.org/10.1212/WNL.0b013e3182872ded
Witten, D. M., Tibshirani, R., & Hastie, T. (2009). A penalized matrix decomposition, with
applications to sparse principal components and canonical correlation analysis.
Biostatistics, *10*(3), 515-534. https://doi.org/10.1093/biostatistics/kxp008
Wray, N. R., Ripke, S., Mattheisen, M., Trzaskowski, M., Byrne, E. M., Abdellaoui, A.,
Adams, M. J., Agerbo, E., Air, T. M., Andlauer, T. M. F., Bacanu, S. A., Baekvad-
Hansen, M., Beekman, A. F. T., Bigdeli, T. B., Binder, E. B., Blackwood, D. R. H.,
Bryois, J., Buttenschon, H. N., Bybjerg-Grauholm, J., . . . Major Depressive Disorder
Working Group of the Psychiatric Genomics, C. (2018). Genome-wide association
analyses identify 44 risk variants and refine the genetic architecture of major
depression. *Nat Genet*, *50*(5), 668-681. https://doi.org/10.1038/s41588-018-0090-3
Xia, C. H., Ma, Z., Ciric, R., Gu, S., Betzel, R. F., Kaczkurkin, A. N., Calkins, M. E., Cook, P.
A., Garcia de la Garza, A., Vandekar, S. N., Cui, Z., Moore, T. M., Roalf, D. R.,
Ruparel, K., Wolf, D. H., Davatzikos, C., Gur, R. C., Gur, R. E., Shinohara, R. T., . . .
Satterthwaite, T. D. (2018). Linked dimensions of psychopathology and connectivity
in functional brain networks. *Nat Commun*, *9*(1), 3003.
https://doi.org/10.1038/s41467-018-05317-y
Xu, L., Skoularidou, M., Cuesta-Infante, A., & Veeramachaneni, K. (2019). Modeling tabular
data using conditional gan. *Advances in neural information processing systems*, *32*.

1063

1064

Supplementary Materials

Details on Genotype Data

The saliva DNA samples of study participants were collected, and 733,293 single nucleotide polymorphisms (SNPs) were genotyped at Rutgers University Cell and DNA Repository (RUCDR) with Affymetrix NIDA Smoke Screen Array. Using PLINK 1.90, we excluded SNPs with genotype call rate <95%, sample call rate <95%, and minor allele frequency (MAF) <1% before imputation. The genotypes were imputed using the Michigan Imputation Server (Das et al., 2016) using the 1000 Genome phase3 version5 panel (Genomes Project et al., 2015) with Eagle v2.4 phasing (Loh et al., 2016). Then, the imputed variants with INFO score > .3 that did not meet our quality control criteria (i.e., call rate <95%, MAF <1%, and Hardy–Weinberg equilibrium p-value <1e-6) were additionally filtered out. To address potential bias derived from genetically diverse and related family members in the ABCD study, we employed PC-Air (Conomos et al., 2015) and PC-Relate (Conomos et al., 2016) to obtain genetically unrelated individuals beyond 4th-degree relatives (i.e., kinship coefficient >0.022) and to remove outliers beyond 6 SD limits from the center of ancestrally informative principal component (PC) space. After quality control procedures, we included a total of 11,301,999 variants in 10,199 multiethnic participants. From these, we excluded first-, second-, or third-degree related samples, resulting in 8,620 unrelated multiethnic participants, among whom 6,555 were of European ancestry.

Details on Polygenic Scores (PGSs)

Polygenic scores (PGSs) analyzed in this study are the same PGSs data

used in (Joo et al., 2024). PGSs were derived using summary statistics from publicly

available genome-wide association studies (GWAS). Thirty traits were selected

based on relevance to cognitive, psychiatric, and behavioral outcomes. The chosen

GWAS include attention-deficit/hyperactivity disorder (ADHD) (Demontis et al.,

2019), cognitive performance (CP) (Lee et al., 2018), educational attainment (EA)

(Lee et al., 2018), major depressive disorder (MDD) (Wray et al., 2018), insomnia

(Jansen et al., 2019), snoring (Jansen et al., 2019), intelligence quotient (IQ)

(Savage et al., 2018), post-traumatic stress disorder (PTSD) (Nievergelt et al., 2019),

depression (DEP) (Howard et al., 2019; Shen et al., 2020), body mass index (BMI)

(Akiyama et al., 2017; Locke et al., 2015), alcohol dependence (ALCDEP) (Walters

et al., 2018), autism spectrum disorder (ASD) (Grove et al., 2019), automobile

speeding propensity (ASP) (Akiyama et al., 2017), bipolar disorder (BIP) (Stahl et al.,

2019), cannabis during lifetimes (Cannabis) (Pasman et al., 2019), ever smoker

(Karlsson Linner et al., 2019), shared effects on five major psychiatric disorder

(CROSS) (Cross-Disorder Group of the Psychiatric Genomics, 2013), alcoholic

drinks consumption per week (Drinking) (Karlsson Linner et al., 2019), anorexia

nervosa (Watson et al., 2019), neuroticism (Nagel et al., 2018), obsessive-

compulsive disorder (OCD) (International Obsessive Compulsive Disorder

Foundation Genetics & Studies, 2018), first principal components of four risky

behaviors (Risky Behav) (Karlsson Linner et al., 2019), general risk tolerance

(RiskTol) (Karlsson Linner et al., 2019), schizophrenia (SCZ) (Bipolar et al., 2018;

Lam et al., 2019), worrying (Nagel et al., 2018), anxiety (Otowa et al., 2016),

subjective well-being (SWB) (Okbay et al., 2016), general happiness (UK Biobank

GWAS. Neale Lab. http://www.nealelab.is/ukbiobank/), and general happiness for

health (happiness-health) (UK Biobank GWAS. Neale Lab.

http://www.nealelab.is/ukbiobank/) and meaningful life (happiness-meaning) (UK

Biobank GWAS. Neale Lab. http://www.nealelab.is/ukbiobank/).

For 25 of these traits, where only European-based GWAS were available, we

calculated polygenic scores using European-based GWAS summary statistics. For

five traits where multiethnic GWAS results were available (PTSD, DEP, BMI,

ALCDEP, SCZ), we calculated polygenic scores using both European-based and

multiethnic-based GWAS summary statistics. In the European-only analysis, we

used polygenic scores for all 30 traits calculated based on European-based GWAS.

The GWAS summary statistics were used as input for PRS-CS (Ge et al.,

2019), a Bayesian regression method, to estimate the posterior effect sizes of SNPs.

The final scores were calculated using PLINK v1.9. To optimize the scores, we

followed the suggestion of the original PRS-CS paper and chose the optimal global

shrinkage hyperparameter (phi, φ) from among four possible values: 1, 1e-2, 1e-4,

and 1e-6. The validation procedure was carried out within 14 PGSs (i.e., DEP, MDD,

ADHD, general happiness, happiness-health, happiness-meaning, SWB, insomnia,

snoring, BMI, PTSD, CP, EA, IQ) that had related measures in the ABCD study. For

each PGS, we performed linear regression of the phenotype variable with each of

the four scores and covariates (sex, age, and the first ten genetic PCs), and then,

based on R^2 and beta coefficient of PGS, selected one of the four PGSs. The

remaining 16 PGSs was automatically validated by PRS-CS-auto (Ge et al., 2019),

which select the optimal value of global shrinkage parameter employing a Bayesian

approach. Finally, to minimize the bias from population stratification, we residualized

the final PGSs with the first ten genetic PCs.

Supplementary Table 1. Demographic information of the supplementary analysis participants (multiethnic analysis and complete data samples).

		PGS – BNM (n=7,297)		ENV – BNM (n=4,215)		Pheno – BNM (n=6,598)	
		train	test	train	test	train	test
N	total	5,837	1,460	3,372	843	5,278	1,320
Sex	Male	3,087	765	1,790	440	3,031	758
	Female	2,747	695	1,581	403	2,247	562
	Other	3	0	1	0	0	0
Race	White	3,251	780	1,702	451	3,036	765
	Black	798	207	474	103	674	161
	Hispanic	1,176	302	781	183	933	228
	Asian	18	6	86	21	94	19
	Other	594	165	329	85	541	147

 Supplementary Figure 1. SCCA Loadings of Polygenic Scores in

 European Samples.

Supplementary Figure 2. SCCA Loadings of Polygenic Scores and

Brain Network Measures in Multiethnic Samples.

1157

1158

1159 **Supplementary Figure 3. SCCA Loadings of Environmental**

1160 **Variables in the Main Analysis.**

1161

1162

1163

1164

1165

1166

1167

1168

1169

Supplementary Figure 4. SCCA Loadings of Environmental Variables and Brain Network Measures in Non-Imputed Samples.

1174 **Supplementary Figure 5. SCCA Loadings of Phenotype Variables in**

1175 **the Main Analysis Mode 1.**

1176

1177

1178

1179

1180

1181

1182

1183

1184

Supplementary Figure 6. SCCA Loadings of Phenotype Variables

and Brain Network Measures in Non-Imputed Samples Mode 1.

Supplementary Figure 7. SCCA Loadings of Phenotype Variables in

1189

the Main Analysis Mode 2.

1190

1191

1192

1193

1194

1195

1196

1197

1198

1199

 Supplementary Figure 8. SCCA Loadings of Phenotype Variables

 and Brain Network Measures in Non-Imputed Samples Mode 2.

1202