The plasma proteome of plant-based diets: analyses of 2920 proteins in 49,615 people.

Tammy Y.N. Tong[1], Karl Smith-Byrne[1], Keren Papier[1], Joshua R. Atkins[1], Mahboubeh Parsaeian[1], Timothy J. Key[1] and Ruth C. Travis[1]

1. Cancer Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, Oxford, UK

Corresponding author: Tammy Y.N. Tong

Postal address: Cancer Epidemiology Unit, Nuffield Department of Population Health,

University of Oxford, Richard Doll Building, Old Road Campus, Oxford, OX3 7LF

Tel: +44 (0) 1865 289653

Email: tammy.tong@ndph.ox.ac.uk

Abstract

Background & Aims: Circulating proteins are integral to many biological processes and could

be influenced by diet. We aimed to assess differences in the plasma proteome between people

of different dietary groups, defined by degree of animal food consumption.

Methods: The UK Biobank recruited middle-aged adults (mostly 40 to 69 years) throughout

the UK between 2006-2010. Relative concentrations of 2920 plasma proteins were quantified

using the Olink Proximity Extension Assay on blood samples from 49,615 participants, who

were also asked to report their ethnicity and consumption of red and processed meat, poultry,

fish, dairy and eggs. We defined six diet groups among the white British participants (23,243

regular meat eaters, 23,472 low meat eaters, 486 poultry eaters, 1081 fish eaters, 721

vegetarians, and 54 vegans), and two diet groups among the British Indians (391 meat eaters

and 167 vegetarians). We used multivariable-adjusted linear regressions to assess the cross-

sectional differences in protein concentrations between diet groups, with correction for

multiple testing.

Results: We observed significant differences in many plasma proteins by diet group (920

proteins in white British participants, 2 in British Indians). Of the biggest differences,

compared with regular meat eaters, the non-meat eaters had significantly higher FGF21 (e.g.

+0.40 SD in vegetarians), CKB (+0.34), GUCA2A (+0.33), FOLR1 (+0.32), IGFBP2 (+0.31)

and DSG2 (+0.30); all groups except the vegans had lower HAVCR1 (-0.38 in vegetarians).

Vegetarians also had significantly lower SELENOP (-0.46), while the vegans had lower

FGFBP2 (-0.68). The observed differences were generally similar in direction in both

ethnicities.

Conclusions: In this first comprehensive assessment of plasma proteins by diet group, we

identified many differences in proteins between vegetarians, vegans and meat eaters; this

variation in protein levels suggests differences in various biological activities, including

gastrointestinal tract and kidney function, which may relate to differences in future disease

risk.

Keywords: vegetarians, vegans, proteomics

Introduction

Proteins are essential for many bodily functions including supporting cell and tissue growth and structural integrity, and for enabling a range of enzymatic, biochemical, signalling and transport functions across different systems in the human body. Proteomics describes the large-scale study of multiple proteins and their interconnected pathways, the investigation of which allows a more comprehensive understanding of disease mechanisms [1]. Protein expression can be altered by environmental factors, including diet [2]. Previous studies have shown that differences in dietary habits with varying levels of animal foods can substantially impact dietary intakes of protein, which supplies the nine essential amino acids that cannot be synthesised endogenously, as well as both dietary intakes and circulating concentrations of amino acids [3]. However, there is scant evidence on how diet may affect the proteome, partly due to the limited availability of proteomics data in large scale studies, and no prior studies have investigated how long-term vegetarian and vegan diets which limit or exclude animal food products may influence the proteome. Circulating proteins have an important role in the aetiology of multiple diseases [4,5], and hence the examination of dietary influences on the proteome may offer unique insights into understanding how vegetarian and vegan diets may affect future disease risk.

The aim of this study is to provide a detailed description of circulating protein concentrations in people with varying degrees of animal food consumption, using data from the UK Biobank.

Methods

Study Population

The UK Biobank is a prospective cohort study of around 500,000 middle-aged people

(recruitment target 40-69 years), recruited from across the United Kingdom between 2006

and 2010. The scientific rationale and design of the UK Biobank have been described in

detail previously [6]. In brief, participants were identified from National Health Service

registers, and were invited to join the study if they lived within travelling distance (~25 km)

of one of the 22 assessment centres in England, Wales and Scotland. People who consented

to participate in the study attended a baseline visit at the assessment centre where they

completed a touchscreen questionnaire which asked about their lifestyle (including diet,

alcohol consumption, smoking status, physical activity), socio-demographic characteristics

and general health and medical history. All participants were also given a verbal interview,

and had their physical measurements and blood samples taken by trained staff. The UK

Biobank study was approved by the National Information Governance Board for Health and

Social Care and the National Health Service Northwest Multicentre Research Ethics

Committee (06/MRE08/65), and all participants gave informed consent to participate using a

signature capture device at the baseline visit.

Ethnicity classification

On the touchscreen questionnaire, participants were asked to self-identify their ethnicity as

'White', 'Mixed', 'Asian or Asian British', 'Black or Black British', 'Chinese', 'Other ethnic

group', 'Do not know', or 'Prefer not to answer', with further sub-categories under each

option. Participants were included for our analyses if they self-identified as 'White', or as

'Asian or Asian British' and subsequently as 'Indian', hereafter referred to as 'white British'

and 'British Indian'. The white British population was included as it made up the majority of

the participants in UK Biobank, while the British Indian population was included because of

the large proportion of vegetarians in this population group (25% compared to less than 2%

in the overall population). The number of vegetarians in the other ethnic groups was too small

to allow valid comparisons by diet groups, and therefore people of other ethnic groups were

not included in these analyses.

Diet group classification

Participants were classified into diet groups based on self-reported dietary data from the

touchscreen questionnaire completed at recruitment. Participants were asked to report their

frequency of consumption of processed meat (including processed poultry), unprocessed red

meat (beef, lamb or mutton, pork), unprocessed poultry (such as chicken or turkey), oily fish,

and other types of fish, with the possible responses ranging from "never" to "once or more

daily". Participants were also asked whether they never consumed dairy and eggs or foods

containing eggs. Based on their responses to these questions, the white British participants

were classified into six diet groups: regular meat eaters (red and processed meat consumption

>3 times per week), low meat eaters (red and processed meat consumption ≤3 times per

week), poultry eaters (participants who ate poultry but no red and processed meat), fish eaters

(participants who ate fish, but not red and processed meat, or poultry), vegetarians

(participants who did not eat any meat or fish), and vegans (participants who did not eat any

meat, fish, dairy products or eggs). The British Indian participants were classified into two

diet groups: meat eaters (ate any combination of red or processed meat or poultry) and

vegetarians; the numbers of fish eaters and vegans were low in the British Indians (around 30

participants combined), and thus were not included in these analyses.

Assay for plasma proteins

Non-fasting blood samples were collected from all UK Biobank participants by trained

personnel (either a phlebotomist or a nurse) except in a small proportion (0.3%) of

participants who declined, were deemed unable to, or where the attempt was abandoned for

technical or health reasons. Proteomic profiling was conducted using the Olink Proximity

Extension Assay on blood plasma samples in 54,219 participants, including 46,595

participants randomly selected at baseline, 6,376 participants selected by the UK Biobank

Pharma Proteomics Project (UKB-PPP) consortium members at baseline and 1,268

individuals who participated in the COVID-19 repeat-imaging study (20 of whom are also in

the UKB-PPP subset) [7]. The samples were shipped on dry ice to Olink Analyses Service,

Uppsala, Sweden for analysis; details of the selection procedures and the technical details of

the proteomics assays have been described previously [7]. In brief, the Olink Explore 3072

platform used in this study is an antibody-based assay which measures the relative abundance

of 2,941 protein analytes, including 2,923 unique proteins, distributed across eight 384-plex

panels (cardiometabolic, cardiometabolic II, inflammation, inflammation II, neurology,

neurology II, oncology and oncology II). Measurements of proteins are expressed as

normalized protein expression (NPX) values which are log-base-2 transformed. Protein

values below the limit of detection (LOD) were replaced with the LOD divided by the square

root of 2 [8], and protein concentrations were subsequently inverse rank normal transformed.

As a result of the processing of protein data, the unit differences in protein concentrations

may be interpreted as SD differences. For these analyses, we excluded three proteins with a

high proportion of missing data, including GLIPR1 (99% missing), NPML (74% missing)

and PCOLCE (64% missing), and thus the analyses included 2920 unique proteins.

Inclusion and exclusion criteria

Of the 54,219 participants selected for proteomic profiling, 53,020 participants remained after

the quality control procedures. Participants were further excluded if they were not of white

British or British Indian ethnicity (n=2989), could not be classified into one of the pre-

specified diet groups or had missing information for fasting time (n=416). After the

exclusions, 49,615 participants (49,057 white British and 558 British Indian) were included

in the analyses. A participant flow chart of the inclusion criteria is shown in **Supplementary**

figure 1.

Statistical analyses

Baseline characteristics of UK Biobank participants included in these analyses were tabulated

by six diet groups in the white British population and by two diet groups in the British Indian

population, as mean (SD) or number (%). We used multivariable adjusted linear regressions

to estimate differences in protein concentrations by diet group, separately by ethnicity, using

regular meat eaters as a reference group in the white British participants, and meat eaters as a

reference group in the British Indians. The model was adjusted for age at recruitment (5 year

categories), sex, region (London, North-West England, North-East England, Yorkshire, West

Midlands, East Midlands, South-East England, South-West England, Wales, Scotland),

fasting status (0–1, 2, 3, 4, 5, 6–7, ≥8 hours), body mass index (BMI; <20, 20.0–22.4, 22.5–

24.9, 25.0–27.4, 27.5–29.9, 30.0–32.4, 32.5–34.9, ≥35 kg/m^2, unknown), alcohol

consumption (<1, 1–7, 8–15, ≥16 g/d, unknown), smoking status (never, previous, current

<15 cigarettes/day, current ≥15 cigarettes/d, unknown) and physical activity (<10, 10-49.9,

≥50 excess metabolic equivalent of task, hr/wk, unknown). Wald tests were used to assess

overall heterogeneity between diet group in each ethnicity, and for pairwise comparisons of

each of the other diet groups compared to the reference group (regular meat eaters in white

British participants/meat eaters in British Indians). Heterogeneity between vegetarians and

vegans in the white British population was assessed based on post-estimation linear

combinations of parameters for all proteins.

To account for multiple testing while considering the high correlations between the

circulating proteins [9], we conducted a principal component analysis of the circulating

proteins in the analysis dataset, and determined that the first 1346 principal components

explained 95% of the total variation in the exposure data. Consequently, the effective number

of independent tests was determined to be 1346, and the *p*-value for statistical significance

was set to be 0.05/1346 = 0.000037. For the top proteins identified (the top 10 significant

proteins in each pairwise comparison against the reference group, based on ranking of *p*-

values), we conducted additional sensitivity analyses to evaluate the extent to which the

associations may be influenced by key covariates including BMI, smoking and alcohol

consumption, by presenting models with and without adjustment of these variables.

Additionally, we also conducted sensitivity analyses restricted to people who self-reported to

be in good and excellent health. All analyses were performed using Stata version 18.1

(StataCorp, TX, USA). All figures were generated using R version 4.2.1, the forest plots

using "Jasper makes plots" package version 2-266 [10].

Results

Baseline characteristics

The baseline characteristics of the study population are shown by ethnicity and diet group in

Table 1. Compared with white British regular meat eaters, the non-meat eaters were on

average slightly younger and more likely to be women. They had lower BMI on average,

reported lower alcohol consumption, more likely to have never smoked, and reported more

physical activity. Compared with British Indian meat eaters, the vegetarians were more likely

to be women and reported lower alcohol consumption, were more likely to have never

smoked and reported lower physical activity, but BMI was not noticeably different between

the two groups. Fasting time was not meaningfully different by diet group in both ethnicities.

Differences in plasma proteins by diet group

The plasma proteins that were significantly different in white British vegetarians and vegans

compared with regular meat eaters are shown as volcano plots in **Figure 1**. Significant

differences in proteins in white British low meat eaters, poultry eaters and fish eaters

compared with regular meat eaters, and in British Indian vegetarians compared with meat

eaters are shown as volcano plots in **Supplementary figures 2-6**. Overall, 920 plasma

proteins were significantly different by diet group (based on p-heterogeneity < 0.000037) in

the white British participants, including 632 plasma proteins that were significantly different

in one or more pairwise comparisons between regular meat eaters and at least one of the other

diet groups. This includes 333 proteins (27 higher, 306 lower) in low meat eaters compared

with regular meat eaters, 52 (9 higher, 43 lower) in poultry eaters, 193 (135 higher, 58 lower)

in fish eaters, 266 (235 higher, 31 lower) in vegetarians and 15 (13 higher, 2 lower) in

vegans; and 7 proteins were significantly different between vegetarians and vegans.

In the British Indian population, only 2 proteins (SUMF2 and ATRAID) were significantly

higher in vegetarians when compared with meat eaters, likely owing to the much smaller

number of British Indians in the study. However, the differences in most proteins were

directly consistent between the two ethnicities, as illustrated by the forest plots of top proteins

by diet group and ethnicity, with the six diet groups in the white British participants and the

two diet groups in British Indians (**Figures 2-3, Supplementary figures 7-9**). The complete

results of all proteins are shown in **Supplementary table 1** (as excel file Supp table 1.xlsx)**.**

Generally, in the multivariable-adjusted model, many proteins showed a gradient effect in

magnitude of differences across diet groups by degree of animal food exclusion, from regular

meat eaters to vegans. Figure 2 shows the top 10 proteins in white British vegetarians (based

on the ranking of pairwise p-values in vegetarians compared with regular meat eaters), and

Figure 3 shows the top 10 proteins in white British vegans. Of the top proteins in vegetarians

(Figure 2), compared with regular meat eaters, vegetarians and vegans both had significantly

higher FGF21 (+0.40 SD in vegetarians; +0.74 SD in vegans), GUCA2A (+0.33; +0.79),

CKB (+0.34, 0.60) and IGFBP2 (+0.31; +0.55) after correction for multiple testing, whereas

vegetarians but not vegans had significantly higher ART3 (+0.34), LAYN (+0.32), and lower

SELENOP (-0.46), CNDP1 (-0.39) and HAVCR1 (-0.38), though the differences in vegans

were directionally similar. In addition to the aforementioned proteins, vegans also had

substantially higher FOLR2 (+0.69), GAS6 (+0.68), DSG2 (+0.68), THY1 (+0.62), FOLR1

(+0.61) than regular meat eaters (Figure 3); the differences in these proteins were

directionally consistent but less extreme in the vegetarians. Vegans, but not the other diet

groups, had significantly lower FGFBP2 (-0.68) compared with regular meat eaters.

The top 10 proteins in white British low meat eaters, poultry eaters and fish eaters compared

with regular meat eaters are shown in Supplementary figures 8, 9 and 10 respectively. Of

these top proteins, none was uniquely different in one diet group only when compared with

regular meat eaters. Similar to the observations in vegetarians, all three groups had

substantially higher DSG2 and lower HAVCR1 compared with regular meat eaters; in both

low meat eaters and poultry eaters HAVCR1 is the protein that exhibited the largest

magnitude in difference across all proteins, while in fish eaters FGF23 showed the biggest

difference. In addition, low meat eaters and poultry eaters both had lower OSM and CD99L2,

the latter of which was higher in vegetarians. In sensitivity analyses of the top proteins with

different levels of covariate adjustment and limited to people of good and excellent health,

results were similar (**Supplementary table 2,** as excel file Supp table 2.xlsx).

Discussion

Summary of main findings

In this large British cohort, we observed differences in the levels of many proteins by diet

group. This is the first large and comprehensive study on vegetarian diet groups and plasma

proteomics, and therefore many of the substantial differences reported here are novel. In

particular, we saw substantial differences in some proteins potentially associated with disease

states such as HAVCR1 and DSG2 (associated with kidney disease [5,11] and inflammatory

bowel disease, IBD [12], respectively), as well as proteins associated with nutritional status

including IGFBP2 [13,14], and possibly FGF21 [15] and the folate receptors FOLR1 and

FOLR2 [16]. The magnitudes of differences in many proteins showed a gradient effect across

the diet groups, and were directionally consistent between the white British and British Indian

populations.

Comparison with previous studies

No previous studies were found that examined plasma proteomics in habitual vegetarians and

vegans. A small trial of 20 participants reported minimal differences in 1306 proteins after

being on a vegan diet for two weeks [17], but these results based on temporary vegan diets

are not comparable to our findings among habitual vegans and vegetarians. Another study

reported 837 proteins associated with different plant-based diet indices in a predominantly

non-vegetarian population [18], while other previous studies have reported on proteomic

profiles by other dietary patterns, including the Dietary Approaches to Stop Hypertension

(DASH) diet, the Healthy Eating Index and the Mediterranean diet, as well as other data

derived dietary patterns [19–22]. Whilst vegetarian and vegan diets share some common

features with these other healthy dietary patterns, such as higher intakes of fruit, vegetables

and whole grains, the diets are also inherently different in many other aspects. For example,

while both the DASH diet and Mediterranean diet recommend reducing red and processed

meat consumption, they generally do not pose any restrictions on consumption of fish and

fish products, or dairy products (particularly low-fat versions) [23,24] , and participants with

the highest adherence to the plant-based diet index still regularly consumed animal products

[25]. In contrast, vegetarian and vegan diets are defined by the complete exclusion of animal

food groups. Taking into account these differences in diet as well as differences in the

proteomics platforms used, it is not surprising that we have not found substantial similarities

in the results; however, the previous studies also reported that people with higher adherence

to the DASH diet had higher FOLR2 [19] and DSG2 [20], while those with higher adherence

to the healthful plant-based diet index also had higher IGFBP2 [18],. There are only a limited

number of studies examining vegetarian and vegan diets with individual proteins that were

also of particular interest in this study, which are described below.

SELENOP

Two previous small studies including 36 vegans [26] and 54 vegetarians/vegans [27] both

found lower circulating levels of SELENOP in vegans/vegetarians than omnivores.

SELENOP is a transporter protein for serum selenium, and previous studies have shown that

vegetarians and vegans both had lower intakes of selenium compared with omnivores who

include meat and fish in their diets [28]. Therefore, lower levels of SELENOP in response to

lower selenium intakes in these diet groups may be expected. However, the soil type in which

crops were grown has been shown to influence both selenium intake and status, thus the

specific selenium content of plant foods may vary by their production region [29]. Some

previous observational studies have suggested that lower circulating selenium might be

associated with higher risks of some types of cancers [30,31] as well as lower bone mineral

density [32,33], though a small selenium supplementation trial in the UK (39 or fewer

participants in each arm) reported no clinically relevant differences in bone mineral density

after 26 weeks [34].

FGF21

In support of our findings, higher concentrations of FGF21 in vegetarians and vegans have

been shown previously in a small study of 36 each of omnivores, vegetarians and vegans

[35]. FGF21 has been hypothesised to be high in vegans as a downstream effect of low intake

of methionine [15], an essential amino acid that is known to be limited in vegan diets [3].

Studies of FGF21 administration in rodents have found favourable effects on adiposity, lipid

profiles, and non-alcohol fatty liver disease, but possible adverse effects on bone homeostasis

[36]; randomised trials of FGF21 analogues also support the reduction in triglycerides [37]

and improvement of fibrosis in patients with non-alcoholic steatohepatitis [38]. Consistent

with these observations, we have previously reported lower BMI and body fat [39], more

favourable lipid profiles [40] and lower heart disease risk [41], but also lower bone mineral

density [39] and higher fracture risk [42] in vegetarians and vegans when compared with

meat eaters; another study has reported lower odds of non-alcohol fatty liver disease in

vegetarians than non-vegetarians [43]. Based on these differences in FGF21, as well the

higher FGF23 in fish eaters and vegetarians, and lower FGFBP2 in vegans but not the other

diet groups, the potential relevance of FGF signalling pathways in relation to both diet and

disease risk deserve further in-depth investigation.

GUCA2A

GUCA2A was substantially higher in all diet groups compared to regular meat eaters, with

the exception of low meat eaters. GUCA2A activates the Guanylate Cyclase C receptor,

which has a role in the maintenance of gut physiology including the increase of water

movement into the intestinal lumen [44]. Consistent with the observations in this study, a

previous EPIC-Oxford study reported a higher frequency of bowel movements in vegetarians

and vegans [45]. Previous studies have found that fasting plasma levels of GUCA2A are

significantly lower in patients who have been diagnosed with Crohn's disease compared with

healthy controls [46], and that loss of GUCA2A expression may be an important determinant

of colorectal cancer via silencing of the GUCY2C tumour suppressor [47].

FOLR1 and FOLR2

We found both FOLR1 and FOLR2 to be higher in non-meat eaters, with vegans having

particularly high levels. FOLR1 and FOLR2 are both folate receptors which bind to and

import folic acid into cells, and are usually downregulated with repletion of folate, however,

this process may also be mediated by homocysteine in a positive direction [16]. While

vegetarians and vegans typically have high dietary and serum folate, they may also have high

homocysteine due to low vitamin B12 intakes [48,49], but no other studies have reported on

folate receptor expression in vegetarians and vegans for comparison. Overexpression of folate

receptors has been reported in patients with some types of cancers [50,51], and previous

prospective analyses in UK Biobank have also found a positive association of both FOLR1

and FOLR2 with kidney cancer [5]. However, previous studies have not shown a higher risk

of kidney cancer in vegetarians or vegans compared with meat eaters [52], and thus it is

possible the upregulation of folate receptors in these groups mainly reflect differences in diet

and homocysteine levels.

IGFBP2

Similar to our findings, a previous study in EPIC-Oxford has found that compared with meat-

eaters, both vegetarians and vegans had higher concentrations of IGFBP2, a binding protein

for IGF-1 [13]. The mechanism for higher IGFBP2 in vegetarians and vegans is not well

established but has been suggested to be related to lower dietary intakes of protein (especially

from dairy) [14], essential amino acids [13], or lower total energy intake [53], which are all

characteristic of vegetarian and vegan diets [3,28,39]. Insulin-like growth factors and their

associated binding proteins, including IGFBP2, are involved in the regulation of cell

proliferation, differentiation, and apoptosis [54,55]. However, previous studies, including

from UK Biobank, have not clearly established an independent association of IGFBP2 with

different cancer types [5,56–58].

CKB

CKB or Creatine kinase B-type, is a class of creatine kinases which are cytoplasmic enzymes

involved in energy homeostasis, and often elevated in cases of musculoskeletal trauma or

myocardial injury [59]. CKB is the main isoform of creatine kinase in the brain, and plays a

critical role in the regulation of ATP levels in neural cells; previous studies have shown that

CKB levels are reduced in brain regions affected by neurodegenerative diseases such as

Alzheimer's disease and dementia [60,61]. On the other hand, elevated levels of CKB have

also been reported in patients with acute brain injury or stroke [62,63], or failure or

suppression of osteoclasts [64]. Given these potentially contradictory reports, the implications

and relevance of higher CKB in non-meat eaters compared with meat eaters require further

investigation.

DSG2

Low and non-meat eaters, particularly vegans, had higher DSG2 in the current study. DSG2

belongs to the family of desmosomal cadherins, which includes both desmogleins (DSGs)

and desmocollins (DSCs), with DSG2 and DSC2 being the major isoforms in humans, and

the only isoforms expressed in the intestinal epithelium [12]. Consistently, the current study

also showed that DSC2 was higher in non-meat eaters particularly vegans. Though the two

proteins are structurally and functionally similar, laboratory evidence in mouse models

suggest that DSG2 may have a more prominent role in the maintenance of the integrity of the

intestinal epithelial barrier than DSC2 [65]. In humans, several small studies on IBD patients

have found lower intestinal protein levels of DSG2 in IBD patients, suggesting a role of the

protein in IBD pathogenesis [12].

HAVCR1

HAVCR1, also known as KIM-1, has been recognised as a biomarker of kidney injury and

possibly an early diagnostic marker of kidney disease [11], as well as being associated with

kidney cancer [5,66]. In mice, high-fat diets or the saturated fatty acid palmitate have been

shown to upregulate expression of HAVCR1 in the proximal tubules of the kidney [67–69].

In humans, vegetarians and vegans tend to have lower intake of saturated fat, which is

consistent with the observation of lower HAVCR1 in the non-meat eaters (though not

significantly so in vegans) in the current study. Additionally, high protein diets, especially

diets high in protein from animal sources, have been suggested to have adverse effects on

kidney health [70]. A previous study suggested a lower risk of chronic kidney disease in

vegetarians after accounting for baseline risk of diabetes and hypertension [71]. Furthermore,

as HAVCR1 is expressed on the surface of immune cells, it is also believed to have a role

regulating immune responses via activation and proliferation of immune cells [11].

Other proteins

The functions of many other proteins identified are not well established. CNDP1 is a

dipeptidase, and one previous study reported higher CNDP1 in people on high protein diets

[72], which would be consistent with our observation of lower CNDP1 in non-meat eaters

(not statistically significant in vegans) who generally have lower protein intake. Several of

the proteins identified such as ART3, GAS6, THY1 which were higher in vegetarians and/or

vegans are involved in a range of signalling pathways and progression of tumour tissues [73–

76], while other proteins including LAYN and CDHR2 are believed to have roles in cell

adhesion [77,78]. AHSP, which was significantly higher in vegans than vegetarians, has a

role in stabilising haemoglobin and its loss of function has been linked to inherited anaemia

[79], but its relevance for nutritional anaemia is unclear. Overall, the reasons that these

proteins appear to be influenced by diet and their implications require further research. As the

first study on habitual vegetarian diet group and a large panel of circulating proteins, the

current study focused on the biggest differences by diet groups, and many other differences

were observed that could not be described in detail, but still deserve further investigation. The

study demonstrates the utility of investigating circulating protein differences by diet groups;

future research should measure these and other proteins using multiple technologies in

relation to dietary factors, to replicate and expand on our findings.

Strengths and limitations

The key strength of this study was that it showed for first time differences in circulating

proteins between people of different habitual vegetarian diet groups in a large population, and

many of the findings were completely novel. We have defined six diet groups in the white

British population and two diet groups in the British Indian population, which allowed a

detailed comparison by both dietary habits and ethnicity. Of the limitations, as with all

observational studies, some level of self-selection bias may be present, which limits the

generalisability of the findings. While we have adjusted for several important confounders,

residual confounding by other dietary and non-dietary factors may still be present, though

sensitivity analyses showed that results were consistent across all adjustment models. We are

also unable to infer causality due to the cross-sectional nature of the study.

Conclusions

In this large population-based cohort in the United Kingdom, many differences in circulating

protein concentrations were observed between different diet groups. These proteins are

involved in a range of different biological pathways and processes, including nutritional

status, and functions related to gastrointestinal tract and kidney function. These differences

likely reflect physiological differences between the different diet groups, and the implications

of these differences for future disease risk, as well as specific diet-protein associations of

relevant foods and nutrients, require further investigation.

Acknowledgements: This research has been conducted using UK Biobank Resource under application 67506.

Author's contributions: TYNT, TJK and RCT conceived and designed the research question. KSB conducted quality control of the proteomics data. TYNT analysed the data and wrote the first draft of the manuscript. All authors provided input on data analysis and interpretation of results, revised the manuscript critically for important intellectual content, and read and approved the final manuscript.

Data sharing: The UK Biobank is an open-access resource. Bona fide researchers can apply to use the UK Biobank data set by registering and applying at http://www.ukbiobank.ac.uk/register-apply/.

Funding: The work is supported by a UK Research and Innovation Future Leaders Fellowship (MR/X032809/1), UK Medical Research Council (MR/M012190/1), Wellcome Trust Our Planet Our Health (Livestock, Environment and People, LEAP 205212/Z/16/Z) and Cancer Research UK (C8221/A29017 and C8221/A29186). The funders had no role in study design, data collection, analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors had no conflicts of interest.

References

[1]   Cui M, Cheng C, Zhang L. High-throughput proteomics: a methodological mini-review. Lab Invest 2022;102:1170–81. https://doi.org/10.1038/s41374-022-00830-7.

[2]   Al-Amrani S, Al-Jabri Z, Al-Zaabi A, Alshekaili J, Al-Khabori M. Proteomics: Concepts and applications in human medicine. World J Biol Chem 2021;12:57–69. https://doi.org/10.4331/wjbc.v12.i5.57.

[3]   Schmidt JA, Rinaldi S, Scalbert A, Ferrari P, Achaintre D, Gunter MJ, et al. Plasma concentrations and intakes of amino acids in male meat-eaters, fish-eaters, vegetarians and vegans: a cross-sectional analysis in the EPIC-Oxford cohort. Eur J Clin Nutr 2016;70:306–12. https://doi.org/10.1038/ejcn.2015.144.

[4]   Pietzner M, Wheeler E, Carrasco-Zanini J, Cortes A, Koprulu M, Wörheide MA, et al. Mapping the proteo-genomic convergence of human diseases. Science (1979) 2021;374. https://doi.org/10.1126/science.abj1541.

[5]   Papier K, Atkins JR, Tong TYN, Gaitskell K, Desai T, Ogamba CF, et al. Identifying proteomic risk factors for cancer using prospective and exome analyses of 1463 circulating proteins and risk of 19 cancers in the UK Biobank. Nat Commun 2024;15:4010. https://doi.org/10.1038/s41467-024-48017-6.

[6]   Fry A, Littlejohns TJ, Sudlow C, Doherty N, Adamska L, Sprosen T, et al. Comparison of sociodemographic and health-related characteristics of UK Biobank participants with those of the general population. Am J Epidemiol 2017;186:1026–34. https://doi.org/10.1093/aje/kwx246.

[7]   Sun BB, Chiou J, Traylor M, Benner C, Hsu Y-H, Richardson TG, et al. Plasma proteomic associations with genetics and health in the UK Biobank. Nature 2023;622:329–38. https://doi.org/10.1038/s41586-023-06592-6.

[8]   Albanes D, Alcala K, Alcala N, Amos CI, Arslan AA, Bassett JK, et al. The blood proteome of imminent lung cancer diagnosis. Nat Commun 2023;14:3042. https://doi.org/10.1038/s41467-023-37979-8.

[9]   Gao X, Starmer J, Martin ER. A multiple testing correction method for genetic association studies using correlated single nucleotide polymorphisms. Genet Epidemiol 2008;32:361–9. https://doi.org/10.1002/gepi.20310.

[10]   Arnold M. Jasper: Jasper makes plots. 2020 2022.

[11]   Liu S, Tang W, Cao J, Shang M, Sun H, Gong J, et al. A Comprehensive Analysis of HAVCR1 as a Prognostic and Diagnostic Marker for Pan-Cancer. Front Genet 2022;13:1–11. https://doi.org/10.3389/fgene.2022.904114.

[12]   Schlegel N, Boerner K, Waschke J. Targeting desmosomal adhesion and signalling for intestinal barrier stabilization in inflammatory bowel diseases—Lessons from experimental models and patients. Acta Physiologica 2021;231:1–15. https://doi.org/10.1111/apha.13492.

[13]   Allen NE, Appleby PN, Davey GK, Kaaks R, Rinaldi S, Key TJ. The associations of diet with serum insulin-like growth factor I and its main binding proteins in 292 women meat-eaters, vegetarians, and vegans. Cancer Epidemiology Biomarkers and Prevention 2002;11:1441–8.

[14]   Crowe FL, Key TJ, Allen NE, Appleby PN, Roddam A, Overvad K, et al. The association between diet and serum concentrations of IGF-I, IGFBP-1, IGFBP-2, and IGFBP-3 in the European Prospective Investigation into Cancer and Nutrition. Cancer Epidemiology Biomarkers & Prevention 2009;18:1333–40. https://doi.org/10.1158/1055-9965.EPI-08-0781.

[15]   McCarty MF. The moderate essential amino acid restriction entailed by low-protein vegan diets may promote vascular health by stimulating FGF21 secretion. Horm Mol Biol Clin Investig 2017;30:1–9. https://doi.org/10.1515/hmbci-2015-0056.

[16]   Antony AC, Tang YS, Khan RA, Biju MP, Xiao X, Li QJ, et al. Translational upregulation of folate receptors is mediated by homocysteine via RNA-heterogeneous nuclear ribonucleoprotein E1 interactions. Journal of Clinical Investigation 2004;113:285–301. https://doi.org/10.1172/JCI200411548.

[17]   Link VM, Subramanian P, Cheung F, Han KL, Stacy A, Chi L, et al. Differential peripheral immune signatures elicited by vegan versus ketogenic diets in humans. Nat Med 2024;30:560–72. https://doi.org/10.1038/s41591-023-02761-2.

[18]   Kim H, Chen J, Prescott B, Walker ME, Grams ME, Yu B, et al. Plasma proteins associated with plant-based diets: Results from the Atherosclerosis Risk in Communities (ARIC) study and Framingham Heart Study (FHS). Clinical Nutrition 2024;43:1929–40. https://doi.org/10.1016/j.clnu.2024.07.005.

[19]   Kim H, Lichtenstein AH, Ganz P, Du S, Tang O, Yu B, et al. Identification of Protein Biomarkers of the Dietary Approaches to Stop Hypertension Diet in Randomized Feeding Studies and Validation in an Observational Study. J Am Heart Assoc 2023;12:e028821. https://doi.org/10.1161/JAHA.122.028821.

[20]   Du S, Chen J, Kim H, Walker ME, Lichtenstein AH, Chatterjee N, et al. Plasma Protein Biomarkers of Healthy Dietary Patterns: Results from the Atherosclerosis Risk in Communities Study and the Framingham Heart Study. J Nutr 2023;153:34–46. https://doi.org/10.1016/j.tjnut.2022.11.008.

[21]   Walker ME, Song RJ, Xu X, Gerszten RE, Ngo D, Clish CB, et al. Proteomic and metabolomic correlates of healthy dietary patterns: The framingham heart study. Nutrients 2020;12:1–20. https://doi.org/10.3390/nu12051476.

[22]   Warensjö Lemming E, Byberg L, Stattin K, Ahmad S, Lind L, Elmståhl S, et al. Dietary Pattern Specific Protein Biomarkers for Cardiovascular Disease: A Cross-Sectional Study in 2 Independent Cohorts. J Am Heart Assoc 2019;8. https://doi.org/10.1161/JAHA.118.011860.

[23]   Kim H, Lichtenstein AH, Ganz P, Du S, Tang O, Yu B, et al. Identification of Protein Biomarkers of the Dietary Approaches to Stop Hypertension Diet in Randomized

Feeding Studies and Validation in an Observational Study. J Am Heart Assoc 2023;12:e028821. https://doi.org/10.1161/JAHA.122.028821.

[24]   Du S, Chen J, Kim H, Walker ME, Lichtenstein AH, Chatterjee N, et al. Plasma Protein Biomarkers of Healthy Dietary Patterns: Results from the Atherosclerosis Risk in Communities Study and the Framingham Heart Study. J Nutr 2023;153:34–46. https://doi.org/10.1016/j.tjnut.2022.11.008.

[25]   Sullivan VK, Kim H, Caulfield LE, Steffen LM, Selvin E, Rebholz CM. Plant-Based Dietary Patterns and Incident Diabetes in the Atherosclerosis Risk in Communities (ARIC) Study. Diabetes Care 2024;47:803–9. https://doi.org/10.2337/dc23-2013.

[26]   Menzel J, Abraham K, Stangl GI, Ueland PM, Obeid R, Schulze MB, et al. Vegan Diet and Bone Health—Results from the Cross-Sectional RBVD Study. Nutrients 2021;13:685. https://doi.org/10.3390/nu13020685.

[27]   Hoeflich J, Hollenbach B, Behrends T, Hoeg A, Stosnach H, Schomburg L. The choice of biomarkers determines the selenium status in young German vegans and vegetarians. British Journal of Nutrition 2010;104:1601–4. https://doi.org/10.1017/S0007114510002618.

[28]   Sobiecki JG, Appleby PN, Bradbury KE, Key TJ. High compliance with dietary recommendations in a cohort of meat eaters, fish eaters, vegetarians, and vegans: Results from the European Prospective Investigation into Cancer and Nutrition-Oxford study. Nutrition Research 2016;36:464–77. https://doi.org/10.1016/j.nutres.2015.12.016.

[29]   Hurst R, Siyame EWP, Young SD, Chilimba ADC, Joy EJM, Black CR, et al. Soil-type influences human selenium status and underlies widespread selenium deficiency risks in Malawi. Sci Rep 2013;3:1425. https://doi.org/10.1038/srep01425.

[30]   Hughes DJ, Duarte-Salles T, Hybsier S, Trichopoulou A, Stepien M, Aleksandrova K, et al. Prediagnostic selenium status and hepatobiliary cancer risk in the European Prospective Investigation into Cancer and Nutrition cohort. Am J Clin Nutr 2016;104:406–14. https://doi.org/10.3945/ajcn.116.131672.

[31]   Fedirko V, Jenab M, Méplan C, Jones JS, Zhu W, Schomburg L, et al. Association of Selenoprotein and Selenium Pathway Genotypes with Risk of Colorectal Cancer and Interaction with Selenium Status. Nutrients 2019;11:935. https://doi.org/10.3390/nu11040935.

[32]   Hoeg A, Gogakos A, Murphy E, Mueller S, Köhrle J, Reid DM, et al. Bone Turnover and Bone Mineral Density Are Independently Related to Selenium Status in Healthy Euthyroid Postmenopausal Women. J Clin Endocrinol Metab 2012;97:4061–70. https://doi.org/10.1210/jc.2012-2121.

[33]   Xie H, Wang N, He H, Yang Z, Wu J, Yang T, et al. The association between selenium and bone health: a meta-analysis. Bone Joint Res 2023;12:423–32. https://doi.org/10.1302/2046-3758.127.BJR-2022-0420.R1.

[34]   Walsh JS, Jacques RM, Schomburg L, Hill TR, Mathers JC, Williams GR, et al. Effect of selenium supplementation on musculoskeletal health in older women: a randomised,

double-blind, placebo-controlled trial. Lancet Healthy Longev 2021;2:e212–21. https://doi.org/10.1016/S2666-7568(21)00051-9.

[35] Castaño-Martinez T, Schumacher F, Schumacher S, Kochlik B, Weber D, Grune T, et al. Methionine restriction prevents onset of type 2 diabetes in NZO mice. FASEB J 2019;33:7092–102. https://doi.org/10.1096/fj.201900150R.

[36] Geng L, Lam KSL, Xu A. The therapeutic potential of FGF21 in metabolic diseases: from bench to clinic. Nat Rev Endocrinol 2020;16:654–67. https://doi.org/10.1038/s41574-020-0386-0.

[37] Bhatt DL, Bays HE, Miller M, Cain JE, Wasilewska K, Andrawis NS, et al. The FGF21 analog pegozafermin in severe hypertriglyceridemia: a randomized phase 2 trial. Nat Med 2023;29. https://doi.org/10.1038/s41591-023-02427-z.

[38] Loomba R, Sanyal AJ, Kowdley K V., Bhatt DL, Alkhouri N, Frias JP, et al. Randomized, Controlled Trial of the FGF21 Analogue Pegozafermin in NASH. New England Journal of Medicine 2023:1–11. https://doi.org/10.1056/nejmoa2304286.

[39] Tong TY, Key TJ, Sobiecki JG, Bradbury KE. Anthropometric and physiologic characteristics in white and British Indian vegetarians and nonvegetarians in the UK Biobank. Am J Clin Nutr 2018;107:909–20. https://doi.org/10.1093/ajcn/nqy042.

[40] Tong TYN, Perez-Cornago A, Bradbury KE, Key TJ. Biomarker Concentrations in White and British Indian Vegetarians and Nonvegetarians in the UK Biobank. J Nutr 2021;151:3168–79. https://doi.org/10.1093/jn/nxab192.

[41] Tong TYN, Appleby PN, Bradbury KE, Perez-Cornago A, Travis RC, Clarke R, et al. Risks of ischaemic heart disease and stroke in meat eaters, fish eaters, and vegetarians over 18 years of follow-up: results from the prospective EPIC-Oxford study. BMJ 2019;366:l4897. https://doi.org/10.1136/bmj.l4897.

[42] Tong TYN, Appleby PN, Armstrong MEG, Fensom GK, Knuppel A, Papier K, et al. Vegetarian and vegan diets and risks of total and site-specific fractures: results from the prospective EPIC-Oxford study. BMC Med 2020;18:353. https://doi.org/10.1186/s12916-020-01815-3.

[43] Chiu T, Lin M-N, Pan W-H, Chen Y-C, Lin C-L. Vegetarian diet, food substitution, and nonalcoholic fatty liver. Tzu Chi Med J 2018;30:102. https://doi.org/10.4103/tcmj.tcmj_109_17.

[44] Dye FS, Larraufie P, Kay R, Darwish T, Rievaj J, Goldspink DA, et al. Characterisation of proguanylin expressing cells in the intestine – evidence for constitutive luminal secretion. Sci Rep 2019;9:1–20. https://doi.org/10.1038/s41598-019-52049-0.

[45] Sanjoaquin MA, Appleby PN, Spencer EA, Key TJ. Nutrition and lifestyle in relation to bowel movement frequency: a cross-sectional study of 20 630 men and women in EPIC–Oxford. Public Health Nutr 2004;7:77–83. https://doi.org/10.1079/phn2003522.

[46] von Volkmann HL, Brønstad I, Tronstad RR, Dizdar V, Nylund K, Hanevik K, et al. Plasma levels of guanylins are reduced in patients with Crohn's disease. Scand J Gastroenterol 2020;55:449–53. https://doi.org/10.1080/00365521.2020.1748224.

[47] Wilson C, Lin JE, Li P, Snook AE, Gong J, Sato T, et al. The paracrine hormone for the GUCY2C tumor suppressor, guanylin, is universally lost in colorectal cancer. Cancer Epidemiology Biomarkers and Prevention 2014;23:2328–37. https://doi.org/10.1158/1055-9965.EPI-14-0440.

[48] Gilsing AMJ, Crowe FL, Lloyd-Wright Z, Sanders TAB, Appleby PN, Allen NE, et al. Serum concentrations of vitamin B12 and folate in British male omnivores, vegetarians and vegans: results from a cross-sectional analysis of the EPIC-Oxford cohort study. Eur J Clin Nutr 2010;64:933–9. https://doi.org/10.1038/ejcn.2010.142.

[49] Hung CJ, Huang PC, Lu SC, Li YH, Huang H Bin, Lin BF, et al. Plasma homocysteine levels in Taiwanese vegetarians are higher than those of omnivores. Journal of Nutrition 2002;132:152–8. https://doi.org/10.1093/jn/132.2.152.

[50] Shiode Y, Kodama T, Sato Y, Takahashi R, Matsumae T, Shirai K, et al. Folate receptor 1 is a stemness trait-associated diagnostic and prognostic marker for hepatocellular carcinoma. Biomark Res 2025;13:37. https://doi.org/10.1186/s40364-025-00752-8.

[51] Bax HJ, Chauhan J, Stavraka C, Santaolalla A, Osborn G, Khiabany A, et al. Folate receptor alpha in ovarian cancer tissue and patient serum is associated with disease burden and treatment outcomes. Br J Cancer 2023;128:342–53. https://doi.org/10.1038/s41416-022-02031-x.

[52] Key TJ, Appleby PN, Crowe FL, Bradbury KE, Schmidt JA, Travis RC. Cancer in British vegetarians: Updated analyses of 4998 incident cancers in a cohort of 32,491 meat eaters, 8612 fish eaters, 18,298 vegetarians, and 2246 vegans. American Journal of Clinical Nutrition 2014;100:378–85. https://doi.org/10.3945/ajcn.113.071266.

[53] Kaaks R, Bellati C, Venturelli E, Rinaldi S, Secreto G, Biessy C, et al. Effects of dietary intervention on IGF-I and IGF-binding proteins, and related alterations in sex steroid metabolism: The Diet and Androgens (DIANA) Randomised Trial. Eur J Clin Nutr 2003;57:1079–88. https://doi.org/10.1038/sj.ejcn.1601647.

[54] Travis RC, Appleby PN, Martin RM, Holly JMP, Albanes D, Black A, et al. A Meta-analysis of Individual Participant Data Reveals an Association between Circulating Levels of IGF-I and Prostate Cancer Risk. Cancer Res 2016;76:2288–300. https://doi.org/10.1158/0008-5472.CAN-15-1551.

[55] Kunhiraman H, McSwain L, Shahab SW, Gershon TR, MacDonald TJ, Kenney AM. IGFBP2 promotes proliferation and cell migration through STAT3 signaling in Sonic hedgehog medulloblastoma. Acta Neuropathol Commun 2023;11:62. https://doi.org/10.1186/s40478-023-01557-2.

[56] Watts EL, Perez-Cornago A, Fensom GK, Smith-Byrne K, Noor U, Andrews CD, et al. Circulating insulin-like growth factors and risks of overall, aggressive and early-onset prostate cancer: a collaborative analysis of 20 prospective studies and Mendelian

randomization analysis. Int J Epidemiol 2023;52:71–86. https://doi.org/10.1093/ije/dyac124.

[57] Cust AE, Allen NE, Rinaldi S, Dossus L, Friedenreich C, Olsen A, et al. Serum levels of C☐peptide, IGFBP☐1 and IGFBP☐2 and endometrial cancer risk; Results from the European prospective investigation into cancer and nutrition. Int J Cancer 2007;120:2656–64. https://doi.org/10.1002/ijc.22578.

[58] Jenab M, Riboli E, Cleveland RJ, Norat T, Rinaldi S, Nieters A, et al. Serum C☐peptide, IGFBP☐1 and IGFBP☐2 and risk of colon and rectal cancers in the European Prospective Investigation into Cancer and Nutrition. Int J Cancer 2007;121:368–76. https://doi.org/10.1002/ijc.22697.

[59] Leverenz D, Zaha O, Crofford LJ, Chung CP. Causes of creatine kinase levels greater than 1000 IU/L in patients referred to rheumatology. Clin Rheumatol 2016;35:1541–7. https://doi.org/10.1007/s10067-016-3242-9.

[60] Aksenov M, Aksenova M, Butterfield DA, Markesbery WR. Oxidative Modification of Creatine Kinase BB in Alzheimer's Disease Brain. J Neurochem 2000;74:2520–7. https://doi.org/10.1046/j.1471-4159.2000.0742520.x.

[61] Burbaeva GSh, Aksenova MV, Makarenko IG. Decreased Level of Creatine Kinase BB in the Frontal Cortex of Alzheimer Patients. Dement Geriatr Cogn Disord 1992;3:91–4. https://doi.org/10.1159/000107000.

[62] Capocchi G, Tassi C, Ricci S, Zampolini M, Fausti R, Rossi A. Creatine kinase BB activity in serum of patients with acute stroke: correlation with the severity of brain damage. The Italian Journal of Neurological Sciences 1987;8:567–70. https://doi.org/10.1007/BF02333663.

[63] Kaste M, Hernesniemi J, Somer H, Hillbom M, Konttinen A. Creatine kinase isoenzymes in acute brain injury. J Neurosurg 1981;55:511–5. https://doi.org/10.3171/jns.1981.55.4.0511.

[64] D'Eufemia P, Finocchiaro R, Villani C, Zambrano A, Lodato V, Palombaro M, et al. Serum brain-type creatine kinase increases in children with osteogenesis imperfecta during neridronate treatment. Pediatr Res 2014;75:626–30. https://doi.org/10.1038/pr.2014.20.

[65] Gross A, Pack LAP, Schacht GM, Kant S, Ungewiss H, Meir M, et al. Desmoglein 2, but not desmocollin 2, protects intestinal epithelia from injury. Mucosal Immunol 2018;11:1630–9. https://doi.org/10.1038/s41385-018-0062-z.

[66] Scelo G, Muller DC, Riboli E, Johansson M, Cross AJ, Vineis P, et al. KIM-1 as a Blood-Based Marker for Early Detection of Kidney Cancer: A Prospective Nested Case–Control Study. Clinical Cancer Research 2018;24:5594–601. https://doi.org/10.1158/1078-0432.CCR-18-1496.

[67] Yu Y, Mo H, Zhuo H, Yu C, Liu Y. High Fat Diet Induces Kidney Injury via Stimulating Wnt/β-Catenin Signaling. Front Med (Lausanne) 2022;9:1–9. https://doi.org/10.3389/fmed.2022.851618.

[68] Zhao X, Chen X, Zhang Y, George J, Cobbs A, Wang G, et al. Kidney injury molecule-1 is upregulated in renal lipotoxicity and mediates palmitate-induced tubular cell injury and inflammatory response. Int J Mol Sci 2019;20. https://doi.org/10.3390/ijms20143406.

[69] Mori Y, Ajay AK, Chang JH, Mou S, Zhao H, Kishi S, et al. KIM-1 mediates fatty acid uptake by renal tubular cells to promote progressive diabetic kidney disease. Cell Metab 2021;33:1042-1061.e7. https://doi.org/10.1016/j.cmet.2021.04.004.

[70] Ko GJ, Rhee CM, Kalantar-Zadeh K, Joshi S. The effects of high-protein diets on kidney health and longevity. Journal of the American Society of Nephrology 2020;31:1667–79. https://doi.org/10.1681/ASN.2020010028.

[71] Liu H-W, Tsai W-H, Liu J-S, Kuo K-L. Association of Vegetarian Diet with Chronic Kidney Disease. Nutrients 2019;11:279. https://doi.org/10.3390/nu11020279.

[72] Franzke B, Bileck A, Unterberger S, Aschauer R, Zöhrer PA, Draxler A, et al. The plasma proteome is favorably modified by a high protein diet but not by additional resistance training in older adults: A 17-week randomized controlled trial. Front Nutr 2022;9. https://doi.org/10.3389/fnut.2022.925450.

[73] Tan L, Song X, Sun X, Wang N, Qu Y, Sun Z. ART3 regulates triple-negative breast cancer cell function via activation of Akt and ERK pathways. Oncotarget 2016;7:46589–602. https://doi.org/10.18632/oncotarget.10306.

[74] Tanaka M, Siemann DW. Gas6/Axl signaling pathway in the tumor immune microenvironment. Cancers (Basel) 2020;12:1–14. https://doi.org/10.3390/cancers12071850.

[75] Hagood JS. Thy-1 as an Integrator of Diverse Extracellular Signals. Front Cell Dev Biol 2019;7:1–5. https://doi.org/10.3389/fcell.2019.00026.

[76] Wee P, Wang Z. Epidermal growth factor receptor cell proliferation signaling pathways. Cancers (Basel) 2017;9:1–45. https://doi.org/10.3390/cancers9050052.

[77] Pan J hua, Zhou H, Cooper L, Huang J lian, Zhu S bin, Zhao X xu, et al. LAYN is a prognostic biomarker and correlated with immune infiltrates in gastric and colon cancers. Front Immunol 2019;10:1–12. https://doi.org/10.3389/fimmu.2019.00006.

[78] Pinette JA, Mao S, Millis BA, Krystofiak ES, Faust JJ, Tyska MJ. Brush border protocadherin CDHR2 promotes the elongation and maximized packing of microvilli in vivo. Mol Biol Cell 2019;30:108–18. https://doi.org/10.1091/mbc.E18-09-0558.

[79] Yu X, Kong Y, Dore LC, Abdulmalik O, Katein AM, Zhou S, et al. An erythroid chaperone that facilitates folding of α-globin subunits for hemoglobin synthesis. Journal of Clinical Investigation 2007;117:1856–65. https://doi.org/10.1172/JCI31664.

Table 1: Baseline characteristics of white British and British Indian participants by diet groups in UK Biobank.

	White British participants						British Indian participants	
	Regular meat eaters N=23,243	Low meat eaters N=23,472	Poultry eaters N=486	Fish eaters N=1,081	Vegetarians N=721	Vegans N=54	Meat eaters N=391	Vegetarians N=167
Age at recruitment	57.2 (8.2)	57.2 (8.0)	56.6 (8.1)	54.4 (8.2)	52.9 (8.1)	55.0 (7.8)	53.9 (8.6)	54.6 (7.8)
Women	9,852 (42.4%)	14,962 (63.7%)	390 (80.2%)	762 (70.5%)	483 (67.0%)	28 (51.9%)	171 (43.7%)	105 (62.9%)
Men	13,391 (57.6%)	8,510 (36.3%)	96 (19.8%)	319 (29.5%)	238 (33.0%)	26 (48.1%)	220 (56.3%)	62 (37.1%)
Fasting time (hours)	3.8 (2.5)	3.7 (2.3)	3.8 (2.2)	3.6 (2.3)	3.7 (2.4)	3.8 (3.0)	4.1 (2.4)	4.1 (2.5)
Body mass index (kg/m^2)	28.0 (4.8)	27.1 (4.7)	25.3 (4.6)	25.3 (4.2)	25.3 (4.4)	25.2 (6.3)	27.1 (4.5)	27.0 (4.5)
Alcohol consumption <1g/d	3,618 (15.6%)	4,573 (19.5%)	152 (31.3%)	255 (23.6%)	171 (23.7%)	27 (50.0%)	157 (40.2%)	143 (85.6%)
Never smoked	11,785 (50.7%)	13,015 (55.4%)	269 (55.3%)	603 (55.8%)	421 (58.4%)	29 (53.7%)	291 (74.4%)	152 (91.0%)
Physical activity (MET hrs/wk)	34.9 (38.6)	33.8 (34.6)	41.5 (38.3)	37.9 (38.8)	34.5 (32.8)	38.9 (41.7)	29.7 (35.8)	22.2 (28.9)

Numbers shown are mean (SD) or number (%).

Figure 1: Volcano plots of proteins in white British vegetarians (top) and vegans (bottom) compared with regular meat eaters.

The red dotted line signifies *p*-value threshold for statistical significance. Each dot represents one protein, which were colour-coded by whether the protein is majority expressed (>50%) in one tissue type. Results were based on the multivariable model adjusted for age at recruitment, sex, region, fasting status, body mass index, alcohol consumption, smoking status and physical activity.

Figure 2: Top 10 proteins in vegetarians by diet group and ethnicity.

The top 10 proteins were selected by ranking the *p*-values of proteins comparing white British vegetarians with regular meat eaters, and sorted by betas in white British vegetarians, where the betas represent SD differences. Results were based on the multivariable model adjusted for age at recruitment, sex, region, fasting status, body mass index, alcohol consumption, smoking status and physical activity.

Figure 3: Top 10 proteins in vegans by diet group and ethnicity.

The top 10 proteins were selected by ranking the *p*-values of proteins comparing white British vegans with regular meat eaters, and sorted by betas in white British vegans, where the betas represent SD differences. Results were based on the multivariable model adjusted for age at recruitment, sex, region, fasting status, body mass index, alcohol consumption, smoking status and physical activity. The results for GUCA2A, FGF21, FOLR1, CKB and IGFBP2 are the same as shown in Figure 2.