# Somatomotor-Visual Resting State Functional Connectivity Increases After Two Years in the UK Biobank Longitudinal Cohort

Anton Orlichenko[a], Kuan-Jui Su[b], Qing Tian[b], Hui Shen[b], Hong-Wen Deng[b], and Yu-Ping Wang[a]

[a]Department of Biomedical Engineering, Tulane University, New Orleans, LA, USA

[b]Center for Biomedical Informatics and Genomics, School of Medicine, Tulane University, New Orleans, LA, USA

## ABSTRACT

Functional magnetic resonance imaging (fMRI) and functional connectivity (FC) have been used to follow aging in both children and older adults. Robust changes have been observed in children, where high connectivity among all brain regions changes to a more modular structure with maturation. In older adults, prior work has identified changes in connectivity associated with the default mode network (DMN); other work has used brain age to predict pre-clinical Alzheimer's disease. In this work, we find an increasing connectivity between the Somatomotor (SMT) and Visual (VIS) Networks using the Power264 atlas in a longitudinal cohort of the UK Biobank (UKB). This cohort consists of 2,722 subjects, with scans being taken an average of two years apart. The average connectivity increase between SMT-VIS is 6.8% compared to the younger scan baseline (from $\rho = 0.39$ to $\rho = 0.42$), and occurs in male, female, older subject ($> 65$ years old), and younger subject ($< 55$ years old) groups. Among all inter-network connections, this average SMT-VIS connectivity is the best predictor of relative scan age, accurately predicting which scan is older 57% of the time. Using the full FC and a training set of 2,000 subjects, one is able to predict which scan is older 82.5% of the time when using the difference of FC between the two scans as input to a classifier. This previously under-reported relationship may shed light on normal changes in aging brain FC, identifies a potential confound for longitudinal studies, and proposes a new area for investigation, specifically the SMT-VIS connectivity.

**Keywords:** fMRI, functional connectivity, UK Biobank, longitudinal, aging, visual network, somatomotor network, Power atlas

# 1. INTRODUCTION

Functional magnetic resonance imaging (fMRI) is a non-invasive technique that has proven indispensable for investigating human neural processes in vivo.[1] For example, it has been used to localize the areas associated with vision,[2] attention,[34] emotion,[567] and language[8] to specific regions in the cortex, or at least find the regions that are most significantly involved in a specific task. Functional connectivity (FC) is a quantity derived from fMRI that measures the time correlation of blood oxygen level-dependent (BOLD) signal between different regions in the brain.[9] FC has recently been used to predict age,[1011] sex,[1213] race,[14] psychiatric disease status,[1516] and pre-clinical Alzheimer's disease.[17] Efforts to predict general fluid intelligence, although common,[1813] are thought by some to be confounded by differential achievement score distribution among ethnicities and the strong presence of race signal in FC.[14] FC has proven effective in predictive studies because of its simplicity and its robust representation of complex BOLD signal activity, as evidenced by high subject identifiability across different scanner tasks and across time.[192021]

Besides being used as a predictive tool, FC has been observed to undergo changes throughout the lifespan. For example, connectivity in young children is generally very high between all brain regions and decreases while also becoming more modularized during and after puberty.[22] The FC of males and females is also quantitatively different, with females having higher intra-DMN connectivity and males having relatively greater connectivity between the DMN and other networks, although there is a wide degree of individual variation.[2324] Meanwhile, studies have shown that changes occur in the DMN during late middle and old age,[25] although the exact direction of change in FC does not always appear constant.[26] In addition, various studies have examined age-related changes in the cingulum[27] and medial temporal lobe.[28] Given the recent interest in using fMRI to predict pre-clinical Alzheimer's disease,[2617] we believe a knowledge of ordinary changes in FC during old age is essential. This is especially true because it has been shown that a confounder can easily be mistaken for a true signal indicative of, e.g., general fluid intelligence or achievement score.[14]

This study uses the longitudinal cohort of the UKB[29] to examine changes in the FC of individuals after an average of two years, the time between longitudinal scans. The UKB population of subjects with fMRI scans is predominantly (98%) Caucasian, ruling out race as a possible confounding effect. Additionally, we investigate changes in FC in longitudinal sub-populations based on subject age and sex. We find that average FC between SMT-VIS networks increases on average from the first scan to the second, and that SMT and VIS-related connectivities are more predictive of scan age than those of other networks. The complete FC, or a large subset, is still required to attain the best accuracy.

## 2. METHODS

We first describe the UKB dataset and the longitudinal subset used for our analysis. We then describe pre-processing of the fMRI data and conversion into FC. Finally, we discuss prediction of older vs younger scan in the longitudinal cohort and detail our methods for analysis of FC changes.

### 2.1 UK Biobank Longitudinal Cohort

The UKB contains various data of more than 500,000 subjects in the UK, of who more than 40,000 have fMRI scans.[29] We processed two longitudinal resting state scans for 2,722 subjects, taken approximately two years apart. These subjects are approximately equally split between male and female, and have significant numbers of younger and older adults (see Figure 3). The longitudinal cohort is composed of 1,289 genetic males and 1,369 genetic females, with the rest not having genetic sex information. The ethnicity of the subset of the UKB with fMRI scans is 98% Caucasian. Besides the 2,722 subjects we processed, an additional 154 subjects have the second longitudinal scan but not the first, resulting either from missing original source data or a failure in our SPM-based preprocessing pipeline.[*]

### 2.2 fMRI Preprocessing

The original scan acquisition parameters are described elsewhere,[30][31] but consist of both resting state and task fMRI scans with a repetition time of TR = 0.735 sec. For this study, we examined the resting state scans only. All resting state 4D fMRI volumes were processed with SPM12 including co-registration and warping to MNI space.[*] BOLD signal was extracted using the Power264 atlas,[32] which consists of 264 ROIs grouped into 14 functional networks and represented by 5mm radius spheres. The resulting timeseries were bandpass filtered between 0.01 and 0.15 Hz to remove scanner drift, noise, heartbeat, and some breathing signal. Pearson correlation of the filtered timeseries was used to create subject-specific FC matrices, which were reduced to the unique entries in the upper right triangle and vectorized. The entire procedure is summarized in Figure 1.

In contrast to the Power264 atlas-derived FC constructed by us, the original UKB data provided the unique part of 21-region and 55-region FC and partial correlation-based connectivity (PC) matrices based on ICA in vectorized format.[30] These matrices were calculated through the use of PCA on whole cohort fMRI data followed by ICA,[30] meaning that regions overlap in an unpredictable way and are not associated with specific functional networks. Although prediction using 55-component ICA-based FC and PC is often as good as and sometimes better than prediction using Power264 atlas-derived FC, the resulting connectivities are uninterpretable with regards to BOLD signal within specific regions. Additionally, in predicting which scan is older, Power264 asymptotes to a higher predictive accuracy than either of the ICA-derived measures (see Figure 4).

---

[*]http://www.fil.ion.ucl.ac.uk/spm/software/spm12/

Figure 1. Preprocessing pipeline for converting 4D fMRI volumes into FC using the Power264 atlas.[32] Reproduced from Orlichenko et al. (2023).[33]

## 2.3 Prediction of Scan Order and Analysis of FC

Prediction of scan age in the UKB longitudinal cohort was carried out by logistic regression[†] models with 20 bootstrapping repetitions, using the scikit-learn implementation.[34] The regularization parameter was fixed to $C = 1$, which was found to be near the optimal value for all training set sizes using grid search. It was found that a simple difference of scan FCs gave the best prediction results compared to concatenation or difference and concatenation, using either logistic regression or MLP. The training set was created with randomization of whether older scan was subtracted from younger scan or younger scan was subtracted from older scan. Our code for computing prediction accuracy can be found online.[‡] However, UKB data sharing policy precludes us from posting the longitudinal data itself; interested researchers may contact us with any questions.

Analysis of FC was performed by finding the mean (Figure 2) and standard deviation (Figure 7) of older scan FC minus younger scan FC for the longitudinal cohort. Additionally, prediction of scan order was carried out using the average connectivity between each of the Power264 networks, each network consisting of many individual ROIs. As before, logistic regression[†] with 20 bootstrap repetitions and $C = 1$ was used for this purpose. A Bonferroni-corrected two-sided t-test was applied to the 105 average inter-network connectivity differences (from the complete graph of 14 functional networks) of the 2,722 longitudinal subjects to determine if they were significantly different from zero (Figure 5 Bottom).

---

[†]https://scikit-learn.org/stable/

[‡]https://github.com/aorliche/ukb-longitudinal-smt-vis

**Functional Networks**

| Label | ROIs | | Label | ROIs | |
|---|---|---|---|---|---|
| 0 | 0-29 | Somatomotor Hand (SMT) | 7 | 156-180 | Frontoparietal (FRNT) |
| 1 | 30-34 | Somatomotor Mouth (SMT) | 8 | 181-198 | Salience (SAL) |
| 2 | 35-48 | Cinguloopercular (CNG) | 9 | 199-211 | Subcortical (SUB) |
| 3 | 49-61 | Auditory (AUD) | 10 | 212-220 | Ventral Attention (VTRL) |
| 4 | 62-119 | Default Mode (DMN) | 11 | 221-231 | Dorsal Attention (DRSL) |
| 5 | 120-124 | Memory (MEM) | 12 | 232-235 | Cerebellar (CB) |
| 6 | 125-155 | Visual (VIS) | 13 | 236-263 | Uncertain (UNK) |

Table 1. Regions, abbreviations, and labels in the Power264 atlas.

## 3. RESULTS

We first describe trends in FC changes during the average of 2 years between longitudinal scans, summarize the ability of simple machine learning models to identify older vs younger scan, and finally investigate the ability of specific inter-network connectivities to predict scan order. Finally, we summarize the statistical significance of inter-network FC changes with aging.

### 3.1 Inter-Network FC Changes

In Figure 2, we show that, on average, SMT-VIS connectivity increases from younger scan to older scan. The right hand side of Figure 2 displays divisions of the 14 functional networks included in the Power264 atlas. Network labels and abbreviations are listed in Table 1. The increase in connectivity is large and distinct over the majority of SMT-VIS FCs compared to other non-SMT and non-VIS FCs. Many FCs involving the VIS network appear to increase in connectivity from the first scan to the second. The average change in FC in the SMT-VIS connection is 6.8%, corresponding to a mean change $\mu_{\Delta\rho} = +0.03$, compared to a standard deviation of $\sigma_{\Delta\rho} = 0.26$. Figure 5 shows that, although small compared to the standard deviation, this difference is very significant.

Figure 3 displays the same analysis, i.e., the average change from first scan to second, for four subsets of the cohort. These subsets are male subjects, female subjects, young ($< 55$ years old) subjects, and old ($> 65$ years old) subjects. All four subsets observed the same effect as the whole cohort, thus we rule out very old age or gender as confounding factors.

Figure 2. Difference in FC calculated by older scan minus younger scan, averaged over all 2,722 longitunidal cohort subjects. There are significant average differences in SMT-VIS connectivity (labeled 0-6). The same plot is displayed on the left and right, with Power264 network divisions on the right hand side. Network labels can be found in Table 1.

## 3.2 Predicting Older Scan of Pair

In Figure 4, one can see that is possible to predict which scan of a longitudinal pair is older with the Power264 atlas at an accuracy of 82.5%, having 2,000 subjects in the training set and the rest in the test set. This measurement was repeated with 20 bootstrap iterations and averaged. The entire 34,716-feature upper right triangle of the FC matrix was used ot make the prediction. One can also see that the ICA FC/PC matrices provided pre-processed along with the UKB data are also able to predict scan order, although at a slightly reduced accuracy. Prediction is possible at an accuracy of 60-70% using only 100-200 training set subjects.

## 3.3 Prediction of Older Scan Using Specific Inter-Network Connections

In Figure 5, we rank average inter-network FCs in their ability to predict scan order. As expected from the mean change in FC (Figure 2), the SMT-VIS connection is the most predictive of longitudinal scan age. Furthermore, SMT and VIS networks are among the next several most predictive inter-network connections. In Figure 5 bottom, we plot the predictive ability of all 105 inter-network connections, along with a p-value for the inter-network FC change being significantly different from zero. The raw p-value has been multiplied by 105 to account for multiple comparisons. It is highly significant for the first 10 or so most predictive inter-network connections.

Table 2 lists the number of subjects whose FC increased or decreased for the SMT-VIS connection and over the entire brain. The table is divided among the four subsets of the longitudinal cohort mentioned previously.

Figure 3. Significant increase in SMT-VIS connectivity after an average of 2 years in the UKB longitudinal cohort appears in male, female, younger, and older groups, and seems to be an invariant feature of FC change in the longitudinal UKB cohort.

Figure 4. Capability of predicting older scan of a pair based on the difference of FC between the two scans, as a function of the number of training subjects. Three inputs are used: 55 component ICA FC, 55 component ICA PC, and the Power264 atlas FC. Prediction accuracy asymptotes at 82.5%.

| Group | +SMT-VIS FC | -SMT-VIS FC | +Total FC | -Total FC | Total Subjects |
|---|---|---|---|---|---|
| Male | 778 (60.4%) | 511 | 690 (53.5%) | 599 | 1289 |
| Female | 741 (54.1%) | 628 | 671 (49.0%) | 698 | 1369 |
| < 55 years old | 269 (59.0%) | 187 | 249 (54.6%) | 207 | 456 |
| > 65 years old | 577 (56.5%) | 445 | 520 (50.1%) | 502 | 1022 |

Table 2. Number of subjects in the longitudinal cohort increasing and descreasing in average FC within the SMT-VIS connection and within the whole brain.

Additionally, we correlated several dozen subject phenotypes and longitudinally-tracked variables with changes in FC and report the most significant in Supplementary Materials Section B. In that section we find an interesting but small correlation with hand grip strength, body mass index (BMI), and basal metabolic rate.

## 4. DISCUSSION

Farràs-Permanyer et al. (2019)[35] find that mean resting state FC may increase throughout the entire brain for the oldest subject (> 80 years old) group. As shown in Supplementary Section A, we confirm a small, statistically insignificant increase in total longitudinal FC in the healthy controls of the ADNI dataset,[36] another elderly population with multiple longitudinal fMRI scans. Other studies have generally focused on examining connectivity in the DMN associated with aging.[37][38] These studies find areas of increased connectivity as well

Figure 5. Ability to predict which scan of a subject is older based on average connectivity between regions. Top: best and worst inter-network connectivities for prediction. Bottom: Prediction accuracy for all 105 inter-network connectivities. We find that SMT-VIS connectivity has the maximum predictive ability of all regions at 57%. In general, network-level connectivities involving SMT and VIS networks have higher predictive ability compared to other regions. The dashed orange line displays negative log base 10 of the Bonferroni-corrected p-value for significance of FC change between scans.

Power264 Atlas Visual and Somatomotor Regions

Figure 6. SMT and VIS regions in the Power264 atlas.

as areas of decreased connectivity. There are two problems with such studies. First, they are for the most part cross-sectional and do not follow a single subject across a multi-year period. Second, they mostly use small numbers of subjects, the majority of studies enrolling fewer than 50, making it impossible to identify small effects. On the other hand, one study performed on a cohort of more than 2,000 older subjects in Rotterdam found age-related changes in connectivity to be complicated, drawing no firm conclusions.[39] We note that the Rotterdam study was not longitudinal but cross-sectional.

We conjecture the fact that most studies only focus on DMN and report decreased connectivity[37] in aging populations may be related to the large number of ROIs in the DMN and an implicit bias inherent in the word "connectivity." Naturally, as we reach very old age we expect physical connections to degenerate, not become stronger. In fact, FC is really the synchronization of BOLD signal between regions, and does not imply a direct physical link between regions. Young children are known to have higher average FC than young adults;[22,40] thus older subjects may been as reverting to a less optimal state as they age.

On the other hand, as we describe in Supplementary Materials Section B, physical observables such as hand grip strength in the UKB longitudinal cohort are weakly correlated with an increase in FC in SMT-CB and VIS-CB connectivity. Additionally, we find BMI and basal metabolic rate are weakly correlated with the longitudinal increase in SMT-VIS connectivity (see Supplementary Section B). This may suggest a small health related effect that is found throughout the study cohort and includes male, female, younger, and older subjects.

We show in this work that the average connectivity increase in the SMT-VIS connection is small but highly

Figure 7. Standard deviation of difference between older scan FC and younger scan FC in the UKB longitudinal cohort. Note that the magnitude of average standard deviation (0.26) of SMT-VIS connectivity change is large compared to mean SMT-VIS connectivity change (0.03). However, we show in Figure 5 that this connectivity change is highly statistically significant. The smallest average standard deviations are found in SMT-SMT (0.2) and VIS-VIS (0.22) connectivities.

statistically significant. The average change in FC in this connection is only 6.8%, corresponding to a mean change $\mu_{\Delta\rho} = +0.03$, compared to a standard deviation of change from subject to subject of $\sigma_{\Delta\rho} = 0.26$ (see Figure 7). However, using our longitudinal sample of 2,722 subjects, we find the average connectivity change from younger scan to older scan is significant as level of $p < 10^{-15}$ after Bonferroni correction for multiple comparisons (Figure 5). Finding such small effects is helped by the use of large number of subjects and longitudinal data.

## 5. CONCLUSION

In this work, we pre-process a 2,722 subject longitudinal subset of the UK Biobank dataset and examine FC using the Power264 atlas. We find that in scans taken an average of two years apart, the average functional connectivity between SMT and VIS network regions tends to increase. This occurs in male, female, younger ($< 55$ years old), and older ($> 65$ years old) subjects. We verify the ability of this average FC increase to predict scan ordering using simple machine learning models. The identification of an increase in connectivity with non-pathological aging, in a longitudinal cohort and specifically in the SMT-VIS synchronization of BOLD

signal, may lead to novel insights about brain function in old age. Additionally, it identifies an effect that could possibly show up as a confounder in studies of dementia or neurodegenerative diseases.

## 6. ACKNOWLEDGEMENTS

The authors would like to acknowledge the UK Biobank (application ID 61915) for the gathering and curation of a large and important dataset accessible to researchers the world over, as well as the curators of the ADNI dataset for their work on neurodegenerative diseases.

The authors would also like to acknowledge the NIH (grants R01 GM109068, R01 MH104680, R01 MH107354, P20 GM103472, R01 EB020407, R01 EB006841, R56 MH124925) and NSF (grant #1539067) for partial funding support.

This research was supported in part using high performance computing (HPC) resources and services provided by Information Technology at Tulane University, New Orleans, LA.

## REFERENCES

[1] Belliveau, J. W., Kennedy, D. N., McKinstry, R. C., Buchbinder, B. R., Weisskoff, R. M., Cohen, M. S., Vevea, J. M., Brady, T. J., and Rosen, B. R., "Functional mapping of the human visual cortex by magnetic resonance imaging.," *Science* **254 5032**, 716–9 (1991).

[2] Cox, D. D. and Savoy, R. L., "Functional magnetic resonance imaging (fMRI) "brain reading": detecting and classifying distributed patterns of fMRI activity in human visual cortex," *Neuroimage* **19**, 261–270 (June 2003).

[3] Coull, J. T. and Nobre, A. C., "Where and when to pay attention: the neural systems for directing attention to spatial locations and to time intervals as revealed by both PET and fMRI," *J. Neurosci.* **18**, 7426–7435 (Sept. 1998).

[4] Pugh, K. R., Shaywitz, B. A., Shaywitz, S. E., Fulbright, R. K., Byrd, D., Skudlarski, P., Shankweiler, D. P., Katz, L., Constable, R. T., Fletcher, J., Lacadie, C., Marchione, K., and Gore, J. C., "Auditory selective attention: An fMRI investigation," *Neuroimage* **4**, 159–173 (Dec. 1996).

[5] Phan, K. L., Wager, T., Taylor, S. F., and Liberzon, I., "Functional neuroanatomy of emotion: a meta-analysis of emotion activation studies in PET and fMRI," *Neuroimage* **16**, 331–348 (June 2002).

[6] Ochsner, K. N., Bunge, S. A., Gross, J. J., and Gabrieli, J. D. E., "Rethinking feelings: An fmri study of the cognitive regulation of emotion," *Journal of Cognitive Neuroscience* **14**(8), 1215–1229 (2002).

[7] Koelsch, S., Fritz, T., Cramon, V, D. Y., Müller, K., and Friederici, A. D., "Investigating emotion with music: an fMRI study," *Hum. Brain Mapp.* **27**, 239–250 (Mar. 2006).

[8] Hernandez, A. E., Dapretto, M., Mazziotta, J., and Bookheimer, S., "Language switching and language representation in Spanish-English bilinguals: an fMRI study," *Neuroimage* **14**, 510–520 (Aug. 2001).

[9] van den Heuvel, M. P. and Hulshoff Pol, H. E., "Exploring the brain network: a review on resting-state fMRI functional connectivity," *Eur. Neuropsychopharmacol.* **20**, 519–534 (Aug. 2010).

[10] Orlichenko, A., Qu, G., Zhang, G., Patel, B., Wilson, T. W., Stephen, J. M., Calhoun, V. D., and Wang, Y.-P., "Latent similarity identifies important functional connections for phenotype prediction," *IEEE Transactions on Biomedical Engineering* **70**(6), 1979–1989 (2023).

[11] Hu, W., Cai, B., Zhang, A., Calhoun, V. D., and Wang, Y.-P., "Deep collaborative learning with application to the study of multimodal brain development," *IEEE Transactions on Biomedical Engineering* **66**(12), 3346–3359 (2019).

[12] İçer, S., İrem Acer, and Baş, A., "Gender-based functional connectivity differences in brain networks in childhood," *Computer Methods and Programs in Biomedicine* **192**, 105444 (2020).

[13] Sen, B. and Parhi, K. K., "Predicting biological gender and intelligence from fmri via dynamic functional connectivity," *IEEE Transactions on Biomedical Engineering* **68**(3), 815–825 (2021).

[14] Orlichenko, A., Daly, G., Liu, A., Shen, H., Deng, H.-W., and Wang, Y.-P., "ImageNomer: developing an fMRI and omics visualization tool to detect racial bias in functional connectivity," (2023).

[15] Wang, S., Zhan, Y., Zhang, Y., Lyu, L., Lyu, H., Wang, G., Wu, R., Zhao, J., and Guo, W., "Abnormal long- and short-range functional connectivity in adolescent-onset schizophrenia patients: A resting-state fMRI study," *Progress in Neuro-Psychopharmacology and Biological Psychiatry* **81**, 445–451 (Feb. 2018).

[16] Rashid, B., Arbabshirani, M. R., Damaraju, E., Cetin, M. S., Miller, R., Pearlson, G. D., and Calhoun, V. D., "Classification of schizophrenia and bipolar patients using static and dynamic resting-state fMRI brain connectivity," *Neuroimage* **134**, 645–657 (July 2016).

[17] Millar, P. R., Luckett, P. H., Gordon, B. A., and Benzinger, T. L., "Predicting brain age from functional connectivity in symptomatic and preclinical alzheimer disease," *NeuroImage* **256**, 119228 (2022).

[18] Qu, G., Xiao, L., Hu, W., Wang, J., Zhang, K., Calhoun, V. D., and ping Wang, Y., "Ensemble manifold regularized multi-modal graph convolutional network for cognitive ability prediction," *IEEE Transactions on Biomedical Engineering* **68**, 3564–3573 (2021).

[19] Finn, E. S., Shen, X., Scheinost, D., Rosenberg, M. D., Huang, J., Chun, M. M., Papademetris, X., and Constable, R. T., "Functional connectome fingerprinting: identifying individuals using patterns of brain connectivity," *Nat. Neurosci.* **18**, 1664–1671 (Nov. 2015).

[20] Cai, B., Zhang, G., Hu, W., Zhang, A., Zille, P., Zhang, Y., Stephen, J. M., Wilson, T. W., Calhoun, V. D., and Wang, Y.-P., "Refined measure of functional connectomes for improved identifiability and prediction," *Hum. Brain Mapp.* **40**, 4843–4858 (Nov. 2019).

[21] Orlichenko, A., Qu, G., Zhou, Z., Ding, Z., and Wang, Y.-P., "Angle basis: A generative model and decomposition for functional connectivity," (2023).

[22] Dosenbach, N. U. F., Nardos, B., Cohen, A. L., Fair, D. A., Power, J. D., Church, J. A., Nelson, S. M., Wig, G. S., Vogel, A. C., Lessov-Schlaggar, C. N., Barnes, K. A., Dubis, J. W., Feczko, E., Coalson, R. S., Pruett, Jr, J. R., Barch, D. M., Petersen, S. E., and Schlaggar, B. L., "Prediction of individual brain maturity using fMRI," *Science* **329**, 1358–1361 (Sept. 2010).

[23] Mak, L. E., Minuzzi, L., MacQueen, G., Hall, G., Kennedy, S. H., and Milev, R., "The default mode network in healthy individuals: A systematic review and meta-analysis," *Brain Connect.* **7**, 25–33 (Feb. 2017).

[24] Ficek-Tani, B., Horien, C., Ju, S., Xu, W., Li, N., Lacadie, C., Shen, X., Scheinos, D., Constable, T., and Fredericks, C., "Sex differences in default mode network connectivity in healthy aging adults," *Cereb. Cortex* (Dec. 2022).

[25] Fjell, A. M., Sneve, M. H., Grydeland, H., Storsve, A. B., Amlien, I. K., Yendiki, A., and Walhovd, K. B., "Relationship between structural and functional connectivity change across the adult lifespan: A longitudinal investigation," *Hum. Brain Mapp.* **38**, 561–573 (Jan. 2017).

[26] Staffaroni, A. M., Brown, J. A., Casaletto, K. B., Elahi, F. M., Deng, J., Neuhaus, J., Cobigo, Y., Mumford, P. S., Walters, S., Saloner, R., Karydas, A., Coppola, G., Rosen, H. J., Miller, B. L., Seeley, W. W., and Kramer, J. H., "The longitudinal trajectory of default mode network connectivity in healthy older adults varies as a function of age and is associated with changes in episodic memory and processing speed," *J. Neurosci.* **38**, 2809–2817 (Mar. 2018).

[27] Hirsiger, S., Koppelmans, V., Mérillat, S., Liem, F., Erdeniz, B., Seidler, R. D., and Jäncke, L., "Structural and functional connectivity in healthy aging: Associations for cognition and motor behavior," *Hum. Brain Mapp.* **37**, 855–867 (Mar. 2016).

[28] Salami, A., Wåhlin, A., Kaboodvand, N., Lundquist, A., and Nyberg, L., "Longitudinal evidence for dissociation of anterior and posterior MTL resting-state connectivity in aging: Links to perfusion and memory," *Cereb. Cortex* **26**, 3953–3963 (Oct. 2016).

[29] Sudlow, C., Gallacher, J., Allen, N., Beral, V., Burton, P., Danesh, J., Downey, P., Elliott, P., Green, J., Landray, M., Liu, B., Matthews, P., Ong, G., Pell, J., Silman, A., Young, A., Sprosen, T., Peakman, T., and Collins, R., "UK biobank: An open access resource for identifying the causes of a wide range of complex diseases of middle and old age," *PLoS Med.* **12**, e1001779 (Mar. 2015).

[30] Smith, S. M., Alfaro-Almagro, F., and Miller, K. L., "Uk biobank brain imaging documentation," tech. rep., UK Biobank (September 2022).

[31] Alfaro-Almagro, F., Jenkinson, M., Bangerter, N. K., Andersson, J. L. R., Griffanti, L., Douaud, G., Sotiropoulos, S. N., Jbabdi, S., Hernandez-Fernandez, M., Vallee, E., Vidaurre, D., Webster, M., McCarthy, P., Rorden, C., Daducci, A., Alexander, D. C., Zhang, H., Dragonu, I., Matthews, P. M., Miller, K. L., and Smith, S. M., "Image processing and quality control for the first 10,000 brain imaging datasets from UK biobank," *Neuroimage* **166**, 400–424 (Feb. 2018).

[32] Power, J. D., Cohen, A. L., Nelson, S. M., Wig, G. S., Barnes, K. A., Church, J. A., Vogel, A. C., Laumann, T. O., Miezin, F. M., Schlaggar, B. L., and Petersen, S. E., "Functional network organization of the human brain," *Neuron* **72**, 665–678 (Nov. 2011).

[33] Orlichenko, A., Qu, G., Su, K.-J., Liu, A., Shen, H., Deng, H.-W., and Wang, Y.-P., "Identifiability in Functional Connectivity May Unintentionally Inflate Prediction Results," (8 2023).

[34] Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D., Brucher, M., Perrot, M., and Duchesnay, E., "Scikit-learn: Machine learning in Python," *Journal of Machine Learning Research* **12**, 2825–2830 (2011).

[35] Farras-Permanyer, L., Mancho-Fora, N., Montalà-Flaquer, M., Bartrés-Faz, D., Vaqué-Alcázar, L., Peró-Cebollero, M., and Guàrdia-Olmos, J., "Age-related changes in resting-state functional connectivity in older adults," *Neural Regen. Res.* **14**(9), 1544 (2019).

[36] Petersen, R. C., Aisen, P. S., Beckett, L. A., Donohue, M. C., Gamst, A. C., Harvey, D. J., Jack, Jr, C. R., Jagust, W. J., Shaw, L. M., Toga, A. W., Trojanowski, J. Q., and Weiner, M. W., "Alzheimer's disease neuroimaging initiative (ADNI): clinical characterization," *Neurology* **74**, 201–209 (Jan. 2010).

[37] Sala-Llonch, R., Bartrés-Faz, D., and Junqué, C., "Reorganization of brain networks in aging: a review of functional connectivity studies," *Front. Psychol.* **6**, 663 (May 2015).

[38] Malagurski, B., Deschwanden, P. F., Jäncke, L., and Mérillat, S., "Longitudinal functional connectivity patterns of the default mode network in healthy older adults," *NeuroImage* **259**, 119414 (Oct. 2022).

[39] Zonneveld, H. I., Pruim, R. H. R., Bos, D., Vrooman, H. A., Muetzel, R. L., Hofman, A., Rombouts, S. A., van der Lugt, A., Niessen, W. J., Ikram, M. A., and Vernooij, M. W., "Patterns of functional connectivity in an aging population: The rotterdam study," *Neuroimage* **189**, 432–444 (Apr. 2019).

[40] Jolles, D. D., van Buchem, M. A., Crone, E. A., and Rombouts, S. A. R. B., "A comprehensive study of whole-brain functional connectivity in children and young adults," *Cereb. Cortex* **21**, 385–391 (Feb. 2011).

Figure 8. Mean longitudinal changes in FC between first scan and second scan in healthy controls of ADNI dataset.

## APPENDIX A. LONGITUDINAL FC CHANGES IN THE ADNI DATASET

We examined the longitudinal change in FC of healthy controls in the Alzheimer's Disease Neuroimaging Initiative (ADNI)[36] (age matched subjects who do not develop AD pathology). We used scans taken an average of one year apart. We confirm a small, statistically insignificant increase in total FC but fail to find the same SMT-VIS increase relative to the rest of FC as in the UKB. Statistics are given in Table 3 and the average FC change is shown in Figure 8.

| Average FC Increase | Std Dev FC Change | +Total FC | -Total FC | Total Subjects |
|---|---|---|---|---|
| 0.015 | 0.131 | 184 (52.7%) | 165 | 394 |

Table 3. There is a small but positive change in FC in ADNI healthy controls (subjects who do not go on to develop AD).

## APPENDIX B. CORRELATION OF CHANGE IN FC WITH LONGITUDINAL OUTCOMES IN THE UKB

We identified several correlations between longitudinal change in FC and changes in clinical outcomes associated with the two scan timepoints in the UKB dataset. These are presented belowm, along with the UKB field identifiers of the outcomes. All p-values are Bonferroni-corrected with $n = 105$ multiple comparisons (one for each average inter-network connectivity).

## B.1 SMT Hand, VIS, and CB Connectivity and Grip Strength (f.46.2.0, f.46.3.0, f.47.2.0, f.47.3.0)

We find a marginally significant association between change in hand grip strength and VIS-CB and SMT-CB connectivity change (Figure 9).

## B.2 Body Mass Index and Basal Metabolic Rate (f.23104.2.0, f.23104.3.0, f.23105.2.0, f.23105.3.0)

We find a not statistically significant but suggestive association between BMI and basal metabolic rate change and SMT-VIS connectivity change (Figure 10).

Figure 9. Hand grip strength change association with functional connectivity change in the longitudinal cohort. Bonferroni-corrected p-values.

Figure 10. BMI and basal metabolic range change association with functional connectivity change in the longitudinal cohort. Bonferroni-corrected p-values.