Participation Questionnaire for Preschoolers with Autism Spectrum Disorder: Item

Development

Takuto Nakamura[a]*, Sakumi Koyama[b], Hirofumi Nagayama[a], and Satoshi

Sasada[c]

[a]Faculty of Health and Social Services, School of Rehabilitation, Kanagawa University of Human

Services, Yokosuka, Japan

[b]Child Development Support Office, Smile-Kids, Nerima, Japan

[c]Graduate School of Health and Social Services, Kanagawa University of Human Services, Yokosuka,

Japan

*Corresponding author:

Takuto Nakamura

Kanagawa University of Human Services,

1-10-1 Heiseicho, Yokosuka, Kanagawa, Japan 238-8522

E-mail: nakamura-y49@kuhs.ac.jp

Word Count: 5300

Participation Questionnaire for Preschoolers with Autism Spectrum Disorder: Item

Development

Occupational therapists need to comprehensively assess participation of children

with autism spectrum disorder (ASD) in daily activities and evaluate the

effectiveness of relevant interventions. Several participation measurement tools

have been developed for children with ASD, but these tools require expert

involvement, which is a barrier to large-scale surveys. To address these concerns,

a caregiver-administered questionnaire—the Participation Questionnaire for

Preschoolers (PQP)—was developed. However, this tool could be improved due

to its narrow age range of 48–72 months and because the item development

process does not reflect the perspectives of children and caregivers. Therefore,

we expanded the PQP's target age range to 36–83 months and developed new

items that reflect the perspectives of professionals and caregivers. Interviews

were conducted with eight experts in supporting children with ASD and 11

caregivers of children with ASD. The interviews were transcribed, and a content

analysis was performed. The number of questions was reduced from 51 to 36,

and the order of items was changed for clarity. Two of the eight sub-domains

were removed to clarify the conceptual difference between activity and

participation. The updated version of the PQP has two unique features: (1) it can

be administered without expert involvement, and (2) it includes items specific to

the challenges faced by children with ASD. Future development of the scale and

validation of its measurement properties are needed.

Introduction

Autism spectrum disorder (ASD) is a neurodevelopmental disorder characterized by challenges related to social interaction, communication, and repetitive behaviors and interests [1]. Typically, ASD is evident early in an individual's development, particularly in childhood, and can considerably impact later life. Therefore, support in early childhood is vital, and significant financial resources are invested in high-income countries to manage ASD [2].

"Participation" is a construct defined in the International Classification of Functioning, Disability, and Health as "involvement in life situations" [3]. This construct, which implies involvement in academics, family roles, interactions with friends, and community activities, is an important goal for occupational therapists [4]. However, the disability characteristics of ASD create limitations to this participation [5]. Prior research suggests that children with ASD have limited participation in many domains compared to typically developing children, and that this limitation is more prominent in ASD than in other disorders [6, 7].

Occupational therapists have examined the use of standardized methods to assess the participation of children with ASD [8]. Many of the earliest participation measurement tools were generic tools not limited to a specific disability or disorder. These tools have helped to characterize participation across disability status and type and build a knowledge base for participation-focused occupational therapy practice. However, in recent years, there has been

a rapid expansion in the development of disability-specific participation measurement tools

that are highly useful in clinical settings and aptly reflect the effects of interventions. For

example, tools such as the "Activities Scale for Kids performance (ASKp) [9]" for measuring

participation in children with musculoskeletal disorders, and "Pediatric Measure of

Participation short forms [10]" for children with spinal cord injuries, have been developed

and utilized for various disabilities. This trend emphasizes the need to develop

disability-specific participation measurement tools for children with ASD [8].

The oldest existing disability-specific participation measurement tool for children

with ASD is the Matrix for Assessment of Activities and Participation [11]. This tool can be

used by professionals to comprehensively measure the activities and participation of children

with ASD aged 3–6 years. The Structured Preschool Participation Observation for Children

with ASD (SPO-ASD) can specifically measure participation in non-inclusive educational

settings [12]. The SPO-ASD is a valid and reliable tool designed for professionals to conduct

detailed assessments of ASD features in children aged 4-6 years. Additionally, the Picture My

Participation-Chinese version is a valid tool that can be used for children with ASD [13]. This

tool is advantageous because it allows children to self-report their participation through a

photo-based interview and can be used by children with ASD aged 5–9 years.

However, the use of these measurement tools necessitates trained experts to conduct interviews or observations. These requirements pose challenges not only for clinical application [14] but also for conducting large-scale surveys.

Given the constraints of these previously developed tools, the Participation Questionnaire for Preschoolers (PQP) was developed [15]. The PQP, developed through interviews with caregivers and professionals, is a participation measurement tool specifically designed for children with ASD. It is a caregiver-administered questionnaire comprising 54 items that comprehensively measure participation and can be answered without professional involvement. The development of a simple, standardized measurement tool such as the PQP can help clinicians easily screen for and monitor changes in participation. Furthermore, the questionnaire can serve as a basis for building evidence in support of participation of children with ASD in large-scale surveys.

However, development of the PQP items is ongoing. Furthermore, the target age range is limited to 48–72 months. Considering that ASD is often detected at approximately three years of age in high-income countries [2], the PQP's current target age range may not satisfy the needs of actual clinical practice and research. There is also a growing demand for new tools to assess and validate changes in participation and the effectiveness of interventions for children with ASD. Therefore, this study aims to expand the PQP's target age range and develop new items adaptable to children of diverse age groups.

Materials and methods

We used a qualitative methodology to develop the questionnaire items. To ensure that the constructs captured important indicators related to the participation of children with ASD, the item development steps of the COnsensus-based Standards for the selection of health Measurement Instruments (COSMIN) methodology for assessing the content validity of patient-reported outcome measures (PROMs)-user manual were used as a reference [16]. Three main steps were followed to develop the questionnaire. First, we determined the basic design of the questionnaire, including the constructs to be measured, the target population, and the response options. This study followed the PQP design used in previous studies [15], except for the target age group. Next, content experts (described below) reviewed the 54 items of the PQP [15] and participated in interviews to assess content validity. Semi-structured interviews were conducted to collect the perspectives of caregivers of children with ASD. Additionally, cognitive interviews were conducted with caregivers to judge whether the items were interpreted as the developers intended. The institutional review board of the first author's university approved the study, and all participants provided informed consent. This study was conducted between February and October 2021. It was, however, not prospectively registered in any public open access database.

Validation by experts

Questionnaire design

The PQP is based on the International Classification of Functioning (ICF), a comprehensive

framework developed by the World Health Organization (WHO) [3] to assess health and

disability [15]. The PQP was developed for professionals to assess the participation of children

with ASD in real-life situations and provide targeted interventions. A reflective model is

adopted in the PQP, and a diverse range of indicators is established to capture the overall

picture of "participation." The preliminary PQP items were developed in Japanese and

included eight tentative sub-areas based on the International Classification of Functioning,

Children, and Youth (ICF-CY) codes related to activity and participation. These sub-domains

are communication; mobility; self-care; domestic life; interpersonal interactions and

relationships; engagement in play; education; and community, social, and civic life [3]. To

evaluate participation in these eight domains, the PQP included both the dimensions of

performance and capacity, as defined by the ICF. Furthermore, to facilitate a more

comprehensive discussion of participation dimensions during interviews, the initial items

incorporated elements such as a child's enjoyment, frequency of engagement, degree of

involvement, and parental aspirations, based on findings from literature reviews [17, 18, 19].

This approach was intended to ensure that each item reflects various aspects of participation,

ultimately contributing to a comprehensive representation of participation as a whole.

Interviews were conducted with experts with at least five years of experience working

with children with ASD to validate the questionnaire content. We gathered feedback from

four occupational therapists, three researchers in occupational therapy theory, one

"participation" researcher, one physician, one speech and hearing therapist, one clinical

psychologist, one certified social worker, one childcare worker, and one physical therapist.

All participants had clinical experience in Japan. Based on their input, a 51-item draft of the

questionnaire was developed (Table 1). Response options were set on a 5-point scale: "agree"

(5 points), "somewhat agree" (4 points), "undecided" (3 points), "somewhat disagree" (2

points), and "disagree" (1 point). The total score was calculated by summing the scores for

each item. The target population for the PQP is preschool-aged ASD children with an

intelligence quotient (IQ) of 50 or higher who have no comorbidities other than

neurodevelopmental disorders [20]. Although the PQP's target age range was 48–72 months,

in this study, the items were modified to verify the content validity when the target age range

was expanded to 36–83 months.

Participants

Eight participants were included in the first phase of this study: three "participation"

researchers, one occupational therapist, one speech–language–hearing therapist, one clinical

psychologist, one certified social worker, and one childcare worker. The inclusion criteria for

selecting the participation researchers were that they had published peer-reviewed articles

that included the word "participation" in the ICF. Regarding the other professionals, selection

was based on them having had at least five years of experience working with children with

ASD in their practice. Using a convenience sampling method, participants were approached

by e-mail or telephone and informed in advance of the study's content, purpose, and

significance. Of the participants, a total of four were working at universities or other

educational/research institutions, one of whom was also working at a clinic, and three were

working at child development support facilities or centers, which are typical early support

services in Japan. No participant refused to participate in the study. All participants were

native Japanese speakers practicing in Japan. Details of participants are presented in Table 2.

Data collection

Individual semi-structured interviews were conducted with each participant using an online

conferencing tool. This format was chosen because there was concern that due to using an

online conferencing tool, it would not be possible to have smooth communication among

participants in interviews with multiple participants, such as focus group interviews. An

interview guide was used for the interviews, asking (1) what were the signs that ASD

children's participation was facilitated or limited, and (2) whether any items were

inappropriate or missing from the draft items developed. The first author (TN), a male

occupational therapist from a university rehabilitation department with experience in multiple

qualitative studies, conducted the interviews in Japanese. The second author (SK), a female

occupational therapist with experience in conducting multiple qualitative studies, was also

present in the interviews as an interview assistant. The interviews were recorded and

transcribed. The average duration of each interview was 93 minutes.

Data analysis

A qualitative content analysis was employed [21]. The analysis was conducted first, with

coding performed independently by the first and second authors. The unit of analysis was

defined as "mentions of item validity." In this study, validity was defined as any of the

following, based on the COSMIN definition of quality guidelines for PROMs [16]:

(1) "Relevance" indicates the relevance of the construct being measured (participation),

the target population (young children with ASD), the context of use (support for

children with ASD), and the appropriateness of the response options and recall

period.

(2) "Comprehensiveness" indicates the coverage of important concepts for measuring

constructs (participation).

(3) "Comprehensibility" indicates that respondents have an adequate understanding of

the instructions, items, and response options, and that the items are adequately

represented.

After each interview, the first and second authors searched the transcripts

independently for statements regarding item validity. The statement context was checked to

determine whether the statement made a clear reference to the question items. For unclear or

ambiguous references, we used the context before and after the statement as a basis for

judgment. Next, we determined which item the statement referred to, and assigned an item

number to the statement. When a statement referred to more than one question item, multiple

codes were extracted from the statement. For example, if a statement referred to both Items 1

and 2, two codes corresponding to each item were extracted. Furthermore, statements that did

not refer to a specific item but were related to the questionnaire's validity, such as those

suggesting a lack of items or a need to modify the instructions or questionnaire design, were

labeled with the code "other" and extracted. Each code was then deductively labeled for

relevance, comprehensiveness, or comprehensibility, based on the aforementioned definition

of validity. Finally, these codes were tied to the corresponding item number and participant. For

example, a reference to the comprehensibility of the 10^{th} item in the first interview was labeled

"1-10-Comprehensibility;" two references to the missing item in the second interview were

labeled "2-Other 1-Comprehensiveness" and "2-Other 2-Comprehensiveness."

After completing the coding, the first and second authors shared their analyses,

discussed any discrepancies, and finalized the codes. Following the coding decisions, the two

authors read the statements in the generated code; decided whether to modify, delete, or add

items (or directives or layouts); and created a draft of revised items. During this process, items

from existing scales were sometimes referenced [22, 23]. When new items were modified or

added, they were identified as either activity or participation codes with reference to the

ICF-CY. This item draft was used in the subsequent round of interviews.

After the interviews were completed, both the first and second authors inductively

categorized the codes. These categories grouped the reasons that modifications were deemed

necessary, as indicated in the responses. The analysis was conducted separately for codes

related to appropriateness and comprehensibility.

Validation by caregivers

Questionnaire design

The draft PQP items as modified in the first phase were used.

Participants

This study was conducted with participants who were caregivers of children with ASD.

Inclusion criteria were as follows: (1) Participants were caregivers of preschool children aged

36–83 months and with a diagnosis of ASD, (2) the children had no comorbid diagnoses other

than neurodevelopmental disorders in the Diagnostic and Statistical Manual of Mental

Disorders, Fifth Edition, and (3) the children had an IQ of 50 or higher. During study

recruitment, the authors requested referrals from child development support centers and

medical institutions in Japan and approached participants by e-mail and telephone. All

participants provided oral and written informed consent. No one refused to participate.

Participants were selected to cover variables such as the child's sex, age in months, comorbid

diagnoses, IQ, region of residence, family structure, the caregiver's sex, and employment

status. Specifically, we collected these variables based on reports from caregivers, and when

recruiting new participants, we continued to recruit participants in categories that had not yet

appeared in one or more variables. The final total number of participants was 11. Details of

the participants are presented in Table 3.

Data collection

Cognitive and semi-structured interviews were conducted using an online conferencing tool.

The interviews were conducted in Japanese by the first author (TN), who received two

months of training in cognitive interviewing. In the cognitive interviews, participants were

asked to use the "think-aloud method" in combination with the "verbal probing method" [24].

Specifically, the think-aloud method was used for each item in each section of the PQP: "First,

please describe your child's condition;" "Next, please describe your feelings;" and "Finally,

please describe your child's feelings." If the first author (TN) felt that there was a need for

further questions, additional open-ended questions were asked (e.g., "You mentioned that it

was difficult to make a judgment, but what specifically was difficult for you?"). In addition, a

semi-structured interview was conducted after the cognitive interview. To ensure that the

items for understanding participation were covered and that no inappropriate items were

included, participants were asked (1) whether the items covered aspects important to the lives

of the participants' children and their families; and (2) whether the drafted items were

irrelevant, inappropriate, or offensive to their daily lives. The second author (SK) was also

present during the interviews as an interview assistant. The interviews were recorded and

transcribed, and either the first or second author took notes for reference in the analysis. In

the analysis, participants who needed revisions for any item underwent a second interview to

determine if the issues were resolved by the modifications. Data collection was concluded

after the interviews with Participant 11, who followed three consecutive participants who did

not mention needing revisions, indicating that data saturation had been reached. The average

duration of each interview was 33 minutes.

Data analysis

The same analysis was performed as in Phase 1.

Results

Content validation by experts

A total of 120 changes were made to the proposed PQP questions, reducing the final number of

questions from 51 to 35. The changes comprised 80 modified questions, 28 deleted questions,

and 12 added questions. The sub-domain most frequently reported was "education," (35 times;

29.2%), followed by "self-care" (23 times; 19.2%), "interpersonal interaction and relationships"

(15 times; 12.5%), "community, social and civic life" (14 times; 11.7%), "engagement in play"

(13 times; 10.8%), "domestic life" (8 times; 6.7%), "communication" (7 times; 5.8%), and

"mobility" (4 times; 3.3%). One modification (0.8%) was also made to the overall design of the

questionnaire. Table 2 shows the number of occurrences of the relevance, comprehensiveness,

and comprehensibility codes in each interview.

Relevance

There were 30 relevance codes, accounting for 25% of all codes. When the relevance codes

were inductively categorized according to their reasons, the most frequently mentioned code

was "clarifying the difference between activity and participation" (n = 17, 56.7%).

Approximately 82.4% of these codes were attributed to Participant 1 (a "participation"

researcher). It was ascertained that all questions within the communication and mobility

domains pertained more to activity than participation, leading to the deletion of these two

domains. Additionally, Participant 1 referenced "clarifying the difference between

environment and participation" (n = 5) 16.7% of times.

I think the sub-domain of communication often asks about skills, so, well, if you ask if

it's about participation, I will say it's not really. (Participant 1)

In addition, "the meaning of the question changes depending on the context" (n = 7)

was mentioned by Participants 2 ("participation" researcher) and 3 (speech–language–hearing

therapist), accounting for 23.3% of the total. Another code, which indicated items that

highlighted caregivers' concerns, was identified: "raises concerns among caregivers." Codes

for appropriateness appeared in 93.3% of the total responses from Participant 3, after which

only two codes appeared from Participant 7.

Comprehensiveness

There were 14 codes for comprehensiveness, accounting for 11.7% of all codes. Half of the

codes (n = 7) were situated within the education domain.

I think it's important for children of the same age to have opportunities; for example, to

get together in a group like during a music festival or a play or a Christmas party and

enjoy the atmosphere. Whether it's possible to share time and place … I feel like this is

something important for participation. (Participant 5)

In response, a new item was added in the education domain: "Has opportunities to

enjoy reading with children of the same age" (later revised to "Enjoys class activities in

cooperation with children of the same age").

Comprehensibility

There were 76 comprehensibility codes, accounting for 63.3% of all codes. Comprehensibility

codes were inductively classified as "question structure is difficult to understand" (n = 41) for

46.1%, "topic is unclear" (n = 39) for 43.8%, "answer is difficult to determine" (n = 8) for 9.0%,

and "instructions are difficult to understand" (n = 1) for 1.1% (13 codes were overlapped into

two classifications). In particular, in the interview with Participant 2, it was suggested that the

items, which had been categorized into eight domains according to the ICF activity and

participation codes, be presented in the following three question formats: "answer the

following questions about your child," "describe how you feel," and "answer the question

about your child's feelings."

307   The subjects have been repeatedly "this child is" and "you are." This item comprises

308   two types of questions: "A question to answer about the child as understood from the

309   mother's perspective" and "A question for the mother to speculate from the child's

310   perspective." We must consider that changing the subject requires effort for people with

311   tendencies of neurodevelopmental disorders, and it could be burdensome for them to

312   answer. (Participant 2)

313   With this change, the items that had been arranged as eight domains were rearranged

314   into the three question formats described above.

Content validation by caregivers

A total of 49 changes were made to the PQP items, resulting in the final number of questions

changing from 35 to 36. There were 48 modifications, zero deletions, and one addition.

Interpersonal interaction and relationships were modified 20 times (40.8%); education, 13

times (26.5%); engagement in play, eight times (16.3%); community, social, and civic life, four

times (8.2%); self-care, two times (4.1%); and modifications regarding response options, two

times (4.1%). No modifications were made to "domestic life." The number of occurrences of

the relevance, comprehensiveness, and comprehensibility codes in each interview is shown in

Table 3; and the revised items, after the interviews were completed, are shown in Table 4.

Relevance

No issues of relevance were reported in the interviews. All participants stated that the items

were relevant to the child, and the questions were not deemed sensitive for respondents to

answer.

Comprehensiveness

Only one code for comprehensiveness was reported, accounting for 2.1% of the total. A mother

of a child within the highest quartile for age in the study noted that there were few questions

about relationships with friends, commenting:

"I have no problem changing clothes and daily habits, and I am worried about what is

going on with their friends, so I thought it would be okay if I had some more questions

like that." (Participant 3)

A new item—"Has close friends (talks a lot, plays together a lot, etc.)"—was added in

response.

Comprehensibility

Regarding comprehensibility, there were 48 references to question ambiguity and misleading

item order, accounting for 98.0% of the total. The codes were inductively categorized as

follows: "topic is unclear" (41.7%, n = 20), "question is difficult to recall" (18.8%, n = 9),

"answer is difficult to determine" (16.7%, n = 8), "question structure is difficult to understand"

(8.3%, n = 4), and "others" (14.9%, n = 7). The "others" included the content of the instructions,

the order in which the items were presented, and the placement of the response options

(applicable options on the left and non-applicable options on the right). For example, although

Participant 6 talked about her child's challenging behavior at mealtime in the semi-structured

interview, she answered "agree" to the item "family time seems to be fun." However, when she

was probed by the interviewer about whether she rarely had fun during family time, she replied

as follows.

When he doesn't get his way, there are quite a few times when he gets angry … But the

joyful look on my child's face is quite impressionable to me, so I ended up choosing

"agree" for this question. (Participant 6)

This item was judged to be a "question difficult to recall" and was revised to "always

appears to enjoy his time with his family."

Discussion

PQP item development and characteristics

This study aimed to extend the age range targeted by the PQP and to develop new items

suitable for a broader spectrum of children. The newly developed PQP items offer three

primary advantages. First, they enable the comprehensive collection of participation data

through a questionnaire, which is essential. The PQP was validated for its relevance,

comprehensiveness, and comprehensibility based on interviews with eight experts and eleven

caregivers of children with ASD until data saturation was achieved. Looking forward, if the

development of this scale is completed and it allows for assessing comprehensive participation

without requiring trained professionals, it could greatly improve the feasibility of conducting

large-scale surveys. Additionally, this could establish a foundation for gathering evidence

about the participation of children with ASD.

Second, the newly developed items reflect the perspectives of both professionals and

caregivers, highlighting the key challenges faced by children with ASD, such as difficulties

performing activities of daily living and accessing specialized services [2]. Disability-specific

measurement tools may be better suited to capture the information required for intervention

and monitor intervention effectiveness than generic instruments [8, 25]. The PQP could,

therefore, be a useful tool for occupational therapists to assess and monitor the participation of

children with ASD during interventions.

Third, this tool facilitates the assessment of a wider age range of children with ASD

than the previous PQP items enabled. The need for early intervention for children with ASD

has been widely recognized, and intervention implemented from infancy is becoming more

common in high-income countries [2]. Therefore, the PQP's expanded age range increases its

accessibility for practical use. It could also contribute to the feasibility of research on the early

childhood participation of children with ASD, such as by enabling longer-term longitudinal

studies.

Limitations and future research directions

The development of measurement tools requires item development, scale development,

and the verification of reliability and validity [26]. To make the PQP usable in practice and

research, these steps must be implemented in the future. Particularly, the selection of

questionnaire items will require item reduction analysis for scale development, and the

validation of content validity needs to be conducted separately after scale development.

COSMIN recommends validating content validity using qualitative research and also

recommends the use of quantitative approaches such as the Delphi method alongside the

qualitative methods as beneficial [16].

This study has some limitations. Between 2020 and 2021, when the data were collected,

many municipalities in Japan implemented travel restrictions owing to the COVID-19

pandemic [27]. This could have limited many people's diversity of events and travel-related

items. To address this issue, it may be useful to reexamine the content validity of the data by

comparing it with data collected during the non-pandemic period. In addition, the data

collected from caregivers included only participants who used child development support

services and medical institutions. Therefore, the participation status of children who did not use

these services may not have been adequately assessed. In future studies, it will be important to

assess the applicability of the PQP to children who do not use these services. This would enable

confirmation that the PQP adequately reflects the participation status of all children with ASD.

Finally, the content of the newly developed PQP items was validated only in the unique cultural

and institutional environment of Japan. Cross-cultural validation is required to use the PQP in

countries other than Japan.

Conclusion

This study expands the target age range of the PQP—a disability-specific participation

measurement tool for children with ASD—to 38–83 months and develops new items. The

updated version of the PQP has two unique features: (1) it provides a comprehensive

assessment of participation in a questionnaire format, and (2) it includes items specific to the

challenges faced by children with ASD.

Acknowledgments: The authors express their gratitude to medRxiv for hosting our initial

findings, available at https://www.medrxiv.org/content/10.1101/2023.08.22.23294206v1. We

also thank Erik and Satomi Phillips for their assistance with the English translation of the

items.

Funding Details: This research did not receive any specific grant from funding agencies in the

public, commercial, or not-for-profit sectors.

Disclosure Statement: The authors declare that they have no conflicts of interest or financial

ties to disclose.

Data Availability Statement: The data that support the findings of this study are available

upon reasonable request but are not publicly accessible due to privacy or confidentiality

concerns.

References

1. American Psychiatric Association. "Neurodevelopmental disorders" In *Diagnostic and Statistical Manual of Mental Disorders* (2022). https://doi.org/10.1176/appi.books.9780890425787.x01_Neurodevelopmental_Disorders

2. Catherine Lord, Tony Charman, Alexandra Havdahl, Paul Carbone, Evdokia Anagnostou, Brian Boyd, Themba Carr, Petrus J de Vries, Cheryl Dissanayake and Gauri Divan, "The Lancet Commission on the future of care and clinical research in autism." *Lancet* 399 No. 10321 (2022): 271–334, https://doi.org/10.1016/S0140-6736(21)01541-5.

3. World Health Organization. International classification of functioning, disability, and health: Children and youth version: ICF-CY (World Health Organization, 2007)

4. Cheryl Boop, Susan M. Cahill, Charlotte Davis, Julie Dorsey, Varleisha Gibbs and Brian Herr, "Occupational therapy practice framework: Domain and process fourth edition," *American Journal of Occupational Therapy* 74 Suppl. 2 (2020): 1–85.

5. Sorayya Askari, Dana Anaby, Melanie Bergthorson, Annette Majnemer, Mayada Elsabbagh and Lonnie Zwaigenbaum, "Participation of children and youth with autism spectrum disorder: A scoping review," *Review Journal of Autism and Developmental Disorders* 2 No. 1 (2015): 103–114, https://doi.org/10.1007/s40489-014-0040-7.

6. Patti LaVesser and Christine Berg, "Participation patterns in preschool children with an autism spectrum disorder," *OTJR: Occupation, Participation and Health* 31 No. 1 (2011): 33–39, https://doi.org/10.3928/15394492-20100823-01.

7. Verónica Schiariti, Soheil Mahdi, and Sven Bölte, "International classification of functioning, disability and health core sets for cerebral palsy, autism spectrum disorder, and attention deficit–hyperactivity disorder," *Developmental Medicine and Child Neurology* 60 No. 9 (2018): 933–941, https://doi.org/10.1111/dmcn.13922.

8. Tamara Yee, Joyce Magill-Evans, Lonnie Zwaigenbaum, Lori-Ann R. Sacrey, Sorayya Askari and Dana Anaby, "Participation measures for preschool children with autism spectrum disorder: A scoping review," *Review Journal of Autism and Developmental Disorders* 4 No. 2 (2017): 132–141, https://doi.org/10.1007/s40489-017-0102-8

9. Costi, Stefania, Maria Cristina Filippi, Luca Braglia, Laura Beccani, Isabella Corradi, Elena Bruzzi, Chiara Signorelli, and Elisa Pelosin. "Reliability and construct validity of the activities scale for kids in Italian children with cerebral palsy." Disability and rehabilitation 44, no. 21 (2022): 6445-6451. https://doi.org/10.1080/09638288.2021.1966519

10. Mulcahey, M. J., Christina C. Thielen, Mary D. Slavin, Pengsheng Ni, and Alan M.

Jette. "Pediatric measure of participation short forms version 2.0: development and evaluation." Spinal Cord 59, no. 11 (2021): 1146-1154. https://doi.org/10.1038/s41393-021-00625-5

11. Susana Castro and Ana Pinto, "Matrix for assessment of activities and participation: Measuring functioning beyond diagnosis in young children with disabilities," *Developmental Neurorehabilitation* 18 No. 3 (2015): 177–189, https://doi.org/10.3109/17518423.2013.806963.

12. Anat Golos, Sophi Chapani Itkin and Hadas Ben-Zur, "The structured preschool participation observation (SPO) for children with ASD: Adaptation, initial psychometric properties, and children's participation," *Physical and Occupational Therapy in Pediatrics* 42 No. 2 (2022): 198–214, https://doi.org/10.1080/01942638.2021.1968096.

13. Liya Li, Berit Møller Christensen, Marita Falkmer, Yue Zhao and Karina Huus, "Content validity of the instrument 'Picture My Participation' for measuring participation of children with and without autism spectrum disorder in mainland China," *Scandinavian Journal of Occupational Therapy* 30 No. 8 (2023): 1237–1247, https://doi.org/10.1080/11038128.2023.2220911.

14. Kathleen N. Lohr and Bradley J. Zebrack, "Using patient-reported outcomes in clinical practice: Challenges and opportunities," *Quality of Life Research* 18 No. 1 (2009): 99–107, https://doi.org/10.1007/s11136-008-9413-7.

15. Takuto Nakamura, Sakumi Koyama, Hirofumi Nagayama and Satoshi Sasada, "Development and content validation of a participation measurement questionnaire for young children with autism spectrum disorder," *Japanese Journal of Occupational Therapy Research* 25 No. 1 (2022): 7–14.

16. Caroline B Terwee, Cecilia AC Prinsen, Alessandro Chiarotto, Henrica CW de Vet, Lex M Bouter, Jordi Alonso, Marjan J Westerman, Donald L Patrick and Lidwine B Mokkink, *COSMIN methodology for assessing the content validity of PROMs–user manual* (VU University Medical Center, 2018)

17. Gary M. Bedell, Mary A. Khetani, Martha A. Cousins, Wendy J. Coster, and Mary C. Law, "Parent perspectives to inform development of measures of children's participation and environment," *Archives of Physical Medicine and Rehabilitation* 92 No. 5 (2011): 765–773, https://doi.org/10.1016/j.apmr.2010.12.029

18. Wendy Coster and Mary Alunkal Khetani, "Measuring participation of children with disabilities: Issues and challenges," *Disability and Rehabilitation* 30 No. 8 (2008): 639–648, https://doi.org/10.1080/09638280701400375.

19. Christine Imms, Brooke Adair, Deb Keen, Anna Ullenhag, Peter Rosenbaum and Mats Granlund, "'Participation': a systematic review of language, definitions, and constructs used in intervention research with children with disabilities,"

Developmental Medicine and Child Neurology 58 No. 1 (2016): 29–38, https://doi.org/10.1111/dmcn.12932.

20. American Psychiatric Association. *Diagnostic and statistical manual of mental disorders: DSM-5 (Vol. 5)* (American Psychiatric Association, 2013)

21. Klaus Krippendorff, *Content analysis: An introduction to its methodology* (Sage Publications, 2018).

22. Toyojiro Matsuishi, Miki Nagano, Yuko Araki, Yoshiyuki Tanaka, Mizue Iwasaki, Yushiro Yamashita, Schinichiro Nagamitsu, Chiho Iizuka, Takashi Ohya, Kunihiko Shibuya et al., "Scale properties of the Japanese version of the Strengths and Difficulties Questionnaire (SDQ): A study of infant and school children in community samples," *Brain and Development* 30 No. 6 (2008): 410–415, https://doi.org/10.1016/j.braindev.2007.12.003.

23. Sara S. Sparrow, David A. Balla, Domenic V. Cicchetti and Edgar A. Doll, *Vineland-II: Vineland Adaptive Behavior Scales;[VABS]; Survey Forms manual; a revision of the Vineland Social Maturity Scale* (NCS Pearson, 2005)

24. Paul C. Beatty and Gordon B. Willis, "Research synthesis: The practice of cognitive interviewing," *Public Opinion Quarterly* 71 No. 2 (2007): 287–311, https://doi.org/10.1093/poq/nfm006.

25. R. Fitzpatrick, C. Davey, M. J. Buxton and D. R. Jones, "Evaluating patient-based outcome measures for use in clinical trials." *Health Technology Assessment* 2 No. 14 (1998): i–iv, 1–74. https://doi.org/10.3310/hta2140.

26. Lidwine Mokkink, Cecilia Prinsen, Donald Patrick, Jordi Alonso, Lex Bouter, Henrica de Vet and Caroline Terwee, "COSMIN methodology for systematic reviews of patient-reported outcome measures (PROMs)", 2018, https://cosmin.nl/wp-content/uploads/COSMIN-syst-review-for-PROMs-manual_version-1_feb-2018.pdf

27. Maskari Machida, Itaru Nakamura, Takako Kojima, Reiko Saito, Tomoki Nakaya, Tomoya Hanibuchi, Tomoko Takamiya, Yuko Odagiri, Noritoshi Fukushima, Hiroyuki Kikuchi, et al., "Acceptance of a COVID-19 vaccine in Japan during the COVID-19 pandemic." *Vaccines* 9 No. 3 (2021): 210, https://doi.org/10.3390/vaccines9030210.

 Tables

 Table 1. The Participation Questionnaire for Preschoolers: preliminary items and subdomains

■ The first question is about your child's self-care and health

1. The child tries to take care of himself/herself (dressing, eating, etc.)

2. The child has difficulty completing his/her daily routine (becomes distracted while

eating, starts something else while getting dressed, etc.)

3. The child does not like to be well-groomed (is reluctant to wear certain clothing,

wash hair, brush teeth, etc.)

4. The child has a sound he/she does not like (vacuum cleaner noise, sibling crying,

sirens, etc.)

5. The child is easily fatigued

6. You are worried about the child's diet

7. You are worried about the child's defecation

8. You are worried about the child's sleep

9. You are having trouble teaching the child to take care of himself/herself (excretion,

changing clothes, eating, etc.)

10. You have a family member who will help you care for the child

11. You have availed a service to take care of the child, which provides you with peace

of mind (nursery school, babysitting, etc.)

12. You are having difficulty in getting the child to seek medical attention or treatment

■ The following questions are about your child's domestic life

13. The child tries to clean up (toys, drawing materials, etc.)

14. You are glad that the child is willing to help others (carries dishes, throws away garbage, etc.)

■ The next questions are about mobility

15. The child is capable of running smoothly and has no issues with changing speed and direction

16. You feel unsafe when walking outdoors with the child (e.g., he/she jumps out into the road)

17. You have difficulties when you are out with the child (he/she goes away on his/her own, does not try to walk by himself/herself, etc.)

18. You do not have access to necessary public transportation with the child (bus, train, plane, etc.)

■ The following questions are about play:

19. The child is able to spend time playing by himself/herself

20. The child has opportunities for active physical play

21. The child is able to have a good time with his/her friends

22. The child is able to participate in play with rules

23. The child has a safe place to play

24. You do not enjoy playing with the child

25. You do not have many places to have fun with the child (park, children's center, etc.)

■ The following questions are about education at home and in preschools and kindergartens:

26. The child obediently listens to adults

27. The child is not able to follow the teacher's instructions to complete the task

28. The child is active in physical activities (running, gymnastics, dancing, etc.)

29. The child is actively involved in creative activities (drawing, clay, crafts, etc.)

30. The child tries to use a tool that he/she has never used before (pencil, jump rope, cleaning supplies, etc.)

31. The child sometimes does not want to go to educational places (kindergarten, preschool, etc.)

32. You want to use medical and welfare services for children with disabilities, but you cannot

33. You are always anxious when you think about the child's future

■ The following questions are about recreational and cultural activities:

34. The child can enjoy outings with the family (eating out, amusement parks, etc.)

35. You have difficulty traveling with the child on overnight stays

36. You have difficulty taking the child to a family gathering (Buddhist memorial service, New Year's Day, etc.)

37. You do not have the financial means to enjoy holidays with the child

■ The following questions are about interpersonal relationships:

38. The child does something that others may find offensive

39. The child enjoys spending time with family

40. The child has many troubles with other children (friends, siblings, etc.)

41. The child has a particularly close friend

42. The child has adults, other than his parents, who appreciate his/her good qualities (grandparents, teachers, etc.)

43. The child has people in the community who are willing to be involved with him/her (e.g., mother's friends)

44. You sometimes do not enjoy spending time with the child

■ The last questions are about communication

45. The child reacts when a friend calls him/her

46. The child understands and acts on words and gestures of encouragement (continues activity, smiles, etc.)

47. The child helps others in need (offers a hand to a child who has fallen, picks up

something that has fallen, etc.)

48. The child knows his/her written name

49. The child can communicate refusals to the family (not wanting to go, not wanting to

drink, etc.)

50. The child can answer in spoken words when asked

51. You can understand what the child is trying to say

528 Each item was written in Japanese and translated into English by three reviewers. One reviewer

529 was a native English-speaking staff member of the thesis editing company, one was a native

530 English-speaking American, and one was a native Japanese speaker who had lived in the USA

531 for more than 20 years.

532 Items 2–9, 12, 16–18, 24, 25, 27, 31–33, 35–38, 40, and 44 were reverse-scored.

Table 2. Demographics of the expert participants involved in content validation and the number of codes that appeared

Participant	Sex	Profession	Practical experience	Facility	Relevance	Comprehensiveness	Comprehensibility
1	Male	"Participation" researcher	None	Professional school	19	1	3
2	Female	"Participation" researcher	21-30 years	University/clinic	5	4	29
3	Female	Speech–language–hearing therapist	5-10 years	Clinic/child development support center	4	5	8
4	Female	Clinical psychologist	11-20 years	Clinic/child development support center	0	2	2
5	Male	Certified social worker	21-30 years	University	0	1	4
6	Female	Childcare worker	11-20 years	Child development support center	0	1	5
7	Male	Occupational therapist	11-20 years	University	2	3	15
8	Male	"Participation" researcher	21-30 years	University	0	1	10

Practical experience was defined as the time spent working with children with ASD. Among the "participation" researchers,

Participants 2 and 8 were researchers who also had experience supporting children with ASD. Participants 5 and 7 belonged to a

research institute and had experience in ASD research. The other participants worked at medical institutions or welfare facilities

related to supporting children with ASD.

Table 3. Demographics of caregivers involved in content validation

Participant	Relationship	Child's sex	Child's age (months)	Comorbid diagnosis	IQ	Residential areas	Relevance (1st/2nd)	Comprehensiveness (1st/2nd)	Comprehensibility (1st/2nd)
1	Mother	Female	36-41	NONE	SI	Big city A	0/0	0/0	9/2
2	Father	Male	65-71	NONE	SI	Big city A	0/0	0/0	1/1
3	Mother	Male	65-71	NONE	SI	Big city A	0/0	1/0	3/0
4	Mother	Male	65-71	NONE	BI	Big city B	0/0	0/0	4/0
5	Mother	Male	54-59	ADHD	SI	Town C	0/0	0/0	2/0
6	Mother	Male	65-71	NONE	ID	Big city D	0/0	0/0	1/15
7	Mother	Male	54-59	NONE	ID	Town E	0/0	0/0	7/0
8	Mother	Male	48-53	NONE	Sup	City F	0/0	0/0	1/1
9	Mother	Male	60-65	ID	ID	City G	0/NA	0/NA	0/NA

| 10 | Mother | Male | 60-65 | DCD | SI | Big city H | 0/NA | 0/NA | 0/NA |
| 11 | Mother | Female | 70-77 | ADHD | ID | Town I | 0/NA | 0/NA | 0/NA |

ADHD, Attention-deficit/hyperactivity disorder; DCD, Developmental dyscoordination disorder; ID, Intellectual disability. The results of the Wechsler Intelligence Scale for Children or the Tanaka-Binet Intelligence Scale are used for the intelligence quotient. However, as there is a high correlation between the scores of both intelligence tests and the Kyoto Scale of Psychological Development, the developmental quotient is regarded as the intelligence quotient only in the results of the Kyoto Scale of Psychological Development. On this basis, IQ 50-69 was classified as Intellectual Disability (ID), IQ 70-84 as Borderline intelligence (BI), IQ 85-115 as Standard intelligence (SI), and IQ 116 and above as Superior intelligence (Sup). IQ 116 and above are classified as Superior intelligence. Residential areas are classified into government-designated cities (hereafter referred to as "big city," indicating municipalities with a population of over 500,000), other cities (hereafter referred to as "city," indicating municipalities with a population between 50,000 and 500,000), and towns and villages (hereafter referred to as "town," indicating municipalities with a population of under 50,000), based on the classification of local governments by Japan's Ministry of Internal Affairs and Communications. The same region of residence, including alphabetic characters, means that participants reside in the same municipality.

Participants 2 and 3 were fathers and mothers of the same child.

Table 4. Items after caregivers' content validation and the number of codes for each sub-domain

■ First, please answer the following questions about your child:	SC	DOM	IIR	EDU	EIP	CSC	COM	MOB
1. Changes clothes by himself/herself	○							
2. Finishes getting dressed	○							
3. Helps around the home		○						
4. Can play alone for more than 15 minutes					○			
5. Engages in active physical play					○			
6. Plays with friends					○			
7. Participates in competitive activities (relay, *sugoroku*, etc.)					○			
8. Learns social skills with friends				○				
9. Learns how to use school supplies (at home, kindergarten, etc.)				○				
10. Enjoys class activities in cooperation with children of the same age				○				

11. Has favorite lessons for learning   ○

12. Has close friends (talks a lot, plays together a lot, etc.)   ○

13. Fights with others of the same age (biting, hitting, etc.)   ○

14. Always feels safe with friends   ○

15. Has a teacher that he/she likes   ○

16. Always has a teacher by his/her side to help develop strengths (kindergarten, lessons, medical care, etc.)   ○

17. Has someone, other than his parents, who is always there when he/she needs help   ○

18. Plenty of places to go out with family   ○

19. Neighbors outside of the immediate family are always nice to him/her   ○

20. Opportunities to participate in community gatherings (festivals, children's meetings, etc.)   ○

■ Next, please describe how you feel	SC	DOM	IIR	EDU	EIP	CSC	COM	MOB
21. Concerned about your child's diet	○							

22. Concerned about your child's defecation　　　　　　　　　　　　　　　　　　　○

23. Concerned about your child's sleep　　　　　　　　　　　　　　　　　　　　○

24. Concerned about changing your child's clothes　　　　　　　　　　　　　　○

25. Playing with your child can be exhausting　　　　　　　　　　　　　　　　　　　　　　　○

26. Suffering from lack of specialized support for children with disabilities (lack of facilities, reluctance to go to

facilities, etc.)　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　○

27. Difficulty in sending your child to preschool or kindergarten　　　　　　　　　　　　　　　○

28. Time at home with your child is always a pleasure　　　　　　　　　　　　　　　○

29. You can take your child on a trip that involves an overnight stay　　　　　　　　　　　　　　　　　○

30. You can take your child to a gathering of relatives　　　　　　　　　　　　　　　　　　　○

■ Finally, please answer the question about your child's feelings	SC	DOM	IIR	EDU	EIP	CSC	COM	MOB
31. Reluctant to seek medical attention at a hospital	○							

32. Seems to like to keep up appearances (wash hair, brush teeth, etc.)   ○

33. Seems to enjoy mealtimes   ○

34. Seems to like helping out   ○

35. Always seems to enjoy his/her time with the family   ○

36. Always seems to enjoy time out with family   ○

Number of code occurrences (Phase 1: Experts)	SC	DOM	IIR	EDU	EIP	CSC	COM	MOB	OTH
Relevance	5	0	2	9	2	1	7	4	0
Comprehensiveness	2	1	2	7	0	2	0	0	0
Comprehensibility	16	7	15	19	11	7	0	0	1

Number of code occurrences (Phase 2: Foster parent)	SC	DOM	IIR	EDU	EIP	CSC	COM	MOB	OTH
Relevance	0	0	0	0	0	0	0	0	0
Comprehensiveness	0	0	1	0	0	0	0	0	0

| Comprehensibility | | 2 | 0 | 19 | 13 | 8 | 3 | 0 | 0 | 3 |

Each item was written in Japanese and translated into English by three reviewers. One reviewer was a native English-speaking staff member of the

thesis editing company, one was a native English-speaking American, and one was a native Japanese speaker who had lived in the USA for more

than 20 years.

Abbreviations: SC, Self-care; DOM, Domestic life; IIR, Interpersonal interactions and relationships; EDU, Education; EIP, Engagement in play;

CSC, Community, social, and civic life; COM, Communication; MOB, Mobility; OTH, Others.

"▯" indicates the sub-domain to which the item corresponds.

The codes related to "Others" pertain to the change in directives and the change in the placement of items and responses. Items 13, 21–27, and 31

are reverse-scored.