Host behaviour driven by awareness of infection risk amplifies the chance of superspreading events

Kris V Parag[1,2,*] and Robin N Thompson[3]

[1]MRC Centre for Global Infectious Disease Analysis, Imperial College London, London, UK.
[2]NIHR HPRU in Behavioural Science and Evaluation, University of Bristol, Bristol, UK.
[3]Mathematical Institute, University of Oxford, Oxford, UK.

*For correspondence: k.parag@imperial.ac.uk.

Abstract

We demonstrate that heterogeneity in the perceived risks associated with infection within host populations amplifies the chances of superspreading during the crucial early stages of an epidemic. Under this behavioural model, individuals less concerned about the dangers from infection are more likely to be infected and attend larger-sized (riskier) events. For directly transmitted diseases such as COVID-19, this leads to infections being introduced at rates above the population prevalence to the events most conducive to superspreading. We develop an interpretable computational framework for evaluating within-event risks and derive a small-scale reproduction number measuring how the infections generated at an event depend on transmission heterogeneities and the number of introductions. This quantifies how event-scale patterns relate to population-level characteristics and generalises previous frameworks. As event duration and size grow, our reproduction number converges to the basic reproduction number. We illustrate that even moderate levels of heterogeneity in the perceived risks from infection substantially increase the likelihood of disproportionately large clusters of infections occurring at larger events, despite fixed overall disease prevalence. We show why collecting data linking host behaviour and event attendance is essential for accurately assessing the risk posed by an invading pathogen in the emerging stages of an outbreak.

Keywords: infectious diseases; behavioural models; risk awareness; reproduction numbers; importations; superspreading events.

Introduction

The prediction and prevention of superspreading events, which are characterised by primary infected individuals generating disproportionately large numbers of secondary infections [1], is a central challenge in infectious disease epidemiology. For acute, directly communicable diseases such as COVID-19, SARS and Ebola virus disease, superspreading is a major driver of transmission that leads to less frequent but more explosive outbreaks than we might expect

under more classical models that neglect the substantial variability in secondary infections
generated by infected hosts [2]. During early or emergent stages of a potential epidemic, when
there are limited immunity levels in the host population and transmission dynamics are
inherently stochastic, superspreading events have been found responsible for spurring both
the initial growth and eventual persistence of epidemics and for limiting the effectiveness of
non-pharmaceutical interventions [1,3–5].

Consequently, identifying the main factors that underly the risk of superspreading is crucial for
effective disease management [4]. Many of these factors are known, with heterogeneities in
(i) host characteristics (e.g., susceptibility, infectiousness and contact patterns), (ii) pathogen
biology (e.g., transmission routes and viral loads), (iii) environmental effects (e.g., ventilation
and gathering size) and (iv) host behaviours (e.g., social customs and intervention adherence)
all contributing to the risk of superspreading [3,4,6–8]. However, incorporating these factors
in parsimonious modelling frameworks can be difficult because the mechanisms linking them
to superspreading are still not fully understood. This is particularly the case for factors (iii) and
(iv), with recurrent calls for more comprehensive data collection to help study the relationships
among behavioural, environmental and epidemiological trends [9–11]. Here we explore how
a key feature of host behaviour can shape the likelihood of superspreading and provide a
mathematical demonstration of the benefits of collecting and analysing more data to elucidate
the links between human behaviour and infectious disease epidemics.

We consider how heterogeneity in perceptions of the risk associated with infection throughout
a host population impact heterogeneity in the transmission of new infections in the early stages
of an epidemic. Risk awareness is a documented phenomenon in which individuals engage in
self-protective behaviours in response to the perceived health, economic and other dangers
of acquiring infection. The exact relationship between risk-perception and self-protection in a
population exists on a spectrum spanning more risk-averse individuals to those with larger risk
appetites (e.g., risk deniers) [12–14]. During an epidemic of a directly transmitted pathogen,
risk-averse individuals may reduce their number of contacts by limiting their socialising and
mobility, while those with larger risk appetites may increase their relative infection exposure
(e.g., by hosting unregulated gatherings when infections are rising) [15]. Risk awareness can
improve intervention efficacy (e.g., reducing mobility) or negate it (e.g., via deliberate non-
compliance) and substantially change outbreak amplitudes and durations [11,16–19].

Despite its importance, the interplay between risk awareness and superspreading risk has not
been studied in detail, with most research focussing on pathogen and host characteristics
instead of exploring behavioural patterns. We study this interplay under a simple but plausible
hypothesis – that more risk-averse hosts are more likely to avoid events of larger size, due to

the perception of heightened infection risk at those events [14,19]. Here, events are short-
term gatherings (e.g., parties) so that only one generation of infection is possible. While data
directly linking risk perception to event attendance are unavailable, it is known that individuals
modify their behaviour in response to population-level prevalence and that the variability in
individuals' level of acceptable risk relative to this prevalence baseline correlates well with the
extent to which contacts are reduced during epidemics [12–14,20,21]. A likely pathway for
reducing exposure to invading pathogens is by limiting attendance at riskier (voluntary) events.
This logic underlies our hypothesis and subsequent analysis.

This awareness mechanism implies, for a fixed population prevalence, that larger events (e.g.,
concerts, sports matches) are more likely to be attended by individuals who are less risk
averse. Since there is limited infection-induced immunity in the host population during early
epidemic stages, these individuals are also more likely to be infected. This stems from
observations that those with larger numbers of contacts have elevated chances of acquiring
infection, and these individuals are also likely to have larger risk appetites (more risk-averse
individuals tend to reduce contacts) [22]. We posit that this coupling between behaviour and
environment (i.e., modification of event attendance due to risk perception and event size) may
amplify the chances of superspreading occurring at larger events, which have the capacity to
support excessive numbers of infections. To test this hypothesis, we develop a framework to
model the number of infections y generated at an event of size n, given that x initially infected
individuals attend that event. This yields a small-scale reproduction number that extends
recent approaches [23–25] to understanding within-event transmission in three directions.

First, we explicitly model the transmission-reducing effects of finite numbers of susceptible
individuals ($n - x$) and imported infections (x) and at the event. As event size and duration
grow, these finite size effects become less important and our small-scale reproduction number
converges to R_0, the popular basic reproduction number. Second, we embed heterogeneity
in transmission at the event within our small-scale reproduction number by allowing variations
in secondary infections that are controlled by a dispersion parameter, k. This is a within-event
version of the seminal model of superspreading [1,5,26] and includes the broad influence of
factors (i)-(ii) described above. Third, we account for how x changes (stochastically) with n.
This considers factors (iii)-(iv) and depends on the prevalence of infection in the population as
well as the size-biased importation rate of infections into the event, $\epsilon(n)$, which is influenced
by the spectrum of risk appetites about that prevalence.

The functional dependence of ϵ on n serves as a parsimonious model of risk awareness and
allows us to assess how host behaviour shapes the risk of superspreading (e.g., if $\epsilon(n)$ is an

increasing function of n, then this implies a higher infection import rate into larger events). We

explore our central hypothesis by comparing the relative and combined impact of $\epsilon(n)$ and k

on the tail probability of observing a disproportionately large number of secondary infections

y at an event. We demonstrate, for a fixed overall import rate (equalling the wider population

prevalence), that risk awareness can substantially amplify the chances of superspreading at

a large event, compared to the scenario in which all individuals attending the large event are

assumed to have similar perceptions of infection risk. This pattern holds regardless of k and,

in some instances, we find that the increase in superspreading risk from risk-aware behaviour

outweighs that from inherent transmission heterogeneity.

Methods

Event reproduction numbers including import risk and transmission heterogeneity

We develop a framework for quantifying the risk of acquiring infection at an event (e.g., a party,

concert, sports match), based on a small-scale (within-event) reproduction number. We detail

this below but also sketch the main steps of our methodology and list key notation in **Fig 1**.

An event is defined as a short-term grouping of n people and we allow $0 \leq x \leq n$ of the

individuals attending the event to be infectious. Initially, there are x introductions (i.e., imported

infections) at this event and $n - x$ susceptible hosts. We assume no prior immunity in the

population and let $\mathbf{P}(y|n)$ be the probability of $0 \leq y \leq n - x$ new infections being generated

at that event as we describe in **Eq. (1)**.

$$121 \qquad \mathbf{P}(y|n) = \sum_{x=0}^{n} \mathbf{P}(y|x,n)\mathbf{P}(x|n). \quad (1)$$

122 This depends on $\mathbf{P}(y|x,n)$, the probability of y new infections occurring given the x infectious

123 individuals initially (for events of size n) and the prior probability of those x imports, $\mathbf{P}(x|n)$.

124 We define the small-scale reproduction number for this event as $R(x) \overset{\text{def}}{=} x^{-1}\mathbf{E}[y|x,n]$, with

125 the expected number of infections generated by x imports denoted by $\mathbf{E}[y|x,n]$. We expand

126 this to obtain **Eq. (2)** below.

$$127 \qquad R(x) \overset{\text{def}}{=} \frac{1}{x} \sum_{y=0}^{n-x} y\mathbf{P}(y|x,n). \quad (2)$$

Here $R(x)$ measures the expected number of new infections generated by each import when

there are x imports in total. Although intuitive, this reproduction number formulation is novel.

A central idea of this study is the importance of $\mathbf{P}(x|n)$ and its dependence on event size n.

Earlier work assumed that $\mathbf{P}(x|n)$ depends solely on the prevalence of the infection in the

population [25], neglecting how heterogeneities in human behaviour may affect the number of

imported cases at a given event of size n. To our knowledge, alternative models for $\mathbf{P}(x|n)$

informed by human behaviour and the influence of this behaviour on the number of infections

generated at the event have not been explored. The heterogeneity in host behaviour that we

consider relates to the spectrum of risk appetites i.e., the fact that different individuals perceive

different infection risks associated with attending an event of size n. This spectrum alters the

rate of importing infections into events, relative to the prevalence, and so modulates $\mathbf{P}(x|n)$.

Fig 1: Modelling framework for event-level transmission subject to risk awareness. We

outline the central steps and define the main notation underlying our proposed framework for

modelling transmission patterns at small events. We refer to equations defined in the Methods.

This framework accounts for how heterogeneity in infection risk perception among individuals

modulates the number of imported cases x at an event of size n and hence contributes to the

secondary infections generated at that event y.

Our event or small-scale reproduction number also generalises prior research by including the

effects of finite x and n. Since only one generation of infection can occur at an event, this finite

initial condition can strongly shape clustering patterns, underscoring the value of modelling

$\mathbf{P}(x|n)$. The original event reproduction number [24] considers a single imported infection and

relates to (but is not the same as) our $R(1)$, which we later show is always an upper bound

for $R(x)$. By extending the event reproduction number definition, we model the influence of

$\mathbf{P}(x|n)$ on the risk of acquiring infection at any event directly. As we explain below, $R(x)$ also

embeds heterogeneity in transmission from both host characteristics as well as pathogen

biology [1] and is explicitly related to the population-level basic reproduction number, R_0 [27].

To convert **Eqs. (1)-(2)** into a computable form, we draw on characteristics of both the event

and disease. We denote the (frequency-dependent) transmission rate as β and the expected

duration of an individual infection as d, so that $R_0 = \beta d$. We then consider an event that lasts

for time τ, which is assumed to be substantially shorter than d, so that infectiousness outlasts

the event and at most one generation of infection is possible at the event. We also assume

that the event is closed i.e., for any specific event, n takes a constant value. The split of n into

x and $n - x$ completely defines the epidemiologically important states for the event

If there is only one infected individual at the start of the event, then the probability that any

susceptible host gets infected is the secondary attack rate (SAR), $p = 1 - e^{-\frac{\beta\tau}{n}}$, making the

standard assumption that the times to infection are exponentially distributed. When there are

s susceptible individuals, then $\mathbf{E}[y|1, n] = sp$. While this assumes that all the susceptible

individuals are exposed to all infectious ones at the event, we can model more realistic contact

networks as in [27] by modifying s to be the subset of susceptible hosts likely to be exposed

to each infection (this connects network and random mixing models).

We generalise this approach in three main directions. First, we model the effect of variability

in the number of imported infections. If there are x imports to the event, then the SAR becomes

$p = 1 - e^{-\left(\frac{\tau}{d}\right)\left(\frac{x}{n}\right)R_0}$ with $\beta = \frac{R_0}{d}$. Since there are initially $s = n - x$ susceptible individuals, the

expected number of infections generated at the event is $\mathbf{E}[y|x, n] = (n - x)p$. The leads to

the event reproduction number $R(x)$ in **Eq. (3)** below. Note that $R(0) = R(n) = 0$.

$$174 \qquad R(x) = \frac{n-x}{x}\left(1 - e^{-\left(\frac{\tau}{d}\right)\left(\frac{x}{n}\right)R_0}\right) \approx \left(1 - \frac{x}{n}\right)\left(\frac{\tau}{d}\right)R_0 + \mathbf{O}\left(\frac{x}{n}\right). \quad (3)$$

This formulation has interesting limiting behaviour at various x. As the number of susceptibles

grows in excess of imports i.e., $\frac{n}{x}$ increases, the $\mathbf{O}\left(\frac{x}{n}\right)$ terms in the Taylor series approximation

of $R(x)$ in **Eq. (3)** become negligible. As n becomes large, we find $R(x) \rightarrow \left(\frac{\tau}{d}\right)R_0$. If the event

lasts for the duration of infectiousness ($\tau = d$), then $R(x) \rightarrow R_0$. This convergence makes

sense since our formulation is equivalent to a finite or small-scale version of random mixing.

Second, we expand this model to include realistic heterogeneity from host characteristics and

pathogen biology. It is unlikely that every infectious individual has the same transmissibility

and we expect substantial variations in the numbers of infections generated by each infected

individual [1,28]. We therefore allow R_0 to have some distribution from which every import is

randomly sampled and let R_0^j indicate the sample for the j^{th} of the x imports at the event. This

heterogeneous version of $R(x)$ is in **Eq. (4)**, with expected number of infections $\mathbf{E}[y|x,n] =$

$xR(x)$. Note that **Eq. (4)** is of the form $\frac{n-x}{x}p_{\text{het}}$, with p_{het} as a heterogeneous SAR.

$$187 \qquad R(x) = \frac{n-x}{x}\left(1 - e^{-\left(\frac{\tau}{d}\right)\left(\frac{1}{n}\right)\Sigma_{j=1}^{x} R_0^j}\right). \quad (4)$$

We compute the mean of $R(x)$ across the transmission heterogeneity for x infectious imports

in **Eq. (5)**, with $\mathbf{E}_{\text{het}}$ indicating expectation about the distributions of the R_0^j and $\mathbf{M}_b(a)$ as the

moment generating function about b evaluated at a. As the transmissibility of the x imported

infections are independently sampled, $\mathbf{M}_{\Sigma_{j=1}^{x} R_0^j}(a) = \prod_{j=1}^{x} \mathbf{M}_{R_0^j}(a)$. This reduces to $\mathbf{M}_{R_0}(a)^x$

if samples are identically distributed. The expected number of infections under this model as

a function of x is $\mathbf{E}_{\text{het}}\big[\mathbf{E}[y|x,n]\big] = x\mathbf{E}_{\text{het}}[R(x)]$ with $\mathbf{E}_{\text{het}}[R(x)]$ from **Eq. (5)**.

$$194 \qquad \mathbf{E}_{\text{het}}[R(x)] = \frac{n-x}{x}\left(1 - \mathbf{M}_{\Sigma_{j=1}^{x} R_0^j}\left(\left(\frac{\tau}{d}\right)\left(\frac{1}{n}\right)\right)\right). \quad (5)$$

195   Following [28], we evaluate the variance around $R(x)$ as $\mathbf{V}_{\text{het}}[R(x)]$ with $a_n = \left(\frac{\tau}{d}\right)\left(\frac{1}{n}\right)$ in **Eq.**

196   **(6)**. This involves expanding $\mathbf{E}_{\text{het}}[R(x)^2] - \mathbf{E}_{\text{het}}[R(x)]^2$ and applying properties of $\mathbf{M}_b(a)$.

197   The variance on the expected number of infections is $\mathbf{V}_{\text{het}}\big[\mathbf{E}[y|x,n]\big] = x^2\mathbf{V}_{\text{het}}[R(x)]$. All of

198   these statistics remain valid for any model of transmission heterogeneity but we derive analytic

199   relations under the most widely used model of [1] in the subsequent section.

200
$$\mathbf{V}_{\mathrm{het}}[R(x)] = \left(\frac{n-x}{x}\right)^2 \left(\mathbf{M}_{\sum_{j=1}^x R_0^j}(-2a_n) - \mathbf{M}_{\sum_{j=1}^x R_0^j}(-a_n)^2\right). \quad (6)$$

Third, we examine how the likelihood of finding that x infectious individuals have attended the

event impacts the above quantities. This involves evaluating how $\mathbf{P}(x|n)$ weights the formulae

in **Eqs. (4)-(6)**. This weighting may be random, depend on behavioural preferences as we

posit in the next section (i.e., risk awareness) or be assigned using other rules. We propose

that a more informative measure of the risk of acquiring infection from an event of size n and

duration τ is the import-weighted event reproduction number R_{imp} as in **Eq. (7)** below.

207
$$R_{\mathrm{imp}} = \sum_{x=0}^{n} R(x)\mathbf{P}(x|n) . \quad (7)$$

208   While R_{imp} averages over the possible numbers of imports, it is still a random variable with

209   samples taken from the distribution controlling transmission heterogeneity. Accordingly, it has

210   statistics $\mathbf{E}_{\mathrm{het}}\left[R_{\mathrm{imp}}\right]$ and $\mathbf{V}_{\mathrm{het}}\left[R_{\mathrm{imp}}\right]$ that we compute by summing and weighting $\mathbf{E}_{\mathrm{het}}[R(x)]$

211   and $\mathbf{V}_{\mathrm{het}}[R(x)]$ by $\mathbf{P}(x|n)$ and $\mathbf{P}(x|n)^2$ respectively. The expected number of new infections

212   $\mathbf{E}_{\mathrm{imp}}[y|n]$ that is associated with R_{imp} follows as in **Eq. (8)**.

213
$$\mathbf{E}_{\mathrm{imp}}[y|n] = \sum_{x=0}^{n} xR(x)\mathbf{P}(x|n). \quad (8)$$

Similarly, we obtain the heterogeneous statistics $\mathbf{E}_{\mathrm{het}}\left[\mathbf{E}_{\mathrm{imp}}[y|n]\right]$ and $\mathbf{V}_{\mathrm{het}}\left[\mathbf{E}_{\mathrm{imp}}[y|n]\right]$ but

the quantities being weighted by $\mathbf{P}(x|n)$ and $\mathbf{P}(x|n)^2$ are now, respectively, $x\mathbf{E}_{\mathrm{het}}[R(x)]$ and

$x^2\mathbf{V}_{\mathrm{het}}[R(x)]$. These all proceed from the properties of expectations and variances applied to

a linear weighted sum with independent terms.

Statistical models for event reproduction numbers and importation patterns

Having outlined measures of infection risk in **Eqs. (7)-(8)**, we build into our framework some

likely approaches for integrating transmission heterogeneities and import patterns (including

when those imported infections are risk-sensitive). This allows us to parsimoniously model

traditional and behavioural drivers of superspreading. Additionally, we incorporate process

stochasticity and provide a full Bayesian formulation for our framework. We start by including

the seminal heterogeneity model of [1], which describes individual variations in transmissibility

via a gamma distribution with dispersion k and mean R_0. We write this as $R_0^j \sim \mathbf{Gam}\left(k, \frac{R_0}{k}\right)$

with **Gam** as a shape-scale parameterised gamma distribution. Using scaling and summing

properties of these gamma variables, we hence obtain $\sum_{j=1}^{x} R_0^j \sim \textbf{Gam}\left(kx, \frac{R_0}{k}\right)$.

This assumes that samples of the basic reproduction number of individuals are independent

and identically distributed and lets us analytically evaluate the moment generating function as

$\textbf{M}_{\sum_{j=1}^{x} R_0^j}(-a_n) = \left(1 + \frac{R_0 a_n}{k}\right)^{-kx}$. We substitute this into **Eqs. (4)-(6)** to precisely compute the

mean and variance of the infections and event reproduction number conditional on a total of

x introductions as detailed above. We can relax the assumption that the R_0^j are independent

and identically distributed by instead sampling them from different distributions or by applying

alternative dispersion models [28]. The heterogeneous R_0^j constitute a major and traditionally

modelled source of stochasticity underpinning the risk metrics we propose in **Eqs. (7)-(8)**.

A less studied source of stochasticity is variability in the probability that infectious individuals

attend the event. Previous work [25] has treated this deterministically, setting the probability

or rate that an attending individual is infected as equal to the population prevalence ρ (or ρ

adjusted by an exposure factor when it is known that the event draws individuals who are less

or more likely to be infected). This is modelled as $x \sim \textbf{Bin}(n, \rho)$, with **Bin** indicating a binomial

distribution. We generalise this under our behavioural hypothesis. We posit, for a fixed overall

importation level, that this import probability increases with n. This models risk awareness, in

which risk-averse individuals who are less likely to be infected avoid larger events, or equally

the individuals attending larger sized events are less risk-averse and more likely to be infected.

Risk appetites may also depend on event duration τ, but we do not explore this here.

We model event size bias using sorted Dirichlet weightings. We consider m events, the i^{th} of

which has size n_i and import rate $\epsilon(n_i)$. Sizes sequentially span all integers from $n_{\min}$ to $n_{\max}$

uniquely (i.e., $n_{\max} = n_{\min} + m - 1$) but we can relax this to include any distribution across

event sizes of interest. We fix the total importation rate across all m events. This constrains

$\sum_{i=1}^{m} n_i \epsilon(n_i) = \rho \sum_{i=1}^{m} n_i$, conserving the total number of infections introduced across events

so that the mean importation rate equals ρ. We enforce this constraint to allow fair comparison

between the conventional model, in which all imports occur with rate ρ, and our size-biased

variations, which describe how variability in perceived risk by hosts affect their attendance at

events. This constraint causes some event sizes to have importation rates above and others

below ρ and allows us to model a spectrum of risk appetites about the baseline prevalence.

This variability in risk perception aligns well with trends found in behavioural surveys [12–14].

The $\epsilon(n_i)$ values encode our event size biases. We construct them by first sampling a set of

random weights $\{w_i\}$ from a symmetrical Dirichlet distribution i.e., $\{w_i\} \sim \mathbf{Dir}(r)$ with r as a

shape parameter applied to every w_i and $\sum_{i=1}^{m} w_i = \rho$. The $\{w_i\}$ set spans all m weights with

smaller values of r leading to more skewed weightings. At very large r, $w_i \approx \frac{\rho}{m}$ for all i. To

model risk-awareness, where we expect that less risk-averse individuals are more likely to

attend larger n_i events relative to more risk-averse individuals, we sort the $\{w_i\}$ in ascending

order so that w_i increases with n_i. We replicate this procedure across many runs to include

variability from the Dirichlet distribution. For every sampled, sorted $\{w_i\}$, we define $\epsilon(n_i) \stackrel{\text{def}}{=}$

$w_i n_i^{-1}(\sum_{l=1}^{m} n_l)$. This satisfies our benchmarking constraint and parsimoniously models the

spectrum of risk appetites across the host population.

We conceptualise this constraint by observing that in the conventional model $\mathbf{E}[x_i] = n_i \rho$ for

the i^{th} event so that $\sum_{i=1}^{m} \mathbf{E}[x_i] = \sum_{i=1}^{m} n_i \rho$ imports occur into all m events on average. In our

risk-aware model $\mathbf{E}[x_i] = n_i \epsilon(n_i)$ and so we chose the $\epsilon(n_i)$ to ensure $\sum_{i=1}^{m} n_i \epsilon(n_i)$ equals

the $\sum_{i=1}^{m} \mathbf{E}[x_i]$ from the conventional model. However, we can relax this constraint to describe

when risk awareness itself changes the prevalence (provided we use updated values at time

snapshots) and we can generalise the model to allow r to also be size dependent (i.e., $r(n_i)$).

In summary, we generate import rates that are size biased (with this bias accounting for risk

awareness) and use a single parameter r to set the strength of this behavioural effect.

Integrating the above models for heterogeneity and importation, we complete our algorithm

(see **Fig 1**) for sampling import weighted distributions of event size risk using **Eqs. (1)-(2)**. We

formulate this in **Eqs. (9)-(10)** with semi-colons discriminating between probabilities that we

evaluate from a distribution and parameters specifying that distribution. For convenience, we

use $S_x = \sum_{j=1}^{x} R_0^j$ for denoting heterogeneous samples and $\epsilon(n)$ for general size bias.

$$280 \qquad \mathbf{P}(y|n,x) = \int_0^{\infty} \mathbf{Bin}\left(y; n-x, \left(1 - e^{-\frac{\tau S_x}{d\,n}}\right)\right) \mathbf{Gam}\left(S_x;\ kx, \frac{R_0}{k}\right) dS_x. \quad (9)$$

$$281 \qquad \mathbf{P}(y|n) = \int_0^1 \sum_{x=0}^{n} \mathbf{P}(y|n,x)\, \mathbf{Bin}(x; n, \epsilon(n))\, \mathbf{P}(\epsilon(n)|n)\, d\epsilon(n). \quad (10)$$

We use the probability distributions in **Eqs. (9)-(10)** together with the definitions of **Eqs. (1)-**

(2) to compute the measures of event risk that we propose in **Eqs. (7)-(8)**. These marginalise

over the distributions of import rate and transmission heterogeneity, which are degenerate

when $\epsilon(n)$ is constant for all n or all $R_0^j = R_0$, respectively. In the Results, we examine the

properties of our computational framework and apply it to explore how behaviour affects
superspreading. Our framework is freely available at: https://github.com/kpzoo/smallscaleR.

Results

289 In the Methods, we developed a framework to assess the risk of acquiring infection at an event
290 by deriving a small-scale reproduction number and the expected number of infections that will
291 occur at the event. Both measures depend on the levels of heterogeneity in transmission and
292 variability in the rate at which infectious individuals are likely to attend the event (i.e., imports).
293 Here we examine the influence of these two key factors in determining outbreak patterns.

Superspreading risk depends on importations and dispersion

Much research has investigated how heterogeneity in transmission can cause superspreading
and hence increase the number of infections likely to result from a gathering or event [1,23].
Specifically, there has been study of how the dispersion parameter k modulates the risk of
superspreading events [26,28,29]. Generally, smaller values of $k < 1$ are predictive of larger
transmission heterogeneity and superspreading risk. However, the influence of the number of
importations x at an event of size n has received relatively little attention. We examine this by
computing the statistics derived in **Eqs. (4)-(6)**, in which we defined the reproduction number
$R(x)$ as a function of the imports and the resulting number of expected infections $\mathbf{E}[y|x,n]$.

We consider an event of size $n = 30$ over a range of dispersions $0.1 \leq k \leq 10$ with a large-
scale limit (see **Eq. (3)**) of $\left(\frac{\tau}{d}\right) R_0 = 0.3$. We sample $R(x)$ and $\mathbf{E}[y|x,n]$ from heterogeneous
gamma distributions describing the transmissibility of the sum of all imported infections (see
Methods) and compute statistics from these samples using **Eqs. (5)-(6)**. We plot these results
in **Fig 2** to explore the properties of these statistics. Interestingly, we find $R(x)$ is a decreasing
function of x, even though every $R(x)$ has the same limit of $\left(\frac{\tau}{d}\right) R_0$. The single import scenario
of $R(1)$ relates to the event reproduction number proposed in [24]. If we assume, as in several
branching process models, that all imports have reproduction number $R(1)$ instead of $R(x)$,
then $\mathbf{E}[y|x,n]$ and the risk of acquiring infection at the event may be notably overestimated.

Further, increasing heterogeneity (decreasing k) increases the variance of our statistics but
decreases mean risk as we see from the inversion of the rank of blue to red curves between
top and bottom panels in **Fig 2**, with $\mathbf{VM}[.]$ as the ratio of variance to mean. Last, we see that
the dependence of our statistics on the number of imports is substantial and can be as critical
as the value of k for describing spread. The value of x that leads to the largest possible (peak)

number of secondary infections at the event is not obvious (and not inferable from $R(x)$) as

imports both cause infections and reduce the available susceptible individuals. In Supplement

Fig S1 we show how this peak changes and that the mean risk difference can be appreciable

in different settings. This underpins the importance of modelling finite event sizes and signifies

that a crucial factor driving the risk of acquiring infection at an event is the import distribution

$\mathbf{P}(x|n)$, which is rarely studied.

Fig 2: Risk statistics for an event with heterogeneous transmission. We plot the mean

($\mathbf{E}[.]$, top subfigures) and variance to mean ratio ($\mathbf{VM}[.]$, bottom subfigures) of the small-scale

event reproduction number $R(x)$ (panel A) and the mean count of new infections $\mathbf{E}[y|x,n]$

(panel B) as a function of the number of imports x. We compute these via **Eqs. (4)-(6)** and

compile statistics over 10^5 samples from heterogeneous offspring distributions with dispersion

parameter k ranging from 0.1 to 10 (increasing from blue to red, with grey depicting all

intermediate values). For comparison, we show the large-scale reproduction number $\left(\frac{\tau}{d}\right) R_0$

and the number of initial susceptible individuals at the event, $n - x$. We repeat this analysis

at a larger value of $\left(\frac{\tau}{d}\right) R_0 = 3$ in Supplement **Fig S1**.

Population prevalence modulates the superspreading potential at events

Having observed the importance of the number of imports, x when assessing the transmission

risk at events, we explore the influence of the distribution of introductions to the event, $\mathbf{P}(x|n)$.

Conventionally [25], $\mathbf{P}(x|n)$ can be defined as a binomial distribution with the probability of an

import being equal to the prevalence of the infection in the wider population, ρ. This is our null

model, and we explore how it integrates with our proposed event statistics (see **Eqs. (4)-(8)**).

We consider epidemics in their initial stages i.e., there is no vaccination- or infection-acquired

immunity, so ρ is small and there are $n - x$ susceptible individuals at the event. We maintain

parameter settings from **Fig 2** but weight samples of small-scale reproduction numbers and

mean numbers of imported infections using $\mathbf{P}(x|n)$, which is $\mathbf{Bin}(x; n, \rho)$ with ρ ranging from

$0.01 - 0.1$ (1–10%). We compute histograms and statistics of these samples in **Fig 3**.

We examine homogeneous ($k = 10$) and heterogeneous ($k = 0.1$) dispersion levels and plot

the log survival (or tail) probabilities of realised numbers of new infections y and associated

small-scale event reproduction numbers R in panels A-B of **Fig 3** for different values of ρ. We

compute these probabilities via **Eqs. (7)-(8)**. Larger values for these probabilities respectively

indicate that superspreading is more likely (i.e., disproportionately more infections than

$\mathbf{E}_{\text{het}}\left[\mathbf{E}_{\text{imp}}[y|n]\right]$ occur) and that imports have increased potential to cause superspreading

(i.e., transmissibility above $\mathbf{E}_{\text{het}}[R_{\text{imp}}]$). This distinction is rarely explored because it is less

important at population levels, where superspreading models are commonly used [1,26,28].

However, the limiting finite-size effects of events make this distinction crucial. Histograms of

samples of the infections at the event for some of the values of ρ in A-B are shown in **Fig 3C**.

Fig 3: The importation rate magnifies the effects of heterogeneous transmission. We plot the log survival probabilities for the number of new infections y (panel A) and related event reproduction numbers R (panel B). We account for the probability of x imports (distributed as **Bin**$(x; n, \rho)$) at an event of size $n = 30$ with the population prevalence as ρ (increasing from blue to red with grey indicating intermediate values). Larger $\mathbf{P}(y \geq c)$ signifies more realised heterogeneity (higher likelihoods that disproportionate numbers of infections result from the event), while larger $\mathbf{P}(R \geq c)$ signifies more heterogeneity in transmissibility (higher potential for superspreading events). In panels A-B, dashed curves are for $k = 10$ (spread is mostly homogeneous) and solid curves are for $k = 0.1$ (spread is heterogeneous). We compute these quantities from **Eqs. (4)-(8)**. Panel C shows histograms of 10^5 samples of y at two ρ

values underpinning panels A-B. Thicker tails or more rightwards mass in these distributions

indicate a higher chance of a large number of infections at the event. We repeat this analysis

at a larger event size of $n = 100$ in Supplement **Fig S2** for comparison.

We find that increasing prevalence ranks the y survival curves for both k scenarios (panel A)

(at a given threshold tail size c, probabilities increase with ρ) but has limited impact on the R

curves (panel B). The latter trend is expected as ρ does not change transmissibility. However,

the fact that prevalence alone can mediate realised superspreading risk is important and, to

our knowledge, unexplored. We confirm this with the histograms of panel C, which have thicker

tails or at least more rightward probability mass as ρ grows (even at large k). The variances

of the y values (not shown) also rise with ρ. We show equivalent analyses for a larger sized

event ($n = 100$) in Supplement **Fig S2** and recover similar results. The rate at which infections

are introduced is therefore critical to assessing the chances of superspreading at event.

The risk of superspreading is a key determinant of whether cases of disease at the beginning

of an outbreak will lead to a major epidemic because local infection clusters can propagate

forward, snowballing into wider waves of infections. In standard models, the R survival curves

correlate strongly with those of y [1,28]. However, the added variation we see in the y curves

in **Fig 3** highlights that superspreading risk is above that expected from R alone and, further,

that chances of stochastic extinction are reduced (the histograms show $\mathbf{P}(y = 0)$ falling with

ρ). Understanding the interaction between the import rate (determined by the prevalence) and

finite event size effects is therefore essential for accurately inferring the risk of superspreading

at an event and hence the chance of epidemic establishment. Next, we demonstrate that the

realised superspreading risk can further rise if risk awareness affects event attendance.

Risk awareness controls importation rates and amplifies superspreading risk

We previously assumed that the importation rate into an event was small, constant and equal

to the population infection prevalence ρ. However, this is unrealistic as event attendance will

depend on individual preferences. Data have found that individual perceptions of infection risk

can regulate transmission dynamics and that a spectrum of risk appetites exist in a population

[13,30,31]. Many models couple behavioural changes to prevalence [10,17] and prevalence

elasticity, in which self-protective behaviours vary with prevalence, have been observed. We

hypothesise, for a fixed prevalence baseline, that heterogeneity in individual risk perception

(i.e., the risk spectrum) may mean that risk-averse individuals avoid larger events where they

expect higher chances of becoming infected. Events with large numbers of attendees are then

disproportionately likely to be attended by less risk-averse individuals (those with large risk

appetites), who have higher chances of introducing infection to the event.

We explore this idea by altering the null model from the above section in which the probability

that an event attendee is already infected is ρ. We propose a size-biased model where risk

appetite or awareness adjusts the event-scale rate of importation based on event size n. We

realise this using weights that assign a rate $\epsilon(n)$ that scales with n (see Methods) but ensures

the total infections imported into all events is conserved on average i.e., overall transmission

levels are constrained. We consider a set of m events, the i^{th} of which has size n_i. The weight

w_i is set to increase with n_i but satisfies $\sum_{i=1}^{m} w_i = \rho$. The skew of the w_i i.e., strength of the

size-bias, is controlled by the parameter r. We apply this model with differing weight strengths

r using **Eqs. (9)-(10)** and under the parameter settings from **Fig 3**, to obtain **Fig 4**.

In **Fig 4**, we study weight choices characterising two risk-awareness levels (green and red),

in which the probability that an attendee is an imported infection increases with the event size,

a relatively risk-stable case (blue) and a null model (black, dashed) completely neglecting risk-

awareness. We show corresponding import rates in panel A of **Fig 4** and compute a critical

event size n^*, at which $\epsilon(n)$ is closest to ρ. For the risk-aware models (green and red), events

above this size have higher infection risk than assumed by conventional (null) models. In

panels B-C, we illustrate that size-biasing substantially amplifies the mean and variance of the

number of infections y, doubling or tripling the risks at larger events, relative to the null model

(see insets), for the risk strength parameters and constraints we consider. This amplification

outweighs the suppression of infections at smaller events as well as susceptible depletion

caused by imports and signifies that risk awareness can strongly shape infection patterns. Our

$\epsilon(n)$ constraints limit variations in the total mean infections across events (panel D).

Fig 4: Event size bias substantially elevates the risk of infection. We compare the risk of acquiring infection at an event under models with size-biased import rates due to variability in risk perception against a null model with constant importation rate at the prevalence ρ. Panel A shows the size-biased rates $\epsilon(n)$, parametrised by r, for $m = 46$ events with sizes spanning $5:50$. Smaller r, decreasing from blue to green to red, indicates more skewed $\epsilon(n)$ functions but conserves the overall infection import rate. The critical event size n^* demarcates when $\epsilon(n)$ is closest to ρ (risk neutral event sizes). Panels B-C illustrate the resulting mean and variance of the number of infections at an event ($\mathbf{E}[y|\epsilon]$, $\mathbf{V}[y|\epsilon]$) relative to the equivalent quantities from the null model ($\mathbf{E}[y|\rho]$, $\mathbf{V}[y|\rho]$) for dispersion parameter $k = 0.1$. In panels A-C, we show medians with 95% credible intervals as computed using **Eqs. (7)-(10)**. These

Fig 5: Superspreading risk increases with risk awareness. We repeat the analyses in **Fig**

4 but for varying prevalence rates ρ at a given risk-awareness strength $r = 0.5$ in panel A and

for differing strengths at prevalence $\rho = 0.05$ in panel B. These show mean numbers of

infections $\mathbf{E}[y|\epsilon]$ under risk-aware models relative to that from the null model $\mathbf{E}[y|\rho]$ (we plot

only medians of distributions for dispersion parameter $k = 0.1$). We demonstrate how risk

awareness modulates the risk of superspreading at a medium and large sized events (dashed

vertical lines in panels A-B) by exploring tail infection survival probabilities $\mathbf{P}(y \geq c)$ in panels
C-D. See Supplement **Fig S3** and **Fig S4** for further accompanying simulations and statistics.

We show the underlying mean, variance and VM ratios of the small-scale, event reproduction
numbers as well as VM ratios for infections in Supplement **Fig S3**. These support the trends
in **Fig 4**. We also repeat this analysis for epidemics with homogeneous spread ($k = 10$) in
Supplement **Fig S4**. Interestingly, while the variance in the number of infections is smaller, the
ratios of the means and variances among risk aware and the null model (panels B-C of **Fig**
S4) are similar and rise with n, indicating that risk awareness alone can introduce additional
superspreading risk. These results (with **Fig 3**) mean that neglecting the risk spectrum within
host populations, as is done in conventional models where the probability that an attendee is
initially infected is set solely by the prevalence ρ, can lead to substantial underestimation of
the likelihood of superspreading.

We confirm this in **Fig 5**, where we illustrate how log survival or tail probabilities of infections
($\log \mathbf{P}(y \geq c)$) change with the risk awareness strength r and ρ. In panel A, we fix r and find
that the median risk of infections, relative to the null risk-neutral model, is largely unchanged.
This verifies that the skew of the size-bias from risk awareness is a key variable. In panel B,
we see how median relative risk at larger events increases with risk awareness levels (i.e., as
r falls), for fixed ρ. We highlight two event sizes $n = (24, 48)$, which have relative risks below
and above 1 (dashed lines in A-B) and examine their tail probabilities in panels C-D. In C we
find that, for both sizes, superspreading risk rises with prevalence as tail probabilities at any
threshold c scale with ρ. In D we note that risk awareness at a given prevalence can reduce
likelihood of superspreading at smaller events but considerably amplify the superspreading
risk at larger events (seen as an inversion in the ranking of curves from blue to red).

Across our simulations, this amplification from host behaviours can be as much as tenfold (2-
2.5 natural log units in the tail probabilities at $4 \leq c \leq 6$ in panel D at $n = 48$). We reinforce
these conclusions by computing the associated statistics of new infections (panel A) and our
small-scale reproduction numbers (panel B) in **Fig 6**. There we verify that the mean and
variance of the number of infections grow with prevalence and the variability in risk awareness
within the population, but that the realised heterogeneity is not inferable from the heterogeneity
in reproduction numbers. Consequently, the population risk spectrum can independently drive
increased superspreading risk at larger events that may have critical ramifications because
larger events can support more infections and contribute disproportionately to the
establishment of infection in the host population early in an epidemic. As a result, accurate
characterisation of small-scale behavioural patterns, in combination with estimation of both

the wider-scale prevalence of infection in the population and transmission heterogeneities, are
integral to correctly quantifying the risk of superspreading.

Fig 6: **Risk awareness is the key driver of superspreading risk at large events.** We
expand on the results in **Fig 5** (using the same parameter values) and compute the mean and
variance of the small-scale reproduction number R and the number of infections at the event
y. Panel A plots these for differing prevalence ρ at fixed risk-awareness strengths r (smaller
values indicate stronger risk awareness), while B varies r at fixed ρ. Increasing ρ leads to a
higher mean number of infections and more variation in the number of infections. Decreasing
r reduces the mean as well as the variance in the number of infections at smaller events but
amplifies them at larger events, increasing the risk of superspreading. The statistics of the

Discussion

Human behaviour is a key driver of infectious disease outbreaks, yet it is not often considered in detail in epidemiological modelling studies. While variations in individual perceptions of the risks associated with acquiring infection are known to shape important macroscopic properties of an epidemic, such as disease incidence time series or patterns of spread [11,17,32], few studies have explored how human behaviour impacts the chances of superspreading. This phenomenon, while infrequent and seeded at small scales such as events or gatherings, can generate disproportionate numbers of infections, which can substantially influence large-scale epidemic growth and persistence, particularly during early or emergent stages [3,5,26]. Data or even models connecting the spectrum of infection risk perceptions within host populations to attendance of events are rarely available or studied [9,24]. Here we aspired to resolve this gap by investigating the effects of plausible relationships between human behaviour and event attendance on superspreading and by highlighting how small-scale behavioural data collection can be useful for improving the accuracy of epidemic modelling and hence control.

We developed a computational framework (**Eqs. (1)-(10)**) to model small-scale transmission at events (e.g., weddings, parties, sports matches or concerts) where superspreading may arise and individual behaviour can impact pathogen dynamics. Our framework quantifies, under a standard random mixing assumption, how finite event size effects together with heterogeneities in both the transmissibility among hosts and the rate of introductions of infections to an event, contribute to the numbers of infections generated at that event. This generalises several earlier approaches [23–25] and allowed us to define a within-event (small-scale) reproduction number that measures how importations and individual-level variations impact the transmissibility at events. Our event reproduction number $R(x)$ meaningfully links to population-level characteristics through its convergence to R_0 when the event size and duration scale asymptotically (see Methods).

Using $R(x)$, we showed that previous transmissibility metrics, whether derived from branching processes [1] or earlier event-level approaches [24], can overestimate transmission and the number of infections likely to occur at an event (**Fig 2** and Supplement **Fig S1**). This result holds for any model in which the finite supply of susceptible individuals at an event is not accounted for and is exacerbated when there are multiple imported infections, because the number of susceptible individuals that any imported case can infect falls [33]. Further, this

finite-size effect highlighted that it is essential to collect data on the number of infections
introduced into events to accurately quantify superspreading risk (**Fig 3** and Supplement **Fig**
S2), which we found to depend strongly on both the number of imported infections and more
conventionally evaluated heterogeneities [3]. This insight hinted at one potential reason why
behavioural patterns might affect the likelihood of superspreading – if risk awareness alters
the distribution of infections imported into events, then it could also modulate the chances of
superspreading occurring at those events.

We explored this possibility using a parsimonious model of human behaviour. We posited that
variations in infection risk perceptions or awareness in a population might cause more risk-
averse individuals to (probabilistically) avoid larger events, which they believe present a higher
risk of acquiring infection. Our framework allowed us to model this as a size-biased weighting
on the rate of introducing infections to an event that is a function of both the wider population
prevalence and the event size. This draws on real-world observations that there is a spectrum
of self-protective behaviours in host populations that are driven by prevalence baselines and
heterogeneous risk perceptions [10,12,14,19,20]. Across numerous model simulations, we
found, for given event sizes and fixed overall prevalence values, that risk awareness amplifies
the chances of superspreading at large events (**Fig 4** and Supplement **Figs S3-4**) but limits
transmission at smaller events.

Moreover, as either the prevalence or variability in risk awareness increases (characterised
by strength parameter r), the chances of superspreading elevate (**Fig 5** and **Fig 6**). This holds
irrespective of the inherent heterogeneity in transmissibility at the event (characterised by
dispersion parameter k), which describes the impact of conventional superspreading drivers
such as pathogen biology and host characteristics. Further, the mean, variance and probability
of large numbers of infections at the event all support this trend. Since this amplification of
within-event transmission occurs precisely at those events with the capacity to support larger
numbers of infections (i.e., at larger events, where the effect of susceptible depletion when
there are more imports is less), this behavioural mechanism can have major consequences.
This may be especially critical during the sensitive, initial stages of potential epidemics, when
increased superspreading can spur growth and trigger progression from sporadic outbreaks
into sustained waves of infection [3,5].

Although these results underscore the importance of human behaviour in driving infectious
disease outbreaks, our approach, like any mathematical modelling study, involved several
simplifying assumptions. First, we assumed random mixing within events, so any susceptible
individual can interact with any infectious individual with equal probability. In reality, contact

networks form at events, and the structure of these networks may differ with the size and type of event. While frameworks for embedding contact structure in epidemiological models exist (for example, multilayer networks can be used to link risk awareness and infection structure [32–34]), they can be complex and difficult to interpret or require high resolution data that are typically unavailable [9,30,35]. We also note that our inclusion of transmission heterogeneities as in [1], together with our weighting of the risk of introductions based on event size, do reflect some features of real-world transmission networks while preserving interpretability.

Second, we assumed that event sizes and durations were pre-determined and fixed the overall import rate across all events. However, risk awareness could itself reduce event durations, sizes, frequencies and thus the prevalence of infections. Conversely, if events are prevented, due to government policy, then less risk-averse individuals may initiate their own unregulated gatherings, which could increase transmission (a rebound effect) [15]. This feedback between behaviour and environment (risk and event properties) might affect chances of superspreading [11]. Characterising this feedback is difficult due to a lack of data linking event properties and human behaviour [9,36]. Future collection and analysis of such data will be vital for grounding hypotheses and avoiding overly strong or prescriptive modelling assumptions.

Third, and relatedly, we did not attempt to model the causes of or temporal changes in the different levels of perceived infection risks among individuals. During initial epidemic stages and especially for novel pathogens, data can be sparse and erratic [35,37]. The perception of the risks associated with acquiring infection may therefore be affected by unreliable reports and major uncertainty about the true risk posed by the invading pathogen. Moreover, these risk perceptions might change across time as population immunity builds or if pharmaceuticals that reduce the severity of infections are introduced at later epidemic stages. It could even be the case that after the epidemic peak, the less risk averse individuals are actually more likely to harbour at least partial immunity. Modelling these poorly understood effects would require strong assumptions that are hard to validate.

To avoid all of the above issues, we only considered initial epidemic stages and focussed on isolating the risk-awareness induced patterns given a set of events and a prevalence baseline. We made a minimal assumption, supported by survey data on population behaviours [12–14], that there is a spectrum of risk about this baseline. Our aim was twofold – to discover how risk perceptions could impact superspreading and to show why collecting auxiliary data describing variations in epidemic-related human behaviour, such as event attendance, are necessary. Improving understanding of the coupling between risk-sensitive behaviours and epidemiology would create a platform for future investigation of the outstanding problems mentioned above.

While our modelling approach is relatively simple, it provides clear evidence that behavioural patterns can substantially amplify the risk of superspreading. Heterogeneity in infection risk perception within host populations, modelled by an event size-biased importation rate $\epsilon(n)$, translates into the potential for substantial transmission at large events during early stages of infectious disease epidemics. Because superspreading plays a pivotal role in epidemic growth and the chance of pathogen establishment, further data are required to uncover and specify the form of $\epsilon(n)$ and the mechanisms that shape the coupling between epidemiological and behavioural patterns. We bolster calls for enhanced surveillance that collects such data [9,30]. Surveys linking perceptions of infection risk with attendance at events [19] are essential to determine when variability in risk awareness may be a principal driver of superspreading. This is important to inform, design and target public health interventions more effectively [4].

Bibliography

1. Lloyd-Smith JO, Schreiber SJ, Kopp PE, Getz WM. Superspreading and the effect of individual variation on disease emergence. Nature. 2005;438: 355–359. doi:10.1038/nature04153

2. Heesterbeek H, Anderson RM, Andreasen V, Bansal S, De Angelis D, Dye C, et al. Modeling infectious disease dynamics in the complex landscape of global health. Science. 2015;347: aaa4339. doi:10.1126/science.aaa4339

3. Althouse BM, Wenger EA, Miller JC, Scarpino SV, Allard A, Hébert-Dufresne L, et al. Superspreading events in the transmission dynamics of SARS-CoV-2: Opportunities for interventions and control. PLoS Biol. 2020;18: e3000897. doi:10.1371/journal.pbio.3000897

4. Frieden TR, Lee CT. Identifying and Interrupting Superspreading Events-Implications for Control of Severe Acute Respiratory Syndrome Coronavirus 2. Emerging Infect Dis. 2020;26: 1059–1066. doi:10.3201/eid2606.200495

5. Lau MSY, Dalziel BD, Funk S, McClelland A, Tiffany A, Riley S, et al. Spatial and temporal dynamics of superspreading events in the 2014-2015 West Africa Ebola epidemic. Proc Natl Acad Sci USA. 2017;114: 2337–2342. doi:10.1073/pnas.1614595114

6. Goyal A, Reeves DB, Cardozo-Ojeda EF, Schiffer JT, Mayer BT. Viral load and contact heterogeneity predict SARS-CoV-2 transmission and super-spreading events. Elife. 2021;10. doi:10.7554/eLife.63537

7. Sun K, Wang W, Gao L, Wang Y, Luo K, Ren L, et al. Transmission heterogeneities, kinetics, and controllability of SARS-CoV-2. Science. 2021;371. doi:10.1126/science.abe2424

8. Chen PZ, Bobrovitz N, Premji Z, Koopmans M, Fisman DN, Gu FX. Heterogeneity in transmissibility and shedding SARS-CoV-2 via droplets and aerosols. Elife. 2021;10. doi:10.7554/eLife.65774

9. Funk S, Bansal S, Bauch CT, Eames KTD, Edmunds WJ, Galvani AP, et al. Nine challenges in incorporating the dynamics of behaviour in infectious diseases models. Epidemics. 2015;10: 21–25. doi:10.1016/j.epidem.2014.09.005

10. Silk MJ, Carrignon S, Bentley RA, Fefferman NH. Improving pandemic mitigation policies across communities through coupled dynamics of risk perception and infection. Proc Biol Sci. 2021;288: 20210834. doi:10.1098/rspb.2021.0834

11. Weitz JS, Park SW, Eksin C, Dushoff J. Awareness-driven behavior changes can shift the shape of epidemics away from peaks and toward plateaus, shoulders, and oscillations. Proc Natl Acad Sci USA. 2020;117: 32764–32771.

637   doi:10.1073/pnas.2009911117

12. Wise T, Zbozinek TD, Michelini G, Hagan CC, Mobbs D. Changes in risk perception and self-reported protective behaviour during the first week of the COVID-19 pandemic in the United States. R Soc Open Sci. 2020;7: 200742. doi:10.1098/rsos.200742

13. d'Andrea V, Gallotti R, Castaldo N, De Domenico M. Individual risk perception and empirical social structures shape the dynamics of infectious disease outbreaks. PLoS Comput Biol. 2022;18: e1009760. doi:10.1371/journal.pcbi.1009760

14. Wambua J, Loedy N, Jarvis CI, Wong KLM, Faes C, Grah R, et al. The influence of COVID-19 risk perception and vaccination status on the number of social contacts across Europe: insights from the CoMix study. BMC Public Health. 2023;23: 1350. doi:10.1186/s12889-023-16252-z

15. Drury J, Rogers MB, Marteau TM, Yardley L, Reicher S, Stott C. Re-opening live events and large venues after Covid-19 "lockdown": Behavioural risks and their mitigations. Saf Sci. 2021;139: 105243. doi:10.1016/j.ssci.2021.105243

16. Eksin C, Paarporn K, Weitz JS. Systematic biases in disease forecasting - The role of behavior change. Epidemics. 2019;27: 96–105. doi:10.1016/j.epidem.2019.02.004

17. Funk S, Gilad E, Watkins C, Jansen VAA. The spread of awareness and its impact on epidemic outbreaks. Proc Natl Acad Sci USA. 2009;106: 6872–6877. doi:10.1073/pnas.0810762106

18. Chan HF, Skali A, Savage DA, Stadelmann D, Torgler B. Risk attitudes and human mobility during the COVID-19 pandemic. Sci Rep. 2020;10: 19931. doi:10.1038/s41598-020-76763-2

19. Shiina A, Niitsu T, Kobori O, Idemoto K, Hashimoto T, Sasaki T, et al. Relationship between perception and anxiety about COVID-19 infection and risk behaviors for spreading infection: A national survey in Japan. Brain Behav Immun Health. 2020;6: 100101. doi:10.1016/j.bbih.2020.100101

20. Savadori L, Lauriola M. Risk perceptions and COVID-19 protective behaviors: A two-wave longitudinal study of epidemic and post-epidemic periods. Soc Sci Med. 2022;301: 114949. doi:10.1016/j.socscimed.2022.114949

21. Dryhurst S, Schneider CR, Kerr J, Freeman ALJ, Recchia G, van der Bles AM, et al. Risk perceptions of COVID-19 around the world. 2022;

22. Mossong J, Hens N, Jit M, Beutels P, Auranen K, Mikolajczyk R, et al. Social contacts and mixing patterns relevant to the spread of infectious diseases. PLoS Med. 2008;5: e74. doi:10.1371/journal.pmed.0050074

23. Tupper P, Pai S, COVID Schools Canada, Colijn C. COVID-19 cluster size and transmission rates in schools from crowdsourced case reports. Elife. 2022;11. doi:10.7554/eLife.76174

24. Tupper P, Boury H, Yerlanov M, Colijn C. Event-specific interventions to minimize COVID-19 transmission. Proc Natl Acad Sci USA. 2020;117: 32038–32045. doi:10.1073/pnas.2019324117

25. Champredon D, Fazil A, Ogden NH. Simple mathematical modelling approaches to assessing the transmission risk of SARS-CoV-2 at gatherings. Can Commun Dis Rep. 2021;47: 184–194. doi:10.14745/ccdr.v47i04a02

26. Garske T, Rhodes CJ. The effect of superspreading on epidemic outbreak size distributions. J Theor Biol. 2008;253: 228–237. doi:10.1016/j.jtbi.2008.02.038

27. Green DM, Kiss IZ, Kao RR. Parameterization of individual-based models: comparisons with deterministic mean-field models. J Theor Biol. 2006;239: 289–297. doi:10.1016/j.jtbi.2005.07.018

28. Parag KV. Sub-spreading events limit the reliable elimination of heterogeneous epidemics. J R Soc Interface. 2021;18: 20210444. doi:10.1098/rsif.2021.0444

29. Edwards DA, Ausiello D, Salzman J, Devlin T, Langer R, Beddingfield BJ, et al. Exhaled aerosol increases with COVID-19 infection, age, and obesity. Proc Natl Acad Sci USA. 2021;118. doi:10.1073/pnas.2021830118

30. Verelst F, Willem L, Beutels P. Behavioural change models for infectious disease transmission: a systematic review (2010-2015). J R Soc Interface. 2016;13.

doi:10.1098/rsif.2016.0820
31. Tan X, Li S, Wang C, Chen X, Wu X. Severe acute respiratory syndrome epidemic and
change of people's health behavior in China. Health Educ Res. 2004;19: 576–580.
doi:10.1093/her/cyg074
32. Granell C, Gómez S, Arenas A. Dynamical interplay between awareness and epidemic
spreading in multiplex networks. Phys Rev Lett. 2013;111: 128701.
doi:10.1103/PhysRevLett.111.128701
33. Moinet A, Pastor-Satorras R, Barrat A. Effect of risk perception on epidemic spreading
in temporal networks. Phys Rev E. 2018;97: 012313.
doi:10.1103/PhysRevE.97.012313
34. Bauch CT, Galvani AP. Epidemiology. Social factors in epidemiology. Science.
2013;342: 47–49. doi:10.1126/science.1244492
35. Parag KV, Donnelly CA, Zarebski AE. Quantifying the information in noisy epidemic
curves. Nat Comput Sci. 2022;2: 584–594. doi:10.1038/s43588-022-00313-1
36. Cauchemez S, Bosetti P, Cowling BJ. Managing sources of error during pandemics.
Science. 2023;379: 437–439. doi:10.1126/science.add3173
37. Yan P, Chowell G. Quantitative Methods for Investigating Infectious Disease
Outbreaks. Cham, Switzerland: Springer; 2019.
710

Funding

KVP acknowledges funding from the MRC Centre for Global Infectious Disease Analysis

(reference MR/R015600/1), jointly funded by the UK Medical Research Council (MRC) and

the UK Foreign, Commonwealth and Development Office (FCDO), under the MRC/FCDO

Concordat agreement and is also part of the EDCTP2 programme supported by the European

Union. The funders had no role in study design, data collection and analysis, decision to

publish, or manuscript preparation. For the purpose of open access, the author has applied a

'Creative Commons Attribution' (CC BY) licence to any Author Accepted Manuscript version

arising from this submission.

Data availability statement

All data and code to reproduce the analyses and figures of this work and to compute formulae

from the Methods are freely available (in MATLAB) at: https://github.com/kpzoo/smallscaleR.