

1           **Treatment of thoracic outlet syndrome to alleviate chronic migraine headache**

2                           **Short title: TOS treatment to relieve migraine**

3                           L Randall<sup>1</sup>, S. Ahn<sup>2,3</sup>, J. Weber<sup>4</sup>, Y.H. Cha<sup>5</sup>

4  
5   <sup>1</sup> Mayo Clinic Alix School of Medicine, Rochester, MN.

6   <sup>2</sup> Texas Christian University, School of Medicine, Fort Worth, TX.

7   <sup>3</sup> Dallas Fort Worth Vascular Surgery, Dallas, TX.

8   <sup>4</sup> St. Francis Medical Center, Tulsa, OK.

9   <sup>5</sup> Department of Neurology, University of Minnesota, Minneapolis, MN.

10

11 Corresponding Author:  
12 Yoon-Hee Cha  
13 717 Delaware Street SE  
14 Minneapolis, MN. 55414  
15 Email: [ycha@umn.edu](mailto:ycha@umn.edu)

16  
17 Abstract: 306

18 Manuscript: 3836

19 Tables: 4

20 Supplemental: 2

21 Figures: 3

22

23

24

**Keywords:** thoracic outlet syndrome, headache, migraine, brachial plexus, subclavian vein

25 **Conflict of interest:** The authors declared no potential conflicts of interest concerning the  
26 research, authorship, and/or publication of this article.

27 **Funding:** The authors received no financial support for the research, authorship, and/or  
28 publication of this article.

29

30

31

## 1 **ABSTRACT**

2 **Background:** Prior case reports have indicated that treating thoracic outlet syndrome (TOS)  
3 may relieve intractable migraine headaches, but the reported experience has been limited. We  
4 report in a large case series how a strategy of treating concurrent TOS can help relieve  
5 intractable migraine headaches in patients with these dual diagnoses.

6 **Methods:** Retrospective chart review for diagnostic and interventional data on patients with  
7 migraine and TOS followed by a questionnaire to investigate specific migraine features and  
8 changes in headache burden before and after treatment of TOS.

9 **Results:** 50 patients (48 women, 2 men, age = 43.9+/-12.7years) with dual diagnoses of chronic  
10 migraine and TOS were included (20 migraine with aura, 28 migraine without aura, two  
11 hemiplegic migraines). Based on review of available data, headaches had become chronic  
12 within one year of onset in 21 patients (42%) and included these characteristics: side-locked or  
13 greater severity ipsilateral to limb paresthesia (38/50 patients), presence of limb swelling (32/48  
14 patients), and worsened by recumbency (32/38 patients). Interventions included physical  
15 therapy, percutaneous transluminal venoplasty, 1st rib removal, scalenectomy, pectoralis minor  
16 tenotomy, and vein patching. Thirty-two patients needed surgery. Mean patient-reported  
17 improvement of headaches on the treated side was 72+/-26.7%; 12 patients experienced  
18 complete resolution of headaches after surgical treatment of TOS (follow-up 7.2+/-5.2 months).  
19 Questionnaire responders reported significant reductions in headache days (18.3+/-8.6 to  
20 11.1+/-10.8 days/month,  $p<0.0016$ ), severity (7.8+/-2.5 to 5.4+/-2.9,  $p<0.00079$ ), and need for  
21 emergency care (3.6+/-4.0 to 0.71+/-1.3 visits/year,  $p<0.0029$ ) after having had their TOS  
22 treated with surgery. Questionnaire responders and non-responders were not significantly  
23 different in underlying clinical features.

24 **Conclusion:** Chronic migraines can be important manifestations of TOS. Early transition to a  
25 chronic state, headaches worsened by recumbency, and headaches with lateralized myofascial

1 pain are clues to a contribution by TOS pathology. The TOS contribution to migraine has been  
2 under-recognized. Addressing it can significantly improve migraine headache burden.

3

#### 4 **INTRODUCTION**

5 Migraine headaches lead to tremendous disability and loss of productivity, particularly when they  
6 become intractable and therapeutic options have been exhausted <sup>1</sup>. While migraine is considered  
7 to be a primary headache disorder and should be diagnosed when explicit criteria as outlined in  
8 the International Classification of Headache Disorders (ICHD) are met, non-headache-related  
9 symptoms may provide clues that a structural cause may be a contributor <sup>2-4</sup>. An underrecognized  
10 contributor to migraine is compression of the brachial plexus, subclavian artery, and/or subclavian  
11 vein leading to thoracic outlet syndrome (TOS) <sup>5-7</sup>. In patients whose migraines are secondary to  
12 TOS, the associated neck and limb pain should be recognized as clues of a structural contributor.  
13 Previous studies have demonstrated that migraines can be a prominent symptom of TOS but  
14 worldwide experience has been limited <sup>8-11</sup>. In 1985, Raskin reported on 30 patients with TOS  
15 whose migraine headaches preceded the onset of arm symptoms and had significantly improved  
16 headache control after treatment of TOS <sup>8</sup>. Given the era, there had been no imaging performed  
17 to diagnose vascular TOS but contemporary criteria for neurogenic TOS had been met. In 1999,  
18 Saxton and colleagues reported detailed anatomical imaging with MRI and MRA of the brachial  
19 plexus in a 36-year-old woman with childhood-onset migraine headaches whose premonitory  
20 symptoms of the migraine headache were bilateral hand pain and paresthesia <sup>10</sup>. Her imaging  
21 showed bilateral cervical ribs and an aberrant subclavian artery compressing the vagus and  
22 recurrent laryngeal nerves, the axillary-subclavian vein, and the stellate ganglia. The patient's  
23 migraines resolved with unilateral 1<sup>st</sup> rib resection, middle and anterior scalenectomy, and  
24 extensive brachial plexus neurolysis. In 2017, Chahwala detailed a case showing retrograde flow  
25 of contrast from the subclavian veins to the internal jugular vein on arm adduction with enhanced

1 collateralization in a patient with chronic migraine <sup>9</sup>. The migraines were relieved by scalenectomy  
2 and 1<sup>st</sup> rib resection. Despite the prominence of myofascial pain and lateralized motor and sensory  
3 symptoms in migraines, recognition of TOS as an underlying contributor to intractable migraines  
4 has been almost completely overlooked since Raskin's report nearly 40 years ago.

5 TOS is designated according to whether the subclavian vein, subclavian artery, or brachial plexus  
6 is affected. Neurogenic TOS (nTOS) relates to symptoms caused by compression of the brachial  
7 plexus and manifests with paresthesia, numbness, pain, and weakness of the affected arm and  
8 hand <sup>12</sup>. Arterial TOS (aTOS) relates to compression or injury of the subclavian artery leading to  
9 symptoms such as pain and pallor due to ischemia, including the production of emboli to the hand.  
10 It is usually associated with cervical ribs or bony abnormalities in the thorax <sup>13, 14</sup>. Venous TOS  
11 (vTOS) relates to compression or thrombosis of the axillary or subclavian vein; it may present as  
12 McCleery syndrome when there is no thrombosis or as Paget Schroetter's syndrome, i.e. "effort  
13 thrombosis," when there is an associated deep vein thrombosis<sup>15, 16</sup>. Venous compression leads  
14 to swelling, warmth, and erythema. Compression of these thoracic outlet structures frequently  
15 occurs together, however, making it difficult to distinguish symptoms due to each structure <sup>5, 17</sup>.  
16 For example, subclavian vein compressions can occur in as many as 65% of patients with  
17 neurogenic TOS <sup>11, 18</sup>.

18 We took on this study to determine which clinical features should alert the clinician of when TOS  
19 is contributing to migraine headache burden. This is particularly important because many of the  
20 patients in this series had experienced severe migraine headaches for years, including starting in  
21 childhood, and only had headache relief once the TOS was treated. While most data presented  
22 is from chart review, patients with a dual diagnosis of chronic migraine and TOS were asked to  
23 provide more detailed information about treatment effects in the form of a questionnaire so that  
24 more specific features could be ascertained. The recognition of TOS contributing to migraine

1 headaches can be critical in converting intractable headaches to manageable ones and  
2 sometimes in completely relieving them.

3

#### 4 **METHODS**

##### 5 ***Participant inclusion criteria***

6 Patients experiencing chronic migraines defined by ICHD-3 criteria as  $\geq 15$  days of headaches per  
7 month in which at least 8 are migraine headaches lasting more than 4 hours a day for  $\geq 3$  months  
8 were included in the study <sup>2</sup>. Patients could also meet the criteria for New Daily Persistent  
9 Headaches (NDPH) as a secondary diagnosis. The entry criteria for TOS were based on nTOS  
10 since these are the most clearly defined criteria by the Consortium for Education and Outcomes  
11 Research on Thoracic Outlet Syndrome (CORE-TOS) and the Society for Vascular Surgery (SVS)  
12 <sup>19</sup> and can be diagnosed by clinical history and exam <sup>19</sup>. This includes at least 4 of 5 clinical  
13 features being met: 1] pain or paresthesia in the neck, upper back, shoulders, arm, and/or hands,  
14 2] symptoms exacerbated by arm elevation, repetitive use, or radiation of pain/paresthesia from  
15 the supra or infraclavicular regions, 3] a mechanism for injuries such as overuse, or prior trauma  
16 to the head, neck or chest, 4] tenderness over the scalene muscles or subcoracoid areas, 5] a  
17 positive upper limb tension test or elevated arm stress test <sup>19</sup>. In addition, concurrent vTOS was  
18 evaluated with CT chest with contrast, digital subtraction venography, and evaluation for the  
19 presence of hand and arm swelling <sup>20</sup> (**Figure 1**). All patients in this study met criteria for both  
20 nTOS and vTOS.

21 Patients underwent CT chest with contrast injected into the symptomatic arm to determine the  
22 presence of subclavian vein thrombosis, stenosis between the clavicle and 1<sup>st</sup> thoracic rib, cervical  
23 ribs, or structural chest abnormalities such as pectus excavatum and axillary masses. This was  
24 followed by digital subtraction venography in patients who required quantitative assessment

1 **(Figure 2A-C)**. Arterial TOS was evaluated by chest CT ruling out cervical ribs and by  
2 arteriography when there was suspicion of arterial involvement. MRI cervical spine, MRI shoulder,  
3 and EMGs were performed to resolve diagnostic ambiguity when there was a suggestion of  
4 cervical radiculopathy, ulnar, or median neuropathy. MRI or CT of the brain was performed to rule  
5 out other secondary causes of headache.

6 Patients in whom TOS was identified underwent an escalating series of treatments including  
7 physical therapy, percutaneous transluminal venoplasty (PTV), and surgical decompression  
8 including 1<sup>st</sup> rib resection plus scalenectomy, scalenectomy alone, pectoralis minor tenotomy, or  
9 vein patching as recommended by the consulting vascular surgeon. Patients with limb swelling  
10 were treated with diuretics. Medical management with standard headache prevention protocols  
11 continued through the TOS treatment course for as long as the patient presented for follow-up.  
12 Improvement in headaches was attributed to the treatment of TOS by temporal association after  
13 specific intervention for TOS.

14 A chart review of all patients seen by the principal investigator over the course of 3 years for a  
15 dual diagnosis of chronic migraine and TOS was performed with data extracted as the patients  
16 were being followed. Institutional approval to send questionnaires outside of regular clinical care  
17 was obtained from the Saint Francis Health Systems Institutional Research Ethics Board with  
18 protocol number 2250-19. Informed consent was obtained from all participants with research  
19 performed in accordance with the Declaration of Helsinki. Participant surveys were conducted to  
20 assess for more detailed TOS and headache features, therapies attempted, changes in symptoms  
21 following treatment, and general well-being and quality of life. Patients were sent a  
22 SurveyMonkey® weblink with a unique study identifier that only they could access.

23

24

## 1 **Measures**

2 Descriptive data on headache time course and diagnoses are reported on all patients for which  
3 that information could be reliably obtained. All post-treatment values were compared against  
4 pretreatment values (e.g. headache days before surgery vs. after surgery). Statistically significant  
5 changes in metrics such as the average number of headache days, severity of headaches, and  
6 number of yearly urgent care/emergency room visits before and after treatment of TOS were  
7 calculated as two-tailed paired t-tests with a significance of  $p<0.05$ . Data are presented for the  
8 number of participants who had sufficient information for that measure with the number of data  
9 points reported. Therefore, some data reported have different denominators.

10

## 11 **RESULTS**

### 12 *Clinical features*

13 A total of 50 patients from a base of approximately 600 patients referred for headaches met the  
14 criteria for both chronic migraine and TOS between March 2016 and April 2019. Clinical features  
15 of 48 women and two men (age 43.9+/-12.7 years, median 44 years) are detailed in **Table 1**.

16

17 Headaches were unilateral and side-locked in 17 patients, and when bilateral were worse on the  
18 side of the greater arm symptoms in an additional 21 patients (**Table 2**). Thirty-two patients  
19 endorsed swelling in their hands and arms, 16 denied swelling, and two did not comment. Twenty-  
20 two of 50 patients noted being woken up in the middle of the night with headaches; 26 of 50 noted  
21 waking up in the morning with headaches. Positional effects were known in 36 patients. Of these,  
22 30 reported having headaches that were worse being reclined, sometimes requiring them to sleep  
23 with the head elevated.

24

1 All 50 patients underwent chest CT, in which there were 26 cases of narrowed subclavian or  
2 innominate veins equally split between the right and left sides (**Table 3**). Other findings relevant  
3 to the diagnosis included three cases of cervical ribs, three cases of thymic hyperplasia causing  
4 mediastinal crowding, a thoracic schwannoma impinging on the subclavian vein of the  
5 symptomatic arm, and one aberrant 2<sup>nd</sup> rib insertion into the sternum on the symptomatic arm.  
6 MRI of the brain was available in 36 patients and head CT in 26 patients, some of whom had  
7 both. There were four cases of cerebellar ectopia, two strokes that occurred after the onset of the  
8 migraines, and one case of demyelinating disease. Fifteen patients underwent digital subtraction  
9 venography. All confirmed severe large vessel venous stenosis ( $\geq 50\%$ ) in the subclavian vein.  
10 Thirteen of these patients also had the internal jugular vein interrogated; 12 had concurrent  
11 internal jugular vein stenosis of  $\geq 50\%$ ; 6 were stenosed  $\geq 75\%$ . Seven underwent additional  
12 arteriography with two showing mild concurrent subclavian artery stenosis ( $\leq 25\%$ ).

13

14 Of the 50 patients, 15 had a lumbar puncture with CSF opening pressure measurements.  
15 Pressures ranged from 14-36 cmH<sub>2</sub>O, mean 21.8 $\pm$ 6.9 cmH<sub>2</sub>O, median 21.0 cmH<sub>2</sub>O. Only one  
16 patient in the series exhibited papilledema. This patient developed migraine headaches and TOS  
17 after a T1 schwannoma resection. She had a pressure of 36 cmH<sub>2</sub>O.

18

19 Patients had tried a mean of 4.0 $\pm$ 2.4 (median number tried = 4, range 0-11) preventative  
20 migraine medical treatments (antihypertensives, anticonvulsants, antidepressants, Botox, CGRP  
21 inhibitors) before starting treatment for TOS. Patients who eventually had surgery showed a trend  
22 towards having tried more medical treatments for migraines than those who did not have surgery  
23 (4.5 $\pm$ 1.9 vs 3.1 $\pm$ 2.4 medications, two-tailed *p*-value 0.0885).

24



1 All patients had a course of physical therapy with a total of 32 patients eventually undergoing  
2 surgical decompressive procedures during the observation period. The same patient could  
3 undergo more than one procedure. Twenty patients underwent bilateral 1<sup>st</sup> rib resections with  
4 scalenectomy, five right-sided 1<sup>st</sup> rib resections with scalenectomy, six left-sided 1<sup>st</sup> rib resections  
5 with scalenectomy, five additional separate scalenectomies, four pectoralis minor tenotomies, two  
6 vein patches (one internal jugular vein, one subclavian vein), and 13 a trial of PTV. All TOS  
7 surgical treatments were done to improve arm symptoms rather than for headaches despite  
8 improvement in headaches being of primary interest in this study. Patients were followed for an  
9 average of 7.2+/-5.2 months (median 7 months, range 1-19 months) after their last procedure.  
10 The follow-up was done by either a vascular surgeon or a neurologist. One patient in this series  
11 developed subclavian vein thrombosis that led to a pulmonary embolus before she had her 1<sup>st</sup> rib  
12 + scalenectomy surgery. Two patients had posterior circulation strokes with cryptogenic sources  
13 before being diagnosed with TOS.

14

15 Risk factors for the development of TOS are detailed in **(Supplemental Table 1)**. Thirty-seven  
16 patients had a history of trauma to the head, neck, or chest that ranged from overuse injury to the  
17 instrumentation of the neck or chest. Some patients had more than one risk factor.

18

#### 19 *Treatment response*

20 Twenty-eight of 41 patients in whom post-treatment clinical status was known reported  $\geq 50\%$   
21 better headache control after treating their TOS. Twelve of these 28 patients reported 100%  
22 headache relief. Overall headache relief in the 41 patients averaged 72+/-26.7% (range 50-100%)  
23 by their global assessments. Four reported no improvement. Two of these four patients had no  
24 improvement in headache but had resolved or majority improvement in syncopal spells. One  
25 patient had no improvement in headache but had resolved supine chest pain and palpitations.

1 Patients whose symptoms did not improve were typically in the middle of treatment at the time  
2 data were summarized and were waiting for surgery on the contralateral side. Nine patients did  
3 not follow up after treatment was initiated to know their response. The two patients with hemiplegic  
4 migraines had hemiplegic symptoms resolve with surgery (cervical rib removal in one, bilateral 1<sup>st</sup>  
5 rib resections in one). In seven patients, headaches recurred after a period of 1-5 months on the  
6 contralateral side which prompted surgery of the contralateral side.

7

## 8 **Survey participants**

9 Of the 50 patients invited to take the survey, 26 started the survey. Of these, 24 completed enough  
10 sections to allow data analysis. Of the remaining 24 who did not take the survey, one patient  
11 declined to participate, one had an inactive email address, and 22 did not respond. To determine  
12 whether there were any systematic biases in participants (n=26) and non-participants (n=24), the  
13 general characteristics of each group were compared (**Supplemental Table 2**). Demographic  
14 features were similar including sex distribution, age, years of migraine, and years of chronic  
15 migraine, but were different in duration of care (1.5+/-0.9 years participants vs 1.0+/-0.9 years  
16 non-participants,  $p=0.0276$ ) and a greater proportion of participants who had started the survey  
17 who had undergone surgery (19 of 24 participants =79%) than non-participants (11 of 24 =49%).

18

### 19 *Clinical features of survey participants*

20 Eighteen of 24 patients who completed the survey reported baseline chronic daily headaches  
21 with no headache-free days of which 11 patients had headaches that were chronic from the  
22 onset. The most frequent area of maximal pain was the skull base, with the most frequent  
23 description of the pain being, “pressure, head exploding,” followed by “throbbing.” The  
24 relationship between the onset of headache and limb symptoms of TOS was answered by 23

1 participants. Limb symptoms with headaches occurred either “always,” or “sometimes,” with  
2 headaches in 87% of participants **Table 4**.

3

#### 4 *Treatment responses from survey participants*

5 Participants rated the change in their headache burden from before to after treatment for TOS  
6 with reports corroborated with the clinical record. The average number of headache days per  
7 month decreased from 18.3+/-8.6 to 11.1+/-10.8 ( $p<0.0016$ ) with the average severity of  
8 headache on a 0-10 scale decreasing from 7.8+/-2.5 to 5.4+/-2.9 ( $p<0.00079$ ). The average  
9 number of visits to an urgent care or emergency room per year decreased from 3.6+/-4.0 to  
10 0.71+/-1.3, ( $p<0.0029$ ). For the 19 of 24 survey participants with completed data and who had  
11 undergone any surgical procedures, the survey was reported at a mean of 10.0+/-6.1 months  
12 (median 9 months, range 0.5-20.7 months) after their last surgical procedure. Of note, patients  
13 did not have to have completed all planned surgical procedures to participate in the survey.  
14 Eighteen patients made responses on a 5-point Likert scale of “strongly disagree (-2),” to “strongly  
15 agree (+2),” for general statements assessing their well-being and physical capabilities at home  
16 and at work, both before and after their TOS diagnosis and intervention. The mean change for “I  
17 am fully functional at work,” was positive (mean 1.22+/-1.48; range 0-4). Similarly, “I am fully  
18 functional at home,” indicated a positive change (mean 1.17+/-1.34; range 0-3) (**Figure 3**).

19

## 20 **DISCUSSION**

21 This study reports the real-life clinical experience of how treating concurrent TOS can alleviate  
22 chronic migraine headaches in patients with these dual diagnoses. It more than doubles the  
23 number of patients reported in the literature for which this association is important and could thus  
24 buttress the rationale for future controlled trials for this underappreciated association.

1 Thoracic outlet syndrome as a contributor to chronic migraines should be considered in patients  
2 with certain clinical features. Our study indicates that these features include: 1. Headaches  
3 ipsilateral to myofascial symptoms including skull base, neck, and shoulder pain, along with arm  
4 and hand symptoms of numbness, tingling, and weakness 2. Headaches associated with edema  
5 of the face, arm, and/or hand, 3. Symptoms worse in recumbency or that frequently wake a person  
6 from sleep, 4. Headaches developing in the context of neck and chest trauma, including postural  
7 strain, 5. Headaches that are chronic within one year of onset, including NDPH. While no feature  
8 is specific to TOS, the combination of supporting features should raise consideration for an  
9 underlying structural disorder such as TOS contributing to migraine pathology.

10 In this study, all patients met CORE TOS SVS criteria for both nTOS and vTOS through imaging  
11 and clinical exams. The nTOS component served as the signal that the thoracic outlet was  
12 compromised whereas the vTOS component explained the clinical features related to the  
13 migraines. It is possible that vTOS alone could contribute to migraines without leading to arm  
14 problems, but these patients would be difficult to find unless routine screening for TOS was  
15 performed in all migraine patients. It remains for a future study to determine whether this is  
16 reasonable.

17 The anatomy of the venous system of the head, neck, and thoracic outlet is consistent with most  
18 clinical features. Subclavian vein obstruction in vTOS impedes drainage from the external jugular  
19 vein and potentially the internal jugular vein when the obstruction is near the confluence of the  
20 innominate vein. During digital subtraction venography, the external jugular vein was frequently  
21 noted to be overdeveloped with exuberant collaterals (**Figure 2B**). This may lead to elevated  
22 venous pressure in tributaries upstream from the compression that can lead to neck pain and  
23 skull-base pressure that triggers a migraine <sup>21, 22</sup>. The areas drained by these veins include the  
24 skull base, lateral neck, periauricular, and suprascapular areas--all areas in which both swelling  
25 and pain have been noted in these patients. Because emissary veins carrying blood from the

1 internal jugular veins collateralize with the external jugular vein at the skull base, subclavian vein  
2 obstruction may potentially impact flow in the internal jugular vein leading to downstream effects  
3 on drainage from the cavernous sinus around the eye and the superior and inferior petrosal  
4 sinuses that drain the inner ear <sup>23, 24</sup>. The superior petrosal sinus sits over the trigeminal nerve  
5 and sometimes wraps around it creating a potential mechanism for the direct triggering of facial  
6 pain with venous congestion <sup>25</sup>. Moreover, the increasingly recognized role of the jugular venous  
7 system in regulating intracranial pressure may be a mechanism for headaches, particularly those  
8 triggered by postural changes <sup>26-28</sup>. Notably, most patients experienced worse headaches in the  
9 supine position when venous outflow is most dependent on the jugular pathway as opposed to  
10 the paravertebral venous plexus that carries greater venous outflow in the upright position <sup>26-28</sup>.  
11 Almost half of the patients in this series noted being woken up in the middle of the night by  
12 headaches. Further, the re-routing of venous blood through intrathoracic veins in upper extremity  
13 venous compressive disorders could be an important factor in the associated symptoms of chest  
14 pain and palpitations, particularly those worsened by supine positioning when venous pressure is  
15 high <sup>9, 10, 29, 30</sup>.

16

### 17 *Special cases*

18 The two patients with hemiplegic migraines are of particular interest. One mechanism of  
19 hemiplegic symptoms in vTOS relates to the parallel path of venous outflow between the  
20 paravertebral venous plexus and the jugular system which bidirectionally shunts with postural  
21 changes <sup>28, 30-32</sup>. Compression of the jugular system shunts venous blood to the vertebral veins  
22 which drain the cervical spinal cord <sup>33</sup>. Over-shunting of venous blood due to either fixed or  
23 frequent obstruction of the jugular venous system could lead to venous congestion and reduced  
24 perfusion pressure of the ipsilateral spinal cord. This can trigger hemiplegic symptoms. In the

1 patient with the cervical rib, TOS decompression also likely removed a component of subclavian  
2 artery compression causing limb ischemia.

3 Two patients had a history of cryptogenic embolic strokes. While one patient had a history of  
4 hyperlipidemia, she was only in her 30s and otherwise in excellent health when she developed a  
5 caudal cerebellar hemispheric stroke. The other patient was in her 40s and developed an occipital  
6 stroke in the setting of a particularly bad migraine headache. Strokes have been reported in TOS,  
7 generally in the context of obstructed flow through a stenosed subclavian artery and stagnation  
8 of blood in an adjacent aneurysm<sup>34, 35</sup>. Neither of these situations applied to our patients but we  
9 note that one of these patients had complete resolution of her headaches after addressing her  
10 nTOS and vTOS.

11

## 12 *Limitations*

13 This report expands the scope of clinical features reported on patients with both chronic migraine  
14 and TOS. It is not a controlled clinical trial, however, and thus has limitations. Baseline clinical  
15 features were reported from chart review over a limited time period with different durations of  
16 follow-up. There was no natural control arm for this study other than showing that standard  
17 migraine strategies had failed in these patients. Compared to non-participants, there were  
18 proportionately more participants who had responded to the questionnaires who had undergone  
19 surgical procedures, who came to follow-up appointments, and who had reported having better  
20 headache control at the last clinical follow-up. Thus, the true percent improvement may be lower  
21 than if all patients had responded. Mitigating these biases is that there were no significantly  
22 different baseline characteristics in clinical features between responders and non-responders to  
23 the questionnaire. Further, there was a counter bias towards poor response in that some patients  
24 were in the middle of treatment during their reporting period rather than at the end of treatment.

1 Ideally, a longitudinal approach could be taken to establish recurrent surveys over the course of  
2 several years with the caveat that more demanding studies may lead to reduced participation and  
3 still select for patients with better responses. Finally, the sample size is small relative to other  
4 headache and vascular studies but a report of 50 patients with combined TOS and chronic  
5 migraine represents the largest case series presented on this association to date in the world  
6 English literature. With the recognition that these disorders are more related than currently  
7 suspected, these cases may appreciably increase with time. This case series may help identify  
8 patients with TOS and migraine for future trials.

9

## 10 **CONCLUSION**

11 Chronic migraines can be important manifestations of TOS that can improve with the treatment  
12 of TOS. TOS should be suspected when neck pain, shoulder pain, and limb paresthesia or  
13 swelling are present, particularly when ipsilateral to the headache, and when headaches are  
14 described as a pressure that is worse with recumbency. We propose that this association is much  
15 more common than is usually appreciated due to the frequency of head, neck, and chest trauma  
16 from overuse injury including some iatrogenic causes such as central lines or neck surgery. TOS-  
17 associated headaches may be chronic from the onset and should be considered in the differential  
18 in patients who meet the criteria for migraine and NDPH. Identifying secondary causes of  
19 headaches such as TOS is of critical importance to avoid losing a window of opportunity to treat  
20 before fibrosis and venous stenosis become permanent and drastic measures such as vein  
21 patching are needed. The consideration of TOS as a possible contributor to chronic migraines  
22 may help address at least one segment of the chronic migraine population and expedite  
23 appropriate therapy. We hope that this report will help justify research resources to study the  
24 association between TOS and migraine more comprehensively.

1 **Ethics approval and consent to participate:** Institutional approval was obtained from the  
2 Saint Francis Health Systems Institutional Research Ethics Board (Tulsa, OK), protocol number  
3 2250-19. A chart review of all patients seen by the principal investigator over the course of 3.5  
4 years was reviewed for a diagnosis of thoracic outlet syndrome. Data are presented  
5 anonymously. Patients in the questionnaire study provided information with participant IDs.  
6 They were allowed to withdraw from the study at any time. All data were treated as confidential.

7

8 **Consent for publication:** Data from chart review are presented anonymously without personally  
9 identifying information. There are no identifying features in the images. Patients who were in the  
10 questionnaire study provided consent for participation and publication.

11

12 **Availability of data and materials:** The dataset supporting the conclusions of this article is  
13 available in the Harvard Dataverse:

14 <https://dataverse.harvard.edu/dataset.xhtml?persistentId=doi:10.7910/DVN/Y4AEPL>

15 **Authors' contributions:** LR performed data analysis, drafted the paper, and reviewed the  
16 interim and final manuscripts. JW provided patient data and reviewed the interim and the final  
17 manuscript. SA provided patient data and reviewed the interim and the final manuscript. YHC  
18 conceived the study, provided patient data, administered the questionnaire study, and reviewed  
19 all drafts of the manuscript.

20 **Acknowledgments:** The authors thank the patients who participated in the study. The authors  
21 received no compensation outside of routine clinical care for these patients. There was no  
22 external funding for this study and no involvement of 3<sup>rd</sup> party institutions in the collection of data  
23 or development of this manuscript.



1 **FIGURE LEGENDS:**

2 **Figure 1:** Examples of clinical signs of venous obstruction. A1] Left supraclavicular fossa, dashed  
3 arrow showing normal concavity. A2] Right supraclavicular fossa, solid arrow showing loss of  
4 definition of the lower sternocleidomastoid posterior border due to swelling in a patient with right  
5 vTOS. B1] Arms at the sides, dashed arrows showing flat external jugular veins. B2] Arms  
6 abducted to 90 degrees, solid arrows showing dilated external jugular veins in a patient with  
7 bilateral vTOS. C] Flushness and swelling of an arm in a patient with bilateral vTOS (only one  
8 arm shown), compared to the normal skin tone of the leg.

9

10 **Figure 2:** Representative examples of imaging in this cohort of chronic headache and TOS  
11 patients. Relevant abnormalities are marked by either open arrowheads or closed arrowheads.

12 A] Right subclavian vein narrowing between the clavicle and 1<sup>st</sup> rib (open arrowhead) with contrast  
13 reflux into the internal jugular vein and lack of flow into the innominate vein (closed arrowhead),  
14 B] Overdeveloped external vein (closed arrowheads) with extensive collaterals (open  
15 arrowheads) in the setting of a stenosed subclavian vein. C] 3D reconstruction of venous system  
16 showing bilateral subclavian vein narrowing (open arrowheads) as well as a narrowed right-sided  
17 internal jugular vein (closed arrowhead).

18  
19 **Figure 3:** Change in patient participants' rating of statements related to their functional  
20 capabilities before and after treating underlying TOS. Scales go from Strongly Disagree (-2) to  
21 Strongly Agree (+2) for a maximum possible change of 4 points.

22

23

24

## 1 REFERENCES

- 2 1. Serrano D, Manack AN, Reed ML, Buse DC, Varon SF, Lipton RB. Cost and predictors of  
3 lost productive time in chronic migraine and episodic migraine: results from the American  
4 Migraine Prevalence and Prevention (AMPP) Study. *Value Health*. 2013;16(1):31-8.
- 5 2. Headache Classification Committee of the International Headache Society (IHS) The  
6 International Classification of Headache Disorders, 3rd edition. *Cephalalgia*. 2018;38(1):1-211.
- 7 3. Dodick DW. Diagnosing Secondary and Primary Headache Disorders. *Continuum*  
8 (Minneap Minn). 2021;27(3):572-85.
- 9 4. Yancey JR, Sheridan R, Koren KG. Chronic daily headache: diagnosis and management. *Am*  
10 *Fam Physician*. 2014;89(8):642-8.
- 11 5. Illig KA, Donahue D, Duncan A, Freischlag J, Gelabert H, Johansen K, et al. Reporting  
12 standards of the Society for Vascular Surgery for thoracic outlet syndrome. *J Vasc Surg*.  
13 2016;64(3):e23-35.
- 14 6. Grunebach H, Arnold MW, Lum YW. Thoracic outlet syndrome. *Vasc Med*.  
15 2015;20(5):493-5.
- 16 7. Klaassen Z, Sorenson E, Tubbs RS, Arya R, Meloy P, Shah R, et al. Thoracic outlet syndrome:  
17 a neurological and vascular disorder. *Clin Anat*. 2014;27(5):724-32.
- 18 8. Raskin NH, Howard MW, Ehrenfeld WK. Headache as the leading symptom of the thoracic  
19 outlet syndrome. *Headache*. 1985;25(4):208-10.
- 20 9. Chahwala V, Tashiro J, Li X, Baqai A, Rey J, Robinson HR. Venous Thoracic Outlet Syndrome  
21 as a Cause of Intractable Migraines. *Ann Vasc Surg*. 2017;39:285 e5- e8.
- 22 10. Saxton EH, Miller TQ, Collins JD. Migraine complicated by brachial plexopathy as displayed  
23 by MRI and MRA: aberrant subclavian artery and cervical ribs. *J Natl Med Assoc*. 1999;91(6):333-  
24 41.
- 25 11. Ahn SS, Miller TJ, Chen JF, Chen SW, Feldtman R, Hwang W. Internal Jugular Vein Stenosis  
26 in Patients with Thoracic Outlet Symptoms. *Journal of Vascular Surgery*. 2013:100S.
- 27 12. Karam J, Thompson R. Neurogenic thoracic outlet syndrome. In: Hans SS, Shepard AD,  
28 Weaver MR, Bove PG, Long GW, editors. *Endovascular and Open Vascular Reconstruction: A*  
29 *practical approach*. 1st ed: CRC Press, Taylor & Francis Group; 2017. p. 377-82.
- 30 13. Gelabert M, Gelabert H. ATOS Consensus Guidelines. 2nd ed. Illig KA, Thompson RW,  
31 Freischlag JA, Donahue DM, Jordan SE, Lum YW, et al., editors. Switzerland: Springer; 2021.
- 32 14. Karam J, Thompson R. Vascular Thoracic Outlet Syndrome. In: Hans SS, Shepard AD,  
33 Weaver MR, Bove PG, Long GW, editors. *Endovascular and Open Vascular Reconstruction: A*  
34 *practical approach*. 1st ed: CRC Press, Taylore & Francis Group; 2017. p. 371-5.
- 35 15. Illig KA, Doyle AJ. A comprehensive review of Paget-Schroetter syndrome. *J Vasc Surg*.  
36 2010;51(6):1538-47.
- 37 16. Likes K, Rochlin DH, Call D, Freischlag JA. McCleery syndrome: etiology and outcome. *Vasc*  
38 *Endovascular Surg*. 2014;48(2):106-10.
- 39 17. Likes K, Rochlin DH, Call D, Freischlag JA. Coexistence of arterial compression in patients  
40 with neurogenic thoracic outlet syndrome. *JAMA Surg*. 2014;149(12):1240-3.
- 41 18. Ahn SS, Miller TJ, Chen SW, Chen JF. Internal jugular vein stenosis is common in patients  
42 presenting with neurogenic thoracic outlet syndrome. *Ann Vasc Surg*. 2014;28(4):946-50.

- 1 19. Thompson R. Diagnosis of Neurogenic Thoracic Outlet Syndrome: 2016 Consensus  
2 Guidelines and Other Strategies. 2nd ed. Illig KA, Thompson RW, Freischlag JA, Donahue DM,  
3 Jordan SE, Lum YW, et al., editors. Switzerland: Springer; 2021.
- 4 20. Illig KA. Diagnosis of vTOS: 2016 Consensus Guidelines. 2nd ed. Illig KA, Thompson RW,  
5 Freischlag JA, Donahue DM, Jordan SE, Lum YW, et al., editors. Switzerland: Springer; 2021.
- 6 21. Balachandra N, Padmalatha K, Prakash B, Ramesh B. Variation of the veins of the head  
7 and neck-external jugular and facial vein. *Int J Anat Var.* 2012;5:99-101.
- 8 22. Deslaugiers B, Vaysse P, Combes JM, Guitard J, Moscovici J, Visentin M, et al. Contribution  
9 to the study of the tributaries and the termination of the external jugular vein. *Surg Radiol Anat.*  
10 1994;16(2):173-7.
- 11 23. Pekcevik Y, Sahin H, Pekcevik R. Prevalence of clinically important posterior fossa emissary  
12 veins on CT angiography. *J Neurosci Rural Pract.* 2014;5(2):135-8.
- 13 24. Zhou D, Ding JY, Ya JY, Pan LQ, Yan F, Yang Q, et al. Understanding jugular venous outflow  
14 disturbance. *CNS Neurosci Ther.* 2018;24(6):473-82.
- 15 25. Akobo S, Tubbs RS. Superior Petrosal Sinus. In: Tubbs RS, editor. *Anatomy, Imaging and*  
16 *Surgery of the Intracranial Dural Venous Sinuses*: Elsevier; 2020. p. 117-24.
- 17 26. Holmlund P, Johansson E, Qvarlander S, Wahlin A, Ambarki K, Koskinen LD, et al. Human  
18 jugular vein collapse in the upright posture: implications for postural intracranial pressure  
19 regulation. *Fluids Barriers CNS.* 2017;14(1):17.
- 20 27. Holmlund P, Eklund A, Koskinen LD, Johansson E, Sundstrom N, Malm J, et al. Venous  
21 collapse regulates intracranial pressure in upright body positions. *Am J Physiol Regul Integr Comp*  
22 *Physiol.* 2018;314(3):R377-R85.
- 23 28. Kosugi K, Yamada Y, Yamada M, Yokoyama Y, Fujiwara H, Yoshida K, et al. Posture-induced  
24 changes in the vessels of the head and neck: evaluation using conventional supine CT and upright  
25 CT. *Sci Rep.* 2020;10(1):16623.
- 26 29. Collins JD, Saxton EH, Miller TQ, Ahn SS, Gelabert H, Carnes A. Scheuermann's disease as  
27 a model displaying the mechanism of venous obstruction in thoracic outlet syndrome and  
28 migraine patients: MRI and MRA. *J Natl Med Assoc.* 2003;95(4):298-306.
- 29 30. Marini TJ, Chughtai K, Nuffer Z, Hobbs SK, Kaproth-Joslin K. Blood finds a way: pictorial  
30 review of thoracic collateral vessels. *Insights Imaging.* 2019;10(1):63.
- 31 31. Qureshi AI, Saleem MA, Naveed O, Qureshi MA, Wallery SS. Upright Catheter-Based  
32 Cerebral Angiography. *J Vasc Interv Neurol.* 2017;9(6):14-9.
- 33 32. Griessenauer CJ, Raborn J, Foreman P, Shoja MM, Loukas M, Tubbs RS. Venous drainage  
34 of the spine and spinal cord: a comprehensive review of its history, embryology, anatomy,  
35 physiology, and pathology. *Clin Anat.* 2015;28(1):75-87.
- 36 33. Carpenter K, Decater T, Iwanaga J, Maulucci CM, Bui CJ, Dumont AS, et al. Revisiting the  
37 Vertebral Venous Plexus-A Comprehensive Review of the Literature. *World Neurosurg.*  
38 2021;145:381-95.
- 39 34. Angappan D, Garrett M, Henry C, Riddle A, Wilson JL. Pediatric Stroke due to Thoracic  
40 Outlet Syndrome Treated with Thrombolysis and Thrombectomy: A Case Report. *Children (Basel).*  
41 2022;9(6).
- 42 35. Meumann EM, Chuen J, Fitt G, Perchyonok Y, Pond F, Dewey HM. Thromboembolic stroke  
43 associated with thoracic outlet syndrome. *J Clin Neurosci.* 2014;21(5):886-9.
- 44

**Table 1: Timeline and diagnosis of chronic migraine TOS patients**

<b>Headache time frame (n=50)</b>	<i>Mean +/- s.d.</i>	<i>Median (range)</i>
Age at presentation	43.9+/-12.7 years	44 (17-78) years
Age at onset any migraine	23.9+/-14.9 years	23.5 (5-62) years
Age at onset chronic migraine	34.8+/-13.4 years	36 (5-62) years
Duration chronic migraine	9.3+/-12.7 years	5.8 (0.3-73) years
<b>Headache types (n=50)</b>		
Migraine without aura total		28
(+) NDPH*		7
Migraine with aura total		20
(+) NDPH*		1
Hemiplegic migraine		2
Chronic migraine within 1 year of headache onset		21
*New Daily Persistent Headache		

**Table 2: Laterality and associated symptoms with migraine**

<b>Associated symptoms</b>	<b>Skull base or neck pain</b>	<b>Shoulder or shoulder blade pain</b>	<b>Chest pain</b>	<b>Arm-hand numbness, tingling, weakness</b>	<b>Syncope or presyncope</b>	<b>Visual problems</b>	<b>Vertigo</b>	<b>Tinnitus</b>	<b>Ear fullness</b>
Symptom present	39	36	31	50	14	25	24	15	20
Ipsilateral to headache	24 (62%)	26 (72%)	10 (32%)	30 (60%)	na	7 (28%)	na	7 (47%)	9 (45%)
Contralateral to headache	0 (0%)	0 (0%)	0 (0%)	1 (2%)	na	0 (0%)	na	0 (0%)	0 (0%)
Bilateral	8 (21%)	5 (14%)	1 (3%)	15 (30%)	na	18 (72%)	na	4 (27%)	6 (30%)
Unclear laterality	7 (18%)	5 (14%)	20 (65%)	4 (8%)	na	0 (0%)	na	4 (27%)	5 (25%)
Symptom absent	2	3	16	0	22	14	12	19	18
Symptom unqueried	9	11	3	0	14	11	14	16	12



**Table 3: Imaging characteristics of chronic migraine TOS patients**

<b>MRI brain with or without contrast (n=36)</b>		<b>CT chest with contrast (n=50)</b>	
Normal	24	Right SCV or BCV stenosis	13
Small vessel disease	4	Left SCV or BCV stenosis	13
Cerebellar ectopia	4		
Occipital lobe stroke	1	Normal	13
Cerebellar stroke	1	Non-diagnostic (poor contrast)	3
Middle cerebral artery aneurysm	1		
Demyelination	1	<i>Structural abnormalities</i>	
Cervical spine syrinx	1	Cervical rib	3
		Thymic hyperplasia	3
<b>CT head without contrast (n=26)</b>		Thoracic schwannoma	1
Normal	21	Anomalous rib insertion	1
Arachnoid cyst	1		
Parietal cyst	1	<i>*SCV=subclavian vein</i>	
Middle cerebral artery aneurysm coiled	1	<i>*BCV=brachiocephalic vein</i>	
Vertebral occlusion	1		
Cerebellar ectopia	1		
No MRI or CT head	5		

**Table 4: Features of headache quality and temporal qualities in survey responders**

<b>Headache features</b>	<b>Participants (n=24)</b>	<b>%</b>
<b>Laterality</b>		
Predominantly unilateral	15	63
<i>Right</i>	10	42
<i>Left</i>	5	21
Alternating unilateral	4	17
Always bilateral	5	21
Unsure	1	4
<b>Area of maximum pain</b>		
Skull base	11	46
Forehead	5	21
Vertex	4	17
Temples	4	17
Around ear	1	4
<b>Headache description</b>		
"Pressure, head exploding"	10	42
"Throbbing"	5	21
"Pressure, head squeezing"	5	21
"Sharp shooting"	3	12
"Dull"	2	8
<b>Years between onset of migraine and arm symptoms</b>		
	<b>Participants (n=23)</b>	<b>%</b>
within 1 year	10	43
1-5 years	6	26
5-10 years	1	4
>10 years	6	26
<b>Relative onset of migraine to arm symptoms in years</b>		
Migraine onset before arm symptoms	17	74
Arm symptoms onset before migraine	5	22
Migraine and arm symptoms onset same time	1	4



<b>Synchronicity of headache and arm symptoms for each migraine episode</b>		
Always together	4	17
Sometimes together	16	70
Never together	3	13





