Integrative single-cell analysis of epigenomic and transcriptomic states in

patients with systemic sclerosis-associated interstitial lung disease

Shifang Li, Meijiao Gong

Laboratory of Immunology and Vaccinology, FARAH, ULiège, Liège 4000, Belgium.

SF and MJ share senior authorship.

Correspondence:

Shifang Li, fruceslee@gmail.com

Laboratory of Immunology and Vaccinology, FARAH, ULiège, Liège 4000, Belgium

Abstract

Interstitial lung disease is the primary cause of death in individuals who have

systemic sclerosis, one of the autoimmune connective tissue diseases. Understanding

the pathophysiology of the disease is crucial to developing treatment options. Here,

we performed a single-cell multi-omic analysis on lung tissue samples from patients

with systemic sclerosis-associated interstitial lung disease (SSC-ILD), profiling

chromatin accessibility and gene expression in the same samples and discovering

significant cellular heterogeneity. Systemic-venous endothelial cells (ECs) have been

shown to be pro-inflammatory and highly active. In addition, it was shown that the

transcription factor FOSL2 targets the genes involved in response to unfolded proteins

in systemic-venous ECs. Furthermore, we prioritized functional risk variants for

systemic sclerosis using a genome-wide association study. Ligand-receptor analysis

revealed that ECs significantly increased interaction with B cells via

CXCL10-CXCR3 in patients with SSC-ILD. Overall, our analysis emphasizes

epigenetic and transcriptional patterns in systemic-venous ECs, which might be

beneficial in understanding the pathogenesis of SSC-ILD.

Introduction

Systemic sclerosis (SSC) is a multi-organ disease with clinical symptoms such as

skin and internal organ fibrosis and vascular abnormalities. Because of limited

therapy choices, SSC causes significant morbidity and mortality. One of the primary

causes of SSC-related death is interstitial lung disease (ILD) [1]. However, the

processes underlying fibrosis in SSC-associated interstitial lung disease (SSC-ILD)

remain largely unknown. According to research, epithelial cell damage, activation of

innate and adaptive immunity, and recruitment and activation of fibroblasts may all

contribute to the excessive synthesis of extracellular matrix and scar formation in

SSC-ILD, similar to the mechanisms in other fibrotic lung diseases [2]. Furthermore,

in SSC-ILD, myofibroblasts have been demonstrated to be essential effector cells for

extracellular matrix remodeling [2-3]. Recent research has given more insight into the

role of SPP1 macrophages in pro-fibrosis in SSC-ILD [4]. However, the role of the

diversity of other cellular subpopulations found in healthy and SSC-ILD lungs in

shaping SSC-ILD pathophysiology is unclear.

Single-cell sequencing, which includes single-cell RNA sequencing (scRNA-seq)

and the assay for transposase-accessible chromatin sequencing (scATAC-seq), is a

valuable method for investigating the molecular and cellular heterogeneity of cell

types. EC, one of the innermost cell types lined along vessels in all organs and tissues

through hosts, is also in the bloodstream, where it can modulate inflammation by

regulating immune cell transport, activation status, and function [5-7]. Furthermore,

EC in the lungs and liver not only promotes immune homeostasis but also mediates

the balance between tolerance and inflammation [6]. However, the role of endothelial

cells (ECs) in lung fibrosis is not well-defined. A previous study showed that ECs to

mesenchymal phenotypic change was observed in the lung tissues of SSC-ILD

patients [8]. A recent study found that ECs in the lungs regulate pulmonary fibrosis

via FOXF1/R-Ras signaling [9]. However, our understanding of EC heterogeneity in

SSC-ILD at the single-cell level remains limited, particularly in the context of

transcriptome and chromatin accessibility. Here, scRNA-seq and scATAC-seq

methods were employed to investigate the role of ECs involved in SSC-ILD in order

to better understand cellular heterogeneity at the single-cell multiomics level.

Furthermore, we used the scATAC-seq with SSC genome-wide association studies

(GWAS) loci to elucidate mechanisms for functional GWAS loci.

Methods

Data download

scRNA-seq for 11 individuals and matched scATAC-seq for 5 individuals were

conducted in the recent study [4]. Therefore, all these raw fastq files were downloaded

from the Gene Expression Omnibus (accession number: GSE212109)

scATAC-seq processing

scATAC-seq data from the scATAC-seq experiment were aligned to the GRCh19

(hg19) reference genome and quantified using the Cellranger count tool (10×

Genomics, v.2.0). ArchR v1.0.2 [10] was used to construct arrow files by reading in

the available read fragments for each sample and applying the default augments. We

selected cells with less than 1000 unique fragments and enrichment at TSSs below 4

to ensure that each cell had a strong signal and was well-sequenced. Meanwhile,

doublets were inferred and filtered by ArchR. The method scOpen (v.1.0.0) [11] was

used to construct a low-dimensional matrix of the cells for dimensionality reduction.

To correct the batch effects (sample and chemistry) and integrate the data, the

Harmony (v.0.1.1) algorithm [12] was used, and cells were clustered using the Leiden

algorithm with a resolution of 1, respectively. A gene activity score matrix was built

using the method "addGeneScoreMatrix" to annotate the clusters and marker genes

were discovered for each cluster using the function "getMarkerFeatures". We also

utilized the MAGIC [13] imputed weight approach on the obtained gene activity

scores to reduce noise in the data scarcity in scATAC-seq. The clusters were annotated

using the same markers from the scRNA-seq data. Clustering with "FindClusters"

with a default resolution of 0.5 during the reclustering process to better find small

clusters. In two-dimensional space, all data were represented using uniform manifold

approximation and projection (UMAP). MACS2 was utilized to accomplish peak

calling [14], and the peak matrix was then generated using "addReproduciblePeakSet"

and these pseudobulk duplicates. Non-overlapping pseudobulk replicates were

constructed from groups of cells using the "addGroupCoverages" function with

distinct inputs for differential comparisons of cell types and clinical stages. The

pseudobulk peak set was subjected to motif enrichment and motif deviation studies.

The chromVAR [15] deviation scores for these motifs were also estimated using the

ArchR implementation. Using the "getMarkerFeatures" function, the pseudobulked

peak set was employed for differential analysis between different cell types and

disease states. To calculate the *p*-value and false discovery rate (FDR) between any

pair of samples, the Wilcoxon test and the Benjamini-Hochberg (BH) multiple tests

were utilized. FDR<=0.01 and log2-fold change (FC)>=1 were used to determine

differentially accessible distal peaks. Meanwhile, we used the ChIPseeker package's

"annotatePeak" function [16] to annotate the closest genes in the peak region using

default inputs. Tn5 insertions in genome-wide motifs were measured and normalized

by eliminating the Tn5 bias from the footprinting signal. We normalized these

footprints using mean values 200-250 from the motif center. To identify peak-to-gene

links, we used the ArchR "addPeak2GeneLinks" function and set the parameters to

"corCutOff=0.45" and "resolution=10,000", with "reducedDims" as the

dimensionality reduction results after batch correction.

Overlap of SSC GWAS SNPs with cell-type peaks

We downloaded summary statistics from López-Isac E *et al* [17] who performed

GWAS meta-analysis in 26,679 European populations involving 9,095 cases and

17,584 controls for comparison with GWASs of SSC via GWAS catalog

(https://www.ebi.ac.uk/gwas) (accession codes GCST009131). The same strategy as

described by Turner *et al* was used to detect overlap between disease-related SNPs

and cell-type peaks [18]. In brief, the lead variants were utilized as input to PLINK

(version 1.9) [19] first, and then kept variants with $r^2>0.8$ utilizing the European

population. To accommodate for SSC variants located close to a peak, a 100-bp

window with the SNP in the center was evaluated by extending the SNP location 50

bp in each direction using BEDOPS (v.2.4.37) [20]. Finally, bedtools (v.2.26.0) [21]

was used to intersect these SNP with each cell type peak.

scRNA-seq processing

For each sample, the raw fastq files were processed and aligned to the GRCh38

reference using CellRanger software with the default parameters

(https://support.10×genomics.com, version 7.1.0). Following that, the Seurat R

package v4.2.0 [22] was used for data quality control, preprocessing, and dimensional

reduction analysis, as previously described [23]. After generating gene-cell data

matrices from combined datasets, matrices were merged, and poor-quality cells with

200 or more expressed genes and mitochondrial gene percentages greater than 50

were excluded, leaving 48,622 cells for further analysis. Doublets were detected using

the Scrublet package (version 2.0.2) using the "scrub.scrub_doublets()" function [24]

and removed by Seurat. After normalizing the data using log transformation and

scaling, the Harmony R package v1.0 was used to correct batch effects (sample and

chemistry). For downstream UMAP visualization and clustering, the top 30

dimensions of Harmony embeddings were used. Cells were clustered using the

Louvain algorithm with a resolution of 0.2, yielding 16 unique cell clusters. We

utilized the Seurat package's "FindAllMarkers" function with the default parameters

for scRNA-seq differential expression analysis. A BH-adjusted p-value of less than

0.05 was regarded as statistically significant. CellTypist [25] was used to determine

the cell type annotations for each cluster based on the expression patterns of known

marker genes.

Multiomics data processing

An anchoring strategy as implemented in the Archr packages was employed to

carry out an integrated analysis of scATAC-seq and scRNA-seq data. Briefly, a

collection of "anchors" between datasets presumed to be in a comparable biological

state was first found (the "FindTransferAnchors" function) using canonical-correlation

analysis (CCA). These "anchors" are then utilized as a reference to integrate the

datasets, and the cluster labels from the scRNA-seq data can be projected onto the

scATAC-seq data using the "TransferData" function. To assess the agreement between

annotated and predicted cell labels in scATAC-seq data, an adjusted rand index was

computed.

Pseudotime and RNA velocity analysis

In the scATAC-seq and integrated datasets, we created a differentiation trajectory

and aligned single cells along it. Using batch-corrected Harmony embeddings, a

cellular trajectory was built in a low-dimension space based on the trajectory

backbone. The cells were assigned based on pseudotime value estimates, and

differential feature z-scores were used to generate a heatmap. For pseudotime

trajectory analysis in scRNA-seq, R packages Monocle3 (version 1.3.1) [26] were

utilized with their default parameters. The root point was determined as "EC venous

systemic". For RNA velocity analysis, CellRanger produced BAM files in conjunction

with velocyto were used to generate a loom file containing the quantification of

spliced and unspliced RNA [27]. Following that, we concatenated samples and

computed RNA velocity using the packages scvelo with the default settings, as

previously described [23].

Receptor-ligand pair analysis

The Python software CellPhoneDB [28] with database version 4 was used to

infer cell-cell communication networks from single-cell transcriptome data to

measure cellular crosstalk across various cell types. Only receptors and ligands

expressed in more than 5% of the cells in a given cell type were taken into account.

We retained only statistically significant averages ($p<0.05$) of interaction partners

from the CellPhoneDB output, which served as a proxy for mean expression of both

the ligand and the receptor of a given anticipated relationship.

Identifying TF target genes

To identify significantly shared TFs and their directly regulated target genes in

SSC-ILD disease, we used the method described by Kuppe *et al* [29], with minor

modifications. In brief, following pseudotime analysis, the transcription factors and

RNA expression along the trajectory were sorted and a pseudotime label for each

transcription factor/gene was assigned. To associate the selected TF with targets, we

explored the correlation of peak accessibility and gene expression to identify

peak-to-gene links and further subset these links by those containing the TF motif.

Significantly correlated links (FDR<0.0001) with a positive correlation were selected.

To build a quantitative transcription factor-gene-regulatory network, we estimated the

correlation of the transcription factor binding activity between scATAC-seq and target

gene expression from scRNA-seq data, and only those interactions with a Pearson

correlation >0.8 were considered.

Results and Discussion

Single-cell landscape in systemic sclerosis-associated interstitial lung disease

A total of 48,622 cells were investigated using scRNA-seq data (**Supplemental**
Figure 1A), including 18,591 cells from 6 healthy individuals and 30,031 cells from 5
SSC-ILD patients, while 2 samples from SSC-ILD and the remaining 3 samples were
included in scATAC-seq (**Supplemental Figure 1B**). Following that, batch
adjustment for samples and chemistry was conducted using the R package "Harmony"
on both the scATAC-seq and scRNA-seq datasets. Based on the expression of
lineage-specific markers, 15 main cell types in the lung were identified in scRNA-seq
(**Figure 1A-1C**). In detail, macrophages, monocytes, T cells, monocyte-derived Mph,
multiciliated, fibroblasts, ECs, basal resting, NK cells, alveolar type 2 epithelial cells
(AT2), mast cells, smooth muscle, AT1, B cells, lymphatic EC mature, and an
unidentified cell population were reported (**Figure 1C**). We obtained a total of 14,717
high-quality cells after filtering for scATAC-seq and identified 13 clusters in ArchR
(**Figure 1D-1E**). 12 cell types were annotated based on gene activity scores provided
to each cluster (**Supplemental Figure 1C**). Label transfer was accomplished by
generating a gene-activity matrix from scATAC-seq data, which is a measurement of
chromatin accessibility inside the gene body and promoter of protein-coding genes.
Transfer anchors have been identified between the "reference" scRNA-seq dataset and
the "query" gene activity matrix, and anticipated cell types were assigned. When
comparing the annotated label using the scRNA-seq dataset, the 12 cell types
identified for scATAC-seq showed high consistency, demonstrating that scATAC-seq
is equivalent to scRNA-seq in the detection and assignment of cell identities (**Figure**
1F). Following that, the MACS2 was used to evaluate the differences in chromatin
accessibility (DAR) between cell types. A master set of 50,229 peaks was generated
by using a log2-fold-change threshold of 1 and adjusted *p*-values to measure DAR
significance (FDR<=0.01). Furthermore, 26% of DAR were found at a promoter
region less than 3 kb from the nearest transcriptional start point (**Figure 1G**). Intronic
was the second most common site, and DAR distribution was relatively consistent
across cell types. Meanwhile, motif enrichment analysis for these marker peaks was
undertaken to identify the TFs likely driving the regulatory profiles in each cell type
(**Figure 1H**). Top enriched motifs in T cells (ERG, ELK4, and ERF) were discovered.

Furthermore, SOX13 motifs were shown to be enriched in ECs, as well as ZNF238

and ZBTB42 motifs in fibroblasts. Overall, the integrated dataset allows us to detect

cell type heterogeneity at the transcriptional and chromatin accessibility levels

(**Figure 1I**).

Heterogeneity of endothelial cell subset

Several investigations have demonstrated that ECs have a role in the

pathophysiology of SSC [30-31]. The chromatin accessibility of ECs from control and

SSC-ILD lungs, however, remains unclear. To investigate the heterogeneity of ECs

subsets in SSC-ILD, we first examined the differences in ECs subsets between HC

and SSC-ILD patients in scRNA-seq including 11 individuals due to the higher

statistical power. Following the re-clustered ECs (4,492 cells), 5 cell subpopulations

were identified based on particular gene expression (**Figure 2A** and **Supplemental**

Figure 2A-2C). Interestingly, our study identified two cell subpopulations of ECs

with unequal frequencies in SSC-ILD patients and healthy individuals ($p<0.01$)

(**Figure 2B** and **Supplemental Figure 2D**). The frequency of SSC-ILD was found to

be higher in EC venous systemic, which substantially expressed RBP7; however, the

frequency of SSC-ILD was significantly lower in EC aerocyte capillary, which highly

expressed TBX2. Our findings from RNA velocity and pseudotime analysis utilizing

scATAC-seq revealed the differentiation of EC venous systemic to EC aerocyte

capillary, indicating that EC venous systemic could serve as a precursor for EC

aerocyte capillary (**Supplemental Figure 2E-2F**).

The integrated analysis of scRNA-seq and scATAC-seq was performed to better

understand the chromatin structure of ECs subpopulations (**Figure 2C-2D**). Label

transfer from scRNA-seq to scATAC-seq revealed the three cell types, with a

consistent lower frequency for EC venous systemic and a greater frequency for EC

aerocyte capillary, as expected (**Figure 2C-2D** and **Supplemental Figure 2G-2H**). A

pseudotime analysis was undertaken to uncover the dynamics and modulation of cell

fate decisions in EC subpopulations (**Figure 2E**). A lineage route from EC venous

systemic to EC aerocyte capillary was discovered by pseudotime analysis utilizing

scATAC-seq. The examination of *cis*-elements near certain genes and the accessibility

of TF motifs revealed distinct regulatory patterns of accessibility over the pseudotime (**Figure 2F-2H**). Interestingly, *cis*-element accessibility in some loci involved in inflammatory response, such as IL1R1 and SELE, as well as response to unfolded protein, such as SERPINH1 and DNAJB1, gradually decreased from EC venous systemic to EC aerocyte capillary, demonstrating that the expression of inflammatory response-related genes was accurately regulated by a distinct *cis*-element network in the different states (**Figure 2F-2G**). The distinct TF activities across the pseudotime of EC venous systemic to EC aerocyte capillary paths were revealed by integrating chromVAR TF-activity and TF RNA expression studies (**Figure 2H**). For example, by assigning a path of accessibility from differentiated EC venous systemic to EC aerocyte capillary, we identified enriched MEOX1 motifs at the start of the trajectory, followed by enrichment of FOS, FOSL2, and JUNB motifs in EC aerocyte capillary, and finally PBX1, KLF11, and KLF4 motifs. According to the trajectory analysis, the EC venous systemic showed significantly higher occupancy in FOSL2 in the footprint analysis than the EC aerocyte capillary, while the EC aerocyte capillary showed a more pronounced DNA occupancy of KLF4, implying a role for FOSL2 and KLF4 as distinct transcription factors in shaping EC maturation (**Figure 2I**). We used a recently described approach to identify probable TF target genes based on the merged scATAC-seq and scRNA-seq datasets to demonstrate the TFs regulatory program network in the ECs [28]. Only Pearson correlations greater than 0.8 between transcription factor binding activity from scATAC-seq and target gene expression from scRNA-seq data were evaluated. Using this method, we discovered 456 genes controlled by KLF4 and 179 genes regulated by FOSL2 (**Figure 2J**). For example, four genes implicated in the response to unfolded protein, namely HSPA8, HSP90AB1, DNAJA1, and HSPB1, were shown to be regulated by FOSL2, which likewise showed decreased activity along the pseudotime axis (**Figure 2K**). Furthermore, our analysis found that KLF4 regulates the expression of genes involved in immune response regulation, such as BTNL9, ITGA4, and ICAM2, in EC aerocyte capillaries. In the pseudotime analysis, we found increased KLF4 binding activity and SITGA4 expression, supporting the significance of KLF4 as a possible activator of

immune response activation (**Figure 2K**). The chromVAR motif variations in the EC

venous systemic and EC aerocyte capillary between HC and SSC-ILD patients were

also compared (**Figure 2L**). AP-1 factors, including FOS and JUNB, were shown to

be abundant in both EC venous systemic and EC aerocyte capillary of SSC-ILD

patients. Furthermore, transcription factors including as GATA2 and GATA3 that are

involved in the negative regulation of ECs apoptosis were shown to be abundant in

the EC venous systemic of healthy individuals. Overall, our analysis indicated

epigenetic reprogramming in the formation of ECs, with activation characteristics and

compositional and epigenomic abnormalities in SSC-ILD patients' EC aerocyte

capillaries.

Annotation of target cell types at SSC GWAS loci

Indeed, the paired scATAC-seq/scRNA-seq datasets enabled us to deconstruct

the mechanisms underlying previously reported causative risk variations from

GWASs and to uncover disease-relevant cell types associated with these loci. Several

studies have found that noncoding GWAS variants are concentrated in CREs and

frequently function in cell-type-specific ways [32-34]. In this study, a multi-tiered

strategy was employed to prioritize putative functional SSC GWAS variants, as

described by Turner *et al* [18]. In brief, after identifying CRE variants, we compared

SSC lead variants (and variants in high LD ($r^2>0.8$; EUR)) from the most recent SSC

GWAS meta-analyses in a European cohort of 26,679 individuals with scATAC peaks

(**Figure 3A**). This identified a subset of variants (50 bp) that overlapped GWASs loci

and marker peaks, with the majority of SSC SNPs located within macrophage peaks,

followed by monocytes, T cells, AT2/basal resting cells, and B cells peaks (**Figure

3B**). We next identified target cell types for SSC regulatory variants based on these

overlaps. Several top potential SSC variants, for example, map to cell-type-specific

peaks, such as rs3792783 at the TNIP1 locus within an AT2 and basal resting cells

peak and rs1005714 at the NUP85 locus within macrophages (**Figure 3C**). Other SSC

variants, such as rs2305743 at IL12RB1 and rs1378942 at MIR4513, map to peaks

shared by T cells, monocytes, and macrophages. We reclustered AT2 using a

scRNA-seq dataset of 11 individuals to characterize its role in SSC-ILD, and four

clusters were identified (**Supplemental Figure 3A-3C**). Notably, the frequencies of

cluster 3, which highly expressed SERPINF1, were considerably greater in patients

with SSC-ILD (**Supplemental Figure 3D**). RNA velocity analysis suggests a C1 to

C3 lineage trajectory (**Supplemental Figure 3E**). Due to the small number of cells

observed in scRNA-seq for AT2 cluster 3, which may represent a rare cell population,

scATAC-seq did not disclose this subpopulation (data not shown). Furthermore,

because of the small number of samples included in the study, no substantial overlap

with GWAS loci was discovered for the cell-type peaks of fibroblasts, one of the main

effector cells in the extracellular matrix remodelling of the disease [3]. A larger

dataset should be used in the future to explore these subpopulations in depth.

Because noncoding variants cannot always regulate the adjacent gene, we used

co-accessibility and scRNA-seq integration to link candidate variants to target gene

expression (**Supplemental Figure 4A**). When all cell types were aggregated, this

orthogonal method yielded a total of 31,432 Peak2Gene linkages (**Supplemental**

Figure 4B). It is worth noting that the macrophages peak-containing variant,

rs1005714, has a co-accessibility and gene expression relationship with MRPS7, and

is also an MRPS7 lung-specific expression quantitative trait locus (eQTL) in the

Genotype-Tissue Expression project (GTEx) (**Figure 3E**). MRSPS7, but not NUP85,

is differently expressed between healthy individuals and SSC-ILD patients

(uncorrected p=2.9e-05) (**Supplemental Figure 4C**). These results suggested that

MRSPS7 could be a candidate gene associated with SSC, a large dataset as well as

experiments should be performed to validate these results.

Cell-cell communication network

To identify the reciprocal communication between different cell types in the lung,

we investigated the accumulated ligand/receptor interaction database CellPhoneDB to

detect reciprocal communication. The number of cell-cell interactions was increased

in SSC-ILD patients than in healthy individuals, as expected (**Figure 4A**). Fibroblasts,

in particular, demonstrated the greatest capacity for cell-cell interactions and shared

an increased number of projected connections with lymphatic EC mature, ECs, and

fibroblasts in patients with SSC-ILD. Furthermore, with the exception of

monocyte-derived Mph, ECs increased cell-cell interactions with practically all other

cell types in SSC-ILD patients. These findings further suggest that ECs may play an

essential role in disease progression.

In SSC-ILD patients, ECs had enhanced CXCL10-DPP4 and CXCL10-CXCR3

interactions with lymphatic EC mature and B cells (**Figure 4B**). According to gene

ontology enrichment analysis, CXCL10 was substantially enriched in the RIG-I-like

receptor signaling pathway and ECs activation (p.adj<0.01), whereas CXCL10 and

CXCR3 were significantly involved in the inflammatory response (p.adj=0.00081).

CXCL10 and its receptors CXCR3 were shown to be more accessible in SSC-ILD

patients, with increased RNA expression of CXCL10 and its receptor CXCR3 (**Figure**

4C-4D). Interestingly, TNFSF10-RIPK1 and TNF-FFAR2 signaling was anticipated

to be active during AT2 and monocyte interactions with other cell types in SSC-ILD

patients. Furthermore, SOSTDC1-LRP6 interactions were expected to be reduced in

the communication between EC cells with basal resting cells and AT2 in SSC-ILD

patients (**Figure 4B**). In fact, this has been identified as a critical regulatory

interaction in the negative control of the canonical Wnt signaling pathway [35].

Overall, our findings point to a complicated interaction network between healthy

individuals and patients. More research is needed to determine whether ECs are

involved in the regulation that leads to the onset of the inflammatory response in B

cells.

In conclusion, by integrating single-cell multiomics data, our research

demonstrated that systemic-venous ECs may play a significant pro-inflammatory role

in SSC patients, providing insights into the pathophysiology of SSC and therapeutic

alternatives.

Competing interests

None.

Acknowledgments

The authors would like to thank Anna and her colleagues for their nice work and the

generation of the single-cell multiomics dataset.

 Figure Legend

Figure 1 An overview of the epigenetic and transcriptional landscape of cell types

in the lungs of SSC-ILD patients and healthy individuals. (**A**) UMAP projection of

48,622 scRNA-seq profiles of cell types in the lung from 11 individuals. Individual

cells are represented by dots, and different cell types are indicated by different colors.

(**B**) Dot plots of gene expression of the marker genes in the scRNA-seq dataset. The

size of the dots represents the percentage of cells in each cluster that have the gene of

interest. Color intensity denoted the standardized gene expression level. (**C**) Barplots

depicting the cell-type proportions of scRNA-seq data across all samples. The various

colors represent the major cell types. (**D**) UMAP projection of 14,717 scATAC-seq

profiles of lung cell types from 5 individuals. Individual cells are represented by dots,

while different cell types are represented by different colors. (**E**) Barplots of

scATAC-seq cell-type proportions across all samples. The colors represent the several

major cell types. (**F**) Validation of the scATAC-seq cell type annotation using paired 5

scRNA-seq data. Heatmap displaying evaluation findings based on the adjusted rand

index (ARI). (**G**) Annotated DAR location bar plot for each cell type. (**H**) Heatmap of

TF motifs found in peak sequences of cell-type markers. The color indicates the

normalized motif enrichment score in ArchR obtained with HOMER and the

hypergeometric test. (**I**) Multi-omics integration for scATAC-seq and scRNA-seq data

processing.

Figure 2 Epigenomic and transcriptional signatures of EC subsets. (**A**) UMAP

projection of 4,492 EC subpopulation scATAC-seq profiles. Individual cells are

represented by dots, while different cell types are represented by different colors. (**B**)

Differences in chosen population proportions across healthy individuals (n=6) and

SSC-ILD groups (n=5). The Wilcoxon test was used to get the *p*-values. (**C**) UMAP

projection of 1,760 scATAC-seq profiles of lung cell types from 5 individuals.

Individual cells are represented by dots, while different cell types are represented by

different colors. (**D**) Barplots of scRNA-seq data cell-type proportions across all

samples. The colors represent the indicated cell types. (**E**) UMAP depicting the ECs

lineage trajectory. Pseudotime values were superimposed on the UMAP embedding;

the smoothed line and arrow indicate the representation of the spline fit trajectory

route. (**F**) Heatmaps depicting the 323 ordered gene-score and matched RNA

expression trajectory from EC venous systemic to EC aerocyte capillary across

pseudotime. (**G**) Accessibility tracks surrounding the IL1R (up) and SELE (down)

loci. (**H**) Heatmaps of 10 positive TF regulators whose TF motif activity (left) and

corresponding gene expression (right) are positively associated across the ECs

differentiation pseudotime trajectory in (E). (**I**) Comparison of aggregate TF

footprints for FOSL2 and KLF4 in EC venous systemic and EC aerocyte capillary,

respectively. (**J**) A TF regulatory network depicting the TFs (FOSL2 and KLF4) and

probable target genes. (**K**) chromVAR bias-corrected deviation scores, RNA

expression, and selected target gene expression across ECs pseudotime for the

specified TFs. In an individual pseudotime-ordered scATAC-seq profile, each dot

indicates the deviation score or RNA expression. The smoothed fit spanning

pseudotime and chromVAR deviation scores or RNA expression is represented by the

line. (**L**) Volcano plots exhibiting the differential TF motif accessibility in the

chromVAR TF bias-corrected deviation between health and SSC-ILD individuals in

the identified ECs subpopulations using the mean TF motif accessibility. The *p*-values

were obtained using a Wilcoxon test, and the FDR was corrected using the

Benjamini-Hochberg method.

Figure 3 scATAC-seq identifies mechanisms for functional SSC GWAS variants.

(**A**) Prioritizing potential SSC-associated GWAS SNPs using a multi-tiered method,

as described by Turner *et al*. (**B**) LD-expanded ($r^2>0.8$; EUR) SSC GWAS variants

(50 bp) overlap with cell-type peaks. (**C**) Heatmap displaying the number of peaks per

cell type that intersect SSC GWAS loci. (**D**) Normalized chromatin accessibility

landscape for cell-type-specific pseudobulk tracks in the vicinity of the TNIP1 (left)

and NUP85 (right) loci. Below the ATAC-seq tracks are the positions of the

ATAC-seq peaks, the GWAS lead SNP, the SNP candidates in LD with the lead SNP,

and the candidate functional SNP. (**E**) Lung eQTL in GTEx for rs3792783 and

rs1005714. The G allele for rs3792783 is strongly related to lower TNIP1 mRNA

levels in GTEx lung (left two panels). The G allele for rs1005714 is strongly related

to elevated MRPS7 and NUP85 mRNA levels (right two panels). The eQTL *p*-value is

displayed as a result of the GTEx pipeline's linear regression between genotype and

normalized gene expression levels. The boxplot within the violin plot includes the

median and interquartile range (IQR) from 25% to 75%.

Figure 4 The cellular-cellular communications of cell types in lung. (**A**) A dotplot

depicting the differences in ligand-receptor interaction events between cell types in

the healthy and SSC-ILD groups. The dotplot (left) and heatmap (right) show the

number of interactions and fold changes (SSC-ILD vs healthy individuals). (**B**) Circos

plots depicting genes implicated in ligand-receptor interactions between cell types.

Color scale is proportional to interaction strength as assessed by CellPhoneDB

interaction means of statistically significant interactions. (**C**) Dot plots demonstrating

an increase in gene expression for the specified genes. The size of the dots represents

the percentage of cells that express the marker. The color scale shows the average

gene expression levels. The arrows represent ligand-receptor pairings generated from

CellPhoneDB. The X represents signaling that has been disabled. (**D**) Single-cell

chromatin accessibility in the CXCL10 and CXCR3 loci as seen by genome browser

tracks.

Figure S1 shows an overview of the single-cell RNA and ATAC datasets. (**A**)

UMAP embedding of scRNA-seq data from all samples for cell types, samples, and

disease states. (**B**) UMAP embedding of scATAC-seq data from all samples for cell

types, samples, and disease states. CCL18 (macrophages), S100A8 (monocytes),

CD3D (T cells), FCN3 (endothelial cells), TMEM190 (multiciliated), DCN

(fibroblasts), BPIFB1 (basal resting), SFTPA1 (AT2), TPSAB1 (mast cells), ACTA2

(smooth muscle), CD79A (B cells), and CCL21 (lymphatic EC mature).

Figure S2 Sub-clustering of ECs and integration of scRNA and scATAC datasets.

(**A**) UMAP embedding of scRNA-seq data from 11 samples in the SSC group and
health individuals. (**B**) Dot plots showing marker gene expression in the scRNA-seq
dataset. The dot size represents the percentage of cells in each cluster that have the
gene of interest. The color intensity reflected the standardized gene expression level.
(**C**) Barplots of scRNA-seq data cell-type proportions across all samples. The colors
represent the several primary cell kinds. (**D**) Proportions of EC subpopulations differ
between healthy individuals (n=6) and SSC-ILD groups (n=5). The Wilcoxon test was
used to compute the *p*-values. (**E**) Monocle 3 inferred scRNA-seq EC venous
systemic differentiation route towards EC aerocyte capillary. Pseudotime colors the
cells. (**F**) RNA velocity cell trajectory analysis of ECs. (**G**) UMAP plot of
scATACA-seq dataset by gene score for cell-type marker genes, including RPS29 (EC
venous systemic), DKK2 (EC arterial), and APLN (EC aerocyte capillaries). (**H**)
Validation of scATAC-seq cell type annotation using scRNA-seq data. Heatmap
showing evaluation results by adjusted rand index (ARI).

Figure S3 Sub-clustering of AT2 using scRNA datasets. (**A**) UMAP embedding of
scRNA-seq data from 11 SSC and healthy individuals. (**B**) Dot plots showing marker
gene expression in the scRNA-seq dataset. The dot size represents the percentage of
cells in each cluster that have the gene of interest. The color intensity reflected the
standardized gene expression level. (**C**) Barplots of scRNA-seq data cell-type
proportions across all samples. The colors represent the several primary cell kinds. (**D**)
Differences in the proportions of four AT2 clusters among healthy individuals (n=6)
and SSC-ILD groups (n=5). The Wilcoxon test was used to compute the *p*-values. (**E**)
AT2 RNA velocity cell trajectory analysis. AT2, alveolar type 2 epithelial cell.

Figure S4 Co-accessibility of scATAC-seq and integration with scRNA-seq link
putative regulatory elements to target gene. (**A**) Heatmap summary of ArchR
Peak2Gene connections (n=31,432) at a resolution of 10 kb, where chromatin
accessibility is substantially linked with target gene expression. The Z-scores for
scATAC-seq peak accessibility are shown on the left, while the Z-scores for RNA

expression are shown on the right. (**B**) *cis*-regulatory architecture and lung cell types at the MRPS7 GWAS loci. (**C**) Violin plot of gene expression in the indicated illness stages. Each dot represents the expression of a single gene in a single cell.

Supplementary Table 1. The statistics generated in the study.

Reference

Steen VD, Medsger TA. Changes in causes of death in systemic sclerosis, 1972-2002. *Ann Rheum Dis* 2007;66:940-4.

Tamby MC, Chanseaud Y, Guillevin L, et al. New insights into the pathogenesis of systemic sclerosis. *Autoimmun Rev* 2003;2:152-7.

Valenzi E, Bulik M, Tabib T, et al. Single-cell analysis reveals fibroblast heterogeneity and myofibroblasts in systemic sclerosis-associated interstitial lung disease. *Ann Rheum Dis* 2019;78:1379-87.

Papazoglou A, Huang M, Bulik M, et al. Epigenetic Regulation of Profibrotic Macrophages in Systemic Sclerosis-Associated Interstitial Lung Disease. *Arthritis Rheumatol* 2022;74:2003-14.

Shao Y, Saredy J, Yang WY, et al. Vascular Endothelial Cells and Innate Immunity. *Arterioscler Thromb Vasc Biol* 2020;40:E138-52.

Amersfoort J, Eelen G, Carmeliet P. Immunomodulation by endothelial cells-partnering up with the immune system? *Nat Rev Immunol* 2022;22:576-88.

Lu Y, Sun Y, Xu K, et al. Editorial: Endothelial cells as innate immune cells. *Front Immunol* 2022;13:1-5.

Mendoza FA, Piera-Velazquez S, Farber JL, et al. Endothelial Cells Expressing Endothelial and Mesenchymal Cell Gene Products in Lung Tissue from Patients with Systemic Sclerosis-Associated Interstitial Lung Disease. *Arthritis Rheumatol* 2016;68:210-7.

Bian F, Lan YW, Zhao S, et al. Lung endothelial cells regulate pulmonary fibrosis through FOXF1/R-Ras signaling. *Nat Commun* 2023;14.

Granja, JM, Corces MR, Pierce SE. et al. ArchR is a scalable software package

for integrative single-cell chromatin accessibility analysis. *Nat Genet* 2021;53:403-411.

Li Z, Kuppe C, Ziegler S, et al. Chromatin-accessibility estimation from single-cell ATAC-seq data with scOpen. *Nat Commun* 2021;12.

Korsunsky I, Millard N, Fan J, *et al*. Fast, sensitive and accurate integration of single-cell data with harmony. *Nat Methods* 2019;16:1289-96.

van Dijk D, Sharma R, Nainys J, et al. Recovering Gene Interactions from Single-Cell Data Using Data Diffusion. *Cell* 2018;174:716-729.e27.

Zhang Y, Liu T, Meyer CA, et al. Model-based analysis of ChIP-Seq (MACS). Genome Biol 2008;9.

Schep A, Wu B, Buenrostro J. et al. chromVAR: inferring transcription-factor-associated accessibility from single-cell epigenomic data. *Nat Methods* 2017;14:975-978.

Yu G, Wang LG, He QY. ChIP seeker: An R/Bioconductor package for ChIP peak annotation, comparison and visualization. *Bioinformatics* 2015;31:2382-3.

López-Isac E, Acosta-Herrera M, Kerick M, et al. GWAS for systemic sclerosis identifies multiple risk loci and highlights fibrotic and vasculopathy pathways. *Nat Commun* 2019;10:1-14.

Turner AW, Hu SS, Mosquera JV, et al. Single-nucleus chromatin accessibility profiling highlights regulatory mechanisms of coronary artery disease risk. *Nat Genet* 2022;54:804-16.

Purcell S, Neale B, Todd-Brown K, et al. PLINK: A tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 2007;81:559-75.

Neph S, Kuehn MS, Reynolds AP, et al. BEDOPS: High-performance genomic feature operations. *Bioinformatics* 2012;28:1919-20.

Quinlan AR, Hall IM. BEDTools: A flexible suite of utilities for comparing genomic features. *Bioinformatics* 2010;26:841-2.

Butler A, Hoffman P, Smibert P, et al. Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol* 2018;36:411-20.

Li SF & Gong MJ. Single-cell RNA sequencing reveals the heterogeneity of

endothelial cells in human dupuytren's disease. Preprint in *Researchsquare.* https://doi.org/10.21203/rs.3.rs-3005025/v2

Wolock SL, Lopez R, Klein AM. Scrublet: Computational Identification of Cell Doublets in Single-Cell Transcriptomic Data. *Cell Syst* 2019;8:281-291.e9.

Domínguez Conde C, Xu C, Jarvis LB, et al. Cross-tissue immune cell analysis reveals tissue-specific features in humans. *Science* (80-) 2022;376.

Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, et al. The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. *Nat Biotechnol* 2014. 32, 381-386.

La Manno G, Soldatov R, Zeisel A, et al. RNA velocity of single cells. *Nature* 2018;560:494-8.

Garcia-Alonso L, Lorenzi V, Mazzeo CI, et al. Single-cell roadmap of human gonadal development. *Nature* 2022;607:540-7.

Kuppe C, Ramirez Flores RO, Li Z, et al. Spatial multi-omic map of human myocardial infarction. *Nature* 2022;608:766-77.

Perelas A, Silver RM, Arrossi A V., et al. Systemic sclerosis-associated interstitial lung disease. *Lancet Respir Med* 2020;8:304-20.

Di Martino ML, Frau A, Losa F, et al. Role of circulating endothelial cells in assessing the severity of systemic sclerosis and predicting its clinical worsening. *Sci Rep* 2021;11:1-9.

Corces MR, Shcherbina A, Kundu S, et al. Single-cell epigenomic analyses implicate candidate causal variants at inherited risk loci for Alzheimer's and Parkinson's diseases. *Nat Genet* 2020;52:1158-68.

Alexi N, Inge RH, Nicole GC, et al. Brain cell type-specific enhancer-promoter interactome maps and disease-risk association. *Science* (80-) 2019;1139:1134-9.

Maurano MT, Humbert R, Rynes E, et al. Systematic localization of common disease-associated variation in regulatory DNA. *Science* (80-) 2012;337:1190-5.

Ahn Y, Sanderson BW, Klein OD, et al. Inhibition of Wnt signaling by wise (Sostdc1) and negative feedback from Shh controls tooth number and patterning. *Development* 2010;137:3221-31.

Fig 1

Fig 2

Fig 3

Fig 4

Fig S2

Fig S4