1 **Opportunistic PSA-free prostate cancer screening utilising biparametric MRI**

2 **(VISIONING)**

3
4

5 Wetterauer C.[1,2,3*]; Matthias M.[1,3*]; Pueschel H.[1,3]; Deckart A.[1]; Bubendorf L.[3,4];
6 Mortezavi A. [1,3]; Arbelaez E. [1]; Winkel D.J.[3,5], Heye T.[3,5], Boll D.T.[3,5]; Merkle E.[3,5];
7 Hayoz S.[6]; Seifert H.H.[1,3]; Rentsch C.A.[1,3]

8
[1] Department of Urology, University Hospital Basel, Basel, Switzerland
[2] Department of Medicine, Faculty of Medicine and Dentistry, Danube Private
University, Krems, Austria
[3] University of Basel, Switzerland,
[4] Institute of Pathology, University Hospital Basel, Switzerland
[5] Department of Radiology, University Hospital Basel, Basel, Switzerland
[6] Swiss group for clinical cancer research (SAKK)
16
17 *authors contributed equally

18
19
20
21 Word count abstract: 300
22 Word count manuscript: 2499
23
24
25
26

27 Corresponding authors:
28
29 C. A. Rentsch; MD-PhD
30 Department of Urology
31 University Hospital Basel
32 Spitalstr. 21, 4031 Basel
33 Switzerland
34
35 C. Wetterauer, MD
36 Department of Urology
37 University Hospital Basel
38 Spitalstr. 21, 4031 Basel
39 Switzerland

40
41

Abstract

Background

Literature suggests that prostate MRI exhibits better predictive capabilities compared to prostate-specific antigen (PSA) in detecting prostate cancer (PCa). Based on this, our present study investigates biparametric MRI (bpMRI) as a PSA-independent screening tool for PCa.

Objective

The primary endpoint was to assess the efforts and effectiveness of identifying 20 participants with clinically significant prostate cancer (csPCa) using bpMRI.

Design, Setting, and Participants

Biopsy-naïve men aged over 45 years were included. All participants underwent 3 Tesla bpMRI, PSA and digital rectal examination (DRE). Targeted-only biopsy was performed in participants with a suspicious lesion (PI-RADS≥3). Men with a negative bpMRI but suspicious DRE or elevated PSA had template biopsies. Pre-intended protocol adjustments were made post-interim analysis for PI-RADS 3 lesions: follow-up mpMRI after 6 months, biopsy only if lesions persisted or upgraded.

Outcome Measurements and Statistical Analysis

Biopsy results underwent comparison using Fisher's exact test and univariable logistic

regression to pinpoint prognostic factors for positive biopsy.

Results and Limitations

A total of 229 participants were enrolled in this analysis. Among these, 77 displayed suspicious

PI-RADS lesions. A total of 79 participants underwent a biopsy. PCa was detected in 29

participants, of whom 21 had csPCa.

BpMRI detected all 21 csPCa, whereas PSA and DRE missed 66.7%. Protocol adjustment led

to a 54.6 % biopsy reduction in PI-RADS 3 lesions. Overall, 10.9 bpMRIs were needed to

identify one participant with csPCa. A major limitation of the study is the lack of a control

cohort undergoing template biopsies.

Conclusions

Opportunistic screening utilising bpMRI as primary tool reveals participants with csPCa that

traditional methods might overlook, even at low PSA levels.

Patient summary

Screening with bpMRI and targeted biopsy identified csPCa in every 11[th] man, regardless of

PSA levels. Preselecting patients based on PSA > 1 ng/ml and positive family history of PCa

as well as other potential blood tests may further improve the effectiveness of bpMRI in this

setting.

Key Words: MRI, PI-RADS, Prostate biopsy, Prostate cancer, PSA, Screening

Trial registry: clinicaltrials.gov

Registration ID: NCT03749993

Introduction

Prostate cancer (PCa) is the most common cancer and the second cause of cancer death in the

male population in Switzerland. The decline in PCa-related deaths in recent years has been

attributed to opportunistic PCa screening based on prostate-specific antigen (PSA) [1].

Without there being any recommended cancer screening at present, the Council of the European

Union has recently proposed to consider the feasibility and effectiveness of implementing

organised prostate cancer screening with appropriate management on the basis of PSA and

magnetic resonance imaging (MRI)[2].

Large screening programs, such as the European Randomized Study of Screening for Prostate

Cancer (ERSPC) and the US Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial

(PLCO), have shown a reduction of up to 30% in PCa-specific mortality [3]. The ERSPC found

that 570 men had to be screened, and 18 patients had to be diagnosed to prevent one PCa-related

death [4].

The ERSPC and PLCO trials have established a PSA range of 3-4 ng/ml as an indication for

prostate biopsy, although using PSA as a single screening parameter has been found to be

suboptimal. This is supported by reports demonstrating an area under the curve of $0.52 - 0.631$

using a receiver operating characteristic (ROC) curve for PSA as a screening tool for PCa [5],

[6], [7]. Furthermore, the STHLM3 model ($\geq 10\%$ risk) has detected PCa ISUP ≥ 2 in 9% and

23% of men with PSA ranges of 1-2 ng/ml and 2-3 ng/ml, respectively [8]

The literature suggests that prostate MRI exhibits superior performance compared to PSA for

PCa detection with areas under the curve of 0.52-0.631 using ROC. The use of multiparametric

MRI (mpMRI) or biparametric MRI (bpMRI) reduces negative biopsies and leads to the

detection of significantly more csPCa [9], [10]. Moreover, the negative predictive value (NPV)

for PCa detection with MRI ranges between 87.9% and 97.5% [11]–[15]. Despite higher initial

costs compared to PSA measurement alone, screening with MRI has therefore the potential to

improve PCa screening.

In this study, prospectively investigating the performance of a purely bpMRI-based

opportunistic PCa screening programme, our primary outcome was the measurement of its

effectiveness, represented in defined parameters. Moreover, we aimed to identify necessary

adaptations to improve it.

1. Participants and methods

1.1 Ethics

The study has been conducted in compliance with the Swiss Association of Research Ethics Committees (EKNZ Nr. 2018-01965) and has been registered at clinicaltrials.gov under the number NCT03749993.

1.2 Study design

This study was designed as a single-centre, prospective cohort study, recruiting participants from between 01/2019 and 05/2023. The study population consisted of men over 50 years of age (over 45 years for individuals of African ancestry or with a family history of prostate cancer) who had not undergone a previous prostate biopsy and had an estimated life expectancy of more than 10 years. Participants with an acute urinary tract infection (defined as voiding symptoms and pathologic urinalysis upon clinical visit), with an NIH-Chronic prostatitis symptom index (NIH-CPSI) score of $\geq$ 19 or an international prostate symptom score (IPSS) $\geq$ 20 were treated and followed up urologically but were excluded from study participation. In addition, participants with contraindications to MRI were also excluded. All participants underwent biparametric MRI (bpMRI) of the prostate using a single 3T scanner (MAGNETOM Prisma, Siemens Healthineers, Erlangen, Germany) at the Department of Radiology of the University Hospital Basel. The bpMRI protocol used is detailed in Supplementary Table 1. T2-weighted turbo-spin echo (TSE) and diffusion-weighted imaging (DWI) series were used for lesion assessment, with reporting radiologists adhering strictly to the PI-RADS (Prostate Imaging-Reporting and Data System) v2.1 guidelines to ensure equal reporting standards for each report [16].

1.3 Interim analysis

An interim analysis to evaluate the performance of the study protocol was planned after a first

screening round (Phase I) with the intention of being able to make amendments to improve the

protocol. Interim analysis was performed after the detection of clinically significant prostate

cancer in five participants. The effectiveness of the study protocol was assessed by measuring

the detecting rates of PCa for bpMRI in comparison to PSA and DRE.

1.4 Indications for biopsy and protocol amendments

In Phase I, PI-RADS 3 lesions or higher were considered suspicious and were followed by

targeted-only biopsy upon detection. MRI reports classified as PI-RADS 1 and 2 were

considered clinically irrelevant and were not biopsied unless the participant had an abnormal

DRE and/or an elevated PSA $\geq$ 10.0 ng/ml. In that case, a template biopsy was performed.

As a result of the interim analysis, protocol amendments for Phase II included a re-assessment

of initial PI-RADS 3 lesions in bpMRI through a follow-up mpMRI (multiparametric MRI)

after 6 months. In case of persistent or upgraded of lesions, targeted prostate biopsy was

performed. Moreover, in addition to suspicious DRE and or (PSA) exceeding 10 ng/ml, PSA

density $\geq$ 0.15 ng/ml^2 was implemented as an indication for template biopsy in the absence of

a PI-RADS lesion $\geq$ 3 (see Fig. 1). The protocol amendments resulting from our interim analysis

(Phase I) were reviewed and approved by the Swiss Association of Research Ethics

Committees.

1.5 Biopsy procedure

At the commencement of our study, transrectal MRI/US-fusion biopsies were performed under local anaesthesia in 36 participants. A commercially available real-time virtual sonography (RVS) system (Hitachi Medical Corporation, Tokyo, Japan) in combination with magnetic position sensors (3D Guidance Trakstar, Ascension) was used for targeted biopsies.

Subsequently, we transitioned to a transperineal approach using a robotic-assisted MRI/US-fusion biopsy system (Monalisa, Biobot Surgical Ltd, Singapore) under general anaesthesia in 39 participants. For both approaches, at least 2 cores per targeted lesion were obtained.

When indicated, a template biopsy with 12 cores equally distributed over the prostate was performed.

1.6 Histopathologic reporting

Gleason score and International Society of Urological Pathology Grade (ISUP) were reported

for each cancer-positive biopsy. Gleason scores 3+3=6 and ISUP 1 were considered clinically

non-significant PCa (cnsPCa). A Gleason score of 3+4=7 or above and ISUP 2 or above would

be clinically significant PCa (csPCa).

1.7 Analysis and statistical methods

Categorical data were summarised using frequency and percentage, and continuous data using

median and range. Biopsy results were compared between subgroups using Fisher's exact test.

Univariable logistic regression was applied to identify possible prognostic factors for positive

biopsy. The significance level of 5% was used for all statistical tests with no adjustment for

multiple testing, due to the exploratory nature of these analyses. All analyses were performed

using SAS 9.4 (SAS Institute, Cary, NC, USA) and R 4.2.1 (The R Foundation; www.r-

project.org).

2. Results

2.1 Participant enrolment and baseline characteristics

From February 2019 to September 2022, 269 participants were enrolled in our study. Finally,

229 participants were ultimately included in the analysis (Figure 2). Of the 79 participants who

underwent a biopsy, all attended follow-up.

Figure 2: CONSORT/TREND flow diagram of the VISIONING study, follow-up and analysis (according to TREND statement)

(MRI = magnetic resonance imaging; PSA = prostate-specific antigen; DRE = digital rectal examination; UTI = urinary tract infection)

The study participants had a median age of 58 years [IQR, 53–64], median PSA level of 1.26

ng/ml [IQR, 0.72–2.84] and median PSA-density of 0.05 ng/ml^2 [IQR, 0.03–0.08]. A positive

family history for PCa was found in 38 participants while 106 participants had neither a PSA >

1 ng/ml nor a positive family history for PCa (Table 1).

Baseline characteristics (n=229)	median [IQR]
Age (years)	58 [53–64]
Prostate volume (ml)	28 [23–39]
PSA (ng/ml)	1.26 [0.72–2.84]
PSA-D (ng/ml^2)	0.05 [0.03–0.08]
Clinical parameters (n=229)	n (%)
PSA < 1 ng/ml & negative family history for PCa	106 (46.2)
Elevated PSA (≥10.0ng/ml)	6 (2.6)
Abnormal DRE	8 (3.5)
Elevated PSA-D (≥0.15ng/ml^2)	18 (7.9)
Positive family history for PCa	38 (16.6)

Table 1: Baseline characteristics and Clinical parameters of all included participants (PSA = prostate-specific

antigen; PSA-D = PSA-density; PCa = prostate cancer).

2.2 Interim analysis (Phase I)

In the first round of opportunistic screening (n=120), we excluded 12 participants leading to

108 participants included in the analysis. Forty-five participants underwent prostate biopsy of

whom 17.8 % (8/45) had a PCa ISUP ≥ 2 (Supplement figure 1). The pre-defined threshold of

detecting n=5 PCa ≥ ISUP 2 in Phase I has been exceeded to n=8 because of the lead time

between bpMRI, biopsy and histopathological report while the opportunistic screening had

continued. Half of the ISUP ≥ 2 detected in Phase I would have been missed by PSA & DRE

alone.

2.3 Overall results (Phase I & II)

Indication for biopsy was given in 89 of the analysed participants. Seventy-nine participants

received prostate biopsy after refusal of biopsy by ten participants.

Cancer was detected in 29 of the 229 (12.7%) recruited men, of whom eight (3.5%) were ISUP

1 and 21 (9.2%) were ISUP ≥ 2. Median PSA level and PSA-density were 2.02 ng/ml [IQR,

1.5–3.21] and 0.067 ng/ml^2 [IQR, 0.049–0.104] for the participants with ISUP 1 cancer and

3.88 ng/ml [IQR, 2.77–7.53] and 0.111 ng/ml^2 [IQR, 0.075–0.294] for the participants with

ISUP 2 or higher, respectively (Table 2).

	n (%)	median PSA (ng/ml) [IQR] {Range}*	median PSA-D (ng/ml^2) [IQR]
Index lesion at participant level (n=229)**			
PI-RADS 1 or 2	152 (66.4)	1.09 [0.67-2.2]	0.042 [0.027-0.065]
PI-RADS 3***	13 (5.7)	1.84 [1.15-3.96]	0.056 [0.033-0.083]
PI-RADS 4	55 (24.0)	2.02 [0.89-3.59]	0.056 [0.031-0.099]
PI-RADS 5	9 (3.9)	5.67 [2.18-20.0]	0.151 [0.054-0.612]
Biopsy intervention (n=229)			
Targeted Biopsy	77 (33.6)	2.06 [0.91-3.88]	0.057 [0.031-0.111]
Template biopsy	2 (0.9)	9.4 [9.05-9.75]	0.234 [0.196-0.272]
No biopsy	150 (65.5)	1.08 [0.67-2.17]	0.042 [0.026-0.064]
Biopsy result (n=79)			
ISUP 1	8 (10.1)	2.02 [1.5–3.21] {0.81-4.69}	0.067 [0.049–0.104]
ISUP ≥ 2	21 (26.6)	3.88 [2.77–7.53] {0.8-39.7}	0.111 [0.075–0.294]
No cancer	50 (63.3)	1.44 [0.79–3.6] {0.21-12.7}	0.051 [0.026–0.087]

*Table 2: Index lesions, performed biopsy and biopsy results. *The range was specifically added in this section to*
*display the lowest PSA values in men with prostate cancer. **In case a participant had more than one suspicious*
*PI-RADS lesion, the highest-graded lesion would count as index lesion. ***The biopsied PI-RADS 3 lesions were*
from Phase I (before the interim analysis) or persistent PI-RADS 3 lesions in follow-up MRI from Phase II
(PSA = prostate-specific antigen; DRE = digital rectal examination; PSA-D = PSA-density; PI-RADS = prostate
imaging-reporting and data system, ISUP = International Society of Urological Pathology).

All 29 PCas were detected by bpMRI screening, while 22 would have been missed by PSA and

DRE. Of these 22 cancers that classical screening would have missed, 14 (63.6%) were ISUP

≥ 2 (Figure 4).

243
244
Figure 4: Histopathologic results (absolute numbers) overall, indicated by bpMRI, PSA-D and PSA &/or DRE.

(ISUP = International Society of Urological Pathology; bpMRI = biparametric MRI; PSA = prostate-specific antigen;

DRE = digital rectal examination; PSA-D = PSA-density)

As a result of the protocol amendments following the interim analysis, we could reduce the

number of negative biopsies in Phase I from 73.3% (33/45) to 50% (17/34) in Phase II (Table

5). More specifically, the rate of negative biopsies indicated by MRI was reduced in Phase II

from initially 72.7% (32/44) to 48.5% (16/33) by not performing an immediate biopsy in men

with a PIRADS 3 score on bpMRI. The number of opportunistic screening MRIs to detect one

ISUP ≥ 2 decreased from 13.6 to 9.3 while increasing the biopsy detection rate for ISUP ≥ 2

from 17.8% (8/45) to 38.2% (13/34).

We calculated detection rates for the scenario of participants with a PSA below 1 ng/ml and

negative family history for PCa not being subjected to further assessment with bpMRI.

Notably, by doing so, the detection of any PCa ISUP ≥ 2 would not have been compromised.

We found that the number of MRIs required to detect one ISUP ≥ 2 could have decreased from

10.9 (229/21) to 5.9 (123/21). Consequently, for the entire cohort, 2.9 (61/21) targeted biopsies

would have been necessary to detect one ISUP ≥ 2 (Table 4).

	Phase I (n=108)	Phase II (n=121)	Overall (n=229)	
	Any PSA	Any PSA	Any PSA	PSA ≥ 1*
Biopsy performed	45	34	79	61
csPCa	8	13	21	21
cnsPCa	4	4	8	6
Neg. biopsy	33	17	50	34
MRI	108	121	229	123
MRI / ISUP ≥ 2	**13.6**	**9.3**	**10.9**	**5.9**
MRI / PCa	9.1	7.1	7.9	4.6
Biopsy / ISUP ≥ 2	**5.6**	**2.6**	**3.8**	**2.9**
Biopsy / PCa	3.8	2	2.7	2.3

265   *Table 4: Absolute numbers (n) of biopsies performed, biopsy results, MRIs carried out and number of MRIs needed*

266   *to detect one ISUP ≥ 2 (MRI/ISUP ≥) in our cohorts of Phase I, Phase II and overall.*

267   **and/or positive family history of prostate cancer*

3. Discussion

In this study, we present the outcomes of a PSA-independent, bpMRI-based, opportunistic screening programme for PCa, comparing its effectiveness in contrast to classical screening methods.

The study encompassed 229 participants in the final analysis, and detected 21 men with ISUP $\geq$ 2 in the biopsy group of 79 individuals, resulting in a 26.7% detection rate for ISUP $\geq$ 2.

The PSA-independent screening strategy using bpMRI detected all 21 men with ISUP $\geq$ 2, whereas classical screening would have missed 66.6% (14/21) of ISUP $\geq$ 2.

While adaptations were made to our study protocol after the interim analysis, the negative biopsy rate indicated by bpMRI could be reduced by 24.2% (from 72.7% down to 48.5%). However, an even more restrained strategy when indicating biopsy for PI-RADS 3 lesions does not appear to be particularly effective, as the detection rates of ISUP $\geq$ 2 in Phase II for PI-RADS 3 and PI-RADS 4 lesions were 25% (1/4) and 36% (9/25), respectively. As a computed result, 11 (229/21) participants had to partake in our opportunistic screening to detect one ISUP $\geq$ 2. The performance of opportunistic screening by bpMRI could be further improved by omitting bpMRI screening in patients with a PSA < 1 ng/ml and a negative family history for PCa. This would have led to a reduction of MRIs by 46.3% (106/229) and to a reduction of the negative biopsy rate by 7.6% (down to 55.7%) and a reduction of the diagnosis of ISUP 1 by 32% (34/50) in the whole cohort. Analysing Phase II of the study under the same assumption, only 5.9 bpMRIs and 2.9 biopsies would have been needed to detect one ISUP $\geq$ 2. Of note, the median PSA-density of ISUP $\geq$ 2 in our study was 0.111 ng/ml2, and as such, below the guidelines-suggested yet debated cut-off of 0.15 ng/ml^2 [17].

By comparison, in the adjusted STHLM3 cohort (adjusted for age, PSA, PSA-naivety, family history and previous negative biopsy using propensity scores to match the Current Prostate Cancer Testing cohort), 53.7 participants had to be screened to detect one ISUP $\geq$ 2 with a biopsy rate without cancer detection of 62% [19].

While the performance of bpMRI in our setting seems very promising, the price and availability

of a 3 Tesla MRI vary strongly between countries. Finally, each individual health system will

have to weigh the availability and costs of MRI against more easily available and probably less

expensive blood tests.

Moreover, we emphasize the financial and temporal benefits associated with bpMRI in

comparison to multiparametric magnetic resonance imaging (mpMRI) when used for screening

purposes with a reduced MRI scan time and omission of gadolinium-containing contrast agents.

Related calculations conducted by Porter K. et al. have demonstrated that bpMRI outperforms

mpMRI in terms of time and cost considerations, while still ensuring high standards of

diagnostic precision [20]. Additionally, other studies have reported similar diagnostic accuracy

between the two techniques [14], [21].

Regarding a pontential selection bias, as we conducted an opportunistic screening and not a

screening-by-invitation programme, men with a positive family history of PCa might have been

more likely to participate in this study. However, the number of participants with a positive

family history of PCa was limited to 16.6%. Another evident limitation of our study protocol is

the absence of a control cohort. Considering the high NPV of MRI for the detection of ISUP $\geq$

2, we weighed the biopsy-related morbidity in a control cohort against the informational benefit.

Therefore, we decided against the template biopsy control group. Moreover, the non-blinding

of the PSA value for participants might have led to the decision to not undergo biopsy in some

men with very low PSA values and PI-RADS 3 and 4 lesions.

In a recent study conducted by Moore C.M. et al. [22], a protocol similar to ours was utilized,

but in a different setting. While we conducted our research under an opportunistic screening

framework, they employed a screening-by-invitation approach. Remarkably, despite these

differing contexts, our findings converged. This consistency between both studies underscores

the robustness and potential generalizability of our results, irrespective of the screening method

employed. Their conclusions, parallel to our own, highlight the significance of MRI in

screening for prostate cancer, further strengthening the validity and relevance of our research.

In order to answer the question whether the good performance of bpMRI also translates into

better survival of men with csPCa, larger and maybe population-based screenings are required.

While csPCa so far equals ISUP $\geq$ 2, new definitions for csPCa might be required as ISUP

grade group II PCa patients with a low PSA may still profit from active surveillance instead of

undergoing definitive radiotherapy or surgery.

328 **4. Conclusions**

This pilot study applying non-PSA triggered biparametric MRI as a sole opportunistic screening

tool for prostate cancer demonstrates superior and promising effectiveness for bpMRI followed

by a targeted biopsy to detect ISUP ≥ 2 compared to classical screening methods. Encompassing

229 participants overall, approximately 11 and 4 men had to undergo bpMRIs and targeted

biopsies, respectively in order to detect one ISUP ≥ 2. The screening protocol may profit from

adjustments leading to a further reduction of the diagnosis of ISUP 1 cancers and negative

biopsies. In a first step, omitting screening in men with a PSA below 1 ng/ml and negative

family history of prostate cancer would reduce the number of bpMRIs and biopsies needed to

detect one ISUP ≥ 2 to approximately 6 and 3, respectively.

5. Take home message

340 BpMRI of the prostate is a promising and highly sensitive tool for PCa screening and improves

341 the detection of ISUP ≥ 2 by 200% as compared to the current screening tools PSA and DRE.

Author contributions:

Data access and responsibility: Marc Matthias had full access to all the data in the study and took responsibility for the integrity of the data and the accuracy of the data analysis.
Study concept and design: Rentsch, Wetterauer, Matthias
Acquisition of data: Matthias
Analysis and interpretation of data: Matthias, Rentsch Wetterauer, Hayoz
Drafting of the manuscript: Matthias, Wetterauer, Rentsch
Critical revision of the manuscript for important intellectual content: Matthias, Wetterauer, Rentsch, Seifert, Pueschel, Bubendorf, Winkel, Boll, Merkle, Heye, Hayoz, Deckart, Arbealez, Mortezavi
Statistical analysis: Matthias, Hayoz
Obtaining funding: Rentsch, Seifert, Boll, Merkle
Administrative, technical or material support: Pueschel
Supervision: Rentsch, Wetterauer, Seifert
Other: none

Financial disclosures:
Prof. Cyrill Rentsch MD-PhD certifies that all conflicts of interest, including specific financial interests and relationships and affiliations relevant to the subject matter or materials discussed in the manuscript (e.g. employment/affiliation, grants or funding, consultancies, honoraria, stock ownership or options, expert testimony, royalties, or patents filed, received, or pending) are the following: none.

Funding/Support and role of the sponsor:
This study is entirely funded by the Department of Urology and Radiology at the University Hospital of Basel, Switzerland. No other sponsors funded this trial.

Previous presentation or release information:
The results of the interim analysis of this study were presented in the form of an unmoderated poster at the annual conference of the swiss association for Urology (SGU) held from the 29th of September to the 1st of October 2021 in Interlaken, Switzerland, as well as in the form of an abstract at the EAU 2021. Furthermore, the presentation was held during the virtual EAU 2021 conference by Marc Matthias on the 11th of July 2021.
The current results of the primary endpoint analysis were presented as an abstract and a live presentation held by Marc Matthias at the EAU 2023 in Milano, Italy as well as at the AUA 2023 in Chicago, USA.

6. References

[1] Bundesamt für Statistik, "Prostatakrebs: zeitliche Entwicklung 2020." Accessed: Jun. 02, 2021. [Online]. Available: https://www.bfs.admin.ch/bfs/de/home/statistiken/gesundheit/gesundheitszustand/kran kheiten/krebs/spezifische.assetdetail.14816169.html.

[2] Council of the European Union, "Council Recommendation on strengthening prevention through early detection: A new EU approach on cancer screening replacing Council Recommendation 2003/878/EC." Accessed: Jul. 30, 2023. [Online]. Available: https://data.consilium.europa.eu/doc/document/ST-14770-2022-INIT/en/pdf

[3] A. Tsodikov *et al.*, "Reconciling the effects of screening on prostate cancer mortality in the ERSPC and PLCO trials," *Ann Intern Med*, vol. 167, no. 7, pp. 449–455, Oct. 2017, doi: 10.7326/M16-2586.

[4] J. Hugosson *et al.*, "A 16-yr Follow-up of the European Randomized study of Screening for Prostate Cancer(Figure presented.)," *Eur Urol*, vol. 76, no. 1, pp. 43–51, Jul. 2019, doi: 10.1016/j.eururo.2019.02.009.

[5] W. J. Catalona *et al.*, "A multicenter study of [-2]pro-prostate specific antigen combined with prostate specific antigen and free prostate specific antigen for prostate cancer detection in the 2.0 to 10.0 ng/ml prostate specific antigen range," *Journal of Urology*, vol. 185, no. 5, pp. 1650–1655, May 2011, doi: 10.1016/j.juro.2010.12.032.

[6] F. A. Distler *et al.*, "The Value of PSA Density in Combination with PI-RADS^{TM} for the Accuracy of Prostate Cancer Prediction," *Journal of Urology*, vol. 198, no. 3, pp. 575–582, Sep. 2017, doi: 10.1016/j.juro.2017.03.130.

[7] J. J. Tosoian *et al.*, "Prostate Health Index density improves detection of clinically significant prostate cancer," *BJU Int*, vol. 120, no. 6, pp. 793–798, Dec. 2017, doi: 10.1111/bju.13762.

[8] H. Grönberg *et al.*, "Prostate cancer screening in men aged 50-69 years (STHLM3): A prospective population-based diagnostic study," *Lancet Oncol*, vol. 16, no. 16, pp. 1667–1676, 2015, doi: 10.1016/S1470-2045(15)00361-7.

[9] F. Giganti, C. Allen, M. Emberton, C. M. Moore, and V. Kasivisvanathan, "Prostate Imaging Quality (PI-QUAL): A New Quality Control Scoring System for Multiparametric Magnetic Resonance Imaging of the Prostate from the PRECISION trial," *Eur Urol Oncol*, vol. 3, no. 5, pp. 615–619, Oct. 2020, doi: 10.1016/j.euo.2020.06.007.

[10] L. Boesen *et al.*, "Assessment of the Diagnostic Accuracy of Biparametric Magnetic Resonance Imaging for Prostate Cancer in Biopsy-Naive Men: The Biparametric MRI for Detection of Prostate Cancer (BIDOC) Study," *JAMA Netw Open*, vol. 1, no. 2, Jun. 2018, doi: 10.1001/jamanetworkopen.2018.0219.

[11] P. C. Moldovan *et al.*, "What Is the Negative Predictive Value of Multiparametric Magnetic Resonance Imaging in Excluding Prostate Cancer at Biopsy? A Systematic Review and Meta-analysis from the European Association of Urology Prostate Cancer Guidelines Panel," *Eur Urol*, vol. 72, no. 2, pp. 250–266, Aug. 2017, doi: 10.1016/j.eururo.2017.02.026.

[12] H. U. Ahmed *et al.*, "Diagnostic accuracy of multi-parametric MRI and TRUS biopsy in prostate cancer (PROMIS): a paired validating confirmatory study," *The Lancet*, vol. 389, no. 10071, pp. 815–822, Feb. 2017, doi: 10.1016/S0140-6736(16)32401-1.

[13] O. Rouvière *et al.*, "Use of prostate systematic and targeted biopsy on the basis of multiparametric MRI in biopsy-naive patients (MRI-FIRST): a prospective, multicentre, paired diagnostic study," *Lancet Oncol*, vol. 20, no. 1, pp. 100–109, Jan. 2019, doi: 10.1016/S1470-2045(18)30569-2.

[14] C. K. Kuhl, R. Bruhn, N. Krämer, S. Nebelung, A. Heidenreich, and S. Schrading, "Abbreviated Biparametric Prostate MR Imaging in Men with Elevated Prostate-specific Antigen," *Radiology*, vol. 285, no. 2, pp. 493–505, Nov. 2017, doi: 10.1148/radiol.2017170129.

[15] V. Kasivisvanathan *et al.*, "MRI-Targeted or Standard Biopsy for Prostate-Cancer Diagnosis," *New England Journal of Medicine*, Mar. 2018, doi: 10.1056/nejmoa1801993.

[16] B. Turkbey *et al.*, "Prostate Imaging Reporting and Data System Version 2.1: 2019 Update of Prostate Imaging Reporting and Data System Version 2," *European Urology*, vol. 76, no. 3. Elsevier B.V., pp. 340–351, Sep. 01, 2019. doi: 10.1016/j.eururo.2019.02.033.

[17] Y. S. Ha *et al.*, "Prostate-specific antigen density toward a better cutoff to identify better candidates for active surveillance," *Urology*, vol. 84, no. 2, pp. 365–372, 2014, doi: 10.1016/j.urology.2014.02.038.

[18] T. Nordström *et al.*, "Prostate cancer screening using a combination of risk-prediction, MRI, and targeted prostate biopsies (STHLM3-MRI): a prospective, population-based, randomised, open-label, non-inferiority trial," *Lancet Oncol*, vol. 22, no. 9, pp. 1240–1249, Sep. 2021, doi: 10.1016/S1470-2045(21)00348-X.

[19] M. Eklund *et al.*, "The Stockholm-3 (STHLM3) Model can Improve Prostate Cancer Diagnostics in Men Aged 50–69 yr Compared with Current Prostate Cancer Testing," *Eur Urol Focus*, vol. 4, no. 5, pp. 707–710, Sep. 2018, doi: 10.1016/j.euf.2016.10.009.

[20] K. K. Porter, A. King, S. J. Galgano, R. L. Sherrer, J. B. Gordetsky, and S. Rais-Bahrami, "Financial implications of biparametric prostate MRI," *Prostate Cancer Prostatic Dis*, vol. 23, no. 1, pp. 88–93, Mar. 2020, doi: 10.1038/s41391-019-0158-x.

[21] L. Xu *et al.*, "Comparison of biparametric and multiparametric MRI in the diagnosis of prostate cancer," *Cancer Imaging*, vol. 19, no. 1, Dec. 2019, doi: 10.1186/s40644-019-0274-9.

[22] C. M. Moore *et al.*, "Prevalence of MRI lesions in men responding to a GP-led invitation for a prostate health check: a prospective cohort study," *BMJ Oncology*, vol. 2, no. 1, p. e000057, Aug. 2023, doi: 10.1136/bmjonc-2023-000057.