

# Determination of glomerular filtration rate, a spin off after contrast-enhanced computed tomography among critically ill patients – proof of concept.

Bertil Kågedal<sup>1\*</sup>, Anders Helldén<sup>2</sup>, Dženeta Nezirević Dernroth<sup>1</sup>, Anders Andersen<sup>3</sup>, Andreas Ekman<sup>4</sup>, Mats Haglund<sup>5</sup>, Bharti Kataria<sup>6</sup>, Frida Oskarsson<sup>3</sup>, Lovisa Tobieson<sup>7</sup>, Åse Östholm Balkhed<sup>8</sup>, and Håkan Hanberger<sup>8</sup>

<sup>1</sup>Department of Clinical Chemistry and Clinical Pharmacology and Department of Biomedical and Clinical Sciences, Linköping University, Linköping, Sweden. E-mail: [bertil.kagedal@liu.se](mailto:bertil.kagedal@liu.se); [dzeneta.dernroth@liu.se](mailto:dzeneta.dernroth@liu.se)

<sup>2</sup>Department of Laboratory Medicine, Division of Clinical Pharmacology, Karolinska Institutet, Karolinska University Hospital, Huddinge, Stockholm, Sweden. E-mail: [anders.hellden@regionstockholm.se](mailto:anders.hellden@regionstockholm.se)

<sup>3</sup>Department of Anesthesia and Intensive Care, Växjö County Hospital, Sweden. E-mail: [anders.andersen@kronoberg.se](mailto:anders.andersen@kronoberg.se); [frida.oskarsson@kronoberg.se](mailto:frida.oskarsson@kronoberg.se)

<sup>4</sup>Department of Anesthesia and Intensive Care, Kalmar County Hospital, Sweden. E-mail: [andreas.ekman@regionkalmar.se](mailto:andreas.ekman@regionkalmar.se)

<sup>5</sup>Department of Infectious Diseases, Kalmar County Hospital, Sweden. E-mail: [mats.haglund@regionkalmar.se](mailto:mats.haglund@regionkalmar.se)

<sup>6</sup>Centre of Medical Image Science and Visualization (CMIV), Department of radiology and Department of Health, Medicine and Caring Sciences, Linköping University, Linköping, Sweden. E-mail: [bharti.kataria@liu.se](mailto:bharti.kataria@liu.se)

<sup>7</sup>Department of Neurosurgery and Department of Biomedical and Clinical Sciences, Linköping University, Linköping, Sweden. E-mail: [lovisa.tobieson@regionostergotland.se](mailto:lovisa.tobieson@regionostergotland.se)

<sup>8</sup>Department of Infectious Diseases, and Department of Biomedical and Clinical Sciences, Linköping University, Linköping, Sweden. E-mail: [ase.ostholm.balkhed@regionostergotland.se](mailto:ase.ostholm.balkhed@regionostergotland.se); [hakan.hanberger@liu.se](mailto:hakan.hanberger@liu.se)

<sup>1,2</sup>Anders Helldén and Bertil Kågedal contributed equally to the manuscript

## Running headline

Iohexol plasma clearance determined from radiology dosing

## Abstract

**Background.** Recently, Gong *et al.* (Gong *et al.* 2022) showed, in nine healthy subjects, that plasma clearance of high doses of iohexol given as contrast enhanced computed tomography (CT) could be used for determination of glomerular filtration rate (GFR). We utilized high doses of iohexol from angiographic or other contrast enhanced CT given to critical ill patients for calculation of  $GFR_{\text{iohexol}}$  and compared these data with standard low dose iohexol GFR determinations.

**Method.** Patients at intensive care units (ICUs) in Southeast Sweden intended for radiographic investigations that included injection of 45-120 ml of iohexol (Omnipaque) were included, and the concentration of iohexol in plasma was measured by HPLC. Iohexol clearance was calculated by the

method of Bröchner-Mortensen. The following days was iohexol clearance determined using the standard low dose of 5 mL of iohexol. Sixteen patients admitted to ICUs were included in this pilot study.

**Results.** GFR after high dosing of iohexol at contrast enhanced CT could be measured for all sixteen critically ill patients. Patients with normal or increased renal function had neglectable iohexol concentrations the day following the CT scan. There was excellent correlation between GFR determination with high and standard low iohexol dosing among these 6 patients. Ten patients had decreased renal function and delayed elimination of iohexol, thus was not GFR measurement with low dose iohexol possible to analyse the day after CT scan with high dose iohexol.

**Conclusion.** This pilot study showed that GFR can be measured after high doses of iohexol at enhanced CT and compare well with the standard low dose of iohexol clearance determinations.

## Introduction

When dosing of renally excreted drugs, the patient's glomerular filtration rate (GFR) must be considered. Current estimated GFR (eGFR) methods, which are based on plasma concentrations of creatinine and cystatin C, are too uncertain to be relied on in patients needing intensive care (Delanaye, Melsom, et al. 2016).

Critical ill patients are frequently investigated by iohexol contrast enhanced computed tomography (CT). Iohexol plasma clearance has for years been the reference method for determination of GFR, but the iohexol dose used is only 5-10% of that used for contrast enhanced CT.

The reference method, low dose iohexol clearance ( $GFR_{\text{iohexol}}$ ), is seldom used as support for drug dosing in patients admitted to ICUs. The method is considered complicated, takes time to implement, requires accurate information on amount of injected iohexol, and exact time schedule for blood sampling (Delanaye, Ebert, et al. 2016; Delanaye, Melsom, et al. 2016; Levey and Inker 2016). According to guidelines, the earliest an examination with standard low dose dosing of iohexol can be carried out is 3-4 days after contrast X-ray, which prevents early knowledge from a more accurate GFR than that from plasma creatinine or cystatin C. However, we hypothesized that clearance of the iohexol given at the radiology department can be used for more rapid information of  $GFR_{\text{iohexol}}$ . A prerequisite is that the pharmacokinetics of iohexol is not dose-dependent. Interestingly, this has been shown in earlier studies (Bäck, Krutzen, and Nilsson-Ehle 1988; Hu et al. 2013).

Recently, Gong *et al.* (Gong et al. 2022) showed that ten times higher dosages of iohexol gave the same clearance rates as the standard dosage of 5 mL iohexol in nine healthy subjects. The aim of this study was to investigate if the administration of iohexol in contrast enhanced CT, on clinical indication, in critically ill patients could be used for determination of GFR by plasma iohexol clearance determination.

## Material and Methods

We conducted a study on ICU patients in collaboration with the Radiology departments at three hospitals in southeast of Sweden. The iohexol preparation used at these departments were Omnipaque 350 mg iodine/mL (GE Healthcare). The radiology departments provided the information on iohexol dose and time for administration to the ICU where four blood samples were taken 2-4 hours after the injection for analysis of plasma iohexol concentration.

Standard clearance investigations were performed the following days with low dose iohexol of 5 mL Omnipaque 300 mg iodine/mL.

We analysed the concentration of iohexol in plasma by HPLC and calculated iohexol clearance by the method of Bröchner-Mortensen (Bröchner-Mortensen 1972). Patients with remaining high iohexol concentration in the pre-study sample (zero sample) day 2 and later would impact on the calculation of the standard low dose iohexol clearance studies and were excluded in the present report. This study

was approved by the Swedish Ethical Review Authority No 2020-04197. The patients gave informed consent to the study.


## Results

Iohexol clearance after CT in sixteen critically ill patients were median (range) 102 mL/min (28 – 152). The correlation between iohexol clearance determined after CT (high dosing day 1; 33975-90600 mg iohexol) and low dosing days 2 to day 8 (3232 mg iohexol) was excellent,  $y = 1.0255x - 0.8807$  mL/min,  $R^2 = 0.8687$  (N=6), where  $x$  = clearance with low dose iohexol and  $y$  = iohexol clearance after CT (Figure 1). Patients with normal or increased renal function had neglectable pre-injection iohexol concentrations (zero sample) the day after the CT scan, and further on the day after the 5 mL injections. For patients with low GFR was GFR measured the day after CT scan based on elimination of the high dose iohexol, but for these patients was repeated GFR measurement the following day with the standard method not feasible due to remaining high concentration of iohexol.

## Discussion

This pilot study shows that the results obtained after high dosing of iohexol at enhanced CT compare well with standard investigation with low dosing. Practically, the results obtained with the high iohexol dosing at radiology can be used for decision of dosing of drugs that are eliminated by glomerular filtration in ICU patients.

**Figure 1.** This figure shows the correlation between iohexol clearance measured with a low dose of 5 mL iohexol and iohexol clearance with high dose iohexol after computed tomography (CT) in six critically ill patients. Four patients had more than one low dose measurements. The figure contains data from patients with no concentrations of iohexol in the zero sample.



## References

- Bröchner-Mortensen, J. 1972. 'A simple method for the determination of glomerular filtration rate', *Scand J Clin Lab Invest*, 30: 271-4.
- Bäck, S. E., E. Krutzen, and P. Nilsson-Ehle. 1988. 'Contrast media and glomerular filtration: dose dependence of clearance for three agents', *J Pharm Sci*, 77: 765-7.
- Delanaye, P., N. Ebert, T. Melsom, F. Gaspari, C. Mariat, E. Cavalier, J. Bjork, A. Christensson, U. Nyman, E. Porrini, G. Remuzzi, P. Ruggenenti, E. Schaeffner, I. Soveri, G. Sterner, B. O. Eriksen, and S. E. Back. 2016. 'Iohexol plasma clearance for measuring glomerular filtration rate in clinical practice and research: a review. Part 1: How to measure glomerular filtration rate with iohexol?', *Clin Kidney J*, 9: 682-99.
- Delanaye, P., T. Melsom, N. Ebert, S. E. Back, C. Mariat, E. Cavalier, J. Bjork, A. Christensson, U. Nyman, E. Porrini, G. Remuzzi, P. Ruggenenti, E. Schaeffner, I. Soveri, G. Sterner, B. O. Eriksen, and F. Gaspari. 2016. 'Iohexol plasma clearance for measuring glomerular filtration rate in clinical practice and research: a review. Part 2: Why to measure glomerular filtration rate with iohexol?', *Clin Kidney J*, 9: 700-4.
- Gong, K., Y. Zou, H. Zhu, X. Peng, D. Qin, B. Sun, and X. Wang. 2022. 'Comparative study of plasma clearance of iohexol at different injection doses', *Clin Chim Acta*, 525: 34-39.
- Hu, S. L., M. Igari, N. L. Walle, M. W. Steffes, M. D. Beland, S. A. Collins, and R. Y. Gohh. 2013. 'Kidney transplant donor glomerular filtration rate by iohexol clearance during computerized tomographic angiography of the kidneys', *Transplant Proc*, 45: 3229-33.
- Levey, A. S., and L. A. Inker. 2016. 'GFR as the "Gold Standard": Estimated, Measured, and True', *American journal of kidney diseases : the official journal of the National Kidney Foundation*, 67: 9-12.