Adverse Long-Term Outcomes and an Immune Suppressed Endotype in Sepsis Patients with Reduced Interferon-γ ELISpot: A Multicenter, Prospective Observational Study

Evan A, Barrios[1], Monty B. Mazer[5], Patrick McGonagill[6] Christian B. Bergmann[3,*], Michael D. Goodman[3], Robert W. Gould[11], Mahil Rao[10], Valerie Polcz[1], Ruth Davis[1], Drew Del Toro[4], Marvin Dirain[1], Alexandra Dram[4], Lucas Hale[11], Mohammad Heidarian[7], Tamara A. Kucaba[12], Jennifer P. Lanz[1], Ashley McCray[1], Sandra Meszaros[4], Sydney Miles[4], Candace Nelson[11], Ivanna Rocha[1], Elvia E Silva[7], Ricardo Ungaro[1], Andrew Walton[4], Julie Xu[12], Leilani Zeumer-Spataro[1], Anne M. Drewry[4] Muxuan Liang[2,], Tyler Loftus[1], Isaiah Turnbull[4], Philip A. Efron[1], Kenneth E. Remy[5], Scott Brakenridge[15] , Vladimir P Badovinac[7,8,9], Thomas S. Griffith[12,13,14], Lyle L. Moldawer[1,16], Richard S. Hotchkiss[4], Charles C. Caldwell[3]

[1]Sepsis and Critical Illness Research Center, Department of Surgery, University of Florida College of Medicine, Gainesville, Florida 32610 USA

[2]Department of Biostatistics, University of Florida College of Medicine, Gainesville, Florida 32610 USA

[3]Department of Surgery, University of Cincinnati College of Medicine, Cincinnati, Ohio 45267, USA.

[4]Department of Anesthesiology, Washington University School of Medicine, St. Louis, Missouri 63110 USA

[5]Department of Pediatrics, Case Western Reserve University School of Medicine, Cleveland, Ohio 44106 USA

[6]Department of Surgery, University of Iowa Carver College of Medicine, Iowa City, Iowa 52242 USA.

[7]Department of Pathology, University of Iowa Carver College of Medicine, Iowa City, Iowa 52242 USA.

[8]Interdisciplinary Program in Immunology, University of Iowa Carver College of Medicine, Iowa City, Iowa 52242 USA.

[9]Experimental Pathology Ph.D. Program, University of Iowa Carver College of Medicine, Iowa City, Iowa 52242 USA.

[10]Department of Pediatrics, University of Iowa Carver College of Medicine, Iowa City, Iowa 52242 USA.

[11]Department of Anesthesiology, University of Minnesota Medical School, Minneapolis, Minnesota 55455 USA.

46
47 [12]Department of Urology, University of Minnesota Medical School, Minneapolis, Minnesota
48 55455, USA.
49
50 [13]Center for Immunology, University of Minnesota Medical School, Minneapolis, Minnesota
51 55455 USA.
52
53 [14]Minneapolis VA Healthcare System, Minneapolis, Minnesota 55417, USA
54
55 [15]Department of Surgery, Harborview Medical Center, University of Washington School of
56 Medicine, Seattle, Washington 98133 USA
57
58 *current address: Department of Anesthesiology and Critical Care Medicine, University of Ulm,*
59 *Ulm Germany.*
60
61 Running head: ELISpot Use in Critical Illness
62
63 Key words: critical illness, late mortality, procalcitonin, IL-6, prediction modeling.
64
65 Correspondence should be directed to:
66
67 [16]Corresponding Author Responsible for Content:
68
Lyle L. Moldawer, Ph.D.
Sepsis and Critical Illness Research Center
Department of Surgery
room 6116, Shands Hospital
University of Florida College of Medicine
Gainesville, Florida 32610-0019
moldawer@surgery.ufl.edu
352, 265-0494
77
78

Conflict of Interest Statements:

Evan A, Barrios, M.D. reports institutional support for salary from a National Institute of
General Medical Sciences training grant in burns, trauma and sepsis (T32 GM-008721) (PAE).

Monty B. Mazer, M.D. is a member of Immune Functional Diagnostics, LLC and receives no
direct financial compensation. Immune Functional Diagnostics, LLC is developing predictive
metrics in critical illness and this technology is evaluated in this research.

Patrick McGonagill M.D. reports no conflicts of interest.

Christian B. Bergmann, M.D. is supported by grants from the Deutsche
Forschungsgemeinschaft (German Research Foundation) (BE 7016/1-1).

Michael D. Goodman, M.D. reports institutional support for salary from the National Institute of

General Medical Sciences (R01 GM-124156) and the Department of Defense (G102983-
6263608307-1).

Robert W. Gould, M.D. reports no conflicts of interest.

Mahil Rao M.D., Ph.D. reports no conflicts of interest.

Valerie Polcz, M.D. reports institutional support for salary from a National Institute of General
Medical Sciences training grant in burns, trauma and sepsis (T32 GM-008721) (PAE).

Ruth Davis, B.S.N, R.N. reports no conflicts of interest.

Drew Del Toro, M.D. reports no conflicts of interest.

Marvin Dirain, M.S. reports no conflicts of interest.

Alexandra Dram reports no conflicts of interest.

Lucas Hale, B.A. reports no conflicts of interest.

Mohammad Heidarian, M.S. reports no conflicts of interest.

Tamara A. Kucaba, B.S. reports no conflicts of interest.

Jennifer P. Lanz, M.S.N., R.N. reports no conflicts of interest.

Ashley McCray, R.N. reports no conflicts of interest.

Sandra Meszaros reports no conflicts of interest.

Sydney Miles, B.S. reports no conflicts of interest.

Candace Nelson, B.A. reports no conflicts of interest.

Ivanna Rocha, M.P.H. reports no conflicts of interest.

Elvia E Silva, M.S. reports no conflicts of interest.

Ricardo Ungaro, B.S. reports no conflicts of interest.

Andrew Walton, M.S. reports no conflicts of interest.

Julie Xu, B.S. reports no conflicts of interest.

Leilani Zeumer-Spataro, B.S. reports no conflicts of interest.

Muxuan Liang, Ph.D. reports no conflicts of interest.

Isaiah Turnbull, M.D is a member of Immune Functional Diagnostics, LLC and receives no
direct financial compensation. Immune Functional Diagnostics, LLC is developing predictive
metrics in critical illness and this technology is evaluated in this research. He is also supported
by R35 GM-133756 from the National Institute of General Medical Sciences.

Tyler Loftus, M.D. is supported by R01 GM-149657 from the National Institute of General
Medical Sciences.

Philip A. Efron, M.D. is supported by grants R35 GM-140806, T32 GM-008721 and RM GM-
139690 awarded by the National Institute of General Medical Sciences.

Kenneth E. Remy, M.D., M.S. is a member of Immune Functional Diagnostics, LLC and
receives no direct financial compensation. Immune Functional Diagnostics, LLC is developing

predictive metrics in critical illness and this technology is evaluated in this research.

Scott Brakenridge, M.D., M.S.C.S. is funded by R35 GM-134880 from the National Institute of
General Medical Sciences. Dr. Brakenridge and the University of Florida may receive royalty
income based on a technology developed by Dr. Brakenridge and others and licensed by
Washington University in St. Louis to IFDx LLC. That technology is evaluated in this research.

Vladimir P. Badovinac Ph.D.is funded by R35 GM-134880 from the National Institute of
General Medical Sciences.

Thomas S. Griffith, Ph.D. is funded by R35 GM-140881 from the National Institute of General
Medical Sciences and the recipient of a Research Career Scientist award (IK6BX006192) from
the Department of Veteran Affairs.

Lyle L. Moldawer, Ph.D. is supported by NIH grants RM1 GM-139690, R01GM-132364 and
RF1 NS128626. Dr. Moldawer and the University of Florida may receive royalty income based
on a technology developed by Dr. Moldawer and others and licensed by Washington University
in St. Louis to IFDx LLC. That technology is evaluated in this research.

Richard S. Hotchkiss, M.D. Dr. Hotchkiss and Washington University in St. Louis may receive
royalty income based on a technology developed by Dr. Hotchkiss and others and licensed by
Washington University in St. Louis to IFDx LLC. That technology is evaluated in this research.
He is also supported by R35 GM-126928, awarded by the National Institute of General Medical
Sciences.

Charles C. Caldwell, Ph.D. Dr. Caldwell and the University of Cincinnati may receive royalty
income based on a technology developed by Dr. Caldwell and others and licensed by
Washington University in St. Louis to IFDx LLC. That technology is evaluated in this research.

Abbreviations

IFNγ, interferon-γ

ELISpot, enzyme-linked immunospot assay

SOFA, sequential organ failure assessment score

ALC, absolute lymphocyte count

IL-6, interleukin 6

sPD-L1, soluble programmed death ligand-1

IL-10, interleukin 10

SEPSIS, critically ill, sepsis cohort of patients

CINS, critically ill, nonseptic cohort of patients

LTAC, long-term acute care facility

IPR, in-patient rehabilitation

SNF, specialized nursing facility

AMA, discharged against medical advice

HLA-DR, human leukocyte antigen DR isotype

PBMC, peripheral blood mononuclear cell

SS, ELISpot spot size

SFU, ELISpot spot forming units

TE, ELISpot total expression

CTSI, UF Clinical and Translational Science Institute,

mAb, monoclonal antibody

AUROC, area under the receiver-operator curve

Abstract

Background: Sepsis remains a major clinical challenge for which successful treatment requires greater precision in identifying patients at increased risk of adverse outcomes requiring different therapeutic approaches. Predicting clinical outcomes and immunological endotyping of septic patients has generally relied on using blood protein or mRNA biomarkers, or static cell phenotyping. Here, we sought to determine whether functional immune responsiveness would yield improved precision.

Methods: An *ex vivo* whole blood enzyme-linked immunosorbent (ELISpot) assay for cellular production of interferon-γ (IFN-γ) was evaluated in 107 septic and 68 non-septic patients from five academic health centers using blood samples collected on days 1, 4 and 7 following ICU admission.

Results: Compared with 46 healthy subjects, unstimulated and stimulated whole blood IFNγ expression were either increased or unchanged, respectively, in septic and nonseptic ICU patients. However, in septic patients who did not survive 180 days, stimulated whole blood IFNγ expression was significantly reduced on ICU days 1, 4 and 7 (all $p<0.05$), due to both significant reductions in total number of IFNγ–producing cells and amount of IFNγ produced per cell (all $p<0.05$). Importantly, IFNγ total expression on day 1 and 4 after admission could discriminate 180-day mortality better than absolute lymphocyte count (ALC), IL-6 and procalcitonin. Septic patients with low IFNγ expression were older and had lower ALC and higher sPD-L1 and IL-10 concentrations, consistent with an immune suppressed endotype.

Conclusions: A whole blood IFNγ ELISpot assay can both identify septic patients at increased risk of late mortality, and identify immune-suppressed, sepsis patients.

192   (word count: 249)

194   **Trial Registry:** Because the study is a prospective observational study, and not a clinical trial,

195   registration with *clinical trials.gov* is not required.

Introduction

Sepsis remains one of the most common causes of critical illness and too often leads to death and morbidity (1-3). Importantly, sepsis is a pathophysiologic host response to microbial infection associated with organ injury and dysfunction (4). However, the nature and magnitude of the host response to sepsis is highly variable depending upon the subject's age, comorbidities and source and severity of microbial infection. Although sepsis is frequently associated with an early exaggerated inflammatory response (5), persistent inflammation (6,7), coagulopathy (8), prolonged immune suppression (3,6,9-12), and lean tissue wasting (13,14), the contribution of these individual responses to the overall outcome of the patient is still unresolved (15,16). Precision medicine has been proposed as a tool to identify which immunologic endotype drives organ injury and is an appropriate target for therapeutic intervention (17). Biomarkers, based on static blood cell phenotypes, protein and transcriptomic metrics have been commonly used to endotype critically ill patients with and without sepsis (15,17-20).

Enzyme-Linked ImmunoSpot assay (ELISpot) is a widely used immunological technique that enables the detection and quantification of individual cells responding to external receptor-specific and nonspecific stimulants and secreting specific proteins, particularly cytokines (21). This method is important for studying the immune response at the single-cell level, offering valuable insights into immune cell function, and immune-related diseases. Its ability to analyze immune responses at the cellular level makes it particularly suitable for monitoring immune system functionality in sepsis. In the current report, we examined the extent to which whole blood ELISpot production of interferon γ (IFNγ) can identify immune suppressed, critically ill patients at increased risk of death.

Materials and Methods

This multi-center, prospective diagnostic and prognostic study conducted between February

23, 2021, and July 22, 2022, enrolled two cohorts of critically ill patients at the time of ICU

admission. The first cohort included patients with a suspected diagnosis of sepsis admitted to

the ICU (SEPSIS). The second cohort included critically ill patients admitted to the ICU without

currently suspected sepsis (Critically Ill, Non-Septic; CINS), but considered at high risk for

subsequent infection (e.g., postoperative, severe trauma). Patient enrollment is shown in

Figure 1, consistent with Enhancing the Quality and Transparency of Health Research

Standards for Reporting of Diagnostic Accuracy (STARD) reporting guidelines (22). All patients

were managed under institutional clinical management protocols. Centralized ethics approval

was obtained from the University of Florida Institutional Review Board (#IRB 202000924)

which served as the sponsoring institution. Written informed consent was obtained from each

patient or their proxy decision-maker at individual clinical sites.

A heparinized blood collection tube (Becton Dickinson) was obtained within the first three days

of ICU admission (labeled as day 1), and on subsequent days 3 through 5 (labeled as day 4),

and weekly thereafter (± 2 days). Self- or proxy-reported race and ethnicity category data were

collected as per National Institutes of Health reporting guidelines and requirements.

Inclusion criteria consisted of ICU admission from the emergency department for community-

acquired sepsis or severe trauma (patients with Injury Severity Scores >15, hemorrhagic

shock, and/or severe chest trauma), as well as non-trauma, postoperative ICU admission, ICU

transfer from the emergency department and inpatient transfer from ward to ICU.

Sepsis was defined according to Sepsis-3 criteria (4), and all subjects were clinically

adjudicated at the individual participating sites. Patients admitted to the ICU for critical illness,

non-sepsis (CINS) were also adjudicated to rule out sepsis. A detailed summary of inclusion

and exclusion criteria is provided in the **Supplemental Materials and Methods.**

Healthy control subjects were recruited at each of the clinical sites. Efforts were made to age,

sex and race-ethnicity match the healthy control subjects to the SEPSIS cohort. Individuals

with autoimmune diseases being treated with biologic immune modulators were excluded, as

were individuals who had received anti-neoplastic therapies or diagnosed with cancer within

the past six months. Vulnerable populations were also excluded.

Primary Outcomes and Clinical Adjudication

The primary clinical outcome for ELISpot was 180-day mortality, determined via clinical

records and telephone follow-up with the patient, their proxy, or their designated contact, and

cross-checked through the US Social Security Death Index. We analyzed temporal trends of

ELISpot in both SEPSIS and CINS patients but compared estimated performance of predictive

models primarily in the SEPSIS patients, as 180-day mortality in the CINS cohort was <4%.

Final sepsis or CINS adjudication was performed by physician-investigator determination at

each clinical site at completion of each patient's hospital course.

Secondary clinical outcome variables included all-cause (in hospital, 30-day) mortality,

development of chronic critical illness (CCI), secondary infections, and poor discharge

disposition. Inpatient clinical trajectory was defined as "early death", "rapid recovery", or "CCI".

CCI was defined as an ICU length of stay $\geq$14 days with evidence of persistent organ

dysfunction (SOFA score $\geq$ 2) (23). Hospitalized patients who died after an ICU length of stay

>14 days from the index hospitalization were also classified as CCI. Poor disposition was

defined as discharge to a skilled nursing facility (SNF), long-term acute care facility (LTAC), or

hospice. Secondary infections were defined as per the US Centers for Disease Control and

Prevention criteria.

ELISpot

ELISpot assays were conducted using the human IFNγ Immunospot[®] kit (CTL Inc., Cleveland,

OH) with several important modifications including the use of diluted whole blood as previously

described (21). Specifically, 100 μl of heparinized whole blood was diluted 1:10 with kit buffer

and 50 μl of the diluted sample was added to each well. Samples were incubated in wells

containing either buffer alone or a soluble anti-CD3/anti-CD28 (125 ng/ml/1.25 μg/ml) mAb

agonist (BioLegend, San Diego, CA). Samples were assayed in duplicate. Optimal

concentrations of agonist were determined in preliminary studies (see **Supplemental**

Materials and Methods).

Samples were quantitated using a CTL S6 Entry or S6 FluoroCore™ ELISpot reader at each

clinical site. To assure comparable results, the instruments were harmonized by CTL Inc. prior

to study using an external standard across all five clinical sites. Results are presented as the

number of spot-forming units (SFU), spot size (μm^2; SS) and total expression (μm^2; TE) a

product of the number of spots and mean spot size using the Immunospot[®] SC software suite

(version 7.0.30.4). Spot forming units represent individual blood cells expressing IFNγ and spot

size is an indication of the amount of IFNγ produced per cell. In subsequent analyses, the

number of IFNγ-producing cells was adjusted for each individual patient's absolute lymphocyte

count to yield the percentage of total lymphocytes expressing IFNγ.

Additional Laboratory Analyses

Whole blood total leukocyte and absolute lymphocyte counts were determined on EDTA-anticoagulated whole blood at the individual clinical sites either using their hospital's Clinical and Diagnostics Laboratory or a research Beckman-Coulter Dx500 or Dx900 hemocytometer (Brea, CA). Cytokine and additional plasma protein analyses were conducted at the University of Florida Sepsis and Critical Illness Research Center (SCIRC) where they were determined in batch using the Luminex MagPix® (Austin, TX) platform using commercial reagents.

Excess sample was stored at the Biorepository of the Clinical and Translational Science Institute (https://www.ctsi.ufl.edu/research/laboratory-services/ctsi-biorepository-2/) where it is available to the scientific community under guidelines promulgated by the National Institute of General Medical Sciences.

Data Collection and Analysis

Clinical data collection was conducted at each site and entered into a web-based electronic case report form created on the REDCap™ platform managed by the University of Florida Clinical and Translational Science Institute (CTSI). Access to the case report form was password protected and limited to only approved research staff, and all interactions with the database were recorded. Peer-to-peer communication allowed approved individuals at all five sites access to their own data and de-identified data from the other four clinical sites. Research data, including ELISpot, total leukocyte and absolute lymphocyte counts, and plasma protein and cytokine data were uploaded into the case report forms from the University of Florida SCIRC. Data managers at the SCIRC were responsible for creating final locked datasets for subsequent analysis.

Statistical Analysis

Descriptive data are presented as frequencies and percentages or medians and IQRs. The

Fisher exact test and Mann-Whitney or Kruskal-Wallis ANOVA tests were used for categorical

and continuous variables, respectively. Area under the receiver operating characteristics curve

(AUROC) values with 95% confidence intervals (computed with 2000 stratified bootstrap

replicates) were used to assess discrimination. Univariable and multivariable logistic

regression were performed to assess whether the combination of metrics improved overall

performance. Post hoc tests were performed for continuous outcomes using the Dunn test. For

post hoc analyses of categorical outcomes, separate 2 × 2 Fisher exact tests were performed.

All significance tests were 2-sided, with a raw $p \leq 0.05$ considered statistically significant.

Analyses were performed using the R Project statistical package, version 4.1.0 (R Project for

Statistical Computing).

Results

Patient Characteristics

Demographic characteristics of the 175 enrolled patients (99 men [57%] and 76 women [43%])

and 46 healthy control subjects (16 men [36%], 30 women [64%]) are summarized in **Table 1**.

The overall cohort included 107 and 68 patients in the SEPSIS and CINS cohorts, respectively.

Patient characteristics were similar across all three cohorts with the exception that the healthy

control subjects were more predominantly female and younger, and SEPSIS patients had a

higher Charlson comorbidity index than CINS subjects (**Table 1**). Within the CINS cohort

([n = 68]), reason for ICU admission was identified in **Supplemental Table 1**.

Table 2 shows clinical outcomes for the SEPSIS and CINS patients. Hospital length of stay

(p<0.02), incidence of secondary infections (p<0.001), development of CCI (p<0.001), and in-

hospital mortality (p<0.01) were all significantly higher in SEPSIS than in CINS patients.

Disposition at discharge also significantly differed between SEPSIS and CINS patients, as did

30- and 180-day mortality (both p<0.01).

IFNγ Production by Unstimulated and anti-CD3/CD28 mAb-Stimulated Whole Blood

Comparison of the first sample collected (days 1-3 post ICU admission) among the three

cohorts revealed considerable heterogeneity in the individual subject response, irrespective of

the cohort. Surprisingly, as a group, spontaneous IFNγ production in *unstimulated* whole blood

was significantly increased from SEPSIS and CINS patients than from healthy subjects at all

sampling intervals (days 1, 4 and 7), despite a significant reduction in lymphocyte numbers

(**Figure 2, Supplemental Figure 1**). This was reflected generally by an increased number of

IFNγ-producing cells (SFU) (all p<0.05;), although the amount of IFNγ produced by each cell

(SS) was increased on day 4 (**Figure 3**). When the total number of IFNγ producing cells was

adjusted for the absolute lymphocyte count, the percentage of lymphocytes expressing IFNγ

was further increased significantly in both SEPSIS and CINS (p<0.001; **supplemental Figure**

2). There was no difference between unstimulated IFNγ expression between the SEPSIS and

CINS cohorts.

Ex vivo stimulation of the whole blood from the three cohorts with agonist anti-CD3/CD28 mAb

resulted in expected increases in the total expression of IFNγ produced when compared with

unstimulated samples. There were also increase in the total amount of IFNγ produced per unit

volume of blood on days 4 and 7 when comparing CINS to healthy subjects (**Figure 2**). This

increased IFNγ expression was only seen on day 4 in SEPSIS patients.

Influence of Outcome on ELISpot Responses

SEPSIS subjects had a greater in-hospital, 30- and 180-day mortality when compared to CINS

patients (**Table 2).** In addition, the incidence of secondary infections, development of CCI and

an adverse discharge disposition were all significantly greater in SEPSIS than CINS patients

(all $p<0.05$).

SEPSIS subjects who died within 180 days of ICU admission did not differ from surviving

patients based on their admission or day 1 SOFA scores or total leukocyte numbers, even

though non-surviving SEPSIS patients were significantly older and had higher Charlson

comorbiditiy scores (both $p<0.05$; **Supplemental Table 1**). Interestingly, there were marked

differences in the IFNγ production from stimulated whole blood between sepsis subjects who

survived or did not survive 180 days. Both the absolute number (SFU) and percentage of IFNγ

producing cells, and the spot size (SS) were significantly lower, and therefore, total expression

(TE) was reduced in non-surviving versus surviving SEPSIS patients (all $p<0.05$; **Figure 4**).

This reduction in expression was, in general, sustained through day 7 in non-survivors,

although significance varied (**Figure 4**). After 7 days, there were too few ICU-remaining

patients to continue the comparison (*data not shown*). Surprisingly, IFNγ production from

unstimulated whole blood did not differ between surviving and non-surviving patients at any

time point, although it only remained elevated in surviving SEPSIS patients when compared

with healthy control subjects at later time points ($p<0.05$; **Figure 4**). There also did not appear

to be any significant changes in ELISpot responses (both stimulated and unstimulated) over

time in either surviving or non-surviving SEPSIS patients.

Univariate and Multivariate Prediction Models for Long-term Survival and Secondary

Outcomes.

Because of the differences in IFNγ expression between surviving and non-surviving SEPSIS

patients, ELISpot AUROCS were evaluated for their discriminatory prediction of long-term

survival (180 day), as well as secondary outcomes, and compared to clinical indices (SOFA,

Charlson comorbidity scores), total white blood cell and absolute lymphocyte counts, and

plasma protein markers (selected cytokines, procalcitonin, and sPD-L1) in the SEPSIS

patients. Similar combined SEPSIS and CINS analyses could not be performed due to the low

mortality in CINS patients. Results are presented in **Table 3** and **Figure 5**.

Similar to data reported by ourselves and others (24-26), the most consistent discriminator of

180-day mortality was the Charlson comorbidity score (AUROC; 0.824 [CI 0.700-0.948], which

also discriminated in-hospital mortality (0.762 [0.611-0.912]) and development of CCI (0.713

[0.594-0.832]), but not the incidence of secondary infection (0.549 [0.426-0.673]. Importantly,

stimulated total IFNγ expression at day 1 and day 4 did not significantly differ from the

Charlson comorbidity score with AUROCs of 0.730 (0.601-0.859) and 0.794 (0.691-0.897),

respectively. Total IFNγ expression on day 4 also discriminated in-hospital mortality with an

AUROC of 0.743 (0.615-0.830), but it was not a strong discriminator of either development of

CCI or incidence of secondary infections.

For total stimulated IFNγ expression, the components contributing to its discriminative power

for 180-day mortality were both the number of IFNγ producing cells (SFU; Day 1, 0.707 [0.568-

0.846] and the amount of IFNγ produced by individual cells (SS; Day 1, 0.707 [0.568-0.846]).

The discriminatory power of the ELISpot total expression (TE) for predicting both in-hospital

and 180-day mortality was much greater than seen for either absolute lymphocyte count

(ALC), IL-6, or procalcitonin at either day 1 or day 4, or for changes in these parameters

between days 1 and 7 (**Table 3**). In addition, the last ELISpot measurement obtained from the

patient prior to discharge or death (usually day 7 or later) was also found to be not as

discriminatory as the earlier day 1 and 4 measurements (*data not shown*).

Setting the threshold for day 1 and day 4 stimulated total IFNγ expression at approximately

80% sensitivity to discriminate 180-day survival, it was possible to assess the

immunosuppressive endotype of those SEPSIS patients with reduced ELISpot total expression

(**Table 4**). These individuals were older and had absolute lymphocyte counts significantly lower

at both days 1 and 4 than SEPSIS patients above the ELISpot threshold. In addition, plasma

sPD-L1 concentrations were significantly higher on day 4. Patients below the ELISpot

threshold also had in-hospital and 180-day mortality, and development of CCI at a markedly

higher frequency than those individuals above the threshold (all p<0.05).

Finally, to examine whether ELISpot could improve the discriminatory power of standard

clinical indices (SOFA and Charlson comorbidity scores), single and multivariate logistic

regression analyses were performed and the AUROCs of the models were calculated by a

four-fold cross-validation procedure (**Table 5**). Model I was built on baseline Charlson

comorbidity data and day 1 SOFA scores yielding AUROCs for 180-day mortality (0.911

[0.858-0.965]); the odds ratios show baseline Charlson comorbidity data and day 1 SOFA

scores are significant predictors for 180-day mortality (**Table 5**). Model II was built on

stimulated total IFNγ expression from Day 4 yielding AUROCs for 180-day mortality (0.794

[0.732-0.855]); the odds ratios show stimulated total IFNγ expression is a significant predictor

for 180-day mortality (**Table 5**). Although ELISpot data show significant odds ratios in Model II,

the addition of stimulated total IFNγ expression to the model built on standard score indices

(Model III) did not significantly increase either the AUROC or the odds ratio for 180-day (0.915

[0.851-0.979] or in-hospital mortality (0.883 [0.801-0.964]), or development of CCI (0.861

[0.830-0.892]) (**Table 5**).

Discussion

Key Findings

This prospective, multi-center observational study has demonstrated that the adaptive immune

response to critical illness, as defined by *ex vivo* whole blood production of IFNγ in response to

T cell receptor stimulation, varied in response to critical illness (**Figures 2, 3**) and could also

discriminate long-term outcomes (**Figure 5, Table 3**). Spontaneous IFNγ production by diluted

whole blood was significantly increased in critically ill patients, irrespective of whether the

critically ill patients were septic. In addition, when whole blood was stimulated *ex vivo* with a T

cell receptor agonist*,* IFNγ production increased dramatically in both healthy and critically ill

cohorts (both SEPSIS and CINS). However, stimulated ELISpot IFNγ total expression early in

the admission to the ICU significantly differed between septic patients who survived 180 days

and those who died, and this reduction in total expression in non-surviving septic patients was

due to both reductions in the total number of IFNγ producing blood cells and the amount of

IFNγ produced by individual cells (**Figure 4**). Importantly, day 1 and day 4 measurements were

more discriminatory than later measurements. Using univariate modeling, stimulated ELISpot

total expression measured in the first week of admission (sampling days 1 and 4) could

differentiate 180-day mortality as well as SOFA and Charlson comorbidity scores, and

markedly better than blood ALC, procalcitonin, IL-6 and sPD-L1 concentrations (**Table 3**).

Septic patients who had low stimulated ELISpot total IFNγ expression at days 1 and 4 had an

immunosuppressed endotype, as reflected in being older, and having lower ALC counts,

higher plasma sPD-L1, and increased incidence of chronic critical illness, in-hospital and late

mortality (**Table 4**). However, in multivariate models, stimulated ELISpot total IFNγ expression

did not significantly improve the discrimination between 180-day survival in models built with

SOFA and Charlson comorbidity scores (**Table 5**).

Context

ELISpot has emerged as a powerful method to assess immunological status in a variety of

clinical disorders (27-30), including sepsis and critically ill patients (20,21,31,32). It offers

several theoretical advantages over other current metrics – for example, cell phenotypes (such

as ALC (33-36) and HLA-DR expression on CD14[+] blood cells (37,38)), plasma protein

concentrations (such as procalcitonin (39-42), IL-6 (43-45) and sPD-L1 (46,47)) or blood

transcriptomics (26,48,49) – used to predict both the severity of the host response and the

immunosuppressed endotype. ELISpot, unlike these static measures, assesses one

component of the functional status of the host protective immune response. In the present

study, ELISpot revealed the capacity of T cells in the blood to produce IFNγ with and without

stimulation through the T cell receptor.

This study is not the first to demonstrate reduced IFNγ ELISpot expression in sepsis patients,

especially in those with adverse clinical outcomes (20,21,50). However, in contrast to these

previous studies, we used diluted whole blood in our ELISpot assay instead of isolated PBMCs

and observed increased IFNγ production in both unstimulated and stimulated whole blood from

critically ill patients. There are two key advantages of using diluted whole blood in the ELISpot.

First, the use of whole blood permits the assay to take place with the entire blood composition

(i.e., all leukocytes, erythrocytes, platelets, and plasma proteins and metabolites) maintained.

Responses to critical illness and *ex vivo* stimulation may be either direct or be mediated via

cell-cell communication and/or plasma mediators. Traditional processing of blood by density

gradient centrifugation separates the PBMC from the neutrophils, platelets, and plasma.

Second, it is a simpler and more 'rapid' assay to set up because there is no required cell

isolation step.

On days 1-7, both the CINS and SEPTIC patients had reduced ALC as compared to healthy

controls (**Supplemental Figure 1**). Surprisingly, despite this lymphopenia, the number of

lymphocytes spontaneously producing IFNγ was higher in both CINS and SEPSIS patients

versus healthy controls at all three time points (**Figure 2**). Day 1 stimulated IFNγ SFU, spot

size (SS), and total expression did not differ between cohorts. However, on Day 4, both the

number of cells producing IFNγ as well as total expression of IFNγ was higher in both the

CINS and SEPSIS cohorts as compared healthy subjects. On Day 7, the number of cells

producing IFNγ remained higher in the CINS and SEPSIS cohorts. Of note, the amount of IFNγ

produced on a per cell basis (reflected by spot size) was lowest in the SEPSIS cohort with total

expression similar to healthy controls.

Given the considerable amount of data showing sepsis can evolve into an immunosuppressed

state, it was surprising to see both the spontaneous and stimulated IFNγ production increase

in septic patients (early after admission) versus healthy subjects. There are several likely

explanations for this apparent paradox. The first potential explanation relates to timing;

specifically, the data presented herein came from blood samples collected within the first 7

days after ICU admission. It is difficult to determine exactly when the sepsis-induced

hyperinflammation transitions to a state of immunoparalysis, but it is tempting to speculate that

our assessment of immune fitness was still within the window of hyperinflammation and

exacerbated immune cell activity.

A second explanation has been termed "*bystander activation*" (51,52). The inflammatory

response that develops during infection has a capacity to trigger antigen-experienced effector

and/or memory CD8$^+$ T cells present in a T cell receptor-independent and cytokine-dependent

manner. A number of cytokines including IL-12, IL-15, TNFα, and IL-18 induce CD8$^+$ T cell

activation and resultant IFNγ production (53). Thus, the sepsis cytokine milieu likely primes

pre-existing, effector and memory CD8$^+$ T cells to produce IFNγ in a cognate antigen-

independent fashion (**Figure 2**). In addition, these cytokine-primed effector/memory CD8$^+$ T

cells will also respond with IFNγ production to a myriad of cytokines produced *ex vivo* during

anti-CD3/CD28 stimulation.

A third explanation may lie in the differences in the lymphocyte subsets present in the

peripheral blood of healthy subjects versus CINS and SEPSIS patients at the time of blood

collection. *De novo* clonal expansion of pathogen-specific effector CD8$^+$ T cells in response to

sepsis-inducing pathogens and resultant inflammation leads the potential for a preponderance

of active response effector cells – especially early in the septic timeline. In contrast, healthy

control volunteers are more likely to have 'resting' naïve and memory T cells and a minimal (if

any) increase in inflammatory cytokines. Consequently, the number of T cells capable of

rapidly responding to polyclonal and/or bystander cytokine stimulations and produce IFNγ in

the ELISpot assay is increased in SEPSIS patients compared with healthy subjects.

Current Work

The current studies add to the body of information suggesting ELISpot examining whole blood

production of IFNγ can both discriminate long-term mortality and identify those patients who

may benefit from therapeutic interventions targeting adaptive immunity.

Limitations

This study has several limitations. Despite multicenter enrollment, sample sizes were still

relatively small for discriminative modeling. Over the past two decades, improved in-hospital

management has reduced the number of adverse events and in-hospital mortality to sepsis

and critical illness (25,54). Discriminatory analyses could only be conducted in the SEPSIS

cohort as CINS patients had very low in-hospital (1%) and 180-day mortality (4%) (**Table 2**).

Second, every effort was made to match healthy control subjects to the SEPSIS and CINS

cohorts, but the healthy donors used in this study were, as a group, significantly younger and

more frequently female (**Table 1**). Median ages in the healthy control cohort were greater than

45 years, a break point often determined to be associated with increased adverse outcomes in

critically ill patients (25). Despite the multicenter nature of the study, the cohorts still were also

predominantly Caucasian.

Future Directions

While the findings presented herein suggest assessing IFNγ production by ELISpot can be

useful in identifying sepsis patients at risk of long-term mortality and the immune suppressed

endotype, its discriminative ability is similar to SOFA and Charlson comorbidity indices and

does not add significantly to their discriminative power. With that said, SOFA and Charlson

comorbidity indices are rarely used for clinical decision making because they provide no

insights or therapeutic directions into the immunological disturbances associated with sepsis

and adverse outcomes. Application of ELISpot to the clinical armamentarium has the potential

to provide important information regarding which sepsis patients would benefit from targeted

therapy (precision medicine). For example, septic patients who have profound suppression of

stimulated IFNγ production may be harmed by therapy with corticosteroids, but might be good

candidates for immune adjuvants therapies to boost their ability to combat invading pathogens.

ELISpot is an FDA-cleared approach for assessing functional immune status to prior

tuberculosis infection and the ELISpot reader used in these studies (CTL S6 Entry) is FDA

510(k) cleared. However, to make these results more actionable, ELISpot results will need to

be obtained within hours, instead of days. Currently, ELISpot results take at least 24 hours to

return, although preliminary data from our consortium suggests the assay can be modified to

produce results in less than 12 hours (*Griffith, T and Caldwell, C., manuscript in preparation*).

In addition, ELISpot can be readily used to assess other components of the blood innate and

adaptive immune response simply by varying the stimulant and the readout metric. For

example, innate immune responses have been readily assessed using endotoxin or other TLR

ligands as a whole blood stimulant and TNFα as the readout (20,21). Furthermore, underlying

mechanisms of adaptive or innate immune responses can be explored using alternative

stimulants (30,31,55), simultaneous adjuvants or inhibitors (56), or different readout metrics

(21,55).

Conclusions

ELISpot can assess functional immune status in critically ill patients, predict adverse long-term

outcomes, and identify subsets of patients who may benefit from immunostimulant therapy.

Contributions:

University of Florida

EAB –Investigation, Visualization, Writing-Review and Editing; **PAE**- Resources, Funding Acquisition, Supervision, Writing-Review and Editing; **ML**-Investigation, Visualization; **TL**-Supervision, Investigation, Writing-Review and Editing; **VP**- Investigation, Visualization, **LLM**-Conceptualization, Funding Acquisition, Resources, Supervision, Project Administration, Writing-Review and Editing; **RU**- Supervision, Investigation, Visualization, **JPL**- Supervision, Investigation, **MD**-Investigation, Visualization, **LZS**-Investigation, Visualization, **RD**-Investigation, **AM**-Investigation

University of Iowa

EES– Investigation, Visualization; **MH**– Investigation, Visualization; **PMcG** – Conceptualization, Project Administration; **MR**– Investigation, Visualization; **VPB**– Conceptualization, Funding Acquisition, Resources, Supervision, Writing-Review and Editing

University of Minnesota

RWG – Conceptualization, Project administration; **TSG**– Conceptualization, Funding, Acquisition, Resources, Supervision, Writing–Review and Editing; **LH**– Project Administration; **TAK**– investigation, visualization; **CN**– project administration; **JX**– investigation, visualization.

Washington University in St. Louis

RSH- Conceptualization, Funding, Acquisition, Resources, Supervision, Writing–Review and Editing; **DDT-** Investigation, **AD**- Investigation, **SM1**-Investigation, **SM2**-Investigation, **IT**-Conceptualization, Supervision, Writing-Review and Editing

University of Washington

SB- Conceptualization, Funding, Writing–Review and Editing

University of Cincinnati

CCC- Conceptualization, Funding, Acquisition, Resources, Supervision, Writing–Review and Editing; **MDG**- Acquisition, Investigation, Visualization, Supervision. **CBB** – Acquisition, Investigation

Case Western Reserve University

MBM- Conceptualization**,** Supervision, Investigation, Writing-Review and Editing; **KR**-Conceptualization, Investigation, Acquisition, Writing-Review and Editing.

Acknowledgements: This work was directly supported by R01 GM-132364, awarded by the National Institute of General Medical Sciences, including a supplement (R01 GM-132364-03S1) in 2022-2024.

Additional support was provided by grants T32 GM-008721 (EAB, VP, PAE), RM1 GM-139690 (LLM, PAE), R01 GM-149657 (TL), R01 GM-124156 (MDG), R35 GM-140806 (PAE), R35 GM-133756 (IT), R35 GM-134880 (VB), R35 GM-134880 (SB), R35 GM1-40881 (TSG), R35 GM-12698 (RSH), all from the National Institute of General Medical Sciences, G102983-6263608307-1 (MDG) from Department of Defense, IK6BX006192 from the Department of Veteran Affairs (TSG), and BE 7016/1-1 (CD) from the Deutsche Forschungsgemeinschaft (German Research Foundation) .

The authors also wish to gratefully acknowledge the patients and their families for their willingness to participate in an observational clinical study with no direct benefit to them.

References

1.   Bauer M, et al. Mortality in sepsis and septic shock in Europe, North America and

Australia between 2009 and 2019- results from a systematic review and meta-analysis. *Crit*

Care. 2020;24(1):239.

2.   Cecconi M, et al. Sepsis and septic shock. *Lancet*. 2018;392(10141):75-87.

3.   Hotchkiss RS, et al. Sepsis and septic shock. *Nat Rev Dis Primers*. 2016;2:16045.

4.   Singer M, et al. The Third International Consensus Definitions for Sepsis and Septic

Shock (Sepsis-3). *JAMA*. 2016;315(8):801-10.

5.   Silva EE, et al. The Calm after the Storm: Implications of Sepsis Immunoparalysis on

Host Immunity. *J Immunol*. 2023;211(5):711-719.

6.   Darden DB, et al. Biomarker Evidence of the Persistent Inflammation,

Immunosuppression and Catabolism Syndrome (PICS) in Chronic Critical Illness (CCI) After

Surgical Sepsis. *Ann Surg*. 2021;274(4):664-673.

7.   Efron PA, et al. The Persistent Inflammation, Immunosuppression, and Catabolism

Syndrome (PICS) Ten Years Later [published online August 8, 2023]. *J Trauma Acute Care*

Surg. doi: 10.1097/TA.0000000000004087.

8.   de Nooijer AH, et al. Complement activation in severely ill patients with sepsis: no

relationship with inflammation and disease severity. *Crit Care*. 2023;27(1):63.

9.   Roth WJ, et al. Characterization of two distinct transglutaminases of murine bone

marrow-derived macrophages: effects of exposure of viable cells to cigarette smoke on

enzyme activity. *J Leukoc Biol*. 1987;42(1):9-20.

10. van Oosterom A, Huiskamp GJ. The effect of torso inhomogeneities on body surface

potentials quantified using "tailored" geometry. *J Electrocardiol*. 1989;22(1):53-72.

11. van der Poll T, et al. The immunopathology of sepsis and potential therapeutic targets.

Nat Rev Immunol. 2017;17(7):407-420.

12. van der Poll T, et al. The immunology of sepsis. *Immunity*. 2021;54(11):2450-2464.

13. Cox MC, et al. The impact of sarcopenia and acute muscle mass loss on long-term

outcomes in critically ill patients with intra-abdominal sepsis. *J Cachexia Sarcopenia Muscle*.

2021;12(5):1203-1213.

14. Zhang W, et al. CMV Status Drives Distinct Trajectories of CD4+ T Cell Differentiation.

Front Immunol. 2021;12:620386.

15. Rincon JC, et al. Immunopathology of chronic critical illness in sepsis survivors: Role of

abnormal myelopoiesis. *J Leukoc Biol*. 2022;112(6):1525-1534.

16. Maslove DM, et al. Redefining critical illness. *Nat Med*. 2022;28(6):1141-1148.

17. Fenner BP, et al. Immunological Endotyping of Chronic Critical Illness After Severe

Sepsis. *Front Med (Lausanne)*. 2020;7:616694.

18. Scicluna BP, et al. Classification of patients with sepsis according to blood genomic

endotype: a prospective cohort study. *Lancet Respir Med*. 2017;5(10):816-826.

19.   Balch JA, et al. Defining critical illness using immunological endotypes in patients with

and without sepsis: a cohort study. *Crit Care*. 2023;27(1):292.

20.   Loftus TJ, et al. Overlapping but Disparate Inflammatory and Immunosuppressive

Responses to SARS-CoV-2 and Bacterial Sepsis: An Immunological Time Course Analysis.

Front Immunol. 2021;12:792448.

21.   Mazer MB, et al. A Whole Blood Enzyme-Linked Immunospot Assay for Functional

Immune Endotyping of Septic Patients. *J Immunol*. 2021;206(1):23-36.

22.   Bossuyt PM, et al. The STARD statement for reporting studies of diagnostic accuracy:

explanation and elaboration. *Ann Intern Med*. 2003;138(1):W1-12.

23.   Gardner AK, et al. The Development of Chronic Critical Illness Determines Physical

Function, Quality of Life, and Long-Term Survival Among Early Survivors of Sepsis in Surgical

ICUs. *Crit Care Med*. 2019;47(4):566-573.

24.   Jouffroy R, et al. Relationship between prehospital modified Charlson Comorbidity Index

and septic shock 30-day mortality. *Am J Emerg Med*. 2022;60:128-133.

25.   Brakenridge SC, et al. Current Epidemiology of Surgical Sepsis: Discordance Between

Inpatient Mortality and 1-year Outcomes. *Ann Surg*. 2019;270(3):502-510.

26.   Brakenridge SC, et al. Evaluation of a Multivalent Transcriptomic Metric for Diagnosing

Surgical Sepsis and Estimating Mortality Among Critically Ill Patients. *JAMA Netw Open*.

2022;5(7):e2221520.

27.   Albareda MC, et al. Distinct Treatment Outcomes of Antiparasitic Therapy in

Trypanosoma cruzi-Infected Children Is Associated With Early Changes in Cytokines, Chemokines, and T-Cell Phenotypes. *Front Immunol.* 2018;9:1958.

28.   Shin KH, et al. CMV specific T cell immunity predicts early viremia after liver transplantation. *Transpl Immunol.* 2018;51:62-65.

29.   Remy KE, et al. Severe immunosuppression and not a cytokine storm characterizes COVID-19 infections. *JCI Insight.* 2020;5(17)

30.   Puertas MC, et al. VIP-SPOT: an Innovative Assay To Quantify the Productive HIV-1 Reservoir in the Monitoring of Cure Strategies. *mBio.* 2021;12(3):e0056021.

31.   Bonavia AS, et al. Comparison of Rapid Cytokine Immunoassays for Functional Immune Phenotyping. *Front Immunol.* 2022;13:940030.

32.   Beckmann N, et al. Staging and Personalized Intervention for Infection and Sepsis. *Surg Infect (Larchmt).* 2020;21(9):732-744.

33.   Podd BS, et al. Early, Persistent Lymphopenia Is Associated With Prolonged Multiple Organ Failure and Mortality in Septic Children. *Crit Care Med.* 2023;

34.   Denstaedt SJ, et al. Blood count derangements after sepsis and association with post-hospital outcomes. *Front Immunol.* 2023;14:1133351.

35.   Drewry AM, et al. Persistent lymphopenia after diagnosis of sepsis predicts mortality. *Shock.* 2014;42(5):383-91.

36.   Zidar DA, et al. Association of Lymphopenia With Risk of Mortality Among Adults in the

US General Population. *JAMA Netw Open*. 2019;2(12):e1916526.

37.   Bidar F, et al. Concomitant Assessment of Monocyte HLA-DR Expression and Ex Vivo

TNF-alpha Release as Markers of Adverse Outcome after Various Injuries-Insights from the

REALISM Study. *J Clin Med*. 2021;11(1)

38.   Bodinier M, et al. Monocyte Trajectories Endotypes Are Associated With Worsening in

Septic Patients. *Front Immunol*. 2021;12:795052.

39.   Uzzan B, et al. Procalcitonin as a diagnostic test for sepsis in critically ill adults and after

surgery or trauma: a systematic review and meta-analysis. *Crit Care Med*. 2006;34(7):1996-

2003.

40.   Prkno A, et al. Procalcitonin-guided therapy in intensive care unit patients with severe

sepsis and septic shock--a systematic review and meta-analysis. *Crit Care*. 2013;17(6):R291.

41.   Wirz Y, et al. Effect of procalcitonin-guided antibiotic treatment on clinical outcomes in

intensive care unit patients with infection and sepsis patients: a patient-level meta-analysis of

randomized trials. *Crit Care*. 2018;22(1):191.

42.   Andriolo BN, et al. Effectiveness and safety of procalcitonin evaluation for reducing

mortality in adults with sepsis, severe sepsis or septic shock. *Cochrane Database Syst Rev*.

2017;1(1):CD010959.

43.   Tocu G, et al. The Contribution of Procalcitonin, C-Reactive Protein and Interleukin-6 in

the Diagnosis and Prognosis of Surgical Sepsis: An Observational and Statistical Study. *J

Multidiscip Healthc*. 2023;16:2351-2359.

44.   Patel RT, et al. Interleukin 6 is a prognostic indicator of outcome in severe intra-

abdominal sepsis. *Br J Surg*. 1994;81(9):1306-8.

45.   Casey LC, et al. Plasma cytokine and endotoxin levels correlate with survival in patients

with the sepsis syndrome. *Ann Intern Med*. 1993;119(8):771-8.

46.   Zeng C, et al. Predictive Value of Soluble Programmed Cell Death Ligand-1 in the

Progression of Septic Patients to Chronic Critical Illness in the Intensive Care Unit: A

Prospective Observational Clinical Study. *Shock*. 2023;60(2):163-171.

47.   Derigs M, et al. Soluble PD-L1 in blood correlates positively with neutrophil and

negatively with lymphocyte mRNA markers and implies adverse sepsis outcome. *Immunol*

Res. 2022;70(5):698-707.

48.   Kostaki A, et al. A 29-Mrna Host Response Whole-Blood Signature Improves Prediction

of 28-Day Mortality and 7-Day Intensive Care Unit Care in Adults Presenting to the Emergency

Department with Suspected Acute Infection and/or Sepsis. *Shock*. 2022;58(3):224-230.

49.   Galtung N, et al. Prospective validation of a transcriptomic severity classifier among

patients with suspected acute infection and sepsis in the emergency department. *Eur J Emerg*

Med. 2022;29(5):357-365.

50.   Haem Rahimi M, et al. Fully automated interferon-gamma release assay to monitor

antigen-independent T cell functionality: A proof of concept study in septic shock. *Cytokine*.

2023;169:156263.

51.   Kim TS, Shin EC. The activation of bystander CD8(+) T cells and their roles in viral

infection. *Exp Mol Med*. 2019;51(12):1-9.

52.   Martin MD, et al. Bystander responses impact accurate detection of murine and human

antigen-specific CD8 T cells. *J Clin Invest*. 2019;129(9):3894-3908.

53.   Freeman BE, et al. Regulation of innate CD8+ T-cell activation mediated by cytokines.

Proc Natl Acad Sci U S A. 2012;109(25):9971-6.

54.   Cox MC, et al. Abdominal sepsis patients have a high incidence of chronic critical illness

with dismal long-term outcomes. *Am J Surg*. 2020;220(6):1467-1474.

55.   Malewicz NM, et al. Early suppression of peripheral mononuclear blood cells in sepsis

in response to stimulation with cytomegalovirus, OKT3, and pokeweed mitogen. *J Appl Physiol*

(1985). 2019;127(6):1539-1547.

56.   Mazer M, et al. IL-10 Has Differential Effects on the Innate and Adaptive Immune

Systems of Septic Patients. *J Immunol*. 2019;203(8):2088-2099.

Table 1. Clinical Characteristics of Patient Cohorts and Healthy Control Subjects.
Patients included two cohorts of individuals admitted to the ICU, one presumed to be septic
(SEPSIS), a second presumed to not be septic, but critically ill and at risk of developing sepsis
(CINS). Values are obtained at time of admission to the ICU.

	CINS (n=68)	SEPSIS (n=107)	Healthy Control (n=46)	p value
Male [n (%)]	37 (54%)	62 (58%)	16 (35%)	0.0283
Age, years	59 (37, 67)	63 (51, 71)	47 (30, 62)	<0.0001
BMI, kg/m^2	27.3 (22.8, 33.9)	28.0 (22.7, 34.8)	N/C	0.718
SOFA Score	6 (4, 8)	6 (4, 10)	N/C	0.718
Charlson Comorbidity Score	2 (0, 4)	3 (1, 4)	2 (0, 2); n=19	0.0045
Total Leukocyte Counts				
WBC (x10^3/μl)	9.3 (7.2, 12.6)	12.8 (8.9, 18.6)	6.5 (5.5, 7.5)	<0.0001
Monocytes (%)	6.9 (5.2, 9.2)	4.6 (3.0, 7.1)	6.9 (6.1, 8.0)	<0.0001
Monocytes (x10^3/μl)	0.7 (0.4, 0.9)	0.7 (0.4, 1.0)	0.4 (0.4, 0.6)	0.0210
Neutrophils (%)	78.1 (71.9, 86.3)	86.1 (81.5, 91.4)	10.9 (7.7, 15.9)	<0.0001
Neutrophils (x10^3/μl)	7.2 (5.1, 10.6)	59 (55.2, 64.3)	3.7 (2.8, 4.6)	<0.0001
Lymphocytes (%)	12.4 (7.2, 17.5)	6.0 (4.3, 9.0)	29.4 (25.6, 35.2)	<0.0001
Lymphocytes (x10^3/μl)	1.0 (0.7, 1.4)	0.8 (0.5, 1.2)	1.9 (1.5, 2.3)	<0.0001
Plasma Proteins				
IL-10 (pg/ml)	18.3 (14.7, 20.7)	20.0 (14.8, 26.3)	17.4 (13.6, 20.8)	0.059
IL-6 (pg/ml)	35.5 (14.2, 155.9)	42.2 (19.9, 101.9)	5.83 (5.05, 6.92)	<0.0001
sPD-L1 (pg/ml)	157 (100, 229)	204.7 (125.0, 311.7)	78.5 (66.5, 93.7)	<0.0001
Procalcitonin (ng/ml)				

729   **N/C**, not collected

Table 2. Clinical Course and Clinical Outcomes in SEPSIS and CINS cohorts.

	CINS (n=68)	SEPSIS (n=107)	p value
LOS, days	10 (7, 14), n=66	13 (7, 21), n=106	0.016
ICU LOS, days	5 (3, 9), n=67	4 (2, 10)	0.904
Secondary infection	17 (29%)	52 (59%)	0.0003
Favorable discharge	52 (76%)	61 (57%)	0.0018
CCI	3 (4%)	15 (14%)	0.074
In-Hospital Mortality	1 (1.5%)	14 (13%)	
30-Day Mortality		12 (11%)	
180-Day Mortality		18 (17%)	
Disposition at discharge			
Home	25 (37%)	27 (25%)	0.104
LTAC	0	4 (3.7%)	0.017
IPR	17 (25%)	12 (11%)	0.104
Hospital	0	2 (1.9%)	0.406
Other	2 (2.9%)	2 (1.9%)	0.319
SNF	10 (15%)	21 (20%)	0.007
Residential facility	0	0	
Home with services	10 (15%)	21 (20%)	0.843
AMA	2 (2.9%)	1 (0.93%)	
Death	1 (1.5%)	14 (13%)	
Hospice facility	1 (1.5%)	2 (1.9%)	
Home with hospice	0	1 (0.93%)	

Sample sizes are identified in column headings except where missing values are noted for specific indices.

LOS, length of hospital stay

LICUS, length of ICU stays

CCI, chronic critical illness

LTAC, long term acute care facility

IPR, inpatient rehabilitation

SNF, specialized nursing facility

AMA, discharged against medical advice

Table 3. Selected Area Under the Receiver Operating Curve Discrimination for Primary Outcome Variables in the SEPSIS cohorts. Values represent the mean and 95% confidence intervals for the primary outcome variables: 180-Day Mortality, In-Hospital Mortality, Incidence of Secondary Infection, Development of Chronic Critical Illness (CCI).

Criterion	Time Point	180-day mortality	In-Hospital Mortality	Secondary Infection	CCI
Age	Baseline	0.68 (0.54,0.81)	0.42 (0.28,0.57)	0.48 (0.35,0.60)	0.62 (0.46,0.77)
Charlson Comorbidity Score	Baseline	**0.82 (0.70,0.95)**	**0.76 (0.61,0.91)**	0.55 (043,0.67)	0.71 (0.59,0.83)
SOFA	Baseline	0.72 (0.57,0.88)	0.72 (0.55,0.88)	0.59 (0.46,0.71)	0.65 (0.49,0.81)
	Day 1	**0.76 (0.62,0.90)**	**0.76 (0.60,0.92)**	0.65 (0.53,0.77)	**0.76 (0.66,0.87)**
	Day 4	**0.79 (0.65,0.93)**	**0.83 (0.67,0.98)**	0.68 (0.56,0.80)	**0.84 (0.74,0.93)**
ELIspot Stimulated IFNγ TE	Day 1	0.73 (0.60,0.86)	0.69 (0.55,0.84)	0.56 (0.44,0.69)	0.60 (0.45,0.75)
	Day 4	**0.79 (0.69,0.90)**	**0.74 (0.61,0.87)**	0.51 (0.37,0.65)	0.62 (0.43,0.81)
ELIspot Stimulated IFNγ SFU	Day 4	**0.74 (0.60,0.88)**	0.65 (0.47,0.84)	0.52 (0.39,0.66)	0.64 (0.47,0.82)
ELIspot Stimulated IFNγ SS	Day 1	0.71 (0.57,0.85)	0.68 (0.51,0.85)	0.56 (0.44,0.69)	0.56 (0.41,0.71)
	Day 4	0.66 (0.53,0.79)	0.70 (0.55,0.85)	0.60 (0.47,0.73)	0.52 (0.36,0.68)
% Lymphocytes Stimulated IFNγ TE	Day 1	0.68 (0.53,0.83)	0.62 (0.42,0.81)	0.58 (0.45,0.71)	0.53 (0.38,0.69)
	Day 4	0.60 (0.45,0.76)	0.54 (0.32,0.75)	0.65 (0.52,0.78)	0.58 (0.34,0.71)
Absolute lymphocyte count	Day 1	0.63 (0.45,0.81)	0.53 (0.31,0.75)	0.58 (0.50,0.72)	0.46 (0.27,0.65)
	Day 4	0.67 (0.52,0.81)	0.64 (0.48,0.80)	0.64 (0.51,0.77)	0.65 (0.47,0.84)
sPD-L1	Day 4	0.67 (0.53,0.80)	0.68 (0.52,0.84)	0.59 (0.45,0.73)	0.57 (0.38,0.77)
IL-6	Day 4	0.63 (0.47,0.79)	0.56 (0.37,0. 75)	0.64 (0.50,0.77)	0.62 (0.43,0.8)
IL-10	Day 4	0.62 (0.45,0.79)	0.51 (0.31,0.72)	0.60 (0.46,0.73)	0.57 (0.43,0.71)
IL-8/CXCL8	Day 4	0.60 (0.45,0.75)	0.64 (0.49,0.79)	0.46 (0.33,0.60)	0.68 (0.54,0.83)

Scores in bold are values equal to or greater than 0.74

TE, total expression

SFU, number of spot forming units

SS, spot size

Table 4. Evidence of Immunosuppression in SEPSIS Patients with Reduced Stimulated IFNγ Total Expression (TE) As Discriminated by ELISpot at Days 1 and 4 Post-ICU Admission. The threshold for reduced TE was set at 80% sensitivity. On day 1 the specificity was 42.1% and on day 4 the specificity was 63.2%.

Day 1	ELISpot IFNγ TE <2334/μm^2) N=65	ELISpot IFNγ TE $\geq$2334 (/μm^2) N=40	P-value
Age	64.2 (14.7)	54.6 (15.1)	0.002
Total WBC cell day 1 (x 10^3/ml)	13.9 (7.07)	15.7 (7.77)	0.158
Absolute Lymphocyte Count Day 1 (/μl^3)	980 (249)	1460 (600)	0.028
sPD-L1 day 1 (pg/ml)	315 (333)	263 (246)	0.311
IL-10, day 1 (pg/ml)	32.3 (70.0)	28.6 (37.7)	0.608
Outcome			**Odds-ratio**
180-day mortality	23.1%	7.5%	3.70 (1.00,13.7)
In-Hospital mortality	18.5%	5.0%	4.30 (0.91,20.3)
CCI	18.5%	7.5%	2.79 (0.74,10.6)
Second Infection	59.6% (6 missing)	58.8% (13 missing)	1.03 (0.43,2.49)

Day 4	ELISpot IFNγ TE <1755/μm^2) N=41	ELISpot IFNγ TE $\geq$1755 (/μm^2) N=51	P-value
Age	66.2(14.0)	58.1(15.7)	0.011

Total WBC cell day 1 (x 10^3/ml)	**13.6(6.93)**	**15.4(8.18)**	0.305
Absolute Lymphocyte Count Day 1 (/ml^3)	70.7 (197)	191 (580)	**0.006**
sPD-L1 day 1 (pg/ml)	337(248)	239(290)	**0.007**
IL-10, day 1 (pg/ml)	42.7(89.9)	24.0(27.7)	0.086
Outcome			**Odds-ratio**
180-day mortality	29.3%	5.9%	6.62 (1.72,25.5)
In-Hospital mortality	19.5%	5.9%	3.88 (0.96,15.7)
CCI	22.0%	7.8%	3.30 (0.94,11.7)
Second Infection	58.8% (12 missing)	56.4% (7 missing)	1.10 (0.44,2.80)

760
761

Table 5. Multivariate Logistic Regression to Discriminate Time to Mortality based on Charlson Comorbidity and SOFA Scores and Day 4 ELISpot Stimulated IFNγ Total Expression in SEPSIS Patients.

	Odds ratio	p value
Model I: Standard clinical indices only		
SOFA score Day 4	1.30 (1.12-1.50)	0.0003
Charlson comorbidity score (CCS) baseline	1.48 (1.24-1.77)	<0.0001
Model II: ELISpot indices only		
ELISpot IFNγ TE Day 4 (Negative of log)	1.44 (1.07-1.95)	0.017
Model III: Standard clinical indices + ELISpot indices		
SOFA score Day 4	1.28 (1.11-1.49)	0.0008
Charlson comorbidity score (CCS) baseline	1.48 (1.22-1.79)	<0.0001
ELISpot IFNγ TE Day 4 (Negative of log)	1.29 (0.90-1.85)	0.165

769 **Figure 1. Flow Diagram for Study Enrollment**

Figure 2. Unstimulated and Stimulated IFNγ Expression as Determined by ELISpot in SEPSIS and CINS Patients, and Healthy Control Subjects on Days 1, 4, and 7 Following ICU Admission. Values represent medians and individual subject responses. Panels A, B, C are from unstimulated whole blood while Panels D, E, F are antiCd3/CD28 stimulated whole blood. Note that the scales for unstimulated expression are logarithmic, whereas they are linear for stimulated expression to appropriately reflect the magnitude and heterogeneity of the individual response. * p< 0.05, ** p<0.01, ***p<0.001, ****p<0.0001 as determined by Kruskal-Wallace ANOVA and post-hoc analyses using the Dunn test. Values are two sided and represent raw p values. SFU, spot forming units. SS, spot size. TE, total IFNγ expression.

Figure 3 – ELISpot Spot Forming Units (SFU) (Panels A, B, C) and Spot Size (SS) (Panels D, E, F) from Unstimulated Whole Blood in the Three Cohorts (Healthy Subjects, SEPSIS and CINS). In unstimulated whole blood, SEPSIS and CINS cohorts demonstrated a consistent increase in the number of cells (SFU) producing IFNγ, when compared to healthy subjects. * p<0.05, *** p<0.001, **** p<0.0001, as determined by Kruskal-Wallace ANOVA and post-hoc analyses using the Dunn test. Values are two sided and represent raw p values. SFU, spot forming units. SS, spot size. TE, total IFNγ expression.

Figure 4. Differential IFNγ Expression by ELISpot in Sepsis Patients Measured at 1, 4 and 7 Days after ICU Admission who Survived or Did Not Survive by 180 days. Panels A, B, C reflect spot number, D, E, F, spot size, and G, H, I, total IFNγ expression. Values represent medians and individual subject responses. The number of subjects declines over time as patients are either discharged from ICU or die. * p< 0.05, ** p<0.01, ***p<0.001, ****p<0.0001 as determined by Kruskal-Wallace ANOVA and post-hoc analyses using the Dunn test. Values are two sided and represent raw p values. SFU, spot forming units. SS, spot size. TE, total IFNγ expression.

Figure 5. Area under the Receiver Operator Curves (AUROC) for Physiologic (SOFA, Charlson comorbidity scores) and Stimulated IFNγ ELISpot Responses in Differentiating In-Hospital and 180-day Mortality. Top Row: Panel A provides SOFA and Charlson Comorbidity Index, and Panel B, selected ELISpot parameters discriminating 180 day mortality. Lower Row: Same breakdowns as for the top row discriminating in-hospital mortality. TE, IFNγ ELISpot total expression; SFU, IFNγ ELISpot spot forming units; SS IFNγ ELISpot spot size

A. Physiologic Metrics – 180 day Mortality

AUC
0.759 SOFA Day 1
0.787 SOFA Day 4
0.824 Charlson Index

B. ELISpot – 180 Day Mortality

AUC
0.730 IFNγ TE Day 1
0.794 IFNγ TE Day 4
0.707 IFNγ SFU Day 1
0.663 IFNγ SS Day 4

C. Physiologic Metrics – In Hospital Mortality

AUC
0.759 SOFA Day 1
0.828 SOFA Day 4
0.762 Charlson Index

D. ELISpot – In Hospital Mortality

AUC
0.695 IFNγ TE Day 1
0.743 IFNγ TE Day 4
0.661 IFNγ SFU Day 1
0.696 IFNγ SS Day 4