A simulation-based approach for estimating the time-dependent reproduction number

from temporally aggregated disease incidence time series data

I Ogi-Gittins[1,2], WS Hart[3], J Song[4], RK Nash[5], J Polonsky[6], A Cori[5], EM Hill[1,2], RN

Thompson[3]*

Affiliations:

[1]Mathematics Institute, University of Warwick, Coventry, CV4 7AL, UK

[2]Zeeman Institute for Systems Biology and Infectious Disease Epidemiology Research

(SBIDER), University of Warwick, Coventry, CV4 7AL, UK

[3]Mathematical Institute, University of Oxford, Oxford, OX2 6GG, UK

[4]Communicable Disease Surveillance Centre, Health Protection Division, Public Health

Wales, Swansea, SA2 8QA, UK

[5]MRC Centre for Global Infectious Disease Analysis, School of Public Health, Imperial

College, London, W2 1PG, UK

[6]Geneva Centre of Humanitarian Studies, University of Geneva, Geneva, 1205, Switzerland

*Correspondence to: robin.thompson@maths.ox.ac.uk

17   **Abstract**

Tracking pathogen transmissibility during infectious disease outbreaks is essential for

assessing the effectiveness of public health measures and planning future control strategies. A

key measure of transmissibility is the time-dependent reproduction number, which has been

estimated in real-time during outbreaks of a range of pathogens from disease incidence time

series data. While commonly used approaches for estimating the time-dependent reproduction

number can be reliable when disease incidence is recorded frequently, such incidence data are

often aggregated temporally (for example, numbers of cases may be reported weekly rather

than daily). As we show, commonly used methods for estimating transmissibility can be

unreliable when the timescale of transmission is shorter than the timescale of data recording.

To address this, here we develop a simulation-based approach involving Approximate

Bayesian Computation for estimating the time-dependent reproduction number from

temporally aggregated disease incidence time series data. We first use a simulated dataset

representative of a situation in which daily disease incidence data are unavailable and only

weekly summary values are reported, demonstrating that our method provides accurate

estimates of the time-dependent reproduction number under those circumstances. We then

apply our method to two previous outbreak datasets consisting of weekly influenza case

numbers from 2019-20 and 2022-23 in Wales (in the United Kingdom). Our simple-to-use

approach allows more accurate estimates of time-dependent reproduction numbers to be

obtained during future infectious disease outbreaks.

Keywords: Mathematical modelling, Infectious disease epidemiology, Reproduction number,

Parameter inference, Serial interval, Approximate Bayesian Computation, EpiEstim,

Influenza, Disease incidence

Introduction

An important challenge for policy makers during infectious disease outbreaks is to devise public health measures that limit transmission without placing an undue burden on the population [1–3]. Central to the decision making process is an ability to monitor changes in pathogen transmissibility in real-time during outbreaks, to determine whether current interventions are sufficient or whether additional restrictions are required.

A widely used measure of transmissibility is the time-dependent reproduction number (R_t) [4–10]. The value of R_t represents the expected number of infections generated by someone infected at time t over the course of their entire infectious period. This quantity changes during an outbreak in response to interventions, variations in host behaviour and depletion of susceptible individuals due to infection-induced immunity. If the value of R_t is (and remains) below one, then the outbreak will decline. On the other hand, if the value of R_t is (and remains) above one, then the outbreak will grow.

Two distinct versions of R_t exist. First, the "instantaneous" reproduction number [4,5,11–13] represents the expected number of infections generated by someone infected at time t over their infectious period if transmission conditions do not change in future (i.e. assuming that the control interventions in place at time t, and any other factors that affect transmission, are not altered after time t). Second, the "case" reproduction number [12,14] is an analogous quantity but accounts for changes in transmissibility that occur after time t (due to, for example, changes in public health policy). Methods exist for estimating each of these versions of R_t [15]. However, here we focus on the instantaneous reproduction number as it is more amenable to analyses conducted in real-time during outbreaks when future changes in pathogen transmissibility are unlikely to be known. We therefore refer to the instantaneous reproduction number as R_t in this article.

A commonly used approach for estimating R_t is the method introduced by Cori *et al.* [4]

(hereafter referred to as the Cori method), implemented in the R software package *EpiEstim*

[16] and the online application *EpiEstim App* [17]. This approach is based on a renewal

equation model of pathogen transmission (see Methods) and involves estimation of R_t from

disease incidence time series and an estimate of the serial interval distribution (the probability

distribution characterising the interval between symptom onset times in infector-infectee

transmission pairs), both on a daily timescale. The Cori method has been extended in a range

of ways following its original development [7], including accounting for imported cases

[5,11,18,19], uncertainty in the serial interval distribution [5], superspreading [20,21],

multiple pathogen variants [22] and unobserved generations of transmission [23].

However, a challenge that besets estimation of R_t using the Cori method is temporal

aggregation of disease incidence time series data [7,24]. For COVID-19, for example, many

public health agencies switched from publishing daily numbers of reported cases to weekly

summaries after the height of the pandemic [25]. Often, disease incidence is reported weekly

even for pathogens including influenza [26] for which realised serial intervals and generation

times are typically only a few days [27–29]. A common workaround when using the Cori

method in these scenarios is to match the timescale of the serial interval distribution to the

timescale of the incidence data; for example, by supplying a weekly serial interval

distribution and applying the Cori method to weekly incidence data. This is problematic not

only because it is hard to unpick within-week changes in pathogen transmissibility when data

are reported weekly, but also because an assumption of the transmission model underlying the

Cori method is that all cases arising at timestep t are generated by infectors from earlier

timesteps. In other words, if the Cori method is applied with a weekly timestep, as considered

in this study, then it is assumed that an infector and infectee cannot both appear as cases in

the disease incidence data in the same week. As the timescale of transmission (as

characterised by the serial interval or generation time) of many pathogens is less than one

week, this assumption is often incorrect when considering weekly aggregated disease

incidence time series data.

In this research article, we address this issue by presenting a novel simulation-based method

for estimating R_t from temporally aggregated disease incidence time series data and the serial

interval distribution. Our approach involves repeated simulation of a renewal equation

transmission model for different values of R_t with a timestep that is smaller than that of the

disease incidence data. Using an iterative version of Approximate Bayesian Computation

(ABC), we show how R_t can be estimated in real-time during outbreaks by matching model

simulations exactly to the temporally aggregated outbreak data. We apply our approach to

simulated data, demonstrating its accuracy and comparing results from our method to those

obtained using the common workaround of the Cori method applied to temporally aggregated

data. We go on to apply our method to real-world outbreak data from the 2019-20 and 2022-

23 influenza seasons in Wales in the United Kingdom.

Methods

In our analyses, we consider two possible approaches for estimating R_t from temporally

aggregated disease incidence time series data: a workaround of the widely used Cori method

(Approach 1 in Fig 1) and our novel simulation-based method (Approach 2 in Fig 1). Since

the simulation-based approach uses a shorter timestep than that of data reporting, this method

accounts for the possibility of multiple generations of transmission occurring between dates

of data reporting. To provide a concrete setting in which to compare the two methods, we

focus on a situation in which disease incidence data are aggregated into weekly timesteps.

Below, we describe how the value of R_t each week can be estimated from the weekly data,

first using the Cori method (with a timestep of one week, since the incidence data are

aggregated into weekly values), and then using our simulation-based approach (using a

timestep shorter than one week, again using the weekly incidence data).

Figure 1. Schematic illustrating the approaches for estimating R_t that we consider. Approach 1 (top)

involves the application of the commonly used Cori method to weekly aggregated disease incidence time series.

Approach 2 (bottom) is the novel simulation-based approach, which involves matching simulations run with a

smaller timestep to the weekly aggregated data to estimate R_t. The second approach relaxes the assumption that

individuals appearing in the incidence data cannot have infected other individuals appearing in the same week.

Relaxing this assumption is particularly important during outbreaks in which the timescale of transmission is

shorter than the temporal aggregation of the data (e.g. if disease incidence time series data are aggregated

weekly, but serial intervals or generation times can be shorter than one week).

<u>The Cori method</u>

Following previous descriptions of the Cori method [4,5,11], we assume that the expected

number of cases, I_t, in week t, is given by

129
$$\mathbb{E}(I_t|\{I_k\}_{k=1}^{t-1}, R_t, \boldsymbol{w}) = R_t \sum_{s=1}^{t-1} w_s I_{t-s}, \quad (1)$$

in which w_s is the probability that the (weekly discretised) serial interval takes the value s

weeks. We use the notation $\boldsymbol{w}$ to denote the sequence of values of w_s ($s = 1,2,\dots$).

The goal of the Cori method is to estimate R_t, assuming that it takes a constant value during

the time period from week $t - \tau$ to week t. In our analyses, we set $\tau = 0$ to obtain an

estimate of R_t each week, but we first present the method for general (non-negative integer

value) τ for consistency with previous presentations of this approach. If the number of cases

in week t is drawn from a Poisson distribution, then the probability of observing weekly

incidence $\{I_k\}_{k=t-\tau}^{t}$ over the time window $[t - \tau, t]$ (which consists of incidence data from

$\tau + 1$ weeks) is

139
$$\mathbb{P}(\{I_k\}_{k=t-\tau}^{t}|\{I_k\}_{k=1}^{t-\tau-1}, R_t, \boldsymbol{w}) = \prod_{k=t-\tau}^{t} \frac{\left(R_t \sum_{s=1}^{k-1} w_s I_{k-s}\right)^{I_k} \exp\left(-R_t \sum_{s=1}^{k-1} w_s I_{k-s}\right)}{I_k!}.$$

140　Assuming that the prior for R_t is a gamma distribution with shape parameter α and rate

141　parameter β, then, by applying Bayes' Theorem, the posterior for R_t is

142
$$p(R_t|\{I_k\}_{k=1}^t, \boldsymbol{w}) = \text{gamma}\left(R_t, \alpha + \sum_{k=0}^{\tau} I_{t-k}, \beta + \sum_{k=0}^{\tau} \sum_{s=1}^{t-k-1} I_{t-k-s} w_s\right),$$

in which we use the notation $p(R_t|\{I_k\}_{k=1}^t, \boldsymbol{w})$ to represent the probability density function of

R_t conditional on past incidence data and the weekly discretised serial interval distribution.

The notation $\text{gamma}(x, a, b)$ represents the probability density of a gamma distribution at

value x with shape parameter a and rate parameter b. In all of our analyses, as in previous

studies [4,5,11], we set $\alpha = 1$ and $\beta = 0.2$. The prior for R_t therefore has mean and standard

deviation equal to five. The large standard deviation is chosen so that the prior is relatively

uninformative. The high mean ensures that the outbreak is not evaluated as being under

control ($R_t < 1$) unless this is very likely to be the case, so that interventions are not relaxed

erroneously.

Throughout the manuscript, we consider estimating individual values of R_t each week, based

on the numbers of new cases observed in that week. In other words, as noted above, we

assume that $\tau = 0$, in which case the above expression simplifies to

155
$$p(R_t|\{I_k\}_{k=1}^t, \boldsymbol{w}) = \text{gamma}\left(R_t, \alpha + I_t, \beta + \sum_{s=1}^{t-1} I_{t-s} w_s\right).$$

The Cori method can therefore be used to obtain a posterior for R_t for $t \geq 2$ weeks.

<u>Simulation-based inference of R_t</u>

In the renewal equation model underlying the Cori method, the number of cases arising in

week t depends on the numbers of cases in previous weeks. Implicit in that approach is an

assumption that individuals appearing in the incidence data in any week cannot generate new

cases in the same week. When disease incidence data are temporally aggregated, so that the

timescale of transmission can be shorter than the timestep in the incidence data, this

assumption may be incorrect. To relax this assumption, we consider a novel simulation-based

approach for estimating R_t. The goal of this method is again to estimate the value of R_t for

each week, $t \geq 2$, but using a renewal equation model with a timestep that is shorter than one

week (e.g., a daily timestep).

Modified renewal equation

In this approach, we consider partitioning the cases in each week into P timesteps, where

each new timestep is $1/P$ weeks. If, for example, $P = 7$, then we are using a daily timestep in

the simulation-based method. We introduce the following notation:

• $I_{P(t-1)+i}^{(P)}$ represents the number of cases in the ith (partitioned) timestep within week t

172           $(i = 1,2,\dots,P)$.

173   • $w_s^{(P)}$ represents the probability that the serial interval, discretised into timesteps of

174   length $1/P$ weeks (see below and Supplementary Material), takes the value s

175   timesteps.

176   • $\boldsymbol{w}^{(P)}$ represents the sequence of values of $w_s^{(P)}$ $(s = 1,2,\dots)$.

177   In forward simulations of the corresponding renewal equation model, we assume that the

178   number of cases in the ith timestep of week t is drawn from a Poisson distribution with mean

179
$$\mathbb{E}\left(I_{P(t-1)+i}^{(P)} \mid \left\{I_k^{(P)}\right\}_{k=1}^{P(t-1)+i-1}, R_t, \boldsymbol{w}^{(P)}\right) = R_t \sum_{s=1}^{P(t-1)+i-1} w_s^{(P)} I_{P(t-1)+i-s}^{(P)},$$

for $i = 1,2,\dots,P$. This is analogous to simulating the renewal equation underlying the Cori

method but with a shorter timestep of $1/P$ weeks (rather than with a timestep of length one

week).

Inference of R_t

Inference of R_t under the simulation-based method involves repeated simulation of the

modified renewal equation model, using an iterative version of ABC. In short, the model is

simulated repeatedly in each week t, with a different value of R_t used in each simulation

(these R_t values are sampled independently from the prior, and incidence data for times

before week t are sampled from matching simulations from earlier weeks). This process is

repeated until a fixed number of simulations (denoted M) have been run in which the

simulated number of cases in week t exactly matches the corresponding number of cases in

the data, I_t. The values of R_t used to generate the matching simulations are then combined

into a posterior estimate for R_t. In all of our analyses using the simulation-based method, a

value of $M = 1000$ was used.

This procedure is repeated iteratively, starting with $t = 2$, then $t = 3$, and so on. Since this

approach only involves obtaining matching simulations for a single week at a time, estimates

of R_t can be obtained relatively quickly (compared to attempting to match an entire

simulation run over multiple weeks to the real-world data, as in standard ABC rejection

sampling [30]). For a more detailed description of the simulation-based inference method,

including an explanation of how cases are distributed between timesteps within the first week

in each simulation, see the Supplementary Material. A schematic explaining the steps

involved in the inference procedure is shown in Fig S1.

Outbreak datasets

We consider three outbreak datasets in our analyses. We first test our approach on a simulated

dataset. The use of simulated data not only enables us to compare estimates of R_t obtained

using the simulation-based approach against analogous estimates using the Cori method, but

it also allows us to verify that the simulation-based approach for estimating R_t generates

accurate estimates in a setting in which we know true value of R_t (i.e., the value used to

generate the simulated dataset). We then go on to compare outputs from the simulation-based

approach and the Cori method using weekly aggregated disease incidence time series for

influenza from 2019-20 and 2022-23 in Wales.

Simulated dataset (Fig 2)

We generated simulated data using the modified renewal equation, using a very small

timestep so that the discretised serial interval is a close approximation to the continuous serial

interval. Specifically, a disease incidence time series was generated starting from one initial

case (in the first timestep) using a timestep of 10 minutes ($P = 24 \times 7 \times 6 = 1{,}008$). To

generate a classic epidemic curve, the simulation was run for 11 weeks with $R_t = 1.5$ for $t \leq$

6 weeks and $R_t = 0.75$ for $t > 6$ weeks.

Influenza in Wales, 2019-20 (Figs 3,4) and 2022-23 (Figs 5,6)

To demonstrate our approach on real-world data, we considered two disease incidence time

series datasets provided by Public Health Wales describing estimated numbers of cases of

influenza-like illness (ILI) in Wales each week. The original data comprised the clinical

consultation rate per 100,000 individuals in sentinel practices in Wales each week [31]. The

total number of weekly cases was then estimated by multiplying each value in the original

data by 31.075 (i.e. scaling these values based on the population size of Wales, which is

3,107,500 [32]). Since the dataset is for ILI, it likely contains some cases that were not

influenza. Nonetheless, these data are sufficient to demonstrate and test the methods that we

present in our study, and so we assume that the datasets are representative of numbers of

influenza cases in Wales. Weekly data were provided from 28 October 2019 to 2 February

2020 (Fig 3A) and 31 October 2022 to 5 February 2023 (Fig 5A). These date ranges each

span 14 weeks with high ILI burden.

<u>Serial interval</u>

Since we analyse influenza outbreak datasets in this study, we assume throughout that the

(continuous) serial interval distribution is a gamma distribution with mean 0.37 weeks (2.6

days) and standard deviation 0.19 weeks (1.3 days) [33]. While this estimate was derived

from household data for pandemic influenza, our focus is on demonstrating the application of

the simulation-based method rather than precise estimation of the serial interval, and we

expect this estimate to be in line with the serial interval for seasonal influenza (i.e., a mean

value of less than one week). Denoting the probability density function of the serial interval

distribution by $g(x)$, then $g(x) = \mathrm{gamma}(x, 4, 10.8)$.

We discretise this distribution into timesteps of length $1/P$ weeks to obtain $\boldsymbol{w}^{(P)}$. To do this,

we adapt the method used by Cori $et\ al.$ [4] in which the serial interval distribution is

discretised into timesteps of length one. Specifically, we set

$$w_k^{(P)} = \int_{(k-1)/P}^{(k+1)/P} g(u)\left(1 - P\left|u - \frac{k}{P}\right|\right) \mathrm{d}u, \qquad \text{for } k = 2,3,4,\ldots$$

as derived in the Supplementary Material. We then choose $w_1^{(P)}$ so that $\boldsymbol{w}^{(P)}$ is a valid

probability distribution (i.e., the sum of the entries of $\boldsymbol{w}^{(P)}$ is one). The rationale for

normalising $\boldsymbol{w}^{(P)}$ in this way is that same-timestep cases (i.e. infectors and infectees

appearing in the same timestep) are not possible in the renewal equation model. Our approach

involves assigning all probability density near zero in $g(x)$ to $w_1^{(P)}$, which is the shortest

possible serial interval in the model.

Results

<u>Simulated dataset</u>

We first considered the simulated disease incidence time series dataset in which the incidence data are aggregated into weekly counts (Fig 2A). Since this dataset was generated using a serial interval for influenza, transmission occurred on a timescale less than one week. The discretised serial interval is shown in Fig 2B, both with a weekly timestep for use with the Cori method ($P = 1$; blue) and with a daily timestep for use with the simulation-based method ($P = 7$; red). Since the renewal equation model underlying both the Cori method and our simulation-based approach does not allow individuals appearing in the incidence data to generate new cases in the same timestep, only the simulation-based approach allows within-week realised serial intervals.

We applied both R_t inference methods to the simulated dataset, finding in this scenario that the simulation-based approach generates more accurate estimates of R_t than the Cori method (Fig 2C). The percentage error in the estimated value of R_t each week using the simulation-based approach with $P = 7$ (compared to value of R_t used to generate the dataset) is shown in Fig 2D.

In addition to our main analysis shown in Fig 2, we also conducted other analyses using the simulated dataset. We demonstrated that when the simulation-based method is applied with $P = 1$, the output matches the results obtained when the Cori method is used to estimate R_t (Fig S2A), as would be expected since the assumptions underlying the two methods are identical in this case. We also considered how R_t estimates obtained using the simulation-based method change when different values of P are chosen (Fig S2B-D), finding that the

method can obtain accurate estimates for relatively small values of P (using a value of $P = 3$

led to similar errors compared to using $P = 7$).

Figure 2. Estimation of R_t from the simulated disease incidence time series dataset. A. The simulated

outbreak dataset (blue bars), generated with $R_t = 1.5$ for $t \leq 6$ weeks and $R_t = 0.75$ for $t > 6$ weeks (black

line). The outbreak was simulated with $P = 1,008$ starting from one initial case in the first timestep, and new

cases were then aggregated into weekly case counts. B. The discretised serial interval, for $P = 1$ (as used with

the Cori method; blue) and $P = 7$ (red). C. Estimates of R_t using the Cori method (blue) and the novel

simulation-based approach (with $P = 7$; red). Blue and red lines are the mean estimates, and the shaded regions

represent the 95% credible intervals. The value of R_t underlying the simulation is shown in black. D. The

percentage error in the mean estimate of R_t each week (relative to the true value of R_t used to generate the

dataset) using the simulation-based method with $P = 7$.

 Influenza in Wales, 2019-20 and 2022-23

We then went on to consider the two Wales influenza outbreak datasets, again using both the

Cori and simulation-based methods to estimate R_t. First, we considered the weekly case

counts from the 2019-20 influenza season (Fig 3A). As with the simulated dataset, the

simulation-based approach led to different estimates of R_t than the Cori method; R_t estimates

obtained using the simulation-based approach were typically lower than those from the Cori

method during November and December 2019 (the simulation-based approach generally led

to R_t estimates between one and two, whereas the Cori method generated estimates above

two in multiple weeks), but then higher than those from the Cori method for most of January

2020 when R_t was estimated to be less that one (Fig 3B). We computed the percentage error

in the R_t estimate each week using the simulation-based method with $P = 7$ (Fig 3C). Since

the true underlying value of R_t was unknown, the percentage error was computed relative to

applying the simulation-based method with a very large value of $P = 168$ (this is

representative of the best possible estimate of R_t obtainable from the weekly incidence data;

using a partitioning value of $P = 168$ returns inferred values of R_t estimated with a one-hour

timestep). We also explored how R_t estimates depend on the value of P that is used (Fig 4).

Estimates obtained using the Cori method and using the simulation-based method with $P = 1$

again matched closely (Fig 4A). We found that a value of $P = 7$ is large enough for accurate

inference of R_t (Fig 4D).

The analyses of the Wales influenza data from 2019-20 were then repeated for the data from

2022-23, with similar results (Figs 5,6). Notably, the Cori method generally led to a higher

estimate of R_t than the simulation-based method when R_t was estimated to be greater than

one, and a lower estimate of R_t than the simulation-based method when R_t was estimated to

be less than one (see Discussion).

Figure 3. Estimation of R_t for influenza in Wales, 2019-2020. A. Weekly numbers of ILI cases in Wales from

28 October 2019 to 2 February 2020, estimated from surveillance data collected in sentinel practices. B.

Estimates of R_t using the Cori method (blue) and the novel simulation-based approach (with $P = 7$; red). Blue

and red lines are the mean estimates, and the shaded regions represent the 95% credible intervals. C. The

percentage error in the mean estimate of R_t each week using the simulation-based method with $P = 7$,

compared to using a larger value of $P = 168$ (which corresponds to estimating R_t with a one-hour timestep).

Figure 4. Dependence of R_t estimates using the simulation-based method on the value of P used, for influenza in Wales, 2019-2020. A. Estimates of R_t obtained when the Cori method (blue) and the novel simulation-based approach with $P = 1$ (red) are applied to the 2019-20 influenza dataset (Fig 3A). B. Analogous to panel A, but with $P = 3$ in the simulation-based approach. C. Analogous to panel A, but with $P = 5$ in the simulation-based approach. D. The average weekly absolute error in mean R_t estimates obtained using the simulation-based method with different values of P, compared to using a larger value of $P = 168$ (which corresponds to estimating R_t with a one-hour timestep). For a given value of P, this measure represents the absolute value of the error in the estimate of R_t in week t (compared to using $P = 168$), averaged over all values of t.

Figure 5. Estimation of R_t for influenza in Wales, 2022-2023. A. Weekly numbers of ILI cases in Wales from

31 October 2022 to 5 February 2023, estimated from surveillance data collected in sentinel practices. B.

Estimates of R_t using the Cori method (blue) and the novel simulation-based approach (with $P = 7$; red). Blue

and red lines are the mean estimates, and the shaded regions represent the 95% credible intervals. C. The

percentage error in the mean estimate of R_t each week using the simulation-based method with $P = 7$,

compared to using a larger value of $P = 168$ (which corresponds to estimating R_t with a one-hour timestep).

Figure 6. Dependence of R_t estimates using the simulation-based method on the value of P used, for

influenza in Wales, 2022-2023. A. Estimates of R_t obtained when the Cori method (blue) and the novel

simulation-based approach with $P = 1$ (red) are applied to the 2022-23 influenza dataset (Fig 5A) B. Analogous

to panel A, but with $P = 3$ in the simulation-based approach. C. Analogous to panel A, but with $P = 5$ in the

simulation-based approach. D. The average weekly absolute error in mean R_t estimates obtained using the

simulation-based method with different values of P, compared to using a larger value of $P = 168$ (which

corresponds to estimating R_t with a one-hour timestep). For a given value of P, this measure represents the

absolute value of the error in the estimate of R_t in week t (compared to using $P = 168$), averaged over all

values of t.

Discussion

During infectious disease outbreaks, evaluation of time-varying changes in pathogen

transmission is essential to inform outbreak responses. Different metrics can be tracked,

including incidence of new cases, hospitalisations and deaths, and outbreak growth rates

[34,35]. A key metric that has been estimated in real-time during outbreaks of a range of

pathogens is R_t, in part because of its straightforward interpretation [7,9,15]. Not only is

there a threshold value of $R_t = 1$, below which an outbreak can be inferred as being under

control, but the value of R_t also provides information about the extent to which the level of

transmission must change (relative to current transmission) for an outbreak to grow or

decline. For example, if $R_t = 2$, then more than half of transmissions must be prevented for

the outbreak to decline. Similarly, if $R_t = 0.5$, then up to twice as many transmissions may

occur before the outbreak begins to grow. Precise estimation of R_t is therefore crucial.

Here, we have presented a novel simulation-based approach for estimating R_t in scenarios in

which disease incidence time series data are aggregated temporally (Fig 1). While

epidemiological data may be collected at a fine temporal resolution, it is common for the data

to then be aggregated (e.g., into weekly or monthly counts). Aggregated data may be easier to

report and can be more accurate than data presented at a high temporal resolution when there

is uncertainty in the precise times at which cases occurred. However, as we have shown,

frequently used methods for inferring R_t, such as the Cori method [4,5], may not generate

accurate estimates when applied to temporally aggregated data if transmission occurs more

rapidly than the temporal resolution of the aggregated data. This is because the renewal

equation model underlying the Cori method involves assuming that an individual appearing

in the disease incidence time series data at timestep t cannot have infected other individuals

appearing in the same timestep. Our proposed simulation-based approach addresses this, by

exactly matching simulations of a renewal equation model run with a shorter timestep (P

timesteps for each timestep in the aggregated data) to the temporally aggregated incidence

data. The simulation-based approach not only provides accurate estimates of R_t (Fig 2), but

can also be applied easily to real-world datasets (Figs 3-6). While using a very large value of

P allows the most accurate possible R_t estimates to be obtained from the aggregated data,

even relatively small values of P are sufficient for R_t to be inferred accurately (Figs 4,6).

We found that, while the Cori method did not always provide an accurate estimate of R_t due

to the temporal aggregation of the disease incidence data, it was able to identify whether or

not R_t is below one (i.e., the outbreak is under control). While this is useful, as noted above

precise estimation of R_t is important as it provides information about the number of

transmissions that must be prevented for an outbreak to be controlled (or the number of

transmissions that can occur for an outbreak to remain under control). The difference in R_t

estimates between the two methods can be explained by the assumption of no same-timestep

cases (i.e., infectors and infectees cannot appear in the disease incidence time series in the

same timestep) in the renewal equation. When the Cori method is applied to weekly data, this

then leads to overestimation of the serial interval, which is known in turn to lead to

overestimation of R_t if the true value of R_t is greater than one and underestimation of R_t if

the true value of R_t is less than one [36,37].

A closely related study by Nash *et al.* [24], undertaken at the same time as the analyses

presented here, has also considered estimation of R_t from temporally aggregated disease

incidence time series data. In that approach, an expectation-maximisation (EM) algorithm is

used to reconstruct daily incidence from any aggregation of disease incidence data using the

serial interval (on a daily timescale). The original version of the Cori method is then applied

to the estimated daily data. This EM approach has been integrated into the EpiEstim R

software package [16]. There are several differences between the approach by Nash *et al.* [24]

and the simulation-based method described here. First, the two approaches are

methodologically distinct, relying on entirely different underlying methods (EM or model

simulation). Second, under the approach by Nash *et al.*, only a single estimated daily disease

incidence time series is obtained. In contrast, our method involves matching a range of

simulations to the temporally aggregated data, thereby considering different possible

disaggregated disease incidence time series that could have led to the weekly aggregated data.

Third, our method can be run straightforwardly for a range of values of P, allowing the most

accurate possible estimates of R_t to be inferred from temporally aggregated incidence data

(discretisation into a timestep of less than one day is straightforward). Fourth, our approach

can be applied easily in scenarios in which the serial interval distribution is discrete rather

than continuous (if, for example, the serial interval distribution is constructed directly from

observations of dates on which infector-infectee pairs report symptoms). A rigorous

comparison of estimated values of R_t obtained using the simulation-based method and using

the EM approach of Nash *et al.* [24] is a target for future exploration. However, an initial

investigation involving applying the methods to the Wales influenza datasets considered here

suggests that the two approaches can obtain consistent results (Fig S3).

Our simulation-based method is conceptually straightforward, simply requiring repeated

simulation of a renewal equation model. It is also computationally efficient to run, as

simulations are only required to match the real-world data for one aggregated timestep at a

time. This contrasts with using ABC rejection sampling to estimate all values of R_t

simultaneously, which would involve matching entire simulated time series to the entire real-

world dataset. The efficiency of our approach allowed us to require that the simulations used

to infer R_t match the real-world data exactly. Further computational efficiency could be

achieved by removing this condition, and instead setting a threshold "distance" within which

a simulation is determined to match the real-world data, as is common when using ABC

[30,38]. However, this necessitates that a distance metric is chosen, and resulting estimates of

R_t may be less accurate.

As in any modelling study, our framework in its current form involves assumptions. We

followed previous publications in which the Cori method has been used [4,5] and assumed

that the time series datasets from which we estimated R_t represent numbers of new

symptomatic cases in each timestep. In the disease incidence time series data, it is then

assumed that each infectee appears after their infector following a time period that reflects a

random draw from the serial interval distribution, which is assumed to always take strictly

positive values. However, in reality, realised serial intervals can be negative (if an infectee

develops symptoms before their infector; this is possible, for example, for transmission of

SARS-CoV-2 [39–42]). Rather than using disease incidence time series, it is possible to apply

both the Cori method and the simulation-based method to data describing incidence of

infections, replacing the serial interval distribution as an input with the distribution of the

generation time (the interval between infection times in infector-infectee pairs). This can be

beneficial as realised generation times are always positive. However, since times of infection

are often unknown, infection incidence data are typically not observed directly. Consequently,

further inference would then be required to estimate incidence of infections, as well as to

estimate the generation time distribution [8,43–45].

In our analyses, we assumed that all cases in the disease incidence time series (after the first

timestep) arose because of transmission within the population under consideration, and that

all cases were recorded. In reality, some infected individuals may become infected outside the

local population [5,11,46,47] and under-reporting of cases is likely for many pathogens [48–

51]. Extension of our method to account for these features of real-world outbreaks is a target

for future research. Similarly, our method assumes that a Poisson distributed number of cases

occur in each timestep of the modified renewal equation model. Considering different

possible probability distributions, including accounting for the possibility of superspreading

events on some days [20,21,52], is another possible area for future work.

Further testing of the performance of the simulation-based approach in different scenarios

would also be worthwhile. For example, in settings in which disease incidence time series are

subject to a "day-of-the-week effect" [53] (e.g., if cases occurring at the weekend are

typically reported with a longer reporting delay than those arising during the week), R_t

inference using the simulation-based method applied to weekly aggregated incidence data

may generate more robust estimates than attempting to infer R_t from less accurate daily

incidence data. Our method can also be adapted for scenarios in which the disease incidence

time series data are aggregated into timesteps that are not all of equal length. For example,

when incidence data are derived from World Health Organization reports that are published

irregularly in time, the timestep changes during the outbreak [54], and those irregular

timesteps can be used directly in our simulation-based method.

In summary, we have presented a novel method for estimating R_t from temporally aggregated

disease incidence time series. Going forwards, the ideal scenario is for disease incidence time

series to be recorded accurately at a fine temporal resolution (e.g., daily). If that occurs, then

existing methods for estimating R_t are generally expected to perform well. However, if

disease incidence time series continue to be aggregated temporally for pathogens for which

transmission occurs on a short timescale, then methods allowing accurate R_t inference from

temporally aggregated data are of paramount importance.

COMPETING INTERESTS

We have no competing interests.

AUTHORS' CONTRIBUTIONS

IOG – formal analysis, investigation, visualisation, validation, writing – original draft, writing – review and editing.

WSH – methodology, writing – review and editing.

JS – methodology, writing – review and editing.

RKN – methodology, writing – review and editing.

JP – methodology, writing – review and editing.

AC – methodology, writing – review and editing.

EMH – methodology, supervision, visualisation, writing – review and editing.

RNT – conceptualization, methodology, project administration, supervision, visualisation, writing – original draft, writing – review and editing.

FUNDING

This research was funded by the EPSRC through the Mathematics for Real-World Systems CDT (ZO-G, RNT; grant number EP/S022244/1) and a doctoral prize (WSH; grant number EP/W524311/1). The collaboration between JS and RNT was funded by a grant from Public Health Wales. EMH and RNT would like to acknowledge the help and support of the JUNIPER partnership, funded by MRC (grant number MR/X018598/1), to which they are linked.

ACKNOWLEDGEMENTS

Thanks to the Communicable Disease Surveillance Centre at Public Health Wales for providing us with the data used in this research. Thanks to members of the Zeeman Institute for Systems Biology and Infectious Disease Epidemiology Research at the University of

Warwick, particularly Alex Kaye, and the Wolfson Centre for Mathematical Biology at the

University of Oxford, for useful discussions about this work.

DATA AVAILABILITY

The computing code used to perform the analyses in this article is available in the following

GitHub repository: www.github.com/billigitt/R_Estim_Simulation_Method. All computer

code was written in the MATLAB programming environment (compatible with version

R2022a).

<u>**References**</u>

1. Hollingsworth TD, Klinkenberg D, Heesterbeek H, Anderson RM. Mitigation strategies
for pandemic influenza A: Balancing conflicting policy objectives. PLoS Comput
Biol. 2011;7: e1001076–e1001076.

2. Smith RD, Keogh-Brown MR, Barnett T, Tait J. The economy-wide impact of pandemic
influenza on the UK: a computable general equilibrium modelling experiment. BMJ.
2009;339: b4571–b4571.

3. Tildesley MJ, Vassall A, Riley S, Jit M, Sandmann F, Hill EM, et al. Optimal health and
economic impact of non-pharmaceutical intervention measures prior and post
vaccination in England: a mathematical modelling study. R Soc Open Sci. 2022;9:
211746.

4. Cori A, Ferguson NM, Fraser C, Cauchemez S. A new framework and software to
estimate time-varying reproduction numbers during epidemics. Am J Epidemiol.
2013;178: 1505–12.

5. Thompson RN, Stockwin JE, van Gaalen RD, Polonsky JA, Kamvar ZN, Demarsh PA, et
al. Improved inference of time-varying reproduction numbers during infectious
disease outbreaks. Epidemics. 2019;29: 100356–100356.

6. Nishiura H, Chowell G. The effective reproduction number as a prelude to statistical
estimation of time-dependent epidemic trends. Math Stat Estim App Epidem. 2009.
pp. 103–121.

7. Nash RK, Nouvellet P, Cori A. Real-time estimation of the epidemic reproduction
number: Scoping review of the applications and challenges. PLoS Digit Health.
2022;1: e0000052.

8. Gostic KM, McGough L, Baskerville E, Abbott S, Joshi K, Tedijanto C, et al. Practical
considerations for measuring the effective reproductive number, Rt. PLoS Comput
Biol. 2020.

9. Vegvari C, Abbott S, Ball F, Brooks-Pollock E, Challen R, Collyer BS, et al. Commentary
on the use of the reproduction number R during the COVID-19 pandemic. Stat Meth
Med Res. 2021;1: 1–11.

10. Thompson RN, Hollingsworth TD, Isham V, Arribas-Bel D, Ashby B, Britton T, et al.
Key questions for modelling COVID-19 exit strategies. Proc Roy Soc B. 2020;287:
20201405–20201405.

11. Creswell R, Augustin D, Bouros I, Farm HJ, Miao S, Ahern A, et al. Heterogeneity in
the onwards transmission risk between local and imported cases affects practical
estimates of the time-dependent reproduction number. Phil Trans R Soc A.
2022;380: 20210308.

12. Fraser C. Estimating individual and household reproduction numbers in an
emerging epidemic. PLoS One. 2007;2: e758.

13. Dai C, Zhou D, Gao B, Wang K. A new method for the joint estimation of
instantaneous reproductive number and serial interval during epidemics. PLoS
Comput Biol. 2023;19: e1011021.

14. Wallinga J, Teunis P. Different epidemic curves for severe acute respiratory
syndrome reveal similar impacts of control measures. Am J Epidemiol. 2004;160:
509–516.

15. White LF, Moser CB, Thompson RN, Pagano M. Statistical estimation of the
reproductive number from case notification data. Am J Epidem. 2020; kwaa211.

16. EpiEstim Team. EpiEstim: Estimate time varying reproduction numbers from
epidemic curves. Version 2.2-4. 2021. Available: www.cran.r-
project.org/web/packages/EpiEstim/

17. EpiEstim App Team. EpiEstim App. 2019. Available:
www.shiny.dide.imperial.ac.uk/epiestim/

18. Li W, Bulekova K, Gregor B, White LF, Kolaczyk ED. Estimation of local time-varying
reproduction numbers in noisy surveillance data. Phil Trans Roy Soc A. 2022.

19. Tsang TK, Wu P, Lau EHY, Cowling BJ. Accounting for imported cases in estimating
the time-varying reproductive number of COVID-19 in Hong Kong. J Infect Dis.
2021;224: 783–787.

20. Johnson KD, Beiglböck M, Eder M, Grass A, Hermisson J, Pammer G, et al. Disease
momentum: Estimating the reproduction number in the presence of
superspreading. Infect Dis Model. 2021;6: 706–728.

21.   Ho F, Parag KV, Adam DC, Lau EHY, Cowling BJ, Tsang TK. Accounting for the
potential of overdispersion in estimation of the time-varying reproduction number.
Epidemiology. 2023;34: 201–205.

22.   Bhatia S, Wardle J, Nash RK, Nouvellet P, Cori A. Extending EpiEstim to estimate the
transmission advantage of pathogen variants in real-time: SARS-CoV-2 as a case-
study. Epidemics. 2023;44: 100692.

23.   Brizzi A, O'Driscoll M, Dorigatti I. Refining reproduction number estimates to
account for unobserved generations of infection in emerging epidemics. Clin Infect
Dis. 2022;75: e114–e121.

24.   Nash RK, Bhatt S, Cori A, Nouvellet P. Estimating the epidemic reproduction
number from temporally aggregated incidence data: a statistical modelling
approach and software tool. PLoS Comput Biol. 2023;19: e1011439.

25.   UK Health Security Agency. The COVID-19 dashboard moves to weekly updates.
2022. Available: www.ukhsa.blog.gov.uk/2022/06/28/the-covid-19-dashboard-
moves-to-weekly-updates/

26.   UK Health Security Agency. National Influenza and COVID-19 surveillance report:
Week 29 report (up to week 28 data). 2023. Available:
www.gov.uk/government/statistics/national-flu-and-covid-19-surveillance-
reports-2022-to-2023-season

27.   Cowling BJ, Fang VJ, Riley S, Malik Peiris JS, Leung GM. Estimation of the serial
interval of influenza. Epidemiology. 2009;20: 344–347.

28.   te Beest DE, Wallinga J, Donker T, Van Boven M. Estimating the generation interval
of Influenza A (H1N1) in a range of social settings. Epidemiology. 2013;24: 244–
250.

29.   Biggerstaff M, Cauchemez S, Reed C, Gambhir M, Finelli L. Estimates of the
reproduction number for seasonal, pandemic, and zoonotic influenza: a systematic
review of the literature. BMC Infect Dis. 2014;14: 480.

30.   Minter A, Retkute R. Approximate Bayesian Computation for infectious disease
modelling. Epidemics. 2019;29: 100368.

31.   Public Health Wales. Weekly influenza and acute respiratory infection surveillance
report: Wednesday 22nd February 2023 (covering week 07 2023). 2023. Available:
www.phw.nhs.wales/topics/immunisation-and-vaccines/fluvaccine/weekly-
influenza-and-acute-respiratory-infection-report/october-2022-october-2023-flu-
season-202223/phw-influenza-surveillance-report-for-2023-week-7pdf/

32.   Office for National Statistics. Population and household estimates, Wales: Census
2021. 2021. Available:
www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/popul
ationestimates/bulletins/populationandhouseholdestimateswales/census2021

33.   Cauchemez S, Donnelly CA, Reed C, Ghani AC, Fraser C, Kent CK, et al. Household
592   transmission of 2009 pandemic Influenza A (H1N1) virus in the United States. N
593   Engl J Med. 2009;361: 2619–2627.

594   34.   Pellis L, Scarabel F, Stage HB, Overton CE, Chappell LH, Fearon E, et al. Challenges in
595   control of Covid-19: short doubling time and long delay to effect of interventions.
596   Phil Trans Roy Soc B. 2021;376: 20200264–20200264.

597   35.   Parag KV, Thompson RN, Donnelly CA. Are epidemic growth rates more informative
598   than reproduction numbers? J R Stat Soc Ser A. 2022;1: 1–11.

599   36.   Wallinga J, Lipsitch M. How generation intervals shape the relationship between
600   growth rates and reproductive numbers. Proc R Soc B Biol Sci. 2007;274: 599–604.

601   37.   Knight J, Mishra S. Estimating effective reproduction number using generation time
602   versus serial interval, with application to COVID-19 in the Greater Toronto Area,
603   Canada. Infect Dis Model. 2020;5: 889–896.

604   38.   Toni T, Welch D, Strelkowa N, Ipsen A, Stumpf MPH. Approximate Bayesian
605   computation scheme for parameter inference and model selection in dynamical
606   systems. J R Soc Interface. 2009;6: 187–202.

607   39.   Du Z, Xu X, Wu Y, Wang L, Cowling BJ, Meyers LA. Serial interval of COVID-19 among
608   publicly reported confirmed cases. Emerg Infect Dis. 2020;26: 1341–1343.

609   40.   Hart WS, Maini PK, Thompson RN. High infectiousness immediately before COVID-
610   19 symptom onset highlights the importance of continued contact tracing. eLife.
611   2021;10: e65534.

612   41.   Geismar C, Nguyen V, Fragaszy E, Shrotri M, Navaratnam AMD, Beale S, et al.
613   Bayesian reconstruction of SARS-CoV-2 transmissions highlights substantial
614   proportion of negative serial intervals. Epidemics. 2023;44: 100713.

615   42.   Madewell ZJ, Yang Y, Longini IM, Halloran ME, Vespignani A, Dean NE. Rapid review
616   and meta-analysis of serial intervals for SARS-CoV-2 Delta and Omicron variants.
617   BMC Infect Dis. 2023;23: 429.

618   43.   Abbott S, Hellewell J, Thompson RN, Sherratt K, Gibbs HP, Bosse NI, et al. Estimating
619   the time-varying reproduction number of SARS-CoV-2 using national and
620   subnational case counts. Wellcome Open Res. 2020;5: 112.

621   44.   Hart WS, Abbott S, Endo A, Hellewell J, Miller E, Andrews N, et al. Inference of the
622   SARS-CoV-2 generation time using UK household data. eLife. 2022;11: e70767.

623   45.   Hart WS, Miller E, Andrews NJ, Waight P, Maini PK, Funk S, et al. Generation time of
624   the alpha and delta SARS-CoV-2 variants: an epidemiological analysis. Lancet Inf
625   Dis. 2022;22: 603–610.

626   46.   Daon Y, Thompson RN, Obolski U. Estimating COVID-19 outbreak risk through air
627   travel. J Travel Med. 2020;27: taaa093.

628   47.   Didelot X, Helekal D, Kendall M, Ribeca P. Distinguishing imported cases from
629   locally acquired cases within a geographically limited genomic sample of an
630   infectious disease. Bioinformatics. 2023;39: btac761.

631   48.   Dalziel BD, Lau MSY, Tiffany A, McClelland A, Zelner J, Bliss JR, et al. Unreported
632   cases in the 2014-2016 Ebola epidemic: Spatiotemporal variation, and implications
633   for estimating transmission. Althouse B, editor. PLoS Negl Trop Dis. 2018;12:
634   e0006161.

635   49.   Albani V, Loria J, Massad E, Zubelli J. COVID-19 underreporting and its impact on
636   vaccination strategies. BMC Infect Dis. 2021;21: 1111.

637   50.   Gibbons CL, Mangen M-JJ, Plass D, Havelaar AH, Brooke RJ, Kramarz P, et al.
638   Measuring underreporting and under-ascertainment in infectious disease datasets:
639   a comparison of methods. BMC Public Health. 2014;14: 147.

640   51.   Lovell-Read FA, Funk S, Obolski U, Donnelly CA, Thompson RN. Interventions
641   targeting non-symptomatic cases can be important to prevent local outbreaks:
642   SARS-CoV-2 as a case study. J R Soc Interface. 2021;18: 20201014.

643   52.   Bradbury NV, Hart WS, Lovell-Read FA, Polonsky JA, Thompson RN. Exact
644   calculation of end-of-outbreak probabilities using contact tracing data. medRxiv.
645   2023.

646   53.   Guzmán-Rincón LM, Hill EM, Dyson L, Tildesley MJ, Keeling MJ. Bayesian estimation
647   of real-time epidemic growth rates using Gaussian processes: local dynamics of
648   SARS-CoV-2 in England. J R Stat Soc C. 2023;00: 1–22.

649   54.   Shaman J, Yang W, Kandula S. Inference and forecast of the current West African
650   Ebola outbreak in Guinea, Sierra Leone and Liberia. PLoS Curr. 2014;1: 6.