

SHARPS INJURIES IN GHANA AND LIBERIA

Sharps Injuries Among Healthcare Workers in Liberia and Ghana: A Cross-Sectional Survey

Laura Jean Ridge^{ab}, PhD, RN, ORCID ID – 0000-0002-6926-2370

John Arko-Mensah^c, PhD

Josh Lambert^a, PhD, ORCID ID, 0000-0002-4513-8156

Lydia Aziato^c, PhD

G Clinton Zeantoe^d, MSN

Henry Duah^a, MPH ORCID ID – 0000-0002-4842-6006

Marjorie McCullagh^b, PhD, ORCID ID – 0000-0002-6504-5176

- a. University of Cincinnati College of Nursing, Cincinnati, OH, USA
- b. University of Michigan School of Nursing, Ann Arbor, MI, USA
- c. University of Ghana, Accra, Ghana
- d. United Methodist University Winifred J Harley School of Nursing, Ganta, Liberia

Corresponding author:

Dr. Laura Jean Ridge

3110 Vine Street

Cincinnati, OH 45221-0038

215.872.2604

ridgelj@ucmail.uc.edu

Abstract

Objective: There is little data on sharps injuries among healthcare workers in West Africa, despite the region's high rate of Hepatitis B and HIV. The purpose of this study is to investigate healthcare workers' history of sharps injury in Liberia and Ghana.

Design: An electronic cross-sectional survey disseminated by local nursing, midwifery, physician assistant, and physician associations.

Setting: Healthcare workers in Liberia and Ghana from February to June 2022.

Participants: Participant were registered nurses, physician assistants, physicians, or midwives, and had been working in a patient care role for at least nine of the previous twelve months.

Methods: A link to the survey was texted to participants through their professional association membership lists, including nursing, midwifery, and physician assistant organizations in both Liberia and Ghana and a physician organization in Ghana only.

Results: 509 participants reported an average of 1.8 injuries per year in Liberia and 1.1 in Ghana ($p < 0.01$). 15.1% of healthcare workers reported three or more injuries in the past year. Liberia had a higher proportion of frequently injured workers ($p < 0.01$). Frequently injured workers were evenly distributed across worker types.

Conclusions: Workers in this region are vulnerable to sharps injuries. A frequently injured subset of workers likely has distinctive risk factors and would benefit from further investigation and intervention.

SHARPS INJURIES IN GHANA AND LIBERIA

Introduction

Sharps injuries are a major occupational hazard among healthcare workers worldwide. It is estimated that these injuries cause 16,000 new Hepatitis C infections, 66,000 new Hepatitis B infections, and 1,000 new HIV infections annually (1). Even though reported sharps injury rates are high in low- and middle-income countries (LMICs), surveillance of injury incidence and reporting is low (2,3).

This is particularly true in West Africa (4), despite the enormous occupational health challenges faced by healthcare workers in that region (5), including the highest rate of HBV (~8%) worldwide (6). It is estimated that mean number of sharps injuries in West Africa per HCW per year is 2.1, resulting in 33,000 exposures to HIV and HCV, and 131,000 exposures to HBV, per year (7). Approximately 41.7% of African healthcare workers reported a sharps injury in the past year, although that data came primarily from other regions in Africa (2).

Liberia, a country of about 5 million people, has about 19,000 healthcare workers, while Ghana, a country of about 31 million people, has about 123,000 healthcare workers (8). Data on sharps injuries in both countries is scant, with only a few facility-specific studies in Ghana (9). One of these studies found that 36.3% of emergency department nurses at a single hospital in Ghana had four or more injuries in the past twelve months, raising the possibility of a larger population of frequently injured workers (10). The purpose of this paper is to investigate the incidence and correlates of sharps injuries among healthcare workers in Liberia and Ghana and to identify subsets or categories of workers with high incidence of injuries.

SHARPS INJURIES IN GHANA AND LIBERIA

Methods

A cross-sectional survey was administered via Qualtrics to healthcare workers in Liberia and Ghana from February to June, 2022. Inclusion criteria included: the participant was a registered nurse, physician assistant, physician, or midwife, and that he or she had been working in a patient care role for at least nine of the previous twelve months. Exclusion criterion was healthcare workers, even from those cadres, who worked primarily as administrators, nursing assistants, or mental health clinicians. A link to the survey was texted to participants through their professional association membership lists including nursing, midwifery, and physician assistant organizations in both Liberia and Ghana, and a physician organization in Ghana only.

The survey instrument was developed by the study team, which included Ghanaian, Liberian, and American researchers with experience in public health and occupational exposures in their respective countries. The survey instrument included questions about demographics and sharps injury experience, both overall and within the past twelve months. A seven-item multiple choice sharps knowledge scale was developed from Liberian and Ghanaian national infection prevention and control policies (11,12). The six-item Safety Culture: Evaluation Survey, developed by the Centers for Disease Control's (CDC) National Institute for Occupational Safety and Health (13), was also included.

The instrument was reviewed for face and content validity by the leadership of participating local health professional associations, and suggested revisions were made by the PI. The survey link and a brief explanation of the project was texted to professional association membership

SHARPS INJURIES IN GHANA AND LIBERIA

lists contact lists no more than three times. Liberian participants received \$3 USD and Ghanaian participants received 15 Cedis via remote scratch card top-up as compensation.

Descriptive and analytic statistical analysis was conducted in JMP 17 (JMP®, Version 17 PRO, SAS Institute Inc., Cary, NC, 1989–2023) by a trained biostatistician. Continuous variables were analyzed using *t*-tests and ANOVA; categorical variables were tested using Pearson chi-squared test. The STROBE checklist for cross-sectional studies was used to guide reporting within the manuscript (Figure 1).

This project received in-country approval from the University of Liberia’s Institutional Review Board (Protocol #21-12-296) and the Ghana Health Services Ethics Research Committee (GHS ERC Number: 018/11/2021). It was deemed exempt by the Institutional Review Board of the University of Michigan and the University of Cincinnati.

SHARPS INJURIES IN GHANA AND LIBERIA

Results

Nine hundred and seventy-one people were screened for participation; 618 met inclusion criteria. Of those 618 participants, 509 recorded the number of injuries they had in the previous 12 months and were included in the analysis (Table 1). Of the participants, 458 were from Ghana and 51 were from Liberia. All 16 regions in Ghana and 11 of Liberia's 15 counties were represented in the analysis. Liberian participants were more likely to be female ($p<.01$) and were older than Ghanaian participants ($p<.01$), although for both groups the largest age group was 31-40 years. Nurses were the largest healthcare worker group represented in both countries.

The majority of participants in both countries had between 4 and 15 years of experience as a healthcare worker, with the plurality of Ghanaian healthcare workers reporting 4 to 8 years of work experience, and the plurality of Liberian healthcare workers reporting 9 to 15 ($p=0.07$). Participants' work settings were fairly evenly spread across different facility types (primary, secondary, and tertiary); in both countries between 38.0-40.3% of participants worked in secondary care facilities. The remainder of participants worked mostly in primary care facilities in Ghana (40.3%) and tertiary care facilities in Liberia (38.0%) ($p=0.02$).

The mean number of injuries in the past twelve months was 1.1, with 1.8 the average in Liberia and 1.1 the average in Ghana ($p<0.01$) (Table 2). The healthcare worker groups with the highest average number of injuries were nurses (1.2) and midwives (1.3), but the difference between the healthcare worker groups was not statistically significant ($p=0.57$).

SHARPS INJURIES IN GHANA AND LIBERIA

Injuries occurred in every unit type with no significant difference between the countries ($p=0.46$) (Table 3). The most frequent units with injuries were: maternity unit (20.7%), the emergency department (20.1%), and operating room (12.8%)/medical-surgical unit (12.8%). The most likely sharp to cause an injury in Ghana was a hollow needle (55.3%), followed by a solid needle (25.3%). In Liberia solid needle was likeliest to cause injury (38%), followed by a hollow needle (30.0%) ($p<0.01$).

Table 4 shows results for frequently injured workers, defined as workers who reported three or more injuries in the past twelve months. Overall, 15.1% of participants were in this category. Liberia had a larger proportion of such workers (29.4%) compared to Ghana (13.5%) ($p=0.01$). Across groups of healthcare workers, physician assistants had the smallest percentage of workers in this category (12.4%) while midwives had the largest (17.0%), however the difference between the two groups was not statistically significant ($p=0.83$). A higher proportion of workers who reported three or more injuries in the last year reported never receiving injury training from infection prevention and control specialists than those with fewer than three injuries ($p=0.01$).

The overall average score on the knowledge scale, out of a possible 10, was 3.63 (Table 5). This average was calculated by adding the number of correct responses. No subtractions were made for incorrect responses. Missing responses were counted as zero. Workers who reported 0 to 2 injuries in the last 12 months had an average score of 3.69, while workers who reported 3 or

SHARPS INJURIES IN GHANA AND LIBERIA

more injuries had an average score of 3.30 ($p=0.45$). Further analysis indicated 54% of nurses, physicians, or physician assistants respondents who completed the scale got a score of zero, while 68% of midwives got a zero ($p=0.09$). More women (60%) than men (53%) got a zero, but this was not significant ($p=0.126$) and may be associated with midwifery, as 95% of midwives are women. Of the respondents who received a zero on the scale, 5% either didn't know what they injured themselves with or injured themselves with another sharp object. For those participants who didn't get a 0, that was only 1%. Removing the workers who received a score of zero on the scale resulted in a median score of 9.

The overall average score on the CDC's Safety Culture test, out of a possible 24, was 21.

Average score on the scale was 21.1 for workers who had zero to 2 injuries in the past year and 20.6 for workers who had 3 or more ($p=0.24$).

SHARPS INJURIES IN GHANA AND LIBERIA

Discussion

This study is the first nationwide study of sharps injury incidence among healthcare workers in either Liberia or Ghana. It indicates that sharps injuries are a major occupational risk for the 142,000 healthcare workers working in those countries, with the average worker experiencing between one and two injuries per year. This is slightly lower than earlier estimates of 2.1 among healthcare workers in West Africa (7), but consistent with the estimate of 41.7% workers injured per year (2), and still indicates that these injuries are prevalent. Different worker groups reported similar frequency of injury; however, the frequency of injuries was highest in Liberia, and 66.4% of participants' most recent injuries took place in four types of units: maternity, emergency department, operating room, and medical-surgical.

Within every healthcare worker group, between 12.4% (physician assistants) and 17.0% (midwives) of workers reported 3 or more injuries in the last year; however, the difference in percentage across groups is not statistically significant. The relatively even distribution of this frequently injured group across healthcare worker types is interesting and suggests a root vulnerability not captured in these data. A larger proportion of Liberians were a part of this frequently injured worker group.

This frequently injured population is slightly smaller than the one suggested in a previous study in Ghana (10), but significant in terms of the number of sharps injuries they experience.

Further, participants in the study came solely from an emergency department (10), which we also identified as a high-risk setting. The fact that a larger percentage of these workers reported

SHARPS INJURIES IN GHANA AND LIBERIA

never receiving sharps training from IPC Focal People than those with fewer than three injuries suggests a training intervention may be useful to this subgroup, but further investigation into whether this subgroup has modifiable work processes or environments that increase their risk should come first. Regardless, targeting this group for intervention is promising, since reducing this group's frequency of injury would have meaningfully lower frequency overall.

The results of the knowledge scale were intriguing. Most workers who attempted the scale received scores of zero, but among workers who did not receive a score of zero, the median knowledge score was high (9 of a possible 10). This suggests that a large proportion of healthcare workers have low sharps safety knowledge, but it is not clear what differentiates low-knowledge workers from those who scored high, as analyses on the two groups typically did not find significant differences. Midwives, the professional group with the highest average number of injuries (1.3), also had the highest percentage of participants who attempted the scale and scored zero (68%). However, neither the average number of injuries nor the rate of zero scores were significantly different for midwives than other groups.

These findings do not seem to suggest an association between low knowledge and frequent injuries, which is consistent with previous studies on occupational exposure to biological hazards in low- and middle-income countries (14). The average knowledge score of the frequently injured group barely differed from the average score of the zero to two injuries group, and the difference was not statistically significant. That said, the relative paucity of training among frequently injured workers suggests that offering training opportunities may

SHARPS INJURIES IN GHANA AND LIBERIA

play a valuable role beyond knowledge building, such as giving workers a chance to practice skills that require dexterity.

Overall, our findings indicate that a better understanding of the vulnerability of frequently injured workers, as associated with their work processes or environments, has the potential to be useful in lowering sharps injuries overall. A better understanding of the >50% of healthcare workers who scored zero on our knowledge scale would also be interesting, but since the association between low knowledge and injury is weak, that is a less promising avenue to approach injury reduction. The limitations of this study is the cross-sectional nature of this data limits our ability to establish causality. We cannot calculate the survey's response rate because while the professional associations did have approximate membership numbers, they had not updated their contact information lists in some time, and it is likely some of the surveys were sent to wrong numbers. We also acknowledge the possibility that workers with recent or frequent sharps injuries were more likely to respond to this survey than workers generally. Both limitations affect generalizability.

Funding Sources for ALL Authors: This study was funded by the University of Michigan's Center of Occupational and Safety Engineering. The PI was also funded by as a T32 fellow in Complexity: Innovations in Promoting Health and Safety (CIPHS), National Institute of Nursing Research (T32 NR016914, PI: Titler) at the time of data collection. This study did not obtain open access funding from either source.

Conflict of Interest: NONE DECLARED

Acknowledgments: The authors would like to acknowledge the work of Mr. Anthony Arkoh and the Graduate Physician Assistants Association of Ghana, Ms. Frederica Hanson and the Ghana Registered Midwives Association, Dr. Kofi Mensah Boateng and the Junior Doctors' Association of Ghana, Mr. Benjamin Suamey and the Liberian Nurses Association, Ms. Shirley Fahnbulleh and the West African College of Nurses, Ms. Wilhelmina Flomo and the Liberian Midwives Association, and Mr. Theophilus Fayiah and the Physician Assistant Association of Liberia.

SHARPS INJURIES IN GHANA AND LIBERIA

Table 1: Participants

	Overall		Ghana		Liberia		p
n	509		458		51		
	N	%	N	%	N	%	
Sex	504		455		49		0.0005
Female	252	50.0%	216	47.5%	36	73.5%	
Male	252	50.0%	239	52.5%	13	26.5%	
Missing	5	1.0%	3	0.7%	2	3.9%	
	N	%	N	%	N	%	
How old are you?	506		456		50		<0.0001
18-30	128	25.3%	124	27.1%	4	8%	
31-40	329	65.0%	297	43.2%	32	64%	
41-50	42	8.3%	28	6%	14	28%	
51-60	4	0.8%	4	1%	0	0%	
>60	3	0.6%	3	1%	0	0%	
Missing	3	0.6%	2	0.4%	1	2.0%	
	N	%	N	%	N	%	
How long have you worked as a healthcare worker?	507		457		50		0.0713
Less than one year	13	2.6%	13	2.8%	0	0.0%	
1-3 years	97	19.1%	93	20.4%	4	8.0%	
4-8 years	173	34.1%	157	34.4%	16	32.0%	
9-15 years	179	35.3%	155	33.9%	24	48.0%	
16-25 years	37	7.3%	32	7.0%	5	10.0%	
>25 years	8	1.6%	7	1.5%	1	2.0%	
Missing	2	0.4%	1	0.2%	1	2.0%	
	N	%	N	%	N	%	
What is your current position in healthcare?	504		456		48		<0.0001
Nurse	193	38.3%	165	36.2%	28	58.3%	
Physician	114	22.6%	114	25.0%	0	0.0%	
Physician Assistant	97	19.2%	89	19.5%	8	16.7%	
Midwife	100	19.8%	88	19.3%	12	25.0%	
Missing	5	1.0%	2	0.4%	3	5.9%	
	N	%	N	%	N	%	
Practice Setting	507		457		50		0.0192

SHARPS INJURIES IN GHANA AND LIBERIA

Primary care	187	36.9%	175	38.3%	12	24.0%	
Secondary care	203	40.0%	184	40.3%	19	38.0%	
Tertiary care	117	23.1%	98	21.4%	19	38.0%	
Missing	2	0.4%	1	0.2%	1	2.0%	

Table 2: Injuries by Country and Healthcare Worker Group

	n	mean	median	p
Location				0.0002
All	509	1.1	1.0	
Liberia	51	1.8	1.0	
Ghana	458	1.1	1.0	
Position in Healthcare				0.5685
Nurse	193	1.2	1.0	
Physician Assistant	97	1.0	1.0	
Midwife	100	1.3	1.0	
Physician	114	1.1	1.0	

SHARPS INJURIES IN GHANA AND LIBERIA

Table 3. Location and Setting of Most Recent Injury

	All		Ghana		Liberia		
	N	%	N	%	N	%	
In which department did the needlestick occur? - Selected Choice	492		442		50		0.4566
Intensive care	13	2.6%	11	2.5%	2	4.0%	
Maternity	102	20.7%	91	20.6%	11	22.0%	
Pediatrics	49	10.0%	43	9.7%	6	12.0%	
Medical surgical	63	12.8%	60	13.6%	3	6.0%	
Injection room	39	7.9%	35	7.9%	4	8.0%	
Operating room	63	12.8%	60	13.6%	3	6.0%	
Psychiatric Unit	5	1.0%	5	1.1%	0	0.0%	
Dental Unit	4	0.8%	4	0.9%	0	0.0%	
Emergency department	99	20.1%	85	19.2%	14	28.0%	
Rotation	13	2.6%	10	2.3%	3	6.0%	
Other	42	8.5%	38	8.6%	4	8.0%	
Missing	17	3.3%	16	3.5%	1	2.0%	
	N	%	N	%	N	%	
What did you injure yourself with?	493		443		50		<0.0001
Hollow needle	260	52.7%	245	55.3%	15	30.0%	
Solid bore needle	131	26.6%	112	25.3%	19	38.0%	
Scalpel	26	5.3%	18	4.1%	8	16.0%	
Don't know	17	3.4%	13	2.9%	4	8.0%	
Another sharp object	59	12.0%	55	12.4%	4	8.0%	
Missing	16	3.1%	15	3.3%	1	2.0%	

SHARPS INJURIES IN GHANA AND LIBERIA

Table 4: Frequently Injured Workers

Injuries in the last 12 months	Location				Position in Healthcare					P
	Ghana % (n)	Liberia % (n)	All % (n)	P	RN % (n)	MD % (n)	PA % (n)	RM % (n)	All % (n)	
0-2	86.5 (396)	70.6 (36)	84.9 (432)	0.0027	84.5 (163)	84.2 (96)	87.5 (85)	83.0 (83)	84.7 (427)	0.8264
>=3	13.5 (62)	29.4 (15)	15.1 (77)		15.5 (30)	15.8 (18)	12.4 (12)	17.0 (17)	15.3 (77)	
All	100.0 (458)	100.0 (51)	100.0 (509)		100 (193)	100 (114)	100 (97)	100 (100)	100 (504)	

Table 5: Knowledge and Safety Culture Scale outcomes

	Knowledge Scale Scores (mean)		Safety Culture Scales Scores (mean)	
Overall?	3.63		21.03	
0-2 injuries	3.69	P=0.451	21.11	P=0.2442
>=3 injuries	3.30		20.55	

SHARPS INJURIES IN GHANA AND LIBERIA

References

1. Prüss-Ustün, A., Rapiti, E., & Hutin, Y. Estimation of the global burden of disease attributable to contaminated sharps injuries among health-care workers. *American Journal of Industrial Medicine* 2005; 48(6), 482-490.
2. Bouya, S., Balouchi, A., Rafiemanesh, H., et al. Global prevalence and device related causes of needle stick injuries among health care workers: A systematic review and meta-analysis. *Annals of Global Health* 2020; 86(1), 35. <https://doi.org/10.5334/aogh.2698>
3. Mossburg, S., Agore, A., Nkimbeng, M., et al. Occupational hazards among healthcare workers in Africa: A systematic review. *Annals of Global Health* 2019; 85(1), 78. <https://doi.org/10.5334/aogh.2434>
4. Groneberg, D. A., Braumann, H., Rolle, S., et al. Needlestick injuries: A density-equalizing mapping and socioeconomic analysis of the global research. *International Archives of Occupational and Environmental Health* 2020; 93(8), 995–1006. <https://doi.org/10.1007/s00420-020-01547-0>
5. Centers for Disease Control and Prevention (CDC). (2017). *Lifesaving Training Protects West Africa's Healthcare Workers and Patients* <https://www.cdc.gov/globalhealth/security/stories/lifesaving-training.html>. published 2017. Accessed June 20, 2023.
6. Assih, M., Ouattara, A. K., Diarra, B., et al. Genetic diversity of hepatitis viruses in West-African countries from 1996 to 2018. *World Journal of Hepatology* 2018; 10(11), 807–821. <https://doi.org/10.4254/wjh.v10.i11.807>

SHARPS INJURIES IN GHANA AND LIBERIA

7. Prüss-Üstün, A., Rapiti, E. & Hutin, Y. (2003). Sharps injuries: Global burden of disease from sharps injuries to health-care workers / <https://apps.who.int/iris/handle/10665/42743>
8. World Bank. (2021). *World Bank Country and Lending Groups*.
<https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups>
9. Ridge, L. J., Dickson, V. V., & Stimpfel, A. W. The occupational health of nurses in the Economic Community of West African States: A review of the literature. *Workplace Health & Safety* 2019; 67(11), 554–564. <https://doi.org/10.1177/2165079919859383>
10. Lori, J. R., McCullagh, M. C., Krueger, A. et al. Sharps injuries among emergency department nurses in one tertiary care hospital in Ghana. *International Emergency Nursing*, 2016; 28, 14–19. <https://doi.org/10.1016/j.ienj.2015.11.007>
11. Ministry of Health, Ghana. (2015). *National Policy and Guidelines for Infection Prevention and Control in Health Care*.
https://www.ghanahealthservice.org/downloads/National_Policy_and_Guidelines%20for_Infection_Prevention_and_Control_in_Health_Care_Settings_2015.pdf published 2015. Accessed July 15, 2023.
12. Ministry of Health, Republic of Liberia (2018). National Infection Prevention and Control Guidelines. Retrieved from https://www.nphil.gov.lr/wp-content/uploads/2022/10/MOH_National-IPC-Guidelines_October-2018.pdf published 2018. Accessed July 10, 2023.

SHARPS INJURIES IN GHANA AND LIBERIA

13. Centers for Disease Control and Prevention (CDC). (2010). *Safety Culture: Evaluation Survey*.

<https://www.cdc.gov/nora/councils/hcsa/stopsticks/survey.html> published 2010

[Accessed July 10](#), 2023

14. Rai, R., El-Zaemey, S., Dorji, N. et al. Exposure to occupational hazards among health care workers in low- and middle-income countries: A scoping review. *International Journal of Environmental Research and Public Health* 2021; 18(5).