1 Targeted malaria elimination interventions reduce *Plasmodium falciparum*
2 infections up to 3 kilometers away
3

4 Jade Benjamin-Chung*[1,2], Haodong Li,[3] Anna Nguyen[1], Gabriella Barratt Heitmann[1], Adam
5 Bennett[4,5], Henry Ntuku,[4] Lisa M. Prach,[4] Munyaradzi Tambo,[6] Lindsey Wu[7], Chris Drakeley[7],
6 Roly Gosling[4,8], Davis Mumbengegwi[9], Immo Kleinschmidt[10,11,12], Alan Hubbard[3], Mark van der
7 Laan[3], Michelle S. Hsiang[2,4,13]
8
[1] Department of Epidemiology and Population Health, Stanford University, Stanford, United States
[2] Chan Zuckerberg Biohub, San Francisco, United States
[3] Division of Biostatistics, University of California, Berkeley
[4] Malaria Elimination Initiative, Global Health Group, University of California, San Francisco (UCSF) , San Francisco, United States
[5] PATH, Seattle, United States
[6] Multidisciplinary Research Centre, University of Namibia, Windhoek, Namibia
[7] Department of Infection Biology, London School of Hygiene and Tropical Medicine, London, UK
[8] Department of Disease Control, London School of Hygiene and Tropical Medicine, London, UK
[9] University of Namibia, Windhoek, Namibia
[10] MRC International Statistics and Epidemiology Group, Department of Infectious Disease Epidemiology, London School of Hygiene and Tropical
Medicine, London, UK
[11] Wits Research Institute for Malaria, Wits/SAMRC Collaborating Centre for Multi-Disciplinary Research on Malaria, School of Pathology,
Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa
[12] Southern African Development Community Malaria Elimination Eight Secretariat, Windhoek, Namibia
[13] Department of Epidemiology and Biostatistics, UCSF, San Francisco, United States
24
25 **Corresponding author:** Jade Benjamin-Chung
26 **Email:** jadebc@stanford.edu

Abstract

29
Malaria elimination interventions in low-transmission settings aim to extinguish hot spots and
prevent transmission to nearby areas. In malaria elimination settings, the World Health
Organization recommends reactive, focal interventions targeted to the area near malaria cases
shortly after they are detected. A key question is whether these interventions reduce
transmission to nearby uninfected or asymptomatic individuals who did not receive
interventions. Here, we measured direct effects (among intervention recipients) and spillover
effects (among non-recipients) of reactive, focal interventions delivered within 500m of
confirmed malaria index cases in a cluster-randomized trial in Namibia. The trial delivered
malaria chemoprevention (artemether lumefantrine) and vector control (indoor residual
spraying with Actellic) separately and in combination using a factorial design. We compared
incidence, infection prevalence, and seroprevalence between study arms among intervention
recipients (direct effects) and non-recipients (spillover effects) up to 3 km away from index
cases. We calculated incremental cost-effectiveness ratios accounting for spillover effects. The
combined chemoprevention and vector control intervention produced direct effects and
spillover effects. In the primary analysis among non-recipients within 1 km from index cases,
the combined intervention reduced malaria incidence by 43% (95% CI 20%, 59%). In secondary
analyses among non-recipients 500m-3 km from interventions, the combined intervention
reduced infection by 79% (6%, 95%) and seroprevalence 34% (20%, 45%). Accounting for
spillover effects increased the cost-effectiveness of the combined intervention by 37%. Our
findings provide the first evidence that targeting hot spots with combined chemoprevention
and vector control interventions can indirectly benefit non-recipients up to 3 km away.

Significance Statement

In settings where malaria transmission is declining and approaching elimination, new malaria cases are clustered in space and time. Prior studies have found that targeting prophylactic antimalarial drugs and vector control in the area around newly detected malaria cases reduced community-wide malaria. Here, we found that when antimalarials and vector control were delivered as a combined strategy in the area near recent cases, malaria incidence was reduced up to 3 kilometers away among individuals who did not receive interventions. Accounting for these benefits to non-recipients increased cost-effectiveness of the intervention. Overall, our findings suggest that combined, targeted malaria interventions can reduce local transmission and support their use for malaria elimination.

Introduction

In the past decade, there has been renewed attention towards global malaria eradication, and many countries have set targets for the elimination of local malaria transmission (1). In Southern Africa, eight countries hope to achieve malaria elimination by 2030 as part of the Elimination Eight Initiative (E8). Yet, global progress has plateaued: annual global malaria cases have increased since 2015, and in 2021, there were 59 cases per 1,000 population at risk, an increase from 57 per 1,000 in 2019 (2).

The ideal malaria elimination intervention would not only prevent disease among recipients but would also prevent onward transmission to nearby non-recipients through spillover effects (i.e., "herd effects", "indirect effects") (4, 5), like some vaccines do (4, 6–12). Prior studies have reported spillover effects for mass drug administration for trachoma (13, 14), school-based deworming (15), insecticide treated bed nets (16–18), and chemoprevention and vector control for malaria (19, 20).

When an intervention reduces disease among intervention non-recipients, accounting for spillover effects can substantially increase cost-effectiveness (21, 22). Identifying cost-effective interventions is crucial to the elimination and eradication enterprise because elimination efforts are projected to cost significantly more than existing malaria control programs in the medium term (3). Even after elimination, countries must continue to conduct intensive surveillance and outbreak response for imported cases to prevent re-establishment.

In settings approaching malaria elimination, the World Health Organization recommends interventions that are "reactive" – delivered soon after a confirmed malaria case is detected – and "focal" – delivered to higher risk individuals who reside near the case (23). A recent cluster-randomized trial in Namibia found that reactive, focal chemoprevention and vector control substantially reduced malaria incidence (24). Spillover effects of these interventions are plausible: chemoprevention may reduce gametocyte biomass in recipients (25), and vector control can reduce the mosquito population near malaria cases. To shed light on whether focal interventions reduce transmission to nearby uninfected or asymptomatic individuals who did not receive interventions, we separately estimated direct effects among intervention recipients and spillover effects among nearby non-recipients. Our approach can be used to estimate spillover effects of other interventions, such as malaria vaccines.

Results

Interventions

We analyzed data from a previously reported cluster-randomized trial of focal malaria interventions conducted in Zambezi region of Namibia in 2017 (NCT02610400) (24) (Figure 1). The region has low *Plasmodium falciparum* malaria transmission (26). Using a two-by-two factorial design, the trial randomized 56 clusters to four arms: 1) reactive case detection (RACD), 2) reactive focal mass drug administration (rfMDA) only, 3) reactive vector control (RAVC) + RACD, 4) RAVC + rfMDA. rfMDA included presumptive treatment with artemether-

106 lumefrantrine to individuals in target areas (Table S1). RACD included testing with rapid
107 diagnostic tests and treatment with artemether-lumefrantrine and single-dose primaquine for
108 those who tested positive. RAVC included indoor residual spraying (IRS) with pirimiphosmethyl.
109 The trial delivered interventions in "target areas" within approximately 500 m of confirmed
110 malaria cases detected through passive surveillance.
111
<u>Effects on malaria incidence</u>
Primary analyses estimated effects on malaria incidence. We estimated three types of effects
on the cumulative incidence of locally acquired, confirmed *Pf* malaria: 1) direct effects among
intervention recipients in target areas within 500 m of confirmed malaria index cases, 2)
spillover effects among non-recipients within 1 km of index cases, and 3) total effects among all
individuals within 1 km of index cases (Figure 2a). We created analytic cohorts including
individuals residing within 1 km of each index case to capture the area and time period in which
we expected each intervention to reduce infections (direct effect) and secondary transmission
to nearby individuals (spillover effect). (Figure 1; Figure S1 see details in Materials and
Methods). We measured effects of the chemoprevention intervention comparing arms with
rfMDA vs. RACD, the vector control intervention by comparing arms with RAVC vs. no RAVC,
and the combined intervention by comparing the rfMDA + RAVC vs. RACD only arms.
124
To estimate direct effects and spillover effects, we used hierarchical targeted maximum
likelihood estimation, a doubly-robust, semiparametric method that adjusts for potential
confounders using ensemble machine learning (27). This approach is appropriate for cluster-
level interventions that may result in statistical dependence between outcomes (see Methods)
(28–30). We adjusted for covariates such as baseline malaria incidence and population size to
account for differences in baseline characteristics between study arms (Tables S2-3).
131
132 In analyses of direct effects among intervention recipients, the chemoprevention intervention
133 reduced malaria incidence among intervention recipients within 500m of index cases, but the
134 confidence interval included the null. There was no evidence of a direct effect for the vector
135 control or combined interventions, but precision was low (Figure 2, Table S4).
136
137 In analyses of spillover effects among intervention non-recipients up to 1 km away from
138 interventions, chemoprevention and vector control interventions reduced incidence, but
139 confidence intervals included the null. There was evidence of a spillover effect of the combined
140 intervention, which reduced malaria incidence by 43% (95% CI 21%, 58%) among non-
141 recipients.
142
We evaluated spillover effect heterogeneity by cluster-level malaria incidence and IRS coverage
prior to the trial, surface temperature, rainfall, the enhanced vegetative index, elevation, and
cohort-level treatment coverage, and gender. Across interventions, spillover effects were
consistently more protective when pre-trial incidence was below the median (Figure S2). For
example, the combined intervention reduced incidence by 68% (95% CI 35%, 84%) when
baseline incidence was lower, but there was no effect when baseline incidence was above the
median. Intervention spillover effects were generally stronger when environmental conditions

favored mosquito breeding and survival (higher rainfall, higher enhanced vegetative index, and
lower elevation) (Table S5). Spillover effects of the chemoprevention intervention were present
for men but were null for women.
153
We performed several sensitivity analyses. When we repeated spillover effect analyses using 2
and 3 km radii around index cases to capture mosquito dispersal over longer distances (31, 32),
results were similar (Figure S3). When using a shorter follow-up period in which intervention
effects may have been stronger (see Methods), results were similar for the chemoprevention
and vector control interventions; for the combined intervention, the spillover effect estimate
was closer to the null, and precision was lower (Figure S4). When we estimated direct effects
including the <3% of intervention recipients who resided >500m from index cases, results were
nearly the same (Figure S5).
162
163 Overlap in analytic cohorts may have resulted in statistical dependence between outcomes
164 (Figure 1; Table S6). When using alternative standard errors accounting for dependence (see
165 Methods), confidence intervals were wider, but there was still evidence of spillover effects and
166 total effects for the combined interventions (Table S4). When we excluded overlapping cohorts
167 from the analysis, results were similar overall (Figure S4).
168
<u>Effects on malaria prevalence</u>
We also estimated effects on malaria prevalence measured using qPCR in a cross-sectional
survey at the end of the malaria season (May to August 2017). In contrast to incidence analyses,
which captured any effects within the period immediately after intervention, prevalence
analyses captured effects of cumulative interventions near the end of the trial. Prevalence also
captures symptomatic and asymptomatic malaria cases that did not necessarily present at
health clinics. Analyses of direct effects included individuals who resided within 500 m any
intervention recipients; spillover effects included individuals with no intervention recipients <
500 m and at least one recipient within 500 m-3 km; total effects included individuals with at
least one intervention recipient within 3 km (Figure 2b). We estimated prevalence ratios using
targeted maximum likelihood estimation and adjusted standard errors for enumeration area-
level clustering.
181
There was evidence of direct effects for all interventions, but confidence intervals included the
null (Figure 2, Table S7). There was evidence of spillover effects: among non-recipients near
intervention recipients, the chemoprevention intervention reduced prevalence by 72% (95% CI
31%, 88%), and the combined intervention reduced it by 79% (95% CI 6%, 95%). For the
chemoprevention and combined interventions, spillover effects decreased in magnitude as
distance to the nearest intervention increased (Figure 3). There was also evidence of spillover
effects on prevalence of households with multiple malaria cases for the chemoprevention and
combined interventions (Table S8).
190
<u>Effects on malaria seroprevalence</u>
We also investigated whether there were effects on seroprevalence of early transcribed
membrane protein 5 antigen (Etramp5.Ag1), an indicator of recent malaria infection (33) that

was measured by Luminex in the cross-sectional survey. The chemoprevention intervention
reduced seroprevalence among individuals who resided within 500m of intervention recipients
by 25% (95% CI 14%, 34%), and the combined intervention reduced it by 34% (95% CI 10%,
42%) (Figure 2, Table S9). Among intervention non-recipients, the combined intervention
reduced seroprevalence by 34% (95% CI 20%, 45%).
199
Cost-effectiveness
To inform policy decisions, we assessed cost-effectiveness using estimates of direct effects and
spillover effects on prevalence. We calculated the incremental cost effectiveness ratio (ICER) by
dividing the difference in cost between arms by the difference in prevalent cases averted
between arms. We included cases averted for both individuals within 500m of any
interventions and those with no intervention recipients < 500 m (direct effect population) and
at least one recipient within 500 m-3 km (spillover effect population). Accounting for direct
effects and spillover effects, the incremental cost-effectiveness ratios were $156 (95% CI $141,
$177), $2,105 (95% CI $1,859, $2,430), and $1,142 (95% CI $944, $1,446) for the
chemoprevention, vector control, and combined interventions (Table S10). Compared to the
trial's original incremental cost-effectiveness ratios estimates, accounting for spillover effects
increased cost-effectiveness by 3%, 21%, and 37% for the chemoprevention, vector control, and
combined interventions (34).
213
214   **Discussion**
215
Here, we showed that a combined intervention of reactive focal chemoprevention plus IRS
reduced malaria infections in intervention recipients as well as non-recipients up to 3 km away.
Overall, spillover effects among non-recipients were strongest for the combined intervention,
which was designed to reduce the parasite reservoir in both humans and mosquitoes. When
accounting for spillover effects, the cost-effectiveness of the combined intervention was 37%
higher than the prior estimate (34).
222
Interventions that produce spillover effects yield greater population health benefits at no
additional cost. A prior analysis found that the combined intervention was highly cost-effective,
but it did not account for possible spillover effects (34). When accounting for spillover effects,
interventions were 3-37% more cost-effective (34). Given that malaria elimination requires
substantially larger investments than malaria control (3, 35), evidence about cost savings due to
spillover effects is critical to policy decisions about elimination strategies.
229
We found stronger evidence of spillover effects of the combined chemoprevention and vector
control intervention over larger spatial scales than two prior studies of targeted malaria
interventions. In Kenya, a trial in a low-transmission area found no change in parasite
prevalence within 500m of serologically-defined hot spots that received targeted larviciding,
long-lasting insecticide treated nets, IRS, and focal mass drug administration (19). In Zambia, an
observational study in a high transmission setting found that IRS targeted to subdistricts with
higher malaria incidence and population density reduced parasite prevalence in sprayed and
unsprayed households within target areas; it did not measure spillover effects outside of target

areas (20). The interventions in our study may have been more likely to produce spillover
effects because they were delivered repeatedly in response to subsequent index cases. In this
trial, clusters received interventions up to 17 times per cluster; they were repeated annually
over three years in the Zambia study and once in the Kenya trial. Further, it is possible that
delivering interventions in response to new index cases can more effectively reduce
transmission than targeting interventions based on an area's incidence or seroprevalence.
244
For the chemoprevention intervention, there was evidence of spillover effects on prevalence
and suggestive evidence of spillover effects and direct effects on incidence. Incidence analyses
measured effects shortly after interventions, while prevalence analyses measured them at the
end of the transmission season. Thus, our findings may indicate that reductions in local
transmission accrued as additional rounds of chemoprevention interventions were delivered
and population intervention coverage increased. This may especially be the case for the
chemoprevention intervention since reductions in infectiousness of malaria cases are typically
short-lived following treatment, especially in the absence of concurrent vector control (36).
Overall, these findings suggest that reactive, focal chemoprevention can more effectively
reduce asymptomatic or subclinical infections among nearby non-intervention recipients than
RACD, particularly after repeated rounds.
256
For the vector control intervention, the primary analysis did not find direct effects, and the
spillover effect estimate included the null despite strong biologic plausibility for both types of
effects. We used a 6-month follow-up period to capture longer-term effects of IRS, which
resulted in spatiotemporal overlap between analytic cohorts (Table S6). This overlap may have
induced dependence between outcomes that was not fully accounted for by covariate
adjustment, resulting in residual bias (28). Analyses of current infection prevalence were not
subject to concerns about cohort overlap and were suggestive of direct effects, but confidence
intervals included the null, and there was no evidence of spillover effects. Finally, our pre-
specified subgroup analyses suggested that spillover effects of RAVC were driven by baseline
transmission levels and environmental conditions: spillover effects on incidence were present
in areas with baseline malaria incidence was <14 per 1,000 and when weather conditions
favored mosquito breeding and survival (temperature < 31 °C; monthly rainfall < 27 mm).
269
The combined intervention appeared to have synergist effects, reducing local transmission to
intervention non-recipients via spillover effects in all analyses. This may be because short-lived
reductions in host infectiousness following chemotherapy can be sustained over time when
coupled with IRS' long-lasting reductions in mosquito populations. In effect, each intervention
reduces the parasite reservoirs in hosts and vectors, and the combination of interventions
slows the replenishment of parasite reservoirs (36). This may explain why we found that
spillover effects were larger for prevalence of current infection, which captured effects at the
end of the season, rather than incidence, which captured short-term effects. Our findings are
consistent with two recent studies that found evidence of potential synergistic community-level
effects when combining community-wide chemoprevention or seasonal malaria
chemoprevention with IRS in high transmission settings (37, 38). Results are also consistent
with a modeling study that estimated that the joint effect of chemoprevention and IRS was over

1.5 times higher than effects of intervention alone in low-transmission settings (36). Taken
together, our results suggest that the combined intervention may be particularly effective as a
reactive intervention or outbreak response in low-transmission settings approaching
elimination or possibly following introduction of cases after elimination has been achieved.
286
Our estimates of direct and spillover effects shed light on the mechanism through which these
targeted interventions work in time and space. We found that spillover effect sizes were
generally similar to or stronger than direct effect sizes. It is possible that during the time
between index case detection and intervention delivery (median 13-14 days) (24), transmission
occurred to nearby intervention recipients. Thus, the interventions may not have been rapid
enough to reduce malaria among recipients but may have prevented onward transmission to
non-recipients further from index cases. In addition, our finding that spillover effects on
prevalence were stronger at shorter distances to interventions suggests that the majority of the
spillover effect occurred within 1km of index cases.
296
Our study was subject to several limitations. First, due to rare outcomes, precision was low in
some analyses and may have increased the chance of Type II error, especially for direct effects.
Second, when constructing analytic cohorts, household relocation after baseline could have
resulted in misclassification of households to target areas or spillover zones. Third, incidence
analyses could not fully rely on randomization-based inference due to cohort overlap; it is
possible that covariate adjustment did not fully account for imbalances between arms. In future
studies, using a ring trial design to test focal interventions could improve baseline balance,
increase precision, and minimize overlap between target areas (39).
305
Despite these limitations, the internal consistency between the findings of this secondary
analysis and the original trial, which each used different data structures and statistical methods,
supports the validity of our findings. Estimates of total effects in this analysis, which pooled
across intervention recipients and non-recipients, were consistent overall with those of the
original trial, which included all individuals in study clusters (intervention recipients and non-
recipients) (24, 40). Additional strengths include pre-specification of spillover analysis methods
and use of individual-level, spatially indexed data to measure spillovers.
313
In conclusion, we found that reactive, focal malaria interventions targeting both human and
mosquito parasite reservoirs reduced malaria risk, even among non-intervention recipients up
to 3km from index cases. Further, the combined intervention could be particularly useful in
responding to imported infections, which pose a persistent threat prior to and following
elimination. Our findings suggest that these interventions are an effective strategy for achieving
and maintaining malaria elimination.
320
321
322   **Materials and Methods**
323
<u>Analysis overview</u>

This study was a secondary analysis of a cluster-randomized trial of focal malaria interventions
conducted in Zambezi region of Namibia from January 1 to December 31, 2017 (NCT02610400)
(41, 24). Using a two-by-two factorial design, the trial randomly allocated 56 study clusters to
study arms: reactive case detection (RACD) only, reactive focal mass drug administration
(rfMDA) only, RACD + reactive vector control (RAVC), and rfMDA + RAVC. Here, we separately
estimated effects among intervention recipients and non-recipients to estimate direct effects
and spillover effects. We estimated the effects of the chemoprevention intervention (rfMDA vs.
RACD), the vector control intervention (RAVC vs. no RAVC), or combined interventions (rfMDA +
RAVC vs. RACD only), consistent with the original trial (24). The analysis plan for this study was
pre-specified at https://osf.io/s8ay4/. Deviations from the pre-analysis plan and details about
the study site and trial are in the Supporting Information.
336
337 Ethics statement
338 The trial protocol was approved by the Namibia Ministry of Health and Social Services (17/3/3)
339 and the Institutional Review Boards at the University of California San Francisco (15–17422) and
340 London School of Hygiene & Tropical Medicine (10411). The secondary analysis protocol was
341 approved by the Stanford University Institutional Review Board (60708).
342
Incidence analyses
The unit of intervention (index cases) and the unit of randomization (clusters) differed, so
cluster-level analyses would not have captured fine-scale direct effects and spillover effects. To
capture the person-time in which we expected each intervention to influence incident malaria
infections, we created analytic cohorts in space and time around each index case that triggered
an intervention. The primary analysis used a 1 km radius around each index case because the
majority of mosquito movement occurs within < 1 km.
350
We pre-specified cohort follow-up length based on the period in which we expected each
intervention to reduce malaria among intervention recipients (direct effects) and non-recipients
(spillover effects). For comparisons of rfMDA and RACD interventions, the direct effect follow-
up period was 35 days, the length of intrinsic incubation period for *Pf* malaria (8). The spillover
effect follow-up period was 21 to 56 days; the 3-week lag period allowed for gametocyte
clearance in the treated individual, sporozoite development in mosquitos, and development of
detectable merozoites in humans. For RAVC interventions, the direct effects follow-up period
was 6 months since IRS can remain effective for an entire transmission season (9). The spillover
effects follow-up period was from day 17 to 6 months. We conducted a sensitivity analysis with
alternative follow-up lengths (rfMDA and RACD direct effects: day 0-21; spillover effects: day
21-42; RAVC direct effects day 0-7; spillover effects day 17-90). Additional details are in the
Supporting Information.
363
Statistical models for incidence
To compare incidence between arms, we used hierarchical targeted maximum likelihood
estimation (TMLE), a double robust, semi-parametric approach appropriate for cluster-level
exposures (27). TMLE estimates both an outcome regression and a propensity score (the
probability of treatment conditional on covariates) and updates the initial parameter estimate

using information in the propensity score. Compared to other parametric models for clustered
data (e.g., mixed effects models, generalized estimating equations), hierarchical TMLE imposes
fewer assumptions and may be more efficient for randomized trials (42). We fit outcome and
propensity score models using an ensemble machine learning algorithm (the Superlearner) (43).
We adjusted standard errors to account for potential correlation due to overlap between some
cohorts using a model of cohort-level influence curves analogous to variance-covariance
models used in cross-random effects models (See details in Supporting Information) (44, 45).
376
Because incidence analyses did not rely on cluster randomization, we adjusted for covariates
that were correlated with the outcome (likelihood ratio test p-value < 0.2) (46). Propensity
score models adjusted for the following baseline variables: cluster-level indoor residual spray
coverage, malaria incidence, median monthly rainfall, median enhanced vegetative index, and
median daytime land surface temperature in the season prior to the trial, population size, and
median elevation. Outcome models adjusted for individual- and cluster-level covariates.
Individual-level covariates included sex, age, calendar month of intervention, distance from an
individual's residence to the residence of the index case that triggered an intervention, number
of interventions an individual previously received, number of previously intervention recipients
within 0.5, 1, 2, and 3 km of the individual's residence (from the start of the trial to the start of
the cohort's observation period), and population size within 0.5, 1, 2, and 3 km of the
individual's residence. Cluster-level covariates included those in the propensity score models as
well as mean distance to the nearest neighboring household, mean distance to the nearest
healthcare facility, and mean time from index case detection to intervention.
391

<u>Prevalence analyses</u>
To capture effects of cumulative interventions at the end of the malaria transmission season,
we estimated effects on prevalence using data from a cross-sectional survey. Direct effects
analyses included individuals who resided near any intervention recipients within 500 m of their
residence, spillover effects analyses included individuals with no intervention recipients < 500
m and at least one recipient within 500 m-3 km, and total effects included individuals with at
least one intervention recipient within 3 km.
399

<u>Statistical models for prevalence</u>
Prevalence analyses used data from the cross-sectional survey conducted at the end of the
malaria transmission season in 2017. We used targeted maximum likelihood estimation (47)
with individual-level data with the same learners included in incidence analyses. We accounted
for correlation within enumeration area-level clusters using cluster-level influence curve-based
standard errors. The covariate set included the same cluster-level covariates included in
incidence analyses and the following individual-level covariates: age, sex, occupation, recent
travel, household slept under a bed net, slept outdoors in the past two weeks, and the total
population, number of intervention recipients, number of intervention recipients in the same
study arm, the number of intervention recipients in a different study arm, and the proportion of
intervention recipients with the same treatment within 500 m, 1 km, 2 km, and 3 km of
sampled individuals. We screened for covariates using the same approach described for
incidence analyses.

413
<u>Cost-effectiveness analysis</u>
We estimated incremental cost effectiveness ratios (ICER) incorporating direct effects and
spillover effects. To facilitate comparison with the original cost-effectiveness estimates for the
trial (34), we used estimated effects on prevalence measured by qPCR. ICERs estimated using
incidence would not be directly comparable between this study and the original trial because
we used a cohort-level analysis, but the original trial used a cluster-level analysis. We used
previously published estimates of total intervention costs in 2017 in US dollars (34). To obtain
the number of prevalent cases averted, we multiplied the difference in prevalence between
arms among intervention recipients and non-recipients by the estimated population size
included in direct effects and spillover effects analyses. We calculated the ICER by dividing the
difference in cost between arms by the difference in prevalent cases averted between arms
among individuals who were located within 500m of any intervention recipients and individuals
who resided within 500m to 3 km of interventions.
427
428 <u>Data and Code Availability</u>
429 Data from the original trial is available at
430 https://clinepidb.org/ce/app/workspace/analyses/DS_f559aee789. Replication scripts are
431 available at https://github.com/jadebc/malaria-focal-spillover-public.
432
433
434 **Author Contributions:** JBC and MH conceptualized the study. JBC, AB, AH, MVDL, and MH
435 developed the analysis plan. HN, LMP, MT, LW, CD, RG, DM, IK, and MH contributed data. JBC,
436 HL, AN, and GBH processed data. JBC and HL conducted the analysis. JBC wrote the manuscript,
437 and all authors edited the manuscript.
438
439 **Competing Interest Statement:** The authors have no competing interests to declare.
440
441 **Acknowledgements**
442

The original trial was supported by Novartis Foundation (A122666), the Bill & Melinda Gates
Foundation (OPP1160129), and the Horchow Family Fund (5300375400). Research reported in
this publication was supported in part by the National Institute of Allergy and Infectious
Diseases of the National Institutes of Health under Award Numbers K01AI141616 (PI: Benjamin-
Chung) and the National Heart, Lung, And Blood Institute of the National Institutes of Health
under award number T32HL151323 (Nguyen) and Novartis Foundation Supplement to A122666
(PIs: Gosling, Hsiang). The content is solely the responsibility of the authors and does not
necessarily represent the official views of the National Institutes of Health. Jade Benjamin-
Chung and Michelle Hsiang are Chan Zuckerberg Biohub Investigators. We also acknowledge
the Stanford Research Computing Center for computational resources at the Sherlock high-
performance cluster.

454
455
456

References

1. R. G. A. Feachem, *et al.*, Malaria eradication within a generation: ambitious, achievable, and necessary. *The Lancet* **394**, 1056–1112 (2019).

2. , "World malaria report 2022" (World Health Organization) (May 8, 2023).

3. O. Sabot, *et al.*, Costs and financial feasibility of malaria elimination. *The Lancet* **376**, 1604–1615 (2010).

4. E. Halloran, I. M. Longini Jr., C. J. Struchiner, *Design and Analysis of Vaccine Studies* (Springer, 2010).

5. J. Benjamin-Chung, *et al.*, Spillover effects in epidemiology: parameters, study designs and methodological considerations. *Int J Epidemiol* **47**, 332–347 (2018).

6. J. Benjamin-Chung, *et al.*, Spillover effects on health outcomes in low- and middle-income countries: a systematic review. *Int J Epidemiol* **46**, 1251–1276 (2017).

7. T. H. Kim, J. Johnstone, M. Loeb, Vaccine herd effect. *Scandinavian Journal of Infectious Diseases* **43**, 683–689 (2011).

8. S. L. Pollard, *et al.*, Estimating the herd immunity effect of rotavirus vaccine. *Vaccine* **33**, 3795–3800 (2015).

9. G. Tsaban, S. Ben-Shimol, Indirect (herd) protection, following pneumococcal conjugated vaccines introduction: A systematic review of the literature. *Vaccine* **35**, 2882–2891 (2017).

10. R. Borrow, *et al.*, The Global Meningococcal Initiative: global epidemiology, the impact of vaccines on meningococcal disease and the importance of herd protection. *Expert Review of Vaccines* **16**, 313–328 (2017).

11. D. Mertz, *et al.*, Herd Effect From Influenza Vaccination in Non-Healthcare Settings: A Systematic Review of Randomized Controlled and Observational Studies. *Open Forum Infect Dis* **2**, 1916 (2015).

12. M. Drolet, *et al.*, Population-level impact and herd effects following the introduction of human papillomavirus vaccination programmes: updated systematic review and meta-analysis. *The Lancet* **394**, 497–509 (2019).

13. K. S. O'Brien, *et al.*, Azithromycin distribution and childhood mortality in compliance-related subgroups in Niger: complier average causal effect and spillovers in a cluster-randomized, placebo-controlled trial. *International Journal of Epidemiology* **51**, 1775–1784 (2022).

14. J. I. House, *et al.*, Assessment of herd protection against trachoma due to repeated mass
antibiotic distributions: a cluster-randomised trial. *The Lancet* **373**, 1111–1118 (2009).

15. E. Miguel, M. Kremer, Worms: Identifying Impacts on Education and Health in the Presence
of Treatment Externalities. *Econometrica* **72**, 159–217 (2004).

16. F. N. Binka, F. Indome, T. Smith, Impact of spatial distribution of permethrin-impregnated
bed nets on child mortality in rural northern Ghana. *Am J Trop Med Hyg* **59**, 80–85
(1998).

17. W. A. Hawley, *et al.*, Community-Wide Effects of Permethrin-Treated Bed Nets on Child
Mortality and Malaria Morbidity in Western Kenya. *Am J Trop Med Hyg* **68**, 121–127
(2003).

18. C. I. Jarvis, *et al.*, Spatial Effects of Permethrin-Impregnated Bed Nets on Child Mortality: 26
Years on, a Spatial Reanalysis of a Cluster Randomized Trial. *The American Journal of
Tropical Medicine and Hygiene* **101**, 1434–1441 (2019).

19. T. Bousema, *et al.*, The Impact of Hotspot-Targeted Interventions on Malaria Transmission
in Rachuonyo South District in the Western Kenyan Highlands: A Cluster-Randomized
Controlled Trial. *PLOS Medicine* **13**, e1001993 (2016).

20. M. A. Hast, *et al.*, The Impact of 3 Years of Targeted Indoor Residual Spraying With
Pirimiphos-Methyl on Malaria Parasite Prevalence in a High-Transmission Area of
Northern Zambia. *American Journal of Epidemiology* **188**, 2120–2130 (2019).

21. J. J. Kim, The Role of Cost-Effectiveness in U.S. Vaccination Policy. *New England Journal of
Medicine* **365**, 1760–1761 (2011).

22. S. Ma, T. A. Lavelle, D. A. Ollendorf, P.-J. Lin, Herd Immunity Effects in Cost-Effectiveness
Analyses among Low- and Middle-Income Countries. *Appl Health Econ Health Policy* **20**,
395–404 (2022).

23. , "WHO Guidelines for malaria" (World Health Organization, 2023).

24. M. S. Hsiang, *et al.*, The effectiveness of reactive focal mass drug administration (rfMDA)
and reactive focal vector control (RAVC) to reduce malaria transmission: a cluster-
randomised controlled open label two-by-two factorial design trial from the low-endemic
setting of Namibia. **395**, 1361–1373 (2020).

25. M. Makanga, A review of the effects of artemether-lumefantrine on gametocyte carriage
and disease transmission. *Malar J* **13**, 291 (2014).

26. E. Chanda, *et al.*, An investigation of the Plasmodium falciparum malaria epidemic in
Kavango and Zambezi regions of Namibia in 2016. *Transactions of The Royal Society of
Tropical Medicine and Hygiene* **112**, 546–554 (2018).

27. W. Zheng, M. J. van der Laan, "Cross-Validated Targeted Minimum-Loss-Based Estimation"
in *Targeted Learning: Causal Inference for Observational and Experimental Data*, Springer
Series in Statistics., M. J. van der Laan, S. Rose, Eds. (Springer, 2011), pp. 459–474.

28. M. E. Sobel, What Do Randomized Studies of Housing Mobility Demonstrate? *Journal of the
American Statistical Association* **101**, 1398–1407 (2006).

29. D. B. Rubin, Comment on: 'Randomisation analysis of experimental data in the fisher
randomisation test' by D. Basu. *Journal of the American Statistical Association* **75**, 591–
593 (1980).

30. C. F. Manski, Economic Analysis of Social Interactions. *Journal of Economic Perspectives* **14**,
115–136 (2000).

31. C. Costantini, *et al.*, Density, survival and dispersal of Anopheles gambiae complex
mosquitoes in a West African Sudan savanna village. *Medical and Veterinary Entomology*
10, 203–219 (1996).

32. D. L. Huestis, *et al.*, Windborne long-distance migration of malaria mosquitoes in the Sahel.
Nature **574**, 404–408 (2019).

33. L. L. van den Hoogen, *et al.*, Selection of Antibody Responses Associated With Plasmodium
falciparum Infections in the Context of Malaria Elimination. *Frontiers in Immunology* **11**
(2020).

34. H. Ntuku, *et al.*, Cost and cost effectiveness of reactive case detection (RACD), reactive focal
mass drug administration (rfMDA) and reactive focal vector control (RAVC) to reduce
malaria in the low endemic setting of Namibia: an analysis alongside a 2×2 factorial
design cluster randomised controlled trial. *BMJ Open* **12**, e049050 (2022).

35. R. Shretta, A. L. V. Avanceña, A. Hatefi, The economics of malaria control and elimination: a
systematic review. *Malaria Journal* **15**, 593 (2016).

36. R. C. Elliott, D. L. Smith, D. C. Echodu, Synergy and timing: a concurrent mass medical
campaign predicted to augment indoor residual spraying for malaria. *Malaria Journal* **18**,
160 (2019).

37. D. C. Echodu, *et al.*, Impact of population based indoor residual spraying with and without
mass drug administration with dihydroartemisinin-piperaquine on malaria prevalence in
a high transmission setting: a quasi-experimental controlled before-and-after trial in
northeastern Uganda. *BMC Infect Dis* **23**, 72 (2023).

38. J. Wagman, *et al.*, Combining next-generation indoor residual spraying and drug-based
malaria control strategies: observational evidence of a combined effect in Mali. *Malaria
Journal* **19**, 293 (2020).

39. Z. Butzin-Dozier, T. S. Athni, J. Benjamin-Chung, A review of the ring trial design for
evaluating ring interventions for infectious diseases. *Epidemiologic Reviews* **In Press**
(2022).

40. L. Wu, *et al.*, Serological evaluation of the effectiveness of reactive focal mass drug
administration and reactive vector control to reduce malaria transmission in Zambezi
Region, Namibia: Results from a secondary analysis of a cluster randomised trial.
eClinicalMedicine **44**, 101272 (2022).

41. O. F. Medzihradsky, *et al.*, Study protocol for a cluster randomised controlled factorial
design trial to assess the effectiveness and feasibility of reactive focal mass drug
administration and vector control to reduce malaria transmission in the low endemic
setting of Namibia. *BMJ Open* **8** (2018).

42. L. B. Balzer, W. Zheng, M. J. van der Laan, M. L. Petersen, A new approach to hierarchical
data analysis: Targeted maximum likelihood estimation for the causal effect of a cluster-
level exposure. *Stat Methods Med Res* **28**, 1761–1780 (2019).

43. M. J. van der Laan, E. C. Polley, A. E. Hubbard, Super learner. *Stat Appl Genet Mol Biol* **6**,
Article 25 (2007).

44. H. Goldstein, Multilevel Covariance Component Models. *Biometrika* **74**, 430–431 (1987).

45. S. W. Raudenbush, A Crossed Random Effects Model for Unbalanced Data With Applications
in Cross-Sectional and Longitudinal Research. *Journal of Educational Statistics* **18**, 321–
349 (1993).

46. S. J. Pocock, S. E. Assmann, L. E. Enos, L. E. Kasten, Subgroup analysis, covariate adjustment
and baseline comparisons in clinical trial reporting: current practice and problems.
Statistics in Medicine **21**, 2917–2930 (2002).

47. M. J. van der Laan, S. Rose, *Targeted Learning: Causal Inference for Observational and
Experimental Data* (Springer, 2011).

Figure 1. Map of target areas and spillover zones in the study site

a) All index cases during the study period. The centroid of each circle is the residence location of a treated index case. Inner circles indicate 500m target areas where interventions were delivered. Outer circles indicate the 1km radius around each index case in which spillover effects were estimated in primary analyses. The dashed line indicates insets in panels B-D showing index cases during the follow-up periods with the largest number of treated index cases. For each comparison of study arms, panels b-d depict examples of analytic cohorts from a single follow-up period (i.e., subsample of person-time) in a subset of the study area. **b**) Inset of study area with index cases in the RACD and rfMDA arms (5-week period: April 25, 2017 – May 30, 2017). **C**) Inset of study area with index cases in the no RAVC and RAVC arms (6-month period: January 1, 2017 - June 30, 2017). **d**) Inset of study area with index cases in the RAVC and rfMDA+RAVC arms (6-month period: January 1, 2017 - June 30, 2017).

RACD: reactive case detection
rfMDA: reactive focal mass drug administration
RAVC: reactive focal vector control

3
4
5

Figure 2. Effects of reactive, focal malaria interventions

a) Definition of effects in incidence analyses. **b**) Definition of effects in prevalence analyses. **c**) Effects on incidence estimated with hierarchical TMLE. All incidence outcome models were fit with cohort-level data except for models of spillover effects of rfMDA vs. RACD and rfMDA + RAVC vs. RACD only. Models were adjusted for covariates that were screened separately for each model using a likelihood ratio test. Confidence intervals shown here do not account for potential outcome correlation. For rfMDA and RACD arms, the incidence analysis includes the period from 0-35 days following index case detection for direct effects and 21-56 days for spillover effects. For rfMDA+RAVC and RAVC only arms, the

analysis includes the period from 0-6 months following index case detection for direct effects and 17 days to 6 months for spillover effects. Total

effects analyses include the person-time for the direct effects and spillover effects analyses. For incidence analyses, direct effects include treated

in target zone, spillover effects include intervention non-recipients up to 1km from an index case, and total effects include all individuals

(intervention recipients and non-recipients) up to 1km from index case. **d**) Effects on prevalence estimated with TMLE using individual-level data;

standard errors were adjusted for clustering at the enumeration area level. Models were unadjusted because there were fewer than 10 malaria

cases per variable. Direct effects include individuals who resided within 500m of any intervention recipients, spillover effects include individuals

with no intervention recipients < 500m and any intervention recipients 500m-3km, and total effects include individuals with any intervention

recipients <3km during the study. In **c**) and **d**), % effectiveness was calculated as the ratio of incidence or prevalence between study arms minus

1 x 100%. The upper bound of the 95% CI for the combined intervention direct effect was truncated from its original value of 381%.

Figure 3. Spillover effects of reactive, focal malaria interventions on prevalence by distance to nearest intervention recipient

Spillover effects include individuals with no intervention recipients < 500m and any intervention recipients within different distance radii. **a)** Includes effects of the human intervention by comparing arms with rfMDA vs. those with RACD. **b)** Includes effects of the mosquito intervention by comparing arms with RAVC vs. those without RAVC. **c)** Includes effects of the combined intervention by comparing arms with rfMDA vs. RACD and rfMDA + RAVC vs. RACD only. Effects on prevalence estimated with TMLE using individual-level data; standard errors were adjusted for clustering at the enumeration area level. Models were adjusted for covariates that were screened separately for each model using a likelihood ratio test. Models were unadjusted because there were fewer than 10 malaria cases per variable. % Effectiveness was calculated as the ratio of incidence or prevalence between study arms minus 1 x 100%.

34
35
36

* Some individuals close to cluster boundaries contributed to cohorts in more than one study arm.

37
38 **Figure S1. Diagram of study randomization, index cases, and population by arm**
39 RACD: reactive case detection. rfMDA: reactive, focal mass drug administration. RAVC: reactive vector control.

Figure S2. Spillover effect estimates on cumulative incidence within subgroups
Cumulative incidence ratios estimated with hierarchical TMLE; outcome models were fit with cohort-level data. Models were adjusted for covariates that were screened separately for each model using a likelihood ratio test. Models for rfMDA + RAVC vs. RACD were unadjusted due to data sparsity. Confidence intervals account for cohort overlap. For rfMDA and RACD arms, the analysis includes the period from 0-35 days following index case detection for direct effects and 21-56 days for spillover effects. For rfMDA+RAVC and RAVC only arms, the analysis includes

47   the period from 0-6 months following index case detection for direct effects and 17 days to 6 months for spillover effects. Total effects analyses
48   include the person-time for the direct effects and spillover effects analyses. Direct effect includes treated in target zone. Spillover effect includes
49   intervention non-recipients up to 1km from an index case. Total effect includes all individuals (intervention recipients and non-recipients) up to
50   1km from index case. For the human intervention, confidence interval upper bounds were truncated at 16 for above median distance to the
51   nearest health facility (observed value: 23).
52
53

54
Figure S3. Sensitivity analyses for spillover effects on cumulative incidence of malaria with different distance radii
For rfMDA and RACD arms, the primary analysis includes the period from 0-35 days following index case detection for direct effects and 21-56
days for spillover effects; the alternative observation period analysis includes the period from 0-21 days following index case detection for direct
effects and 21 to 42 days for spillover effects. For rfMDA+RAVC and RAVC only arms, the primary analysis includes the period from 0-6 months
following index case detection for direct effects and 17 days to 6 months for spillover effects; the alternative observation period analysis
includes the period from 0-7 days following index case detection for direct effects and 17 to 90 days for spillover effects. Total effects analyses
include the person-time for the direct effects and spillover effects analyses. Direct effect includes intervention recipients in target zone. Spillover
effect includes intervention non-recipients up to 1km from an index case in the primary analysis and up to 2km or 3km in sensitivity analyses.
Total effect includes all individuals (intervention recipients and non-recipients) up to 1km from index case in the primary analysis and up to 2km
or 3km in sensitivity analyses. Includes cohort-level analyses for all estimates except spillover effects of the combined intervention. All incidence
outcome models were fit with cohort-level data except for models of spillover effects of rfMDA vs. RACD and rfMDA + RAVC vs. RACD only.
66

67
Figure S4. Sensitivity analyses for effects on cumulative incidence of malaria
For rfMDA and RACD arms, the primary analysis includes the period from 0-35 days following index case detection for direct effects and 21-56
days for spillover effects; the alternative observation period analysis includes the period from 0-21 days following index case detection for direct
effects and 21 to 42 days for spillover effects. For rfMDA+RAVC and RAVC only arms, the primary analysis includes the period from 0-6 months
following index case detection for direct effects and 17 days to 6 months for spillover effects; the alternative observation period analysis
includes the period from 0-7 days following index case detection for direct effects and 17 to 90 days for spillover effects. Total effects analyses
include the person-time for the direct effects and spillover effects analyses. Direct effect includes intervention recipients in target zone. Spillover
effect includes intervention non-recipients up to 1km from an index case. Total effect includes all individuals (intervention recipients and non-
recipients) up to 1km from index case. Sensitivity analyses for no overlap of spillover zones excluded any cohorts whose spillover zones
overlapped spatially or temporally with other spillover zones. Sensitivity analyses for no overlap of target areas excluded any cohorts whose
target areas overlapped spatially or temporally with other target areas. Some direct effects models could not be fit due to data sparsity. All
incidence outcome models were fit with cohort-level data except for models of spillover effects of rfMDA vs. RACD and rfMDA + RAVC vs. RACD
only.
81
82

83
Figure S5. Sensitivity analyses for direct effects including all intervention recipients
85 The observation period was 0-35 days for rfMDA and RACD arms and 0-6 months for rfMDA+RAVC and RAVC only arms. Black points indicate
86 estimates from analyses including all intervention recipients, regardless of whether they resided within the target zone within 500m of index
87 cases. Mauve points indicate estimates from analyses restricting to intervention recipients within 500m of index cases that triggered
88 interventions. Analyses were performed at the cohort level.
89

		Human intervention	
		Reactive case detection only (28 clusters)	**Reactive focal mass drug administration** (28 clusters)
Mosquito intervention	**No reactive focal vector control** (28 clusters)	Reactive case detection only (14 clusters)	Reactive focal mass drug administration only (14 clusters)
	Reactive focal vector control (28 clusters)	Reactive case detection plus reactive focal vector control (14 clusters)	Reactive focal mass drug administration plus reactive focal vector control (14 clusters)

Table S1. Two-by-two factorial study design of reactive focal interventions
Reactive case detection (RACD) involved administering rapid diagnostic tests for malaria to individuals living within a 500-m radius of an index case and treating
individuals who tested positive with artemether-lumefantrine and single-dose primaquine. Reactive focal mass drug administration (rfMDA) involved
presumptively treating individuals living within a 500-m radius of an index case with artemether-lumefantrine, without testing for malaria beforehand. Reactive
focal vector control (RAVC) involved spraying the long-lasting insecticide, pirimiphos-methyl, to the interior walls of households located within a seven-
household radius of an index case. The effectiveness of three interventions were compared to three respective controls: (1) rfMDA versus RACD (B and D vs A
and C); (2) RAVC versus no RAVC (C and D vs A and B); and (3) rfMDA plus RAVC versus a RACD only (D vs A). Reproduced from Hsiang et al. 2020 *Lancet* with
permission.

		Human intervention		Mosquito intervention		Human & mosquito intervention	
		RACD	rfMDA	No RAVC	RAVC	RACD only	rfMDA + RAVC
Population characteristics							
Number of cohorts		161	149	152	158	73	70
Mean cohort population size (SE)		26 (1)	27 (1)	26 (1)	27 (1)	26 (1)	29 (1)
Mean cluster population size (SE)		389.6 (1.94)	346.4 (1.96)	358.9 (2.01)	376.9 (1.94)	353.0 (2.05)	328.0 (1.89)
Malaria incidence per 1,000 in 2016 (SE)		27.0 (0.37)	55.8 (1.26)	31.9 (0.60)	50.0 (1.15)	26.6 (0.56)	75.4 (2.25)
Pre-season indoor residual spray coverage 2016 (SE)		76.3 (0.32)	77.1 (0.36)	77.9 (0.37)	75.6 (0.31)	83.6 (0.43)	81.5 (0.42)
Distance to nearest healthcare facility (km) (SE)		5.2 (0.06)	6.7 (0.08)	5.0 (0.06)	6.7 (0.07)	3.5 (0.06)	6.9 (0.12)
Ecological factors (range)							
Median monthly rainfall November 2016-April 2017 (mm)		23.7 (18.4, 26.7)	23.5 (18.4, 26.7)	23.5 (18.4, 26.7)	23.7 (18.4, 26.7)	23.7 (18.4, 26.7)	23.7 (18.4, 26.7)
Median enhanced vegetative index January 2017-July 2017		0.15 (0.09, 0.31)	0.15 (0.09, 0.27)	0.15 (0.09, 0.22)	0.15 (0.09, 0.31)	0.15 (0.10, 0.21)	0.15 (0.09, 0.27)
Median elevation (m)		522 (387, 1021)	541 (412, 1124)	527 (398, 1124)	547 (387, 1021)	522 (398, 921)	576 (412, 984)
Median daytime land surface temperature (C)		30.5 (28.9, 33.4)	31.1 (28.6, 32.5)	30.7 (28.6, 33.4)	30.8 (28.7, 32.5)	30.7 (28.9, 33.4)	31.1 (28.7, 32.5)

102 **Table S2. Baseline characteristics among intervention recipients**

103 Includes data from intervention recipients in target areas located within 500m of an index case.

	Human intervention		Mosquito intervention		Human & mosquito intervention	
	RACD	rfMDA	No RAVC	RAVC	RACD only	rfMDA + RAVC
Population characteristics						
Number of cohorts	161	149	152	158	73	70
Mean cohort population size (SE)	238 (9)	232 (12)	223 (9)	247 (11)	256 (13)	276 (19)
Mean cluster population size (SE)	379.4 (0.63)	355.8 (0.59)	354.0 (0.58)	380.5 (0.63)	349.5 (0.61)	352.8 (0.63)
Malaria incidence per 1,000 in 2016 (SE)	29.2 (0.12)	41.0 (0.35)	28.3 (0.17)	40.4 (0.29)	27.3 (0.16)	50.0 (0.55)
Pre-season indoor residual spray coverage 2016 (SE)	77.1 (0.10)	81.0 (0.12)	78.6 (0.12)	79.2 (0.10)	82.8 (0.14)	86.9 (0.12)
Distance to nearest healthcare facility (km) (SE)	4.9 (0.02)	6.7 (0.03)	4.4 (0.02)	6.9 (0.02)	3.2 (0.02)	7.3 (0.04)
Ecological factors (range)						
Median monthly rainfall November 2016-April 2017 (mm)	23.7 (18.4, 26.7)	23.5 (18.4, 26.7)	23.7 (18.4, 26.7)	23.7 (18.4, 26.7)	23.7 (18.4, 26.7)	23.7 (18.4, 26.7)
Median enhanced vegetative index January 2017-July 2017	0.15 (0.09, 0.31)	0.15 (0.09, 0.27)	0.15 (0.09, 0.22)	0.15 (0.09, 0.31)	0.15 (0.10, 0.21)	0.15 (0.09, 0.27)
Median elevation (m)	522 (387, 1021)	535 (412, 1124)	527 (398, 1124)	547 (387, 1021)	522 (398, 921)	677 (412, 984)
Median daytime land surface temperature (C)	30.5 (28.9, 33.4)	31.1 (28.6, 32.5)	30.7 (28.6, 33.4)	30.8 (28.7, 32.5)	30.6 (28.9, 33.4)	31.1 (28.7, 32.5)

107 **Table S3. Baseline characteristics among non-intervention recipients up to 1km away from index cases**
108 Includes data from intervention non-recipients up to 1km from an index case that triggered interventions.
109
110

	N cohorts	N	Incidence proportion		Incidence ratio (95% CI)		
			Intervention arm	Reference arm	Unadjusted	Adjusted	Adjusted, CI adjusted for cohort overlap
Human intervention (rfMDA vs. RACD)							
Direct effect	310	8,252	3.4	6.5	0.53 (0.25, 1.11)	0.40 (0.11, 1.48)	0.40 (0.10, 1.56)
Spillover effect	310	72,830	9.0	9.9	0.91 (0.60, 1.37)	0.82 (0.52, 1.29)	0.82 (0.44, 1.51)
Total effect	310	81,082	8.4	9.6	0.88 (0.59, 1.31)	0.83 (0.51, 1.35)	0.83 (0.43, 1.60)
Mosquito intervention (RAVC vs. no RAVC)							
Direct effect	310	8,252	8.9	7.6	1.17 (0.62, 2.23)	1.35 (0.54, 3.34)	1.35 (0.43, 4.25)
Spillover effect	310	72,830	12.9	18.5	0.69 (0.47, 1.03)	0.68 (0.46, 1.00)	0.68 (0.36, 1.30)
Total effect	310	81,082	12.5	17.4	0.72 (0.49, 1.06)	0.73 (0.49, 1.08)	0.73 (0.36, 1.45)
Combined intervention (rfMDA + RAVC vs. RACD only)							
Direct effect	143	3,914	6.4	7.4	0.87 (0.32, 2.41)	1.03 (0.22, 4.81)	1.03 (0.19, 5.58)
Spillover effect	143	38,048	11.2	18.1	0.62 (0.34, 1.13)	0.57 (0.41, 0.80)	0.57 (0.42, 0.79)
Total effect	143	41,962	10.8	17.1	0.63 (0.35, 1.12)	0.37 (0.22, 0.63)	0.37 (0.18, 0.79)

Table S4. Direct effect, spillover effect, and total effect estimates on cumulative incidence of malaria infection
For rfMDA and RACD arms, the analysis includes the period from 0-35 days following index case detection for direct effects and 21-56 days for
spillover effects. For rfMDA+RAVC and RAVC only arms, the analysis includes the period from 0-6 months following index case detection for
direct effects and 17 days to 6 months for spillover effects. Total effects analyses include the person-time for the direct effects and spillover
effects analyses. Direct effect includes intervention recipients in the target zone. Spillover effect analyses includes intervention non-recipients up
to 1km from an index case. Total effect includes all individuals (intervention recipients and non-recipients) up to 1km from index case. Models
were fit with hierarchical targeted maximum likelihood. All outcome models were fit with cohort-level data except for models of spillover effects
of rfMDA + RAVC vs. RACD only. Adjusted models were fit if there were fewer than 10 malaria cases per variable. Covariates were screened
separately for each model using a likelihood ratio test. We separately fit individual- and cohort-level outcome models and report the model with
the smaller cross-validated mean squared error. All models except spillover effects of the human and combined interventions were fit on cohort-
level data.

	Below median		Above median	
	Minimum	Maximum	Minimum	Maximum
Malaria incidence per 1,000 in 2016	0.0	13.9	14.9	293.3
Pre-season indoor residual spray coverage 2016 (%)	27.2	77.3	77.9	100
Median daytime land surface temperature (C)	28.6	31.1	31.1	33.4
Median monthly rainfall November 2016-April 2017 (mm)	18.4	23.7	23.7	26.7
Median enhanced vegetative index January 2017-July 2017	0.09	0.15	0.15	0.31
Median elevation (m)	387	541	544	1124
Cohort-level treatment coverage (%)	0.0	8.3	8.3	97.4

125 **Table S5. Range above and below median value in each enumeration area for subgroup variables**

	Primary analysis		Sensitivity analysis with shorter observation period	
	Target areas	Spillover zone	Target areas	Spillover zone
Human intervention (rfMDA vs. RACD)	32.0	28.9	21.2	18.4
Mosquito intervention (RAVC vs. no RAVC)	59.2	47.5	53.8	41.8
Combined intervention (rfMDA + RAVC vs. RACD only)	60.5	28.1	60.2	24.1

Table S6. Percentage of cohorts overlapping with other cohorts

Overlap in target area was defined as index cases that triggered interventions located within <1km of each other and observation periods that temporally overlapped with another cohort's. Overlap in spillover zones was defined as index cases that triggered interventions located within 1-2km of each other and observation periods that temporally overlapped with another cohort's. The denominator was the total cohorts included in each analysis.

	N		Prevalence		Prevalence ratio (95% CI)	
	Intervention arm	Reference arm	Intervention arm	Reference arm	Unadjusted	Adjusted
Human intervention (rfMDA vs. RACD)						
Direct effect	1537	1835	0.029	0.033	0.90 (0.61, 1.31)	0.84 (0.53, 1.32)
Spillover effect	244	229	0.025	0.087	0.28 (0.12, 0.69)	--
Total effect	1781	2064	0.029	0.039	0.74 (0.52, 1.04)	0.79 (0.51, 1.19)
Mosquito intervention (RAVC vs. no RAVC)						
Direct effect	1710	1662	0.026	0.037	0.70 (0.48, 1.03)	0.78 (0.51, 1.21)
Spillover effect	195	278	0.051	0.058	0.89 (0.41, 1.92)	--
Total effect	1905	1940	0.028	0.040	0.71 (0.51, 1.01)	0.64 (0.43, 0.96)
Combined intervention (rfMDA + RAVC vs. RACD only)						
Direct effect	758	883	0.017	0.033	0.52 (0.27, 1.00)	--
Spillover effect	118	152	0.017	0.079	0.21 (0.05, 0.94)	--
Total effect	876	1035	0.017	0.040	0.43 (0.24, 0.78)	--

Table S7. Direct effect, spillover effect, and total effect estimates on malaria prevalence measured by qPCR
Prevalence was measured in a cross-sectional survey in a random sample of households at the end of the malaria season. Analyses were
restricted to individuals located within 3 km of at least one intervention recipient. Direct effects include individuals with any intervention
recipients within 500m, spillover effects include individuals with no intervention recipients < 500m and any intervention recipients 500m-3km,
and total effects include individuals with any intervention recipients <3km during the study. Prevalence ratios were estimated using TMLE with
individual-level data, and standard errors were adjusted for clustering at the enumeration area level. Adjusted analyses were not fit there were
fewer than 30 observations within strata of the intervention and outcome. Adjusted models were not fit if the number of cases within treatment
arm strata was <30.
144

	N households		Prevalence		Unadjusted
	Intervention arm	Reference arm	Intervention arm	Reference arm	Prevalence Ratio (95% CI)
Human intervention (rfMDA vs. RACD)					
Direct effect	456	506	0.018	0.018	0.99 (0.38, 2.54)
Spillover effect	72	69	0.000	0.043	0.00 (0.00, 0.00)
Total effect	528	575	0.015	0.021	0.73 (0.30, 1.76)
Mosquito intervention (RAVC vs. no RAVC)					
Direct effect	481	481	0.012	0.023	0.55 (0.20, 1.46)
Spillover effect	65	76	0.015	0.026	0.58 (0.05, 6.35)
Total effect	546	557	0.013	0.023	0.55 (0.22, 1.37)
Combined intervention (rfMDA + RAVC vs. RACD only)					
Direct effect	219	244	0.005	0.016	0.28 (0.03, 2.48)
Spillover effect	36	40	0.000	0.050	0.00 (0.00, 0.00)
Total effect	255	284	0.004	0.021	0.19 (0.02, 1.53)

Table S8. Direct effect, spillover effect, and total effect estimates on household-level malaria prevalence of measured by qPCR

Prevalence was measured in a cross-sectional survey in a random sample of households at the end of the malaria season. Analyses were run at
the household level. Household-level malaria prevalence was the percentage of households with more than one malaria case detected in the
prevalence survey by qPCR. Direct effects include households with any intervention recipients within 500m, spillover effects include households
with no intervention recipients < 500m and any intervention recipients 500m-3km, and total effects include households with any intervention
recipients <3km during the study. Prevalence ratios were estimated using TMLE with household-level data. Adjusted analyses were not fit there
were fewer than 30 observations within strata of the intervention and outcome. Adjusted models were not fit if the number of cases within
treatment arm strata was <30.

	N		Prevalence		Prevalence ratio (95% CI)	
	Intervention arm	Reference arm	Intervention arm	Reference arm	Unadjusted	Adjusted
Human intervention (rfMDA vs. RACD)						
Direct effect	1316	1611	0.215	0.285	0.75 (0.66, 0.86)	0.84 (0.71, 1.00)
Spillover effect	198	182	0.227	0.225	1.01 (0.69, 1.46)	1.32 (0.73, 2.41)
Total effect	1514	1793	0.217	0.279	0.78 (0.69, 0.88)	0.85 (0.73, 0.99)
Mosquito intervention (RAVC vs. no RAVC)						
Direct effect	1475	1452	0.241	0.267	0.90 (0.80, 1.02)	0.90 (0.79, 1.04)
Spillover effect	133	247	0.188	0.247	0.76 (0.50, 1.15)	--
Total effect	1608	1699	0.236	0.264	0.90 (0.80, 1.01)	0.88 (0.76, 1.01)
Combined intervention (rfMDA + RAVC vs. RACD only)						
Direct effect	634	770	0.194	0.295	0.66 (0.54, 0.80)	--
Spillover effect	81	130	0.136	0.208	0.66 (0.55, 0.80)	--
Total effect	715	900	0.187	0.282	0.65 (0.34, 1.25)	--

158
Table S9. Direct effect, spillover effect, and total effect estimates on Etramp5.Ag1 seroprevalence
Prevalence was measured in a cross-sectional survey in a random sample of households at the end of the malaria season. Analyses were
restricted to individuals located within 3 km of at least one intervention recipient. Direct effects include individuals with any intervention
recipients within 500m, spillover effects include individuals with no intervention recipients < 500m and any intervention recipients 500m-3km,
and total effects include individuals with any intervention recipients <3km during the study. Prevalence ratios were estimated using TMLE with
individual-level data, and standard errors were adjusted for clustering at the enumeration area level. Adjusted analyses were not fit there were
fewer than 30 observations within strata of the intervention and outcome. Adjusted models were not fit if the number of cases within treatment
arm strata was <30.
167

		N individuals		Prevalence		Prevalent cases		Total prevalent cases averted (95% CI)	Incremental cost-effectiveness ratio (95% CI)	% change from original estimate
	Intervention cost	Target area	Spillover zone	Target area	Spillover zone	Target area	Spillover zone			
Human intervention										
RACD	$354,750	8,187	996	0.033	0.087	268	87	(ref)	(ref)	
rfMDA	$368,321	8,060	1,301	0.029	0.025	236	32	87 (77, 96)	$156 ($141, $177)	-3%
Mosquito intervention										
No RAVC	$261,409	7,845	1,290	0.037	0.058	288	74	(ref)	(ref)	
RAVC	$461,661	8,426	980	0.026	0.051	217	50	95 (82, 108)	$2,105 ($1,859, $2,430)	-21%
Combined intervention										
RACD only	$127,312	3,697	626	0.033	0.079	121	49	(ref)	(ref)	
rfMDA+RAVC	$234,223	3,878	635	0.017	0.017	66	11	94 (74, 113)	$1,142 ($944, $1,446)	-37%

Table S10. Cost-effectiveness analysis

Prevalent cases averted were estimated using hierarchical TMLE models for prevalence measured by qPCR. The number of prevalent cases
averted equaled the produce of the difference in prevalence between arms among intervention recipients and non-recipients by the estimated
population size within target areas vs. spillover zones. The incremental cost effectiveness ratio is the ratio of the difference in cost between arms
by the difference in prevalent cases averted in both target area and spillover zones within 3 km of index cases for rfMDA + RAVC vs. RACD.
Original estimates were reported in Ntuku et al., 2022 10.1136/bmjopen-2021-049050.

Supporting Information

Study population
This study analyzed data from a cluster-randomized trial of focal malaria interventions conducted in Zambezi region of Namibia from January 1 to December 31, 2017 (NCT02610400) (1, 2). The region has seasonal malaria transmission that peaks between January and June. *Plasmodium falciparum* is the dominant species, and annual *Pf* incidence was less than 15 per 1,000 from 2010-2015. In 2016, the incidence was 32.5 per 1,000 following an outbreak (3). In 2015, prevalence measured by loop-mediated isothermal amplification was 2.2% (4). In the study site, the Namibia Ministry of Health and Social Services routinely delivered case management and annual preseason household IRS with dichlorodiphenyltrichloroethane, with the exception of a small number of structures that were sprayed with deltamethrin. In addition, they offered reactive case detection (RACD) within 500 m of confirmed malaria cases, which included testing with rapid diagnostic tests and treatment with artemether-lumefrantrine and single-dose primaquine for those who tested positive.

Cluster-randomized trial design
The trial included 56 clusters defined based on census enumeration areas that were within the catchment area of study health care facilities. Enumeration areas were eligible for inclusion in the trial if they 1) were located in the catchment areas of 11 health facilities, 2) had complete incidence data from 2012-13, and 3) had at least one incident case during the trial. Using a two-by-two factorial design, the trial randomized 56 clusters to four arms: 1) RACD only, 2) reactive focal mass drug administration (rfMDA) only, 3) reactive vector control (RAVC) + RACD, 4) RAVC + rfMDA. rfMDA included presumptive treatment with artemether-lumefrantrine to individuals in target areas (Extended Data Table 1). The trial used restricted randomization with the following criteria: mean annual incidence in 2013 and 2014, population size, population density, and mean distance from the household to a health-care facility. It was not practical to blind study participants or field staff to intervention assignment, but laboratory analyses and primary statistical analyses were blinded.

Interventions
Field staff delivered interventions in response to passively detected malaria index cases that were confirmed by rapid diagnostic tests or microscopy if the case had resided in the study cluster at least one night in the prior 4 weeks. The trial delivered interventions in "target areas" within approximately 500 m of confirmed malaria cases detected through passive surveillance. In the RACD arms, individuals were eligible to receive rapid diagnostic tests, and individuals who tested positive were eligible for treatment with artemether-lumefrantrine and single-dose primaquine (Coartem, Novartis Pharmaceuticals, Kempton Park, South Africa; or Komefan 140, Mylan Laboratories, Sinnar, India). In the rfMDA arms, individuals were eligible for presumptive treatment with artemether-lumefrantrine. In the RAVC arms, households were eligible for IRS with pirimiphosmethyl (Actellic 300CS, Syngenta, Basel, Switzerland). In all arms, study teams aimed to deliver interventions within 500 m of a clinical malaria case and within 7 days to 5 weeks of the case report. RACD and rfMDA interventions were delivered to at least 25 people within target areas and RAVC was delivered to at least seven households within target areas.

Over 80% of eligible confirmed malaria cases received interventions, and over 85% of eligible intervention recipients were covered by interventions (2). Since compliance was high, for intervention recipients, we analyzed treatment as randomly assigned. Field staff did not offer repeat interventions in response to subsequent index cases within 5 weeks for rfMDA and RACD and within the same malaria season for RAVC. Field staff recorded the household geocoordinates of the index case and intervention recipients. Additional details about the interventions were previously published (1, 2).

Procedures
Prior to randomization, field staff conducted a geographic census and recorded the latitude and longitude of all households in the study area. During the trial, trial staff extracted data on confirmed incident malaria cases and travel history from the rapid reporting system. At the end of malaria season between May and August 2017, the study team collected an endline cross-sectional survey to measure infection prevalence. Field staff collected dried blood spots on filter paper (Whatman 3 Corporation, Florham Park, NJ, USA) by finger prick from consenting individuals, and qPCR was performed targeting the acidic terminal sequence of the *var* gene.(5) Field staff also collected 250 ml of whole blood in BD Microtainer tubes with EDTA additive (Becton, Dickinson and Corporation, Franklin Lakes, NJ, USA) for serological analyses. Using human plasma, Luminex assays were performed to detect malaria antigens using previously described procedures (6, 7). Field staff recorded the geocoordinates of all sampled households.

Informed consent
In the original trial, written informed consent was obtained from individual participants for rfMDA or RACD, and from heads of households (≥18 years of age) for RAVC. A parent or guardian was required to provide written informed consent for children younger than 18 years receiving rfMDA or RACD, and written assent for receiving these interventions was also obtained from children aged 12–17 years.

Construction of analytic cohorts for incidence analysis
To construct cohorts, we matched index cases and intervention recipients to individuals recorded in the baseline census using household geocoordinates, age, and sex. We required that geocoordinates be < 100m apart to allow for small deviations in the location of geocoordinate recordings. We excluded 32 cohorts from the analysis for which it was not possible to merge intervention recipient geocoordinates with index data geocoordinates. Because clusters were contiguous with no buffer zones between them, to capture potential dependencies across study clusters, we allowed cohorts to include individuals assigned to an adjacent cluster with a different treatment assignment from the triggering index case if it was within 1 km of an index case.

Follow-up periods for analytic cohorts
We pre-specified cohort follow-up length based on the period in which we expected each intervention to reduce malaria among intervention recipients (direct effects) and non-recipients (spillover effects). Day 0 for each cohort was the date of index case detection. For comparisons of rfMDA and RACD interventions, the direct effect follow-up period was 0 to 35 days, the

length of intrinsic incubation period for *Pf* malaria (8). This is the period of time in which we would expect the intervention to interrupt the parasite life cycle in treated, infected individuals, and in turn, prevent symptoms and/or infectiousness. The spillover effect follow-up period was 21 to 56 days; the 3-week lag period allowed for gametocyte clearance in the treated individual, sporozoite development in mosquitos, and development of detectable merozoites in humans. For RAVC interventions, the direct effects follow-up period was 6 months since IRS can remain effective for an entire transmission season (9). The spillover effects follow-up period was from day 17 to 6 months. A mosquito bite could hypothetically be prevented on the day of intervention, so the earliest secondary case could occur after sporozoite development in mosquitos (minimum 10 days), and development of detectable merozoites in humans (minimum 7 days). We conducted a sensitivity analysis with alternative follow-up lengths (rfMDA and RACD direct effects: day 0-21; spillover effects: day 21-42; RAVC direct effects day 0-7; spillover effects day 17-90).

Hierarchical TMLE
We compared incidence between arms using hierarchical targeted maximum likelihood estimation (TMLE) (10). We fit propensity score models at the cohort-level since interventions were delivered to cohorts. Within study clusters and cohorts, we expected individuals' outcomes to be correlated due to interventions, social interactions, and local environmental factors. We fit two types of outcome models that accounted for statistical dependence in different ways (11). Cohort-level models allowed for statistical dependence between individuals in the same cohort without making any assumptions about the nature of the dependency. Individual-level models assumed that cluster-level and individual-level covariates removed any dependence between outcomes of individuals in nearby geographic areas (11). We separately fit individual- and cohort-level models and then chose the outcome model with the smaller cross-validated mean squared error.

We fit outcome and propensity score models using an ensemble machine learning algorithm (the Superlearner) (12). For propensity score models, learners included generalized linear models, least absolute shrinkage and selection operator (LASSO) (13), and elastic net regression (14). For outcome models, we used the same learners as well as extreme gradient boosting (15). We performed 10-fold cross-validation using a loss function at either the individual- or cohort-level (11). Validation samples were constructed from randomly sampled individuals or cohorts. Because comparisons of rfMDA + RAVC vs. RACD had rare outcomes and a smaller sample size, we used 30-fold cross-validation.

Adjusting standard errors for cohort overlap
We adjusted standard errors to account for potential correlation due to overlap between some cohorts using a model of cohort-level influence curves analogous to variance-covariance models used in cross-random effects models (16, 17). Specifically, we fit the model:

$$D_i \times D_j \sim d(i,j) + t(i,j) + C \tag{1}$$

where $D_i \times D_j$ is the product of influence curves of cohorts i and j, $d(i,j)$ is the distance between the location of the index case that triggered the intervention in each cohort, $t(i,j)$ is the start date of the intervention in each cohort, and C is the cluster-level intervention assignment (18). Adjustment for intervention assignment accounted for correlation due to shared exposure to or receipt of the intervention. For cohorts with no overlap, we set $D_i \times D_j$ to zero. The regression was implemented with a simplified SuperLearner library including the generalized linear models and LASSO (13). We calculated the variance accounting for outcome dependence as follows:

$$var(\hat{\psi} - \psi) = var\left(\frac{1}{N}\sum_{i=1}^{N} D_i\right) = \frac{1}{N^2}\left(\sum_{i=1}^{N} var(D_i) + 2\sum_{i<j} cov(D_i, D_j)\right)$$

where $\hat{\psi}$ is the estimator, ψ is the estimand, and N is the number of cohorts.

In both incidence and prevalence analyses, we excluded any categorical covariates with less than 5% prevalence to avoid positivity violations. To minimize empirical positivity violations (19), we only fit models if the number of outcome events per variable was $\geq$10 and only fit adjusted models if the number of observations per strata was $\geq$30 (20).

Deviations from pre-analysis plan

The analysis plan for this study was pre-specified at https://osf.io/s8ay4/. We note the following deviations from the plan:

1. We originally planned to conduct an individual participant data meta-analysis including data from three trials in Namibia, Eswatini, and Zambia. However, after reviewing the data for the Eswatini and Zambia trials, we determined that the geocoding of participants was not sufficient to allow for the planned spillover analyses. Thus, we proceeded with an analysis using data only from the Namibia trial.

2. In primary analyses using incidence data, we did not impose bounds on the mean outcome conditional on treatment and covariates because in initial models using bounds, estimates were very unstable.

3. In secondary analyses using prevalence data, we corrected standard errors at the cluster-level instead of at the household-level as specified in the pre-analysis plan. This better reflected the clustered sampling in the original trial.

References

1. O. F. Medzihradsky, *et al.*, Study protocol for a cluster randomised controlled factorial design trial to assess the effectiveness and feasibility of reactive focal mass drug administration and vector control to reduce malaria transmission in the low endemic setting of Namibia. *BMJ Open* **8** (2018).

2. M. S. Hsiang, *et al.*, The effectiveness of reactive focal mass drug administration (rfMDA) and reactive focal vector control (RAVC) to reduce malaria transmission: a cluster-randomised controlled open label two-by-two factorial design trial from the low-endemic setting of Namibia. **395**, 1361–1373 (2020).

3. E. Chanda, *et al.*, An investigation of the Plasmodium falciparum malaria epidemic in Kavango and Zambezi regions of Namibia in 2016. *Transactions of The Royal Society of Tropical Medicine and Hygiene* **112**, 546–554 (2018).

4. P. McCreesh, *et al.*, Subpatent malaria in a low transmission African setting: a cross-sectional study using rapid diagnostic testing (RDT) and loop-mediated isothermal amplification (LAMP) from Zambezi region, Namibia. *Malar J* **17**, 480 (2018).

5. N. Hofmann, *et al.*, Ultra-Sensitive Detection of Plasmodium falciparum by Amplification of Multi-Copy Subtelomeric Targets. *PLOS Medicine* **12**, e1001788 (2015).

6. L. Wu, *et al.*, Optimisation and standardisation of a multiplex immunoassay of diverse Plasmodium falciparum antigens to assess changes in malaria transmission using sero-epidemiology. *Wellcome Open Res* **4**, 26 (2020).

7. L. Wu, *et al.*, Serological evaluation of the effectiveness of reactive focal mass drug administration and reactive vector control to reduce malaria transmission in Zambezi Region, Namibia: Results from a secondary analysis of a cluster randomised trial. *eClinicalMedicine* **44**, 101272 (2022).

8. S. K. Nilsson, L. M. Childs, C. Buckee, M. Marti, Targeting Human Transmission Biology for Malaria Elimination. *PLOS Pathogens* **11**, e1004871 (2015).

9. World Health Organization, "WHO recommended insecticides for indoor residual spraying against malaria vectors" (2018).

10. W. Zheng, M. J. van der Laan, "Cross-Validated Targeted Minimum-Loss-Based Estimation" in *Targeted Learning: Causal Inference for Observational and Experimental Data*, Springer Series in Statistics., M. J. van der Laan, S. Rose, Eds. (Springer, 2011), pp. 459–474.

11. L. B. Balzer, W. Zheng, M. J. van der Laan, M. L. Petersen, A new approach to hierarchical data analysis: Targeted maximum likelihood estimation for the causal effect of a cluster-level exposure. *Stat Methods Med Res* **28**, 1761–1780 (2019).

12. M. J. van der Laan, E. C. Polley, A. E. Hubbard, Super learner. *Stat Appl Genet Mol Biol* **6**, Article 25 (2007).

13. R. Tibshirani, Regression Shrinkage and Selection via the Lasso. *Journal of the Royal Statistical Society. Series B (Methodological)* **58**, 267–288 (1996).

14. J. Friedman, T. Hastie, R. Tibshirani, Regularization Paths for Generalized Linear Models via Coordinate Descent. *J Stat Softw* **33**, 1–22 (2010).

15. J. H. Friedman, Greedy Function Approximation: A Gradient Boosting Machine. *The Annals of Statistics* **29**, 1189–1232 (2001).

16. H. Goldstein, Multilevel Covariance Component Models. *Biometrika* **74**, 430–431 (1987).

17. S. W. Raudenbush, A Crossed Random Effects Model for Unbalanced Data With Applications in Cross-Sectional and Longitudinal Research. *Journal of Educational Statistics* **18**, 321–349 (1993).

18. M. M. Davies, M. J. van der Laan, Sieve Plateau Variance Estimators: A New Approach to Confidence Interval Estimation for Dependent Data. *U.C. Berkeley Division of Biostatistics Working Paper Series* **Working Paper 322** (2014).

19. M. L. Petersen, K. E. Porter, S. Gruber, Y. Wang, M. J. van der Laan, Diagnosing and responding to violations in the positivity assumption. *Stat Methods Med Res* **21**, 31–54 (2010).

20. P. Peduzzi, J. Concato, E. Kemper, T. R. Holford, A. R. Feinstein, A simulation study of the number of events per variable in logistic regression analysis. *Journal of Clinical Epidemiology* **49**, 1373–1379 (1996).