

1 **Hebbian plasticity induced by temporally coincident BCI enhances post-stroke motor recovery**

2 Johanna Krueger¹, Richard Krauth¹, Christoph Reichert², Serafeim Perdakis³, Susanne Vogt⁴, Tessa
3 Huchtemann⁵, Stefan Dürschmid^{2,6}, Almut Sickert⁷, Juliane Lamprecht^{7,8}, Almir Huremovic^{7,9}, Michael
4 Görtler⁴, Slawomir J. Nasuto¹⁰, I-Chin Tsai¹, Robert T. Knight^{11,12}, Hermann Hinrichs², Hans-Jochen
5 Heinze^{2,13}, Sabine Lindquist¹⁴, Michael Sailer⁷, Jose del R. Millán^{15,16}, Catherine M. Sweeney-Reed^{1,6*}

6

7 ¹Neurocybernetics and Rehabilitation, Dept. of Neurology, Otto von Guericke University, Magdeburg,
8 Germany

9 ²Leibniz Institute for Neurobiology, Magdeburg, Germany

10 ³School of Computer Science and Electronic Engineering, University of Essex, Colchester, UK

11 ⁴Dept. of Neurology, Otto von Guericke University Magdeburg, Germany

12 ⁵Dept. of Neurology, University Hospital Münster, Münster, Germany

13 ⁶Center for Behavioral Brain Sciences (CBBS), Otto von Guericke University Magdeburg, Germany

14 ⁷Neurorehabilitation Centre, Median Hospital, Magdeburg, Germany

15 ⁸Health and Care Sciences, Martin Luther University Halle-Wittenberg, Halle, Germany

16 ⁹Dept. of Neurology, Ingolstadt Hospital, Ingolstadt, Germany

17 ¹⁰Biomedical Engineering, University of Reading, Reading, UK

18 ¹¹Helen Wills Neuroscience Institute, University of California – Berkeley, Berkeley, USA

19 ¹²Dept. of Psychology, University of California, Berkeley, Berkeley, USA

20 ¹³University Hospital Magdeburg, Otto von Guericke University Magdeburg, Magdeburg, Germany

21 ¹⁴Dept. of Neurology, Pfeiffersche Stiftung, Magdeburg, Germany

22 ¹⁵Dept. of Electrical and Computer Engineering, The University of Texas at Austin, Austin, USA

23 ¹⁶Dept. of Neurology, The University of Texas at Austin, Austin, USAs, USA

24

25 *Corresponding author:

26 Catherine Sweeney-Reed: catherine.sweeney-reed@med.ovgu.de

27 **Abstract**

28 Functional electrical stimulation (FES) can support functional restoration of a paretic limb post-stroke.
29 Hebbian plasticity depends on temporally coinciding pre- and post-synaptic activity. A tight temporal
30 relationship between motor cortical (MC) activity associated with attempted movement and FES-
31 generated visuo-proprioceptive feedback is hypothesized to enhance motor recovery. Using a brain-
32 computer interface (BCI) to classify MC spectral power in electroencephalographic (EEG) signals to
33 trigger FES-delivery with detection of movement attempts improved motor outcomes in chronic stroke
34 patients. We hypothesized that heightened neural plasticity earlier post-stroke would further enhance
35 corticomuscular functional connectivity and motor recovery. We compared subcortical non-dominant
36 hemisphere stroke patients in BCI and Sham (FES temporally independent of MC movement attempt
37 detection) groups. The primary outcome measure was the Fugl-Meyer Assessment, Upper Extremity
38 (FMA-UE). We recorded high-density EEG and transcranial magnetic stimulation-induced motor
39 evoked potentials before and after treatment. The BCI group showed greater: FMA-UE improvement;
40 motor evoked potential amplitude; beta oscillatory power and long-range temporal correlation
41 reduction over contralateral MC; and corticomuscular coherence with contralateral MC. These
42 changes are consistent with enhanced post-stroke motor improvement when movement is
43 synchronized with MC activity reflecting attempted movement.

44 **Key words:** brain-computer interface; BCI; functional electrical stimulation; FES; acute stroke;
45 subacute stroke; upper limb rehabilitation; Fugl-Meyer Assessment; FMA; EEG; TMS

46 Introduction

47 Stroke is a leading cause of motor disability¹, with upper limb impairment occurring in over 75% of
48 patients following acute stroke². Despite the successes of thrombolytic therapy in reducing mortality
49 and morbidity, a third or less of patients meet the criteria for this treatment, and over half of those
50 receiving it are left with functional deficits³. Motor recovery depends on neural plasticity and the
51 reorganization of structural and functional motor networks to re-establish corticomuscular
52 connectivity⁴⁻⁸. Neural plasticity is task-specific, time-dependent, and environmentally-influenced⁹.
53 Various approaches to re-establishment and reinforcement of connectivity between paretic
54 musculature and residual motor areas are based on targeting Hebbian plasticity by synchronizing
55 movement-associated visuo-proprioceptive feedback and motor cortical electrophysiological
56 correlates of movement within a narrow time window^{10,11}. Functional electrical stimulation (FES) is an
57 established therapeutic tool for assisting movement attempts and promoting motor recovery. Studies
58 involving chronic and subacute stroke patients have enhanced motor recovery when FES delivery is
59 temporally coupled to movement attempts detected in brain electrical activity, using a brain-
60 computer interface (BCI)¹²⁻¹⁴. Electroencephalographic (EEG) signals recorded over motor cortex
61 provided the input to a classifier, and FES was triggered when features derived from these signals were
62 classified as reflecting attempted movement as opposed to rest. Although starting rehabilitation early
63 post-stroke is associated with better motor outcomes, putatively due to heightened neural
64 plasticity^{4,15,16}, the majority of studies implementing BCI-FES-based rehabilitation focus on patients in
65 the chronic phase^{17,18}. We hypothesized that earlier initiation of BCI-FES would improve
66 corticomuscular functional connectivity, resulting in greater motor recovery. We also aimed to
67 investigate neural correlates of motor recovery in patients receiving BCI-FES to gain a better
68 understanding of potential mechanisms of action. The early phase post-stroke poses challenges in
69 therapy program completion, and heterogenous patient groups with cortical and subcortical stroke,
70 affecting either hemisphere are commonly included. Here we compared outcomes in a BCI and Sham
71 group in a matched lesion subgroup from the Magdeburg patient cohort (German Clinical Trials
72 Register: DRKS00007832; DRKS00011522). BCI and Sham patients had suffered a subcortical stroke
73 affecting the non-dominant hemisphere, and the tight uniformity of the study group enabled group-
74 level comparisons of electrophysiological and behavioral markers over the treatment period.

75 While clinical outcome is the primary focus in the evaluation of rehabilitation measures, understanding
76 the mechanisms underlying recovery is the key to informing further development. Electrophysiological
77 and functional measures of brain activity can provide potential markers of modulation during therapy,
78 which is an essential component of evaluating the potential of this approach in facilitating motor
79 recovery. Brain oscillatory activity¹⁹ and corticomuscular functional connectivity²⁰ have been proposed
80 as biomarkers of post-stroke recovery. Here we compared clinical outcome and neural correlates of
81 motor recovery in patients in the acute and subacute phases post-stroke allocated to BCI-FES therapy
82 (BCI group) or FES delivered without a tight temporal relationship with EEG correlates of movement
83 attempts (Sham group). The patients-underwent a three-week FES, either BCI-controlled or Sham,
84 rehabilitation program, with transcranial magnetic stimulation (TMS)-induced motor evoked potential
85 (MEP) amplitude measurement, and high-density EEG recordings. The EEG analyses included
86 sensorimotor cortical spectral power, corticomuscular coherence, and long-range temporal correlation
87 (LRTC).

88 In the BCI group, movement attempts were detected based on the online classification of EEG signals
89 during the therapy. The sensorimotor rhythm refers to oscillations in brain electrical activity over

90 motor cortical regions in the alpha (8-12 Hz) and beta (13-30 Hz) frequency ranges. Event-related
91 desynchronization and synchronization (ERD/ERS) index reduction/increase of the sensorimotor
92 rhythms, detectable as changes in EEG recordings starting before and changing over the course of
93 movement^{21,22}. They provide well-established indices of actual movement, as well as of imagined
94 movement²³ and movement attempts²⁴ and are commonly used in BCI applications²⁵. EEG recordings
95 of this rhythm were made from each individual patient during a training session during cued movement
96 attempts and rest periods. The electrode locations and specific frequencies at which oscillatory power
97 differences were greatest between movement and rest were selected as features for training the BCI
98 classifier. EEG data were continuously recorded during the subsequent treatment sessions, with
99 repeated online classification, and when a movement attempt was identified, FES to the paretic limb
100 was triggered.

101 The Sham group had the same external set-up at the BCI group, to enable blinding to group allocation.
102 Each Sham group patient was assigned the classifier output stimulation timing and frequency from a
103 randomly selected BCI group participant, to ensure that the only difference between the groups was
104 that the timing of FES in the Sham group was independent of the patient's own cortical activity.

105 Laterality of motor cortical activity plays an important role in post-stroke recovery in two ways. First,
106 handedness has an impact on movement- and imagined-movement-related sensorimotor cortical
107 oscillatory activity as well as fMRI activation in healthy participants²⁶⁻²⁸, and activation patterns during
108 post-stroke rehabilitation differ according to whether the dominant or non-dominant hemisphere is
109 affected^{29,30}. We therefore focused our analyses on the largest possible uniform patient group: right-
110 handed patients with a non-dominant hemisphere stroke. Second, shifts of abnormal bilateral motor
111 area activation during paretic hand movement in the subacute phase toward a more unilateral
112 activation pattern of ipsilesional motor areas in chronic stroke is associated with better motor
113 outcome^{31,32}. On the other hand, while contralesional motor cortical activity is associated with poorer
114 motor outcomes in the chronic phase post-stroke, this activity appears to play an important role early
115 post-stroke³¹. We therefore examined electrophysiological changes over the treatment period both in
116 contralesional and ipsilesional motor cortical regions.

117 The primary outcome measure was change in the Fugl-Meyer Assessment of the Upper Extremity
118 (FMA-UE) score from before to after treatment. We also examined potential neural correlates of a
119 direct effect of BCI-FES on relevant neural processes. The amplitude of TMS-induced MEPs provides an
120 index of the integrity of corticomuscular connectivity, and these were measured before and after the
121 treatment program. Based on delivery of FES in temporal association with movement-associated
122 spectral power changes in the sensorimotor rhythm, we compared spectral power across the alpha
123 and beta frequency ranges after, with that before the treatment program in each group. This included
124 comparison at a group and an individual patient level and evaluating correlation between oscillatory
125 power in the sensorimotor rhythm and the FMA-UE score after treatment. As the aim was re-
126 establishment of corticomuscular functional connectivity, we also assessed change in the EEG-EMG
127 coherence in the same frequency range from before to after treatment in each group, based on the
128 suggestion that this measure could provide a biomarker for motor recovery post-stroke²⁰. We also
129 evaluated a potential impact of BCI-FES on LRTC. LRTC provides an index of correlation between
130 different time periods in a time series, providing an index of the extent to which neuronal systems are
131 at a near-critical state permitting rapid changes in functional connectivity as processing demands
132 change over time³³. LRTC has been postulated to facilitate information transfer in neuronal networks,
133 with physiological memory of a past activity influencing future activity through continuous

134 modification and recurrent interactions between ongoing activity and stimulus-induced changes in
135 activity^{33,34}. Cumulative modification in network functional connectivity, resulting from activity-
136 dependent plasticity, has been proposed to provide the physiological mechanism underlying the power
137 law correlations in ongoing oscillatory neuronal network activity, thus influencing future recruitment
138 of neurons to engage in particular oscillatory activity³³. LRTC observed in EEG signals shows power-law
139 behavior, suggesting that the underlying neurodynamic processes are similar on different time scales³⁴.
140 The amplitude envelope of alpha and beta oscillations displays intermittent fluctuations and power-
141 law decay of the autocorrelation over hundreds of seconds, suggesting a self-organized dynamical
142 critical state³³. Task-relevant neural assemblies, defined by temporal relationships between activity in
143 different brain areas, form and dissolve over time^{35,36}. Sensory stimuli result in reorganization of
144 ongoing endogenous brain dynamics³⁷. As activity is propagated through cortical networks, altering
145 functional connectivity, reflected in changes in LRTC, and influencing future neuronal recruitment,
146 somatosensory stimuli disrupt these transient neural assemblies, degrading ongoing LRTC³³. We
147 hypothesized that tight temporal coupling between motor cortical oscillatory power and the
148 somatosensory stimulus in the BCI group would result in a greater reduction in LRTC than a
149 somatosensory stimulus delivered independently of motor cortical activity corresponding to a
150 movement attempt.

151 **Results**

152 Patients

153 Of the patients recruited in Magdeburg (N = 32), 62.5 % (n = 20) completed the rehabilitation program.
154 The reasons for discontinuing participation were complete recovery (n = 2), finding the therapy too
155 tiring (n = 1), the sequelae of a previously diagnosed psychiatric (n = 4) or physical illness (n = 4), and
156 the patient leaving the region (n = 1). Ten patients were allocated to the BCI group and 10 patients to
157 the Sham group. The analysis was applied to the largest sub-group of patients with similar lesion
158 location, which was those whose non-dominant hemisphere was affected by a subcortical stroke,
159 resulting in equal BCI (n = 6) and Sham (n = 6) group sizes.

160 BCI features

161 The features (electrode locations and spectral power frequencies) that were selected at each re-
162 training of the classifier for the BCI group patients changed over the course of treatment in all patients
163 (Supplementary Fig. 1). Early in the program, bilateral features provided the best classification, with a
164 tendency towards ipsilesional (contralateral) features being selected by the final training of the
165 classifier. While the features included power in both the alpha and beta frequencies throughout, alpha
166 power continued to be relevant by the end of the treatment period. By week 4 or later, all classifiers
167 included an alpha power feature. Only one patient had an ipsilateral beta feature by the end.

168 Clinical evaluation

169 Examining the FMA-UE scores before and after the program, an interaction was observed between
170 *Time* and *Group* ($F(1) = 8.03$, $p = 0.030$) (Fig. 1). No other interactions were significant. A main effect
171 of *Time* was also observed ($F(1,6) = 8.93$, $p = 0.024$). No other within-subject main effects were
172 significant. No between-subject effects were significant. Post hoc pairwise comparisons showed a
173 significant increase in FMA-UE score from pre- to post-treatment in the BCI group ($p = 0.004$) but not
174 in the Sham group ($p = 0.77$). The scores did not differ between the groups pre-treatment ($p = 0.81$),

175 and a trend towards a higher score in the BCI than the Sham group was seen post-treatment ($p =$
176 0.062).

177 Of the secondary clinical outcome measures, a trend towards a *Group x Time* interaction ($F(1) = 4.97,$
178 $p = 0.056$) and a main effect of *Time* ($F(1) = 4.08, p = 0.078$) were only observed for the National
179 Institute of Health (NIHSS) upper limb score.

180 When *Therapy start* (Acute, Subacute) was included as a between-subject factor, the only significant
181 interaction remained *Time x Group* ($F(1) = 6.66, p = 0.049$) (Fig. 2). Post hoc tests showed an increase
182 in FMA-UE score in the BCI group ($p = 0.010$) but not in the Sham group ($p = 0.89$). The FMA-UE score
183 increased in the BCI group from pre- to post-treatment when therapy was started in the acute (within
184 one month of stroke: $p = 0.016$) but not the subacute (one to six months post-stroke: $p = 0.21$) phase,
185 but the increase was not significant in the Sham group, starting in either the acute ($p = 0.94$) or the
186 subacute phase ($p = 0.78$).

187 TMS

188 TMS measurements were available from patients with a subcortical stroke from both groups (BCI: $n =$
189 3; Sham: $n = 3$). An interaction was observed between *Group* and *Time* ($F(1) = 27.69, p = 0.034$) (Fig.
190 3). There was no main effect of *Group* ($F(1,2) = 9.12, p = 0.094$) or *Time* ($F(1,2) = 1.36, p = 0.36$). Post
191 hoc revealed a significant amplitude increase from pre- to post-treatment in the BCI group ($p = 0.012$)
192 only (Sham group: $p = 0.50$).

193 High-density EEG

194 Oscillatory spectral power differed between pre- and post-treatment in the BCI group ($p = 0.036$), with
195 a reduction in upper beta (15-23 Hz) oscillatory spectral power around 0.5 to 1.5 s following the
196 movement cue over the ipsilesional motor cortex (at electrode C2), which was not seen in the Sham
197 group (Fig. 4). Spectral power was compared before and after treatment for each patient on the
198 contralateral (C2) and ipsilateral (C1) side to movement, at the time post-movement cue, at which the
199 pre- to post-movement change was greatest (1.2 to 1.4 s) (Fig. 4). Beta power reduction over the
200 treatment period was most consistent across individuals in the BCI group over the ipsilesional motor
201 cortex, contralateral to movement (Fig. 5). The contralateral beta power after therapy correlated with
202 the FMA-UE score ($r(2) = 0.96, p = 0.044$) (Fig. 5). No significant correlation was observed in the Sham
203 group nor in either group before therapy.

204 LRTC, quantified using the Hurst parameter, was lower after than before the treatment program in the
205 BCI group in the beta frequency range according to pairwise T-tests (Fig. 6A, B). Averaging over the
206 beta frequency range at which power changed over time in the BCI group (15-23 Hz) and over time, a
207 reduction in LRTC was seen in the BCI group only (paired T-tests, BCI: $T = -3.38, p = 0.043$; Sham: $T =$
208 0.19, $p = 0.86$). While LRTC was higher after than before the program in the Sham group in the alpha
209 frequency range (8-12 Hz), averaging over frequency and time, the difference was not significant
210 (paired T-tests, BCI: $T = -0.85, p = 0.46$; Sham: $T = 0.52, p = 0.64$). The reduction in beta-LRTC was
211 consistently observed at an individual patient level in the BCI group only, and the increase in alpha-
212 LRTC was consistently seen in the Sham group only (Fig. 6C, D).

213 We examined EEG–EMG coherence in the time–frequency window in which spectral power changed
214 from pre- to post-treatment in the BCI group (0.5-1.5 s; 15-23 Hz), at the electrode location over the
215 contralateral primary motor cortex at which the power difference was greatest (C2). The EEG–EMG

216 coherence was greater after than before treatment in the BCI group (paired T-test: $T = -3.45$, $p = 0.041$)
217 but not in the Sham group ($T = -0.073$, $p = 0.95$) (Fig. 7).

218 **Discussion**

219 Greater motor recovery, reflected by improved FMA-UE scores, was seen in the group receiving BCI–
220 FES, with stimulation temporally locked to oscillatory spectral power changes in the sensorimotor
221 rhythm, compared to the Sham group, who received FES at times unrelated to oscillatory correlates of
222 movement attempts. Recovery was greater if the BCI–FES therapy was started in the acute phase post-
223 stroke. Neural correlates of improved functional connectivity between contralateral (ipsilesional)
224 motor cortex in the BCI group included greater increases in TMS-induced MEP amplitudes and in
225 corticomuscular coherence in the beta frequency range pre- to post-treatment than in the Sham group.
226 Moreover, movement-associated beta spectral power reduction was more pronounced post-
227 treatment in the BCI than the Sham group, commensurate with a reduction in compensatory activity.
228 Finally, long-range temporal correlation within beta oscillations was also reduced post-treatment in
229 the BCI group, suggesting that a subcritical state could be advantageous to motor recovery. Our
230 findings are consistent with the proposal that FES delivery in a tight temporal window coupled with
231 movement attempts using a BCI could improve post-stroke motor recovery, particularly if started early.
232 Multiple neural correlates of motor recovery were modulated by the treatment program in BCI group,
233 supporting the notion that timing FES delivery according to sensorimotor electrophysiological
234 correlates of movement attempts could have a specific impact on recovery processes.

235 Few studies have investigated the potential impact of using BCI-FES early post-stroke^{13,38,39}. Of the
236 eight patients receiving BCI-FES in a partial crossover design study, four commenced treatment in the
237 subacute phase, from 2-6 months post-stroke, with three in the BCI-FES and one in a control group
238 receiving no FES³⁹. Handedness, hemisphere affected, and lesion location varied. All three BCI patients
239 showed improved motor function after treatment, while the control patient, whose impairment was
240 also the most severe, did not. In another study with a partial crossover design, in which five of the 21
241 patients (mainly with stroke affecting the non-dominant hemisphere, including subcortical and cortical
242 stroke) commenced treatment in the subacute stage, a clinically relevant improvement was seen in
243 three of these patients¹³. A further study, involving seven right-handed patients with mainly subcortical
244 stroke receiving BCI-FES in the acute/subacute phase, also showed greater motor recovery and
245 enhanced sensorimotor rhythm desynchronization on the affected side after BCI-FES, which was not
246 observed in the control group receiving FES unrelated to EEG features³⁸. The improvements following
247 BCI-FES in patients early post-stroke in these studies are consistent with our findings. Moreover, our
248 preliminary analyses indicated that the increases in FMA and in beta desynchronization, as well as the
249 reduction in beta LRTC, were observable in all four BCI group patients individually⁴⁰. The small numbers
250 of patients included at an early stage post-stroke highlights the importance of meta-analyses
251 combining the findings from different studies.

252 Comparing alpha and beta oscillatory power pre- and post-treatment showed an increase in
253 ipsilesional beta desynchronization in the BCI group. On the other hand, alpha power provided more
254 ipsilesional classification features by the end than at the start of the treatment program. Enhanced
255 ipsilateral beta and also alpha desynchronization on motor imagery have been reported following BCI-
256 based neurofeedback training in subacute stroke patients⁴¹. Modulations of alpha and beta power
257 have been postulated to enable selection of task-relevant neural assemblies, with separate roles
258 proposed for alpha and beta oscillations during goal-directed actions⁴². A decrease in contralateral
259 sensorimotor beta power in healthy participants on increasing action selection difficulty was suggested

260 to reflect disinhibition of cortical regions engaged in determining movement parameters, while
261 increased ipsilateral alpha power was proposed to facilitate disengagement of task-unrelated neuronal
262 populations⁴². Reinforcing alpha modulation associated with movement attempts, through providing
263 visual and proprioceptive feedback generated by BCI-FES-induced movement using alpha power as a
264 classifier feature, could have facilitated synaptic strengthening or maintenance of neuronal networks
265 oscillating in the alpha frequency range involved in movement generation. Reducing the selection of
266 beta power features for classification could have reduced the integrity of networks oscillating in the
267 beta range. We note that greater pre-treatment ipsilesional alpha desynchronization has been
268 associated with better outcome in chronic stroke patients, with increased desynchronization over a
269 BCI-training program correlating with greater motor recovery¹⁹. A pre- to post-treatment change in
270 movement-related sensorimotor oscillations in the BCI group here is consistent with a modulatory
271 effect of BCI-FES on the sensorimotor rhythm.

272 We observed a reduction in LRTC in beta oscillations in the BCI group post- compared with pre-
273 treatment. LRTC has been proposed to reflect neuronal systems close to a critical state, allowing fast
274 reorganization of functional neural networks in response to changing demands. Better performance in
275 an attentional task has been found to be associated with lower beta LRTC than at rest, and it was
276 postulated that performance in tasks requiring sustained attention benefits from a sub-critical state⁴³.
277 LRTC in alpha band oscillations is also reduced following perturbation by a stimulus and on
278 movement^{33,44}. LRTC was not examined in the beta band in these studies, however. Our finding of
279 reduced LRTC post-treatment in the BCI group, who had shown better motor recovery than the Sham
280 group, suggests moving to a sub-critical state is associated with improved motor function and could be
281 induced by BCI-FES trained using the sensorimotor rhythm.

282 Pre- with post-treatment comparison of electrophysiological markers differed on the contralateral
283 (ipsilesional) side. An fMRI meta-analysis found that while contralesional motor cortical involvement
284 is common, an eventual predominance of ipsilesional activity is associated with better motor
285 outcome³². While lateralization of sensorimotor activity during post-stroke recovery to the
286 contralesional hemisphere has been associated with better motor outcomes in a cohort including
287 subcortical and cortical stroke patients⁴⁵, better recovery has been reported with ipsilesional
288 lateralization following subcortical stroke^{11,32,46}.

289 The main limitations of this study are associated with the early timing of the intervention and its impact
290 on patient numbers. The most significant recovery post-stroke is seen in the first few weeks³,
291 suggesting that intervention at this time may offer a window period with heightened neural plasticity,
292 potentially enhancing facilitation of motor recovery. However, multiple factors contribute to the
293 limited patient numbers included in BCI studies in early post-stroke patients^{13,38,39}. Extensive
294 investigations and treatments are frequently required on hospital admission, presenting a challenge
295 to study recruitment. Moreover, co-existing medical conditions, often associated with the stroke, in
296 this patient group can impede treatment program completion. Finally, spontaneous post-stroke
297 recovery is most common in the acute phase, in the first days to weeks post-stroke¹³. These limitations
298 are common across centers, underlining the need for multi-center studies and meta-analyses to
299 address the efficacy of rehabilitation approaches in this group.

300 The non-dominant hemisphere was affected in the majority of patients able to participate, due to
301 aphasia being an exclusion criterion. The laterality of brain activity associated with movement depends
302 on whether the dominant or non-dominant side is affected and the handedness of the patient. Group
303 level statistical analyses comparing pre- and post-treatment activity required these factors to be

304 uniform across patients. A tendency to use the non-dominant hand less may impede use-related
305 spontaneous recovery, which could play a role in the benefits seen following BCI-FES in this patient
306 group. Further studies directly comparing groups in whom the non-dominant and dominant
307 hemispheres affected are needed, but again, the group sizes required will necessitate large-scale multi-
308 center patient recruitment to reach the necessary patient numbers in each group.

309 Our findings support the proposal that using a BCI to trigger FES temporally coupled with movement
310 attempts detected in motor cortical oscillations enhances post-stroke motor recovery, especially
311 starting early after stroke. The electrophysiological findings suggest BCI-driven FES supports re-
312 establishment of movement-associated processing on the ipsilesional side and a transition towards a
313 subcritical state as contributing to the mechanism of Hebbian facilitation.

314 **Methods**

315 Patients

316 The patients were a subgroup of the Magdeburg patient cohort in an international, multi-center
317 double-blind, randomized controlled study, which comprised two registered trials with the same study
318 protocol but differing target patient populations. The first trial targeted patients in the acute phase
319 post-stroke (German Clinical Trials Register: DRKS00007832) and the second included patients in the
320 subacute phase (DRKS00011522). Patients were recruited following acute hospital admission post-
321 stroke or on transfer to the rehabilitation center, from the University Hospital Magdeburg stroke ward
322 and the Neurorehabilitation Centre, Median Hospital Magdeburg, Germany, respectively. The study
323 protocol was approved by the Local Ethics Committee of the University Hospital, Magdeburg, Germany
324 and performed in accordance with the principles of the Declaration of Helsinki. All patients discussed
325 study participation and the possibility of withdrawing from the study at any time, without a need to
326 provide a reason, with CMSR, and subsequently provided informed, written consent to participation.

327 *Inclusion criteria*

328 The primary inclusion criterion was upper limb paresis following stroke affecting wrist extension, with
329 a Medical Research Council Power Test score < 3, persisting >24 hours, and still present on recruitment.
330 The acute group was recruited less than 1 month and the subacute group 1-6 months after stroke
331 onset. Patients were required to be a minimum of 18 years of age, with no upper age limit. Diagnosis
332 was confirmed using magnetic resonance imaging (MRI) or computerized tomography (CT), and
333 patients with thrombotic or haemorrhagic stroke were included.

334 *Exclusion criteria*

335 The ability to understand the therapy instructions was a prerequisite, both to fulfill the requirement
336 of provision of informed, written consent, and to enable active participation. Exclusion criteria were
337 therefore a score <25 on the Montreal Cognitive Assessment⁴⁷ or severe aphasia, precluding active
338 discussion of the instructions. Further exclusion criteria were severe hemi-neglect, depression
339 (Hospital Anxiety and Depression Scale: HADS-total >15/21)⁴⁸, fatigue (Fatigue Severity Scale > 36/63,
340 i.e., > 4/7 on 9 items)⁴⁹, pain in the neck/shoulder/arm (Pain Scale > 5/10)⁵⁰, or a history of epilepsy.
341 Other exclusion criteria were medical instability (orthostatic hypotension, sepsis, end-stage renal
342 failure, severe visual impairment, fixed joint contractures, a skin condition that could be worsened
343 through electrode placement), and taking certain regular medication (L-dopa, amantadine).

344

345 BCI-FES

346 On recruitment, patients were pseudorandomly allocated to the BCI or Sham group. The groups were
347 counterbalanced according to the following factors: *Age*, *Sex*, *Lesion Side*, *Lesion Site* (subcortical,
348 cortical), *Lesion Type* (ischaemic, haemorrhagic), and *Pre-treatment FMA*, to control for potential
349 confounding factors. Patients were added sequentially to the database containing these factors and
350 also the factor *Group Allocation*. The first four patients were allocated to the BCI group, so that FES
351 delivery parameters would be available for generating comparable parameters for the Sham group.
352 Frane's algorithm was then applied to the database to determine group allocation. An index of
353 imbalance of each factor among patients so far recruited was calculated, based on each possible group
354 allocation for the next patient. The index was a p-value from testing the hypothesis that the factor did
355 not differ between groups. The Chi-square-goodness-of-fit test was used for *Group Allocation*, the
356 Wilcoxon rank sum test for *Age* and *Pre-treatment FMA*, and the chi-square test for the remaining
357 factors. For each possible group allocation, the largest imbalance was selected and converted to a
358 probability of *Group Allocation* to each group by normalization. With each patient allocation, the most
359 unbalanced factor at that time point was thus considered. The patients, therapists, and evaluating
360 clinicians were blinded to group allocation.

361 Sixteen EEG electrodes were placed bilaterally over motor cortical regions using a customized
362 electrode cap, with electrode positions based on the 10-20 international system as follows: Fz, FC3,
363 FC1, FCz, FC2, FC4, C3, C1, Cz, C2, C4, CP3, CP1, CPz, CP2, and CP4. The reference electrode location
364 was the right mastoid, and the ground electrode was at AFz. Selective electrode coverage was used,
365 as our aim was to base FES timing on motor cortical activity, and the reduced electrode number
366 enabled rapid application, which was important for daily electrode application, to minimize therapist
367 time and maximize compliance. EEG signals were recorded at a sampling rate of 512 Hz using a
368 g.USBamp V2.14.07 amplifier (g.tec, Austria).

369 Patients received a maximum of five sessions per week, with sessions occurring on different days. The
370 total number of possible sessions depended on the length of the patients' stay at the rehabilitation
371 center. All patients received a minimum of three weeks of rehabilitation, and a two-week extension
372 was granted in certain cases by the individual state or private health insurance company. A mean of
373 18.8 [SD 5.7] treatment sessions were performed. Due to the variation, analyses of clinical outcomes
374 were corrected for the number of sessions received.

375 An initial training session was carried out to record EEG data during attempted movement and at rest,
376 which were used to train the classifier. Patients were seated comfortably in front of a computer screen,
377 with a table in front of them on which to rest their forearms, palms down, with flexed elbows. When
378 a green up-arrow was presented, patients were instructed to attempt to extend the wrist of the paretic
379 limb. To provide analogous visual stimulation for both trial types, a red down-arrow was presented
380 when patients were to remain at rest. An upwardly moving bar was presented as visual feedback during
381 movement attempts, and a downwardly moving bar was present during rest. The cue to begin each
382 trial was presented at 0 s. Four to six five-minute blocks were performed. Feature and selection and
383 classification were performed in accordance with the previously reported chronic stroke study¹².
384 Following Laplacian-based spatial filtering, the Welch periodogram was applied to calculate the power
385 spectral density at each electrode in 2 Hz bands from 8-30 Hz in 1 s sliding windows, shifting at 62.5
386 ms intervals (i.e., 16 times per second). Canonical variates analysis was used to identify up to 10
387 features for initial classifier training⁵¹. The trials were labelled as movement attempt or rest to provide
388 input to train the Gaussian classifier using gradient-descent supervised learning. During the therapy,

389 the probability was determined that a particular power spectral density value belonged to the
390 movement attempt or rest trial class. When the classification threshold was not exceeded, a leaky
391 integrator was used to smooth the ongoing output of the classifier. FES was triggered at the time point
392 at which the probabilities integrated over time reached a threshold. If neither class was determined
393 over a maximum 7 s trial, the trial was terminated, and the next one started. EEG data recorded during
394 the therapy sessions were used to retrain the classifier each week, to account for changes over the
395 course of the treatment.

396 Each subsequent therapy session comprised 3-7 blocks, according to fatigue levels, and lasted 10-25
397 minutes, including breaks. Fifteen movement attempts were made per block. For each therapy session,
398 the EEG electrode cap was again applied, and two stimulating electrodes were placed over the
399 extensor digitorum communis of the paretic forearm for inducing or assisting wrist extension by
400 applying FES using a RehaStim stimulation device (Hasomed, Germany). EEG data were recorded
401 continuously, with online classification 16 times per second. When a movement attempt was detected,
402 FES was delivered. To balance the stimulation frequency between the groups, a BCI group patient was
403 arbitrarily selected for each Sham group patient, and the corresponding frequency of stimulation was
404 applied.

405 Clinical evaluation

406 A series of clinical evaluations was made to determine potential differences between the groups in
407 terms of direct physical assessment and also impact on ability to perform daily tasks. The Edinburgh
408 Handedness Inventory (EHI) was used to evaluate handedness.

409 The Fugl-Meyer Assessment upper extremity (FMA-UE) score (max. 66)⁵² was the primary outcome
410 measure. A repeated measures ANOVA with the between-subject factor *Group* (BCI, Sham), the within-
411 subject factor *Time* (pre- and post-treatment), and the covariates *Age*, *Sex*, *Days Post-Stroke*, and *Days*
412 *of Therapy* (i.e., number of treatment sessions) was used to compare the difference between FMA-UE
413 score changes over the program between the groups. A repeated measures ANOVA was also applied
414 including *Therapy start* (acute, subacute) as an additional between-subject factor.

415 A range of secondary endpoints was determined, to enable a detailed exploration of any potential
416 differences between the groups. They included the Medical Research Council Power Test, the
417 Rivermead Test, the Barthel Index, the National Institute of Health Stroke Scale (motor: Arm), the
418 European Stroke Scale, the Modified Ashworth Scale (spasticity), the Goal Attainment Assessment, and
419 the Stroke Impact Scale.

420 TMS

421 TMS was performed as a part of routine clinical monitoring from patients who fulfilled the inclusion
422 and exclusion criteria relating to high magnetic field exposure. Before and after treatment, TMS was
423 delivered to EEG location C4, over the primary motor cortex, while electromyographical signals (EMG)
424 were simultaneously recorded over the affected (left) extensor digitorum communis. TMS was
425 commenced at 70% of capacity and increased repeatedly by 10%, until the maximum motor evoked
426 potential (MEP) amplitude was observed. The change in MEP from before to after treatment was
427 compared between groups using a repeated measures ANOVA, with the between-subject factor *Group*
428 (BCI, Sham) and the within-subject factor *Time* (before, after), correcting for the covariates, *Age* at
429 stroke onset and *Sex*.

430 High-density EEG

431 High-density EEG data were recorded using a BrainAmpDC amplifier (Brain Products GmbH, Germany)
432 from 64 channels (sampling rate: 500 Hz), simultaneously with electromyographic (EMG) data from
433 electrodes placed over extensor digitorum communis of the affected limb during movement attempts,
434 in twelve runs pre- and post-treatment. Each run comprised 10 movement and 5 rest trials in a
435 pseudorandom order. Trials were presented using Presentation software (Version 18.2,
436 Neurobehavioral Systems, Berkeley, CA, USA), analogously to movement cue presentation during the
437 treatment program. The data were analyzed using custom Matlab scripts, EEGLab⁵³, and FieldTrip⁵⁴.
438 Consistent with the clinical analyses, EEG data were analyzed from the patients with a non-dominant
439 hemisphere, subcortical stroke. To enable electrode level comparison, we focused on patients who
440 were purely right-handed (N = 8; BCI: n = 4, Sham: n = 4).

441 The EEG data were checked for internal consistency, then a notch (49–51 Hz) and a bandpass (1-200
442 Hz) filter were applied. The channels from each patient were then visually inspected and marked for
443 ocular, EMG, and other artifacts. If more than 10 % of the data in a given channel were marked, the
444 relevant channel was removed and replaced by spline-interpolated data from neighboring channels.
445 The data were then re-referenced to an average reference, before being epoched according to
446 movement cue presentation (at time = 0 s) with a window of -2 s to 2.998 s (2500 frames). Epochs
447 containing artifacts, determined by visual inspection, were excluded from subsequent analysis by JK
448 and RK, supervised by CMSR. Independent component analysis (ICA) was applied, and components
449 containing eye-blink, eye movement, and muscle artifacts were identified by JK and RK on visual
450 inspection and removed, followed by back-projection of the ICs to the electrode space. The EMG data
451 were epoched with the EEG data but separately notch- and high-pass filtered (10 Hz cut-off), then
452 rectified. The data were further epoched to the times relevant for the subsequent analyses.

453 Time-frequency decomposition was carried out through convolution with 5-cycle Morlet wavelets
454 from 4 to 31 Hz. Change in oscillatory spectral power from pre- to post-treatment was compared for
455 each group. Paired T-tests were applied to each time-frequency point, with a threshold of $p = 0.05$,
456 followed by cluster-based permutation tests with 500 randomizations. We then examined the change
457 in individual patient beta spectral power pre- to post-treatment on an individual level over motor
458 cortex ipsi- and contralateral to movement of the affected hand for each group, followed by calculation
459 of Pearson's correlation coefficient between post-treatment contralateral beta spectral power and
460 FMA-UE.

461 Coherence was calculated between the EMG signal recorded over the extensor digitorum communis
462 during movement attempts and each EEG channel in the time–frequency window (0.5 to 1.5 s, 15-23
463 Hz) at which the pre- to post-treatment spectral power reduction differed between the BCI and sham
464 groups. The EMG and EEG data were Fourier-transformed, with multitaper spectral smoothing, and
465 the cross spectra were calculated based on the phase difference between the EMG and each EEG
466 signal. The change in EEG–EMG coherence from pre- to post-treatment was compared for each group
467 over contralateral motor cortex, at electrode C2, where power modulation was greatest, using paired
468 T-tests.

469 Long-range temporal correlation (LRTC) was calculated using detrended fluctuation analysis (DFA). DFA
470 was developed, because autocorrelation function analyses may yield spurious long-range correlations
471 when the data are non-stationary. Evaluation of the decay in auto-correlation between remote parts
472 of a non-stationary data sequence using DFA⁵⁵ is therefore applicable in EEG data³³. LRTC can be
473 quantified in EEG data in either the time or the frequency domain, the former by fitting the power law

474 to the autocorrelation, and the latter by estimating the slope of the 1/f power spectrum on a log–log
475 scale and computing the scaling exponent. DFA provides a more practical and most common approach
476 to quantifying the degree of temporal dependency in non-stationary signals, captured in the Hurst
477 exponent (H), and has been shown to be consistently related to both of those approaches⁵⁶. In EEG
478 signals, the degree of self-similarity within the time series has previously been quantified based on
479 power law scaling, by applying least squares linear regression to determine the slope of a log–log plot
480 of detrended fluctuations against window size (time scale) to yield $H^{44,55,57}$. LRTC is deemed present
481 when H is between 0.5 and 1.

482 LRTC in alpha and beta oscillations partially overlaps topologically with the distribution of spectral
483 power, and alpha and beta power and LRTC correlate weakly³⁴. We therefore evaluated LRTC at the
484 electrode location at which power differences from pre- to post-treatment differed most between the
485 BCI and Sham groups. The data were time–frequency decomposed using the wavelet transform with 5
486 cycle wavelets, amplitudes were extracted for alpha and beta frequencies (9–30 Hz), and H was
487 calculated in 1 Hz steps. Long signal segments are needed to estimate H in narrowband signals⁵⁸, so
488 we concatenated the movement trials before applying DFA, following the approach of Wairagkar and
489 colleagues⁴⁴, as the DFA scaling exponent is not affected by stitching data together^{57,59}. The minimum
490 available number of trials for a given patient was 20, so 20 sequential trials were concatenated for
491 each patient. The LRTC was then calculated over a 47.5 s sliding window in 50 ms steps, and the LRTC
492 value was assigned to the first time point of each window. Paired T-tests were applied to compare
493 LRTC before and after treatment for each group across time and the alpha and beta frequency ranges,
494 as these frequencies were used as classifier features during the treatment program.

495 **Author contributions statement**

496 CMSR: wrote first draft of manuscript; JK, CR, SP, SV, TH, SD, RTK, HH, HJH, MS, JRM, CMSR: study
497 concept and design; CMSR: study coordination; JK, RK, SV, TH, AS, JL, AH, MG, HJH, SL, MS, CMSR:
498 patient recruitment; CMSR: patient consent; JK, RK, SV, TH, AH, SL, CMSR: clinical evaluation for
499 inclusion; JK, RK, SV, TH, AH, SL: blinded pre- and post-therapy clinical evaluation; SV, TH, AH, MG, HJH,
500 SL, MS: clinical management; AS, JL, AH, SL, MS, CMSR: therapy coordination; CR, SP, JRM, CMSR: group
501 allocation, therapy preparation, and implementation; CR, CMSR: instructed therapists in treatment
502 implementation; SP, JRM: wrote group allocation and classifier software; JK, RK, CMSR:
503 electrophysiological recording; JK, RK, CMSR: electrophysiological and TMS analyses; JK, CR, SJN, ICT,
504 RTK, CMSR: data interpretation. All authors approved the final manuscript.

505 **Competing Interests statement**

506 None declared.

507 **Acknowledgements**

508 The authors would like to thank Anne-Katrin Baum, Manuela Reichwald, and Angelika Klemme for
509 support in data acquisition and Arne Leukert and Luise Ebeling-Jung for carrying out the therapy
510 sessions. We also thank the patients for their participation.

511 **Data availability**

512 The datasets analyzed during the current study are available from the corresponding author on
513 reasonable request.

514 **References**


- 515 1. Kyu, H., Abate, D., Abate, K. & Al., E. Global, regional, and national disability-adjusted life-
516 years (DALYs) for 359 diseases and injuries and healthy life expectancy (HALE) for 195
517 countries and territories, 1990 – 2017: a systematic analysis for the Global Burden of Disease
518 Study 2017. *Lancet* **392**, 1859–1922 (2018).
- 519 2. Lawrence, E. S. *et al.* Estimates of the prevalence of acute stroke impairments and disability in
520 a multiethnic population. *Stroke* **32**, 1279–1284 (2001).
- 521 3. Grefkes, C. & Fink, G. R. Recovery from stroke: current concepts and future perspectives.
522 *Neurol Res Pract* **2**, (2020).
- 523 4. Murphy, T. H. & Corbett, D. Plasticity during stroke recovery: From synapse to behaviour.
524 *Nature Reviews Neuroscience* vol. 10 861–872 Preprint at <https://doi.org/10.1038/nrn2735>
525 (2009).
- 526 5. Takeuchi, N. & Izumi, S. I. Maladaptive plasticity for motor recovery after stroke: mechanisms
527 and approaches. *Neural Plasticity* vol. 2012 Preprint at <https://doi.org/10.1155/2012/359728>
528 (2012).
- 529 6. Su, F. & Xu, W. Enhancing brain plasticity to promote stroke recovery. *Frontiers in Neurology*
530 vol. 11 Preprint at <https://doi.org/10.3389/fneur.2020.554089> (2020).
- 531 7. Soekadar, S. R., Birbaumer, N., Slutzky, M. W. & Cohen, L. G. Brain–machine interfaces in
532 neurorehabilitation of stroke. *Neurobiol Dis* **83**, 172–179 (2015).
- 533 8. Dokkum, L. E. H. Van, Ward, T. & Laffont, I. Brain computer interfaces for neurorehabilitation
534 – its current status as a rehabilitation strategy post-stroke. *Ann Phys Rehabil Med* **58**, 3–8
535 (2015).
- 536 9. Cramer, S. C. *et al.* Harnessing neuroplasticity for clinical applications. *Brain* vol. 134 1591–
537 1609 Preprint at <https://doi.org/10.1093/brain/awr039> (2011).
- 538 10. Takeuchi, N. & Izumi, S. I. Combinations of stroke neurorehabilitation to facilitate motor
539 recovery: Perspectives on Hebbian plasticity and homeostatic metaplasticity. *Front Hum*
540 *Neurosci* **9**, (2015).
- 541 11. Ramos-Murguialday, A. *et al.* Brain-machine-interface in chronic stroke rehabilitation: a
542 controlled study. *Ann. Neurol.* **74**, 100–108 (2014).
- 543 12. Biasucci, A. *et al.* Brain-actuated functional electrical stimulation elicits lasting arm motor
544 recovery after stroke. *Nat Commun* **9**, 1–13 (2018).
- 545 13. Remsik, A. B. *et al.* Behavioral outcomes following brain computer interface intervention for
546 upper extremity rehabilitation in stroke: a randomized controlled trial. *Front Neurosci* **12**, 752
547 (2018).
- 548 14. Wu, Q. *et al.* Brain functional networks study of subacute stroke patients with upper limb
549 dysfunction after comprehensive rehabilitation including BCI training. *Front Neurol* **10**, (2020).
- 550 15. Maulden, S. A., Gassaway, J., Horn, S. D., Smout, R. J. & DeJong, G. Timing of initiation of
551 rehabilitation after stroke. *Arch Phys Med Rehabil* **86**, 34–40 (2005).
- 552 16. Hordacre, B. *et al.* Evidence for a window of enhanced plasticity in the human motor cortex
553 following ischemic stroke. *Neurorehabil Neural Repair* **35**, 307–320 (2021).

- 554 17. Remsik, A. *et al.* A review of the progression and future implications of brain-computer
555 interface therapies for restoration of distal upper extremity motor function after stroke.
556 *Expert Rev Med Devices* **4440**, 17434440.2016.1174572 (2016).
- 557 18. Krueger, J. *et al.* Brain-computer interface-driven functional electrical stimulation for motor
558 rehabilitation following stroke. *Klinische Neurophysiologie* **51**, (2020).
- 559 19. Ray, A. M., Figueiredo, T. D. C., López-Larraz, E., Birbaumer, N. & Ramos-Murguialday, A. Brain
560 oscillatory activity as a biomarker of motor recovery in chronic stroke. *Hum Brain Mapp* **41**,
561 1296–1308 (2020).
- 562 20. Krauth, R. *et al.* Cortico-muscular coherence is reduced acutely post-stroke and increases
563 bilaterally during motor recovery: a pilot study. *Front Neurol* **10**, (2019).
- 564 21. Pfurtscheller, G. & Lopes da Silva, F. Event-related EEG/MEG synchronization and
565 desynchronization: basic principles. *Clin. Neurophysiol.* **110**, 1842–57 (1999).
- 566 22. Sweeney-Reed, C. M. & Nasuto, S. J. Detection of neural correlates of self-paced motor
567 activity using empirical mode decomposition phase locking analysis. *J Neurosci Methods* **184**,
568 54–70 (2009).
- 569 23. Pfurtscheller, G., Brunner, C., Schoegl, A. & Lopes da Silva, F. Mu rhythm (de)synchronization
570 and EEG single-trial classification of different motor imagery tasks. *Neuroimage* **31**, 153–9
571 (2006).
- 572 24. López-Larraz, E. *et al.* Event-related desynchronization during movement attempt and
573 execution in severely paralyzed stroke patients: An artifact removal relevance analysis.
574 *Neuroimage Clin* **20**, 972–986 (2018).
- 575 25. Chaudhary, U., Birbaumer, N. & Curado, M. R. Brain-Machine Interface (BMI) in paralysis. *Ann*
576 *Phys Rehabil Med* **58**, 9–13 (2015).
- 577 26. Vukelić, M., Belardinelli, P., Guggenberger, R., Royter, V. & Gharabaghi, A. Different oscillatory
578 entrainment of cortical networks during motor imagery and neurofeedback in right and left
579 handers. *Neuroimage* **195**, 190–202 (2019).
- 580 27. Stancak Jr., A. & Pfurtscheller, G. The effects of handedness and type of movement on the
581 contralateral preponderance of mu-rhythm desynchronisation. *Electroencephalogr Clin*
582 *Neurophysiol* **99**, 174–182 (1996).
- 583 28. Tzourio Mazoyer, N. *et al.* Between-hand difference in ipsilateral deactivation is associated
584 with hand lateralization: fMRI mapping of 284 volunteers balanced for handedness. *Front*
585 *Hum Neurosci* **9**, (2015).
- 586 29. Gao, J. *et al.* Hemispheric difference of regional brain function exists in patients With acute
587 stroke in different cerebral hemispheres: a resting-state fMRI study. *Front Aging Neurosci* **13**,
588 (2021).
- 589 30. Liew, S. L. *et al.* Laterality of poststroke cortical motor activity during action observation is
590 related to hemispheric dominance. *Neural Plast* **2018**, (2018).
- 591 31. Buetefisch, C. M. Role of the contralesional hemisphere in post-stroke recovery of upper
592 extremity motor function. *Frontiers in Neurology* vol. 6 Preprint at
593 <https://doi.org/10.3389/fneur.2015.00214> (2015).

- 594 32. Rehme, A. K., Eickhoff, S. B., Rottschy, C., Fink, G. R. & Grefkes, C. Activation likelihood
595 estimation meta-analysis of motor-related neural activity after stroke. *Neuroimage* **59**, 2771–
596 2782 (2012).
- 597 33. Linkenkaer-Hansen, K., Nikulin, V., Palva, J., Kalia, K. & Ilmoniemi, R. Stimulus-induced change
598 in long-range temporal correlations and scaling behaviour of sensorimotor oscillations. *Eur. J.*
599 *Neurosci.* **19**, 203–211 (2004).
- 600 34. Nikulin, V. v. & Brismar, T. Long-range temporal correlations in electroencephalographic
601 oscillations: Relation to topography, frequency band, age and gender. *Neuroscience* **130**, 549–
602 558 (2005).
- 603 35. Varela, F., Lachaux, J., Rodriguez, E. & Martinerie, J. The brainweb: phase synchronization and
604 large-scale integration. *Nat. Rev. Neurosci.* **2**, 229–39 (2001).
- 605 36. Sweeney-Reed, C. M. *et al.* Corticothalamic phase synchrony and cross-frequency coupling
606 predict human memory formation. *Elife* **3**, e05352 (2014).
- 607 37. Makeig, S. *et al.* Dynamic brain sources of visual evoked responses. *Science (1979)* **295**, 690–
608 694 (2002).
- 609 38. Li, M. *et al.* Neurophysiological substrates of stroke patients with motor imagery-based brain-
610 computer interface training. *International Journal of Neuroscience* **124**, 403–415 (2014).
- 611 39. Young, B. M. *et al.* Changes in functional brain organization and behavioral correlations after
612 rehabilitative therapy using a brain-computer interface. *Front Neuroeng* **7**, (2014).
- 613 40. Krueger, J. *et al.* Functional electrical stimulation driven by a brain-computer interface in
614 acute and subacute stroke patients impacts beta power and long-range temporal correlation.
615 in *2022 IEEE Workshop on Complexity in Engineering, COMPENG 2022* (Institute of Electrical
616 and Electronics Engineers Inc., 2022). doi:10.1109/COMPENG50184.2022.9905448.
- 617 41. Pichiorri, F. *et al.* Brain-computer interface boosts motor imagery practice during stroke
618 recovery. *Ann Neurol* **77**, 851–865 (2015).
- 619 42. Brinkman, L., Stolk, A., Dijkerman, H. C., De Lange, F. P. & Toni, I. Distinct roles for alpha- and
620 beta-band oscillations during mental simulation of goal-directed actions. *Journal of*
621 *Neuroscience* **34**, 14783–14792 (2014).
- 622 43. Irrmischer, M., Poil, S. S., Mansvelder, H. D., Intra, F. S. & Linkenkaer-Hansen, K. Strong long-
623 range temporal correlations of beta/gamma oscillations are associated with poor sustained
624 visual attention performance. *European Journal of Neuroscience* **48**, 2674–2683 (2018).
- 625 44. Wairagkar, M., Hayashi, Y. & Nasuto, S. J. Dynamics of long-range temporal correlations in
626 broadband EEG during different motor execution and imagery tasks. *Front Neurosci* **15**,
627 660032 (2021).
- 628 45. Young, B. M. *et al.* Changes in functional brain organization and behavioral correlations after
629 rehabilitative therapy using a brain-computer interface. *Front Neuroeng* **7**, 15 (2014).
- 630 46. Tombari, D. *et al.* A longitudinal fMRI study: In recovering and then in clinically stable sub-
631 cortical stroke patients. *Neuroimage* **23**, 827–839 (2004).
- 632 47. Nasreddine, Z. S. *et al.* The Montreal Cognitive Assessment, MoCA: a brief screening tool for
633 mild cognitive impairment. *J Am Geriatr Soc* **53**, 695–699 (2005).


- 634 48. Sagen, U. *et al.* Screening for anxiety and depression after stroke: comparison of the Hospital
635 Anxiety and Depression Scale and the Montgomery and Åsberg Depression Rating Scale. *J*
636 *Psychosom Res* **67**, 325–332 (2009).
- 637 49. Cumming, T. B., Packer, M., Kramer, S. F. & English, C. The prevalence of fatigue after stroke: a
638 systematic review and meta-analysis. *International Journal of Stroke* **11**, 968–977 (2016).
- 639 50. Radbruch, L. *et al.* Validation of the German version of the Brief Pain Inventory. *J Pain*
640 *Symptom Manage* **18**, 180–187 (1999).
- 641 51. Galán, F., Ferrez, P. W., Oliva, F., Guàrdia, J. & Millán, J. D. R. Feature extraction for multi-class
642 BCI using canonical variates analysis. in *2007 IEEE International Symposium on Intelligent*
643 *Signal Processing, WISP* (2007). doi:10.1109/WISP.2007.4447615.
- 644 52. Fugl-Meyer, A., Jääskö, L., Leyman, I., Olsson, S. & Steglind, S. The post-stroke hemiplegic
645 patient. 1. A method for evaluation of physical performance. *Scand J Rehabil Med* **7**, 13–31
646 (1975).
- 647 53. Delorme, A. & Makeig, S. EEGLAB: an open source toolbox for analysis of single-trial EEG
648 dynamics including independent component analysis. *J Neurosci Methods* **134**, 9–21 (2004).
- 649 54. Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J. M. FieldTrip: Open source software for
650 advanced analysis of MEG, EEG, and invasive electrophysiological data. *Comput Intell Neurosci*
651 **156869**, (2011).
- 652 55. Peng, C.-K., Havlin, S., Stanley, H. E. & Goldberger, A. Quantification of scaling exponents and
653 crossover phenomena in nonstationary heartbeat time series. *Chaos* **5**, 82–87 (1995).
- 654 56. Rangarajan, G. & Ding, M. Integrated approach to the assessment of long range correlation in
655 time series data. *Phys. Rev. E Stat. Phys.* **61**, 4991–5001 (2000).
- 656 57. Botcharova, M., Berthouze, L., Brookes, M. J., Barnes, G. R. & Farmer, S. F. Resting state MEG
657 oscillations show long-range temporal correlations of phase synchrony that break down
658 during finger movement. *Front Physiol* **6**, (2015).
- 659 58. Linkenkaer-Hansen, K., Nikouline, V. v., Palva, J. M. & Ilmoniemi, R. J. Long-range temporal
660 correlations and scaling behavior in human brain oscillations. *Journal of Neuroscience* **21**,
661 1370–1377 (2001).
- 662 59. Chen, Z., Ivanov, P. C., Hu, K. & Stanley, H. E. Effect of nonstationarities on detrended
663 fluctuation analysis. *Phys Rev E Stat Phys Plasmas Fluids Relat Interdiscip Topics* **65**, 15 (2002).
- 664

665 **Figure legends**




666

667 **Figure 1.** The Fugl-Meyer Assessment of the upper extremity (FMA-UE) showed greater motor recovery
668 in the BCI group post-treatment than the Sham group. Interaction between *Time* and *Group*: $F(1) =$
669 8.03 , $p = 0.030$, correcting for covariates *Age*, *Sex*, *Days Post-Stroke*, and *Days of Therapy*. Post hoc
670 tests pre- to post-treatment: BCI: $p = 0.004$; Sham: $p = 0.77$. (The maximum score of the FMA-UE is 66
671 points.) Error bars = standard error of the mean.




672

673 **Figure 2.** The improvement in Fugl-Meyer Assessment of the upper extremity (FMA-UE) score was
674 greatest in the BCI group from pre- to post-treatment in patients who started treatment in the acute
675 phase (within one month) post-stroke compared with patients in the Sham group and with patients in
676 either group starting treatment in the subacute phase. Including *Therapy start* as a factor: interaction
677 *Time x Group* ($F(1) = 6.66$, $p = 0.049$) **A.** Patients starting treatment in the acute phase: post hoc $p =$
678 0.016 . **B.** Patients starting treatment in the subacute phase: post hoc $p = 0.78$. Error bars = standard
679 error of the mean.




680

681 **Figure 3.** The amplitude of the transcranial magnetic stimulation (TMS)-induced motor evoked
 682 potentials increased following treatment in the BCI group but not the Sham group. Interaction: *Group*
 683 and *Time* ($F(1) = 27.69$, $p = 0.034$), correcting for covariates. BCI: post hoc $p = 0.012$; Sham: post hoc p
 684 $= 0.050$. TMS was applied at electrode location C2, contralateral to the affected limb. Error bars =
 685 standard error of the mean.



686


687 **Figure 4.** Spectral power pre-treatment minus power post-treatment in each group. Spectral power
 688 reduction was greatest from pre- to post-treatment over contralateral (ipsilesional) primary motor
 689 cortex in the BCI group. Note that the positive T-values indicate a greater in desynchronization post-
 690 than pre-treatment. Black contour = cluster of adjacent time–frequency points at which the post- vs-
 691 pre-treatment power differed according to paired T-tests at threshold $p = 0.05$. **A.** BCI group: at each
 692 electrode. **B.** BCI group: largest cluster observed at electrode C2, over right primary motor cortex. **C.**
 693 Sham group: at each electrode. **D.** Sham group: at electrode C2, over right primary motor cortex.
 694 Cluster-based permutation testing showed a significant difference between spectral power pre- and
 695 post-treatment in the BCI group ($p = 0.036$).




696

697 **Figure 5.** Individual patient beta (15-23 Hz) spectral power at 1.2 to 1.4 s post-movement cue before
 698 before and after treatment. **A.** Over motor cortex ipsilateral to affected hand movement (C1) in the BCI group.
 699 **B.** Over motor cortex contralateral to affected hand movement (C2) in the BCI group. **C.** Over motor
 700 cortex ipsilateral to affected hand movement (C1) in the Sham group. **D.** Over motor cortex
 701 contralateral to affected hand movement (C1) in the Sham group. **E.** Correlation between beta spectral
 702 power ipsilateral to affected hand movement (C2) and FMA-UE after treatment. **F.** Correlation
 703 between beta spectral power contralateral to affected hand movement (C2) and FMA-UE after
 704 treatment ($r(2) = 0.96$, $p = 0.044$). No other correlation was significant.


705



706
 707 **Figure 6.** Changes in long-range temporal correlation (LRTC), quantified using the Hurst parameter, in
 708 high density EEG data recorded after compared with before the therapy program. Averaging over the
 709 beta frequency range at which power decreased post-therapy in the BCI group (15-23 Hz) and over
 710 time, LRTC decreased only in the BCI group (paired T-tests, BCI: $T = -3.38$, $p = 0.043$; Sham: $T = 0.19$, p
 711 $= 0.86$). **A, B.** Significance of the pre- to post-treatment LRTC difference over frequency and time based
 712 on pairwise T-tests. **A.** BCI group. **B.** Sham group. **C, D.** Changes in Hurst parameter in individual
 713 patients. Green: BCI group; Blue: Sham group; Solid lines: significant difference on T-test in this group
 714 and frequency; Dashed lines: difference not significant **C.** At beta (18 Hz). **D.** At alpha (9 Hz).



715
 716 **Figure 7.** EEG-EMG coherence in the beta frequency range (0.5-1.5 s; 15-23 Hz): difference between
 717 pre- and post-therapy. **A.** BCI group (paired T-test: $T = -3.45$, $p = 0.041$). **B.** Sham group (paired T-test:
 718 $T = -0.073$, $p = 0.95$).



719

720

721

722

723

Supplementary figure 1. Classifier features. Features selected for classifiers for patients in the BCI group over the course of the therapy program. Note that classifiers were not trained to determine the timing of functional electrical stimulation in the Sham group. Selected electrodes are highlighted in red, and the frequencies chosen are given below the relevant electrode.