TITLE

Rare variant analyses validate known ALS genes in a multi-ethnic population and
identifies *ANTXR2* as a candidate in PLS

AUTHORS: Tess D. Pottinger[1,2], Joshua E. Motelow[1,3], Gundula Povysil[1], Cristiane A. Martins
Moreno[4], Zhong Ren[1], Hemali Phatnani[5,6,7], The New York Genome Center ALS Sequencing
Consortium, Timothy J. Aitman[8], Javier Santoyo-Lopez[9], Scottish Genomes Partnership, Hiroshi
Mitsumoto[5], ALS COSMOS Study Group, PLS COSMOS Study Group, GTAC Investigators,
David B. Goldstein[1], Matthew B. Harms[1,5,6*]

[1]Institute for Genomic Medicine, Columbia University Irving Medical Center, New York, New
York, United States of America.

[2]Department of Internal Medicine, Columbia University Irving Medical Center, New York, New
York, United States of America.

[3]Department of Pediatrics, Columbia University Irving Medical Center, New York, New York,
United States of America.

[4]Department of Neurology, School of Medicine, Universidade de São Paulo, São Paul, Brazil.

[5]Department of Neurology, Columbia University Irving Medical Center, New York, New York,
United States of America.

[6]Center for Motor Neuron Biology and Disease, Columbia University Irving Medical Center, New
York, New York, United States of America.

[7]New York Genome Center, New York, New York, United States of America.

[8]Centre for Genomic and Experimental Medicine, University of Edinburgh, Edinburgh, Scotland.

[9]Edinburgh Genomics, University of Edinburgh, Edinburgh, Scotland.

34
35

CORRESPONDING AUTHOR
Matthew B. Harms, MD
CUIMC/Neurological Institute of New York
Department of Neurology
710 West 168th Street
New York, NY 10032
Telephone: 212-342-4536
mh3607@cumc.columbia.edu

TOTAL WORD COUNT: 6,704 (including refs/tables/legends)

ABSTRACT

Background: Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disease affecting over 30,000 people in the United States. It is characterized by the progressive decline of the nervous system that leads to the weakening of muscles which impacts physical function. Approximately, 15% of individuals diagnosed with ALS have a known genetic variant that contributes to their disease. As therapies that slow or prevent symptoms, such as antisense oligonucleotides, continue to develop, it is important to discover novel genes that could be targets for treatment. Additionally, as cohorts continue to grow, performing analyses in ALS subtypes, such as primary lateral sclerosis (PLS), becomes possible due to an increase in power. These analyses could highlight novel pathways in disease manifestation.

Methods: Building on our previous discoveries using rare variant association analyses, we conducted rare variant burden testing on a substantially larger cohort of 6,970 ALS patients from a large multi-ethnic cohort as well as 166 PLS patients, and 22,524 controls. We used intolerant domain percentiles based on sub-region Residual Variation Intolerance Score (subRVIS) that have been described previously in conjunction with gene based collapsing approaches to conduct burden testing to identify genes that associate with ALS and PLS.

Results: A gene based collapsing model showed significant associations with *SOD1*, *TARDBP*, and *TBK1* (OR=19.18, p = 3.67×10^{-39}; OR=4.73, p = 2×10^{-10}; OR=2.3, p = 7.49×10^{-9}, respectively). These genes have been previously associated with ALS. Additionally, a significant novel control enriched gene, *ALKBH3* (p = 4.88×10^{-7}), was protective for ALS in this model. An intolerant domain based collapsing model showed a significant improvement in identifying regions in *TARDBP* that associated with ALS (OR=10.08, p = 3.62×10^{-16}). Our PLS protein truncating variant collapsing analysis demonstrated significant case enrichment in *ANTXR2* (p=8.38×10^{-6}).

Conclusions: In a large multi-ethnic cohort of 6,970 ALS patients, rare variant burden testing validated known ALS genes and identified a novel potentially protective gene, *ALKBH3*. A first-ever analysis in 166 patients with PLS found a candidate association with loss-of-function mutations in *ANTXR2*.

Keywords: Amyotrophic lateral sclerosis, ALS, peripheral lateral sclerosis, PLS, Burden testing, Rare-variant analyses

MAIN TEXT

Background

Amyotrophic lateral sclerosis (ALS) is a rare neurodegenerative disease characterized by the progressive loss of upper motor neurons in the cortex and lower motor neurons of the brainstem and spinal cord. Even with FDA-approved disease modifying medication and palliation by artificial nutrition and ventilation, the prognosis is poor and death from accumulating paralysis occurs a median of 32 months after symptoms first manifest(1). Over the last 30 years, genetic study of the 5-10% of ALS patients with family history(2, 3) have securely implicated ~20 monogenic causes and showed possible association to a similar number of genes (https://clinicalgenome.org/affiliation/40096/). Causative mutations in the most prevalent ALS genes (*C9ORF72, SOD1*, *TARDBP*, and *FUS)* explain ~70% of familial ALS(4, 5). Due in part to incomplete penetrance, 10% of simplex ALS cases also carry mutations in these same genes(6).

A paucity of unsolved ALS pedigrees for family studies has intersected with falling sequencing costs for large-scale sequencing to allow gene discovery studies based on rare variant burden or collapsing methods on cohorts using predominantly simplex patients. Since our group first used this approach to implicate *TBK1* and *NEK1,* others have also identified *DNAJC7, TUBA4A* and several candidates(6-9). These analyses utilized the entire gene or recognizable functional domains as the regions for burden testing (6, 7) and were restricted to cohorts with European ancestry, or with less than 5% non-European ALS cases. Recognizing that power for discovery could be improved by a) increasing case and control numbers, b) diversifying the ancestries of participants, and c) collapsing on domains known to be intolerant of variation, we conducted both standard gene and intolerant domain-based collapsing analyses on 6,970 multi-ethnic ALS cases and ancestry-matched controls. Primary lateral sclerosis (PLS) is also a neurodegenerative disease of motor neurons with clinical features, neuropathology, and some genetics that overlaps with ALS(10-12). PLS is nearly always simplex and 20 times

rarer than ALS(13). Because large-scale sequencing studies of ALS often include PLS patients,

we were able to conduct a gene-based collapsing analysis in 166 PLS multi-ethnic cases and

ancestry matched controls.

Methods

Study population

All samples and data came from participants that provided written, informed consent for genetic

studies that had been IRB-approved at each contributing center. The study cohort includes

participants from the Genomic Translation for ALS Care (GTAC study), the Columbia University

Precision Medicine Initiative for ALS, the ALS COSMOS Study Group, the PLS COSMOS Study

Group, the New York Genome Consortium, and the ALS Sequencing Consortium (IRB-

approved genetic studies from Columbia University Medical Center, including the Coriell NINDS

repository), University of Massachusetts at Worchester, Stanford University (including samples

from Emory University School of Medicine, the Johns Hopkins University School of Medicine,

and the University of California, San Diego), Massachusetts General Hospital Neurogenetics

DNA Diagnostic Lab Repository, Duke University, McGill University (including contributions from

Saint-Luc and Notre-Dame Hospital of the Centre Hospitalier de l'Université de Montréal

[CHUM], [University of Montreal]), Gui de Chauliac Hospital of the CHU de Montpellier

(Montpellier University), Pitié Salpêtrière Hospital, Fleurimont Hospital of the Centre Hospitalier

Universitaire de Sherbrooke (CHUS) (University of Sherbrooke), Enfant Jésus Hospital of the

Centre hospitalier affilié universitaire de Québec (CHA) (Laval University), Montreal General

Hospital, Montreal Neurological Institute and Hospital of the McGill University Health Centre, the

University of Edinburgh Scotland, and Washington University in St. Louis (including

contributions from Houston Methodist Hospital, Virginia Mason Medical Center, University of

Utah, and Cedars Sinai Medical Center). Participants were determined to have ALS or PLS by

neuromuscular specialists at tertiary motor neuron disease care centers with expertise in

distinguishing between the two. ALS diagnoses were based on the El Escorial Criteria in all

cases. For PLS, explicit criteria requiring >3 years of symptoms without conversion to ALS were

used for 79 of the 172 PLS participants (14). The criteria used for the remaining PLS diagnoses

were not available.

Controls were selected from >100,000 whole-exome or -genome sequenced individuals

housed in the IGM Data Repository. Individuals with known neurodegenerative disease were

excluded. However, none of the controls were screened for neurodegenerative disease. All

participants consented to the use of DNA in genetic research.

Whole exome and genome sequencing

DNA sequencing was performed at Columbia University, the New York Genome Center, Duke

University, McGill University, Stanford University, HudsonAlpha, and University of

Massachusetts, Worcester. Kits used to conduct whole-exome capture are as follows: Agilent

All Exon kits (50MB, 65MB, and CRE), Nimblegen SeqCap EZ Exome Enrichment kits (V2.0,

V3.0, VCRome, and MedExome), and IDT Exome Enrichment panel. There were 2,185

participants with ALS who were sequenced using Nimblegen SeqCap EZ Exome Enrichment

kits and 51 who were sequenced using the IDT Exome Enrichment panel (**Supplemental Table**

1). While 1,272 controls were evaluated using the Aligent All Exon kits, 8,498 with the IDT

Exome Enrichment panel, and 11,201 with the Nimblegen SeqCap EZ Exome Enrichment kits.

Sequencing was performed using Illumina GAIIx, HiSeq 2000, HiSeq 2500, and NovaSeq 6000

sequencers according to standard protocols. Whole genome sequencing was conducted at the

New York Genome Center and in-house at the IGM. Sample-level BAM files were transferred

from the New York Genome Center to the IGM (n = 3,418). An additional 1,316 genomes were

processed by the IGM. There were 1,553 genomes in our control cohort (**Supplemental Table**

1). Data were aligned to the human reference genome (NCBI Build 37) using DRAGEN (Edico

Genome, San Diego, CA, USA). Picard (http://picard.sourceforge.net) was used to remove

duplicate reads and to process lane-level BAM files to create a sample-level BAM file. GATK

was used to recalibrate base quality scores, realign around indels, and call variants utilizing the

Best Practices recommendations v3.6 (15). Variants were annotated using ClinEff and the

Analysis Tool for Annotated Variants (ATAV), an in-house IGM annotation tool (16). Variants

were annotated with the Genome Aggregation Database (gnomAD) v2.1 frequencies, regional-

intolerance metrics, and the clinical annotations by the Human Gene Mutation Database

(HGMD), ClinVar, and Online Mendelian Inheritance in Man (OMIM). Exonic regions were

retained for downstream statistical analyses.

Sample and variant quality control

Samples reporting >2% contamination according to verifyBamID (17) and those with consensus

coding sequence (CCDS release 20) <90% were excluded from these analyses. KING (18) was

used to test for relatedness. Only unrelated (up to second-degree) individuals were included in

these analyses. For related pairs, samples were chosen to prefer cases. Samples where X:Y

coverage ratios did not match expected sex were excluded.

Only variants within the CCDS or the 2 bp canonical sites were included in these

analyses. These variants were also required to have a quality score of at least 50, a quality by

depth score of at least 5, genotype quality score of at least 20, read position rank sum of at least

−3, mapping quality score of at least 40, mapping quality rank sum greater than −10, and a

minimum coverage of at least 10. SNVs had a maximum Fisher's strand bias of 60, while indels

had a maximum of 200. For heterozygous genotypes, the alternative allele ratio was required to

be greater than or equal to 30%. Only variants with the GATK Variant Quality Score

Recalibration filter "PASS", "VQSRTrancheSNP90.00to99.00", or

"VQSRTrancheSNP99.00to99.90" were included. Variants were excluded if they were marked

by EVS, ExAC, or gnomAD as being failures (http://evs.gs .washington.edu/EVS/).

Clustering, ancestry, and coverage harmonization

A neural network pre-trained on samples of known ancestry was used to calculate probability

estimates for six ancestry groups (African, East Asian, European, Hispanic, Middle Eastern, and

South Asian). Methods for characterizing samples into clusters has been previously described

(19).

To ensure balanced sequencing coverage of evaluated sites between cases and

controls, we imposed a statistical test of independence between the case/control status and

coverage as previously described (20). Sites were removed where the absolute difference in

percentages of cases and controls with at least 10x coverage was greater than 7%. Samples

were then pruned using this method on a cluster-by-cluster basis. Through this approach,

approximately 7- 11% were removed. Clusters with less than 5 participants were not included in

these analyses, thereby removing 6 participants with PLS but none with ALS.

Variant-level statistical analysis

The models that were used to test for associations of nonsynonymous coding or canonical

splice variants with outcome included variants with MAF <0.1% for each population represented

in gnomad and internal AF of <0.1%. Models tested were a standard gene-unit collapsing

analysis, and a domain-unit analysis. The models used for these analyses were previously

described (7). A domain-based approach utilizing sub-region Residual Variation Intolerance

Score (subRVIS) domain percentage(7) with a threshold of 25 was also used to evaluate case

enrichment of rare variants. The full list of 18,653 CCDS genes was analyzed for each model.

Genes with at least one qualifying variant were included for analyses. As we are meta analyzing

across clusters an exact 2-sided Cochran-Mantel-Haenszel test was used (using the statistical

package in R v3.6). Study-wide significance was determined by accounting for 6

nonsynonymous models- multiplicity-adjusted significance threshold $\alpha = 4.9 \times 10^{-7}$

(**Supplemental Table 2**). Model inflation was calculated using empirical (permutation-based)

expected probability distributions as described by Povysil and colleagues (19).

ALS and PLS rare variant burden testing

We conducted both standard gene and intolerant domain-based collapsing analyses on 6,970

multi-ethnic ALS cases (87% European) and 22,534 ancestry-matched controls. Standard gene

collapsing analyses identified case enrichment of rare variants (minor allele frequency of 0.001)

in an ALS cohort with 12 sub-population groups (**Supplemental Figure 1A**) that correspond to

ancestry-based clusters (**Supplemental Figure 1B; Supplemental Table 3**). Analyses were

conducted on clusters with at least 3 cases. Controls were drawn from individuals sequenced

for phenotypes/diseases with no known association with ALS (**Supplemental Table 4**). As

expected, a negative control analysis for rare synonymous variants found no case-enrichment

(**Supplemental Figure 2**). Because gene-based collapsing considers variation across the entire

gene, regions that are tolerant of variation could swamp case-enrichment signals originating

from regions that are intolerant of variation (7). To overcome this limitation, we conducted rare

variant collapsing on domains that are intolerant to variation as defined as a subRVIS domain

score threshold of 25, a cutoff based on threshold testing.

As large-scale sequencing studies of ALS often include PLS patients, we were able to

conduct a gene-based collapsing analysis in 166 PLS multi-ethnic cases (88% European) and

17,695 ancestry matched controls (**Supplemental Figure 3; Supplemental Tables 5**). We

expected the study would be underpowered for securely implicating causative genes but used

this as an opportunity to generate candidates for future study.

ALS gene set enrichment analyses

An ALS gene set enrichment analysis was conducted using the gene strength association list outlined in **Table 1**. We utilized the qualifying variants that were associated with ALS in each gene set category and used the exact two-sided CMH test to analyze burden of ALS genes defined by gene set. These lists were curated using data published by Gregory and colleagues (21). As outlined, "ALS Confirmed" genes were found to have ample published replication evidence, while 'ALS Plus' genes had some replication data and/or functional evidence for an association with ALS. However, 'ALS Replication Needed' genes, required additional replication analyses and/or functional data, and 'ALS Weak Evidence' genes were genes that overlapped with ALS phenotypically.

Results

Rare variant burden testing

Collapsing analysis of all rare functional variants (missense and protein truncating variants) (**Supplemental Table 2**) found genome-wide and study-wide significant (p < 4.9 × 10^{-7}) case-enrichment for known ALS genes *SOD1, TARDBP, TBK1* (OR=19.18, p = 3.67 × 10^{-39}; OR=4.73, p = 2 × 10^{-10}; OR=2.3, p = 7.49 × 10^{-9}, respectively) and control-enrichment for *ALKBH3* (OR=0.26, p = 4.88 × 10^{-7}) (**Figure 1A; Supplemental Data**). Although *SOD1, TBK1* and *TARDBP* are definitive ALS genes, we were intrigued by the identification of control-enriched *ALKBH3.* Control-enrichment was not explained by sequencing methodology, ancestry cluster, or specific phenotype/disease population within the control cohort. Because *ALKBH3* plays a role in DNA repair(22), a mechanism increasingly implicated in ALS pathogenesis(23), we attempted to replicate this novel association by analyzing summary statistics from the Project MinE cohort, which is similar in size to ours (24). None of the available models focused on variation as rare as in our analyses, but at a higher minor allele frequency (MAF) for qualifying variants (0.005), a minor degree of control-enrichment was in

fact observed (OR= 0.56, p = 3.96 × 10^{-4}). This raises the possibility that rare missense and

protein truncating variants (PTVs) in *ALKBH3* could protect from ALS, a finding that requires

validation in large cohorts.

Intolerant domain analyses implicated the same three known ALS genes (*SOD1*,

TARDBP, and *TBK1* at OR=20.63, p = 1.68 × 10^{-38}; OR=10.08, p = 3.62 × 10^{-16}; and OR=3.15,

p = 8.38 × 10^{-11}, respectively) (**Figure 1B; Supplemental Data**). The intolerant domain

analysis did not improve over the gene-based analysis for *SOD1* or *TBK1* (**Figure 2; Figure 3**)

but doubled the odds ratio and significantly lowered the p-value obtained for *TARDBP.* The

improvement of the intolerant domain model (**Figure 1C, 1D**) stemmed from a significant drop

(one-tailed z-score p=0.031) in the number of qualifying variants found in controls dispersed

across tolerant regions, while highlighting qualifying variants in ALS cases predominantly in the

intolerant C-terminal region.

Although most models showed no significant genes, the dominant PTV model showed

significant case enrichment for *ANTXR2* (OR=174.57, p=8.38 x 10^{-6}) (**Figure 4; Supplemental**

Table 6; Supplemental Data), a gene associated with brain connectivity changes and

Alzheimer's disease[25]. Currently, there are no additional large sequencing studies of PLS in

which we could attempt replication.

ALS gene set enrichment analyses

A gene set enrichment analysis of genes that were defined as 'ALS Confirmed' were

significantly associated with the ALS for all dominant models, including PTV only (p = 9.12 × 10^{-}

^{24}), Missense & PTV (p = 6.63 × 10^{-19}), and Missense only (p = 1.03 × 10^{-19}) (**Figure 5**). The

synonymous model, which served as a control, showed no association (p = 0.79) between these

genes and ALS. Genes that are weakly associated with ALS, 'ALS Weak Evidence', showed no

significant enrichment of rare variants for the 4 models that were analyzed. The group of genes

that were described as needing additional replication studies, 'ALS Replication Needed',

showed a significant association with rare variants and ALS for the Missense & PTV model (p =

4.6×10^{-3}). For all other models, rare variants in these genes were not significantly associated

with ALS. An analysis of genes that are characterized as 'ALS Plus' showed no significant

association of rare variants with ALS for the 4 models that were analyzed.

Discussion

Burden testing. Conducting genic and intolerant domain based rare variant burden testing in a

large multi-ethnic population provides insight into novel and established biological mechanisms

in disease manifestations. Additionally, analyzing specific disease subtypes can capture critical

disease pathways that could be targets for clinical intervention. Here we show, that performing

burden testing in multi-ethnic populations and in disease subtypes found novel genetic

associations in individuals diagnosed with ALS and PLS. These analyses implicated ALS genes

that have previously been identified (*SOD1*, *TARDBP*, and *TBK1*). We also identified *ALKBH3*

as a potentially protective gene that warrants further study in additional cohorts. In addition, we

conducted the first rare variant collapsing analysis in PLS, identifying PTVs in *ANTXR2*. This

gene will need to be investigated further in larger PLS cohorts or in targeted functional analyses.

Lastly, gene set enrichment analyses provide evidence that genes known to be associated with

ALS show strong evidence to have a rare variant burden especially for protein truncating

variants.

ALKBH3 associates with ALS. We found that genic burden testing of individuals diagnosed

with ALS identified known risk genes (*SOD1*, *TARDBP*, and *TBK1*) and a novel protective gene

(*ALKBH3*). *ALKBH3* encodes for AlkB homolog 3, Alpha-Ketoglutarate Dependent Dioxygenase

which protects against the cytotoxicity of methylating agents by repair of the specific DNA

lesions (26-28). ALKBH3 potentially acts as a putative hyperactive promotor to suppress

transcription associated DNA damage of highly expressed genes (29). Genes that play a role in

DNA repair and DNA damage response such as *TARDBP*, *FUS*, and *NEK1* (30-33) are known

to play a role in ALS potentially through neuronal death pathways.

ANTXR2 associated with PLS. Genic burden testing of protein truncating variants on

individuals with PLS identified a suggestive gene (*ANTXR2*). *ANTXR2* encodes a receptor for

anthrax toxin that may be involved in extracellular matrix adhesion. Variants in this gene have

been associated with hyaline fibromatosis (34, 35), and has been shown to play a role in

angiogenesis (36). This finding adds to the number angiogenic genes that have been implicated

in ALS including *VEGF* and *ANG* (37).

While we identified a potentially important gene that is associated with PLS, we were

limited in our sample size and will therefore need additional cohorts or functional studies to

further investigate this finding. Additionally, there are potentially more ALS subtypes that could

be investigated to better understand this heterogeneous disease. Lastly, unknown confounders

could be contributing to the signal that are found in these association analyses.

Conclusions

In summary, we performed the largest rare variant analyses of a multi-ethnic population of

patients with ALS to date. Our analysis did not identify new ALS risk genes but demonstrated

that collapsing models informed by regions of intolerance can be useful for identifying genes

where disease-associated variation is limited to regions with low background variation. This

analysis also confirmed the association of the C-terminal domain of *TARDBP*. We also

identified *ALKBH3* as a potentially protective gene that warrants further study in additional and

larger cohorts. Finally, we conducted the first rare variant collapsing analysis in PLS, identifying

PTVs in *ANTXR2* as a candidate disease gene. This association and potential mechanisms for

PTVs in this gene will need to be investigated further in larger PLS cohorts.

It is important to note that this analysis doubled the number of ALS cases and

quadrupled the number of controls from our first study(6) but remained underpowered for the

identification of new ALS genes. A recently published rare variant burden analysis with a similar

number of ALS cases did not identify new genes(24) either, emphasizing the need for

increasingly large genomically characterized ALS cohorts, especially in non-European

populations.

Figure 1. Q-Q plots of gene- and domain-level collapsing of ALL functional coding

variants in ALS cohort. (A) The results for a standard gene-level collapsing of 6,970 ALS

cases and 22,524 controls. P-values were generated using an exact two-sided Cochran-Mantel-

Haenszel (CMH) by gene by cluster. The genes with the top associations that achieved study-

wide significance of $p<4.9 \times 10^{-7}$ (*SOD1* (OR=19.18)*, TARDBP* (OR=4.73)*, TBK1* (OR=2.3)*,* and

ALKBH3 (OR=0.26)) are labeled. *SOD1, TARDBP, TBK1* have been previously implicated in

rare variant association studies of ALS. Yellow and green lines indicate the 2.5[th] and 97.5[th]

percentile of expected p-values, respectively. (B) The results for the domain-based collapsing

restricting qualifying variants to those with subRVIS domain percentage score < 25 of 6,970

cases and 22,524 controls. P-values were generated using an exact two-sided Cochran-Mantel-

Haenszel (CMH) by gene by cluster. The genes with the top associations (*SOD1* (OR=20.63)*,*

TARDBP (OR=10.08)*,* and *TBK1* (OR=3.15)) are labeled. (C) Standard gene-level collapsing

model showed 44 qualifying variants in cases (red circles) and 31 in controls (blue circles) for

TARDBP (D) subRVIS domain collapsing improved association by removing control variants

(cases = 43; controls = 15). Regions with subRVIS domain percentage below 25 are highlighted

in orange while those above this threshold are highlighted in blue. A one tailed z-score showed

that there were significantly less controls in the intolerant domain as indicated by subRVIS

domain percentage score < 25 (p=0.031).

Figure 2. Plot of gene- and domain-level collapsing of ALL *SOD1* functional coding

variants. Standard gene-level collapsing model showed 93 qualifying variants in cases (red

circles) and 18 in controls (blue circles) for *SOD1*. subRVIS domain collapsing improved

association by removing control variants (cases = 90; controls = 16). Regions with subRVIS

domain percentage below 25 are highlighted in orange while those above this threshold are

highlighted in blue. However, a one tailed z-score showed that the differences in the number of

controls in the intolerant domain was not significantly lower than those in the entire gene as

indicated by subRVIS domain percentage score < 25 (p=0.4).

Figure 3. Plot of gene- and domain-level collapsing of ALL *TBK1* functional coding

variants. Standard gene-level collapsing model showed 73 qualifying variants in cases (red

circles) and 143 in controls (blue circles) for *TBK1*. subRVIS domain collapsing improved

association by removing control variants (cases = 47; controls = 72). Regions with subRVIS

domain percentage below 25 are highlighted in orange while those above this threshold are

highlighted in blue. However, a one tailed z-score showed that the differences in the number of

controls in the intolerant domain was not significantly lower than those in the entire gene as

indicated by subRVIS domain percentage score < 25 (p=0.3).

Figure 4. Q-Q plot of gene-level collapsing of protein truncating variants (PTV) in PLS cohort. The results for a standard gene-level collapsing of 166 PLS cases and 17,695 controls. P-values were generated using an exact two-sided Cochran-Mantel-Haenszel (CMH) by gene by cluster. The gene with the top associations that achieved genome-wide significance of p<8.38x10^{-6} (*ANTXR2* (OR=174.57)) is labeled. *ANTXR2* has not been previously implicated in rare variant association studies of PLS. Yellow and green lines indicate the 2.5th and 97.5th percentile of expected p-values, respectively.

Figure 5. Forest plot of ALS genes by model. Rare variants in "ALS Confirmed" genes were

significantly associated with ALS in all gene-based collapsing models except the control

synonymous model. Rare variants in "ALS Plus" genes were associated with ALS in "Missense

& PTV" gene-based collapsing model. There was no association with ALS of rare variants in

"ALS Replication Needed" and "ALS Weak Evidence" genes. Pooled odds ratio, 95%

confidence intervals, and p-values were generated from exact two-sided Cochran-Mantel-

Haenszel (CMH) tests.

Tables

490 **Table 1. ALS gene association strength**

ALS Confirmed	ALS Plus	ALS Replication Needed	ALS Weak Evidence
C9ORF72	*ALS2*	*ANXA11*	*ANG*
FUS	*ATXN2*	*C21ORF2*	*DAO*
NEK1	*CHCHD10*	*CCNF*	*EWSR1*
OPTN	*CHMP2B*	*DNAJC7*	*NEFH*
PFN1	*DCTN1*	*ERBB4*	*PRPH*
SETX	*ERLIN2*	*GLT8D1*	*SQSTM1*
SOD1	*FIG4*	*hnRNPA1*	*SSI8L1*
TARDBP	*GRN*	*hnRNPA2B1*	*TAF15*
TBK1	*KIF5A*	*LGALSL*	
UBQLN2	*MATR3*	*TIA1*	
VAPB	*SIGMAR1*	*TUBA4A*	
	SPG11		
	VCP		

ACKNOWLEDGEMENTS:

We thank the following groups for contributing ALS/PLS samples, sequencing, or clinical data:

New York Genome Center ALS Consortium: J. Kwan, D. Sareen, J.R. Broach, Z. Simmons, X. Arcila-Londono, E.B. Lee, V.M. Van Deerlin, E. Fraenkel, L.W. Ostrow, F. Baas, N. Zaitlen, J.D. Berry, A. Malaspina, P. Fratta, G.A. Cox, L.M. Thompson, S. Finkbeiner, E. Dardiotis, T.M. Miller, S. Chandran, S. Pal, E. Hornstein, D.J. MacGowan, T. Heiman-Patterson, M.G. Hammell, N.A. Patsopoulos, J. Dubnau, and A. Nath.

ALS Exome Sequencing Consortium: S.H. Appel, R.H. Baloh, R.S. Bedlack, R. Brown, W.K. Chung, S. Gibson, J.D. Glass, A. Gitler, D.B. Goldstein, T.M. Miller, R.M. Myers, S.M. Pulst, J.M. Ravits, G. Rouleau, E. Greene, N. Shneider, and W.W. Xin;

*Genomic Translation for ALS Care (GTAC) study***:** S.H. Appel, R.H. Baloh, R.S. Bedlack, S. Chandran, L. Foster, S. Goutman, E. Green, C. Karam, D. Lacomis, G. Manousakis, T.M. Miller, S. Pal, D. Sareen, A. Sherman, Z. Simmons, L. Wang.

ALS COSMOS Study Sites Group: **Columbia University Coordinating Center, NY, NY:** Hiroshi Mitsumoto, MD, DSc, Pam Factor-Litvak, PhD, Regina Santella, PhD, Howard Andrews, PHD; **Texas Neurology, P.A., Dallas, TX:** Daragh Heitzman, MD; **Duke University, Durham, NC:** Richard S. Bedlack, MD, PhD; **California Pacific Medical Center, San Francisco, CA:** Jonathan S. Katz, MD, Robert Miller, MD, Dallas Forshew; **University of Kansas, Kansas City, KS:** Richard J. Barohn, MD, PhD; **Mayo Clinic, Rochester, MN; D**r. Eric J. Sorenson, MD; **University of California - Davis, Sacramento, CA:** Bjorn E. Oskarsson, MD, PhD; **University of Kentucky, Lexington, KY:** Edward J. Kasarskis, MD, PhD; **University of California - San Francisco, San Francisco, CA:** Catherine Lomen-Hoerth, MD, PhD, Jennifer Murphy, PhD; **University of Colorado, Aurora, CO:** Yvonne D. Rollins, MD, PhD; **University of California – Irvine, Orange, CA:** Tahseen Mozaffar, MD; **University of Nebraska, Omaha, NE;** J. Americo M. Fernandes, MD; **University of Iowa, Iowa City, IA:** Andrea J. Swenson, MD; **University of Texas - Southwestern, Dallas:** Sharon P. Nations, MD; **SUNY - Upstate Medical University, Syracuse, NY:** Jeremy M. Shefner, MD, PhD; and **Hospital for Special Care, New Britain, CT:** Jinsy A. Andrews, MD, MS, Dr. Agnes Koczon-Jaremko.

PLS COSMOS Study Group: **Columbia University Irving Medical Center, NY, NY:** Hiroshi Mitsumoto, MD, DSc, Peter L. Nagy, MD, PhD, Pam Factor-Litvak, PhD, PhD, Rejina Santella, PhD, Howard Andrews, PhD, Raymond Goetz, PhD; **Icahn School of Medicine - Mount Sinai, NY, NY:** Chris Gennings, PhD; **University of California - San Francisco, San Fransisco, CA:** Jennifer Murphy, PhD; **National Institute of Neurological Disorders and Stroke, Bethesda, MD:** Mary Kay Floeter, MD, PhD; **University of Kansan Medical Center, Kansas City, KS:** Richard J. Barohn, MD; **University of Texas, Dallus, TX:** Sharon Nations, MD; **Western University, London, Ontario**: Christen Shoesmith, MD; and **University of Kentucky, Louisville, KT**: Edward Kasarskis, MD, PhD.

We thank The Washington Heights–Inwood Columbia Aging Project (WHICAP) for the
contribution of control samples. We also thank the WHICAP study participants and the WHICAP
research and support staff for their contributions to this study: K. Welsh-Bomer, C. Hulette, and
J. Burke; D. Valle, J. Hoover-Fong, and N. Sobriera; A. Poduri; S. Palmer; R. Buckley; K.
Newby; The Murdock Study Community Registry and Biorepository Pro00011196; National
Institute of Allergy and Infectious Diseases Center for HIV/AIDS Vaccine Immunology (CHAVI)
(U19-AI067854); National Institute of Allergy and Infectious Diseases Center for HIV/AIDS
Vaccine Immunology and Immunogen Discovery (UM1-AI100645); CHAVI Funding; R. Ottman;
V. Shashi; S. Berkovic, I. Scheffer, and B. Grinton; The Epi4K Consortium and Epilepsy
Phenome/Genome Project; C. Depondt, S. Sisodiya, G. Cavalleri, and N. Delanty; S. Hirose; C.
Woods, C. Village, K. Schmader, S. McDonald, M. Yanamadala, and H. White; G. Nestadt, J.
Samuels, and Y. Wang; D. Levy; E. Pras, D. Lancet, and Z. Farfel; S. Chen; R. Wapner; C.
Moylan, A. Mae Diehl, and M. Abdelmalek; DUHS (Duke University Health System)
Nonalcoholic Fatty Liver Disease Research Database and Specimen Repository; M. Winn and
R. Gbadegesin; M. Hauser; S. Delaney; A. Need and J. McEvoy; M. Walker; M. Sum;
Undiagnosed Diseases Network; National Institute on Aging (R01AG037212, P01AG007232).

FUNDING SOURCES:

The collection of ALS and PLS samples and data was funded in part by: The Scottish Genomes Partnership (Chief Scientist Office of the Scottish Government Health Directorates (SGP/1) and The Medical Research Council Whole Genome Sequencing for Health and Wealth Initiative (MC/PC/15080); The New York Genome Center ALS Consortium (ALS Association 15-LGCA-234, 19-SI-459, and the Tow Foundation; The GTAC study (ALS Association 16-LGCA-310 and Biogen Idec); ALS Exome Sequencing Consortium (Biogen Idec). Funding for the ALS 561 COSMOS and PLS COSMOS studies was provided by NIEHS R01ES016348, the Muscular 562 Dystrophy Association, MDA Wings Over Wall Street, Spastic Paraplegia Foundation (SPF), private 563 donations from Mr. and Mrs. Marren, the Adams Foundation, and Ride for Life.

The collection of control samples and data was funded in part by: Bryan ADRC NIA P30 AG028377; NIH RO1 HD048805; Gilead Sciences, Inc.; D. Murdock; National Institute of Allergy and Infectious Diseases Center for HIV/AIDS Vaccine Immunology (CHAVI) (U19-AI067854); National Institute of Allergy and Infectious Diseases Center for HIV/AIDS Vaccine Immunology and Immunogen Discovery (UM1-AI100645); Bill and Melinda Gates Foundation; NINDS Award# RC2NS070344; New York-Presbyterian Hospital; The Columbia University College of Physicians and Surgeons; The Columbia University Medical Center; NIH U54 NS078059; NIH P01 HD080642; The J. Willard and Alice S. Marriott Foundation; The Muscular Dystrophy Association; The Nicholas Nunno Foundation; The JDM Fund for Mitochondrial Research; The Arturo Estopinan TK2 Research Fund; UCB; Epilepsy Genetics Initiative, A Signature Program of CURE; Epi4K Gene Discovery in Epilepsy study (NINDS U01-NS077303) and The Epilepsy Genome/Phenome Project (EPGP - NINDS U01-NS053998); The Ellison Medical Foundation New Scholar award AG-NS-0441-08; National Institute Of Mental Health of the National Institutes of Health under Award Number K01MH098126; B57 SAIC-Fredrick Inc. M11-074; OCD Rare 1R01MH097971-01A1. This research was supported in part by funding from Funding from the Duke Chancellor's Discovery Program Research Fund 2014; an American Academy of Child and Adolescent Psychiatry (AACAP) Pilot Research Award; NIMH Grant RC2MH089915; Endocrine Fellows Foundation Grant; The NIH Clinical and Translational Science Award Program (UL1TR000040); NIH U01HG007672; The Washington Heights Inwood Columbia Aging Project; The Stanley Institute for Cognitive Genomics at Cold Spring Harbor Laboratory and the Utah Foundation for Biomedical Research. Data collection and sharing for the WHICAP project (used as controls in this analysis) was supported by The Washington Heights–Inwood Columbia Aging Project (WHICAP, PO1AG07232, R01AG037212, RF1AG054023) funded by the National Institute on Aging (NIA) and by The National Center for Advancing Translational Sciences, National Institutes of Health, through Grant Number UL1TR001873. This manuscript has been reviewed by WHICAP investigators for scientific content and consistency of data interpretation with previous WHICAP Study publications. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

1.   Dorst J, Ludolph AC, Huebers A. Disease-modifying and symptomatic treatment of amyotrophic lateral sclerosis. Ther Adv Neurol Disord. 2018;11:1756285617734734.
2.   Chen S, Sayana P, Zhang X, Le W. Genetics of amyotrophic lateral sclerosis: an update. Mol Neurodegener. 2013;8:28.
3.   Gibson SB, Figueroa KP, Bromberg MB, Pulst SM, Cannon-Albright L. Familial clustering of ALS in a population-based resource. Neurology. 2014;82(1):17-22.
4.   Chia R, Chiò A, Traynor BJ. Novel genes associated with amyotrophic lateral sclerosis: diagnostic and clinical implications. Lancet Neurol. 2018;17(1):94-102.
5.   Boylan K. Familial Amyotrophic Lateral Sclerosis. Neurol Clin. 2015;33(4):807-30.
6.   Cirulli ET, Lasseigne BN, Petrovski S, Sapp PC, Dion PA, Leblond CS, et al. Exome sequencing in amyotrophic lateral sclerosis identifies risk genes and pathways. Science. 2015;347(6229):1436-41.
7.   Gelfman S, Dugger S, de Araujo Martins Moreno C, Ren Z, Wolock CJ, Shneider NA, et al. A new approach for rare variation collapsing on functional protein domains implicates specific genic regions in ALS. Genome Res. 2019;29(5):809-18.
8.   Jih KY, Tsai PC, Tsai YS, Liao YC, Lee YC. Rapid progressive ALS in a patient with a DNAJC7 loss-of-function mutation. Neurol Genet. 2020;6(5):e503.
9.   Smith BN, Ticozzi N, Fallini C, Gkazi AS, Topp S, Kenna KP, et al. Exome-wide rare variant analysis identifies TUBA4A mutations associated with familial ALS. Neuron. 2014;84(2):324-31.
10.   Tartaglia MC, Rowe A, Findlater K, Orange JB, Grace G, Strong MJ. Differentiation between primary lateral sclerosis and amyotrophic lateral sclerosis: examination of symptoms and signs at disease onset and during follow-up. Arch Neurol. 2007;64(2):232-6.
11.   Statland JM, Barohn RJ, Dimachkie MM, Floeter MK, Mitsumoto H. Primary Lateral Sclerosis. Neurol Clin. 2015;33(4):749-60.
12.   Mitsumoto H, Nagy PL, Gennings C, Murphy J, Andrews H, Goetz R, et al. Phenotypic and molecular analyses of primary lateral sclerosis. Neurol Genet. 2015;1(1):e3.
13.   Turner MR, Barohn RJ, Corcia P, Fink JK, Harms MB, Kiernan MC, et al. Primary lateral sclerosis: consensus diagnostic criteria. J Neurol Neurosurg Psychiatry. 2020;91(4):373-7.
14.   Mitsumoto H, Factor-Litvak P, Andrews H, Goetz RR, Andrews L, Rabkin JG, et al. ALS Multicenter Cohort Study of Oxidative Stress (ALS COSMOS): study methodology, recruitment, and baseline demographic and disease characteristics. Amyotroph Lateral Scler Frontotemporal Degener. 2014;15(3-4):192-203.
15.   McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. Genome Res. 2010;20(9):1297-303.
16.   Ren Z, Povysil G, Hostyk JA, Cui H, Bhardwaj N, Goldstein DB. ATAV: a comprehensive platform for population-scale genomic analyses. BMC Bioinformatics. 2021;22(1):149.
17.   Jun G, Flickinger M, Hetrick KN, Romm JM, Doheny KF, Abecasis GR, et al. Detecting and estimating contamination of human DNA samples in sequencing and array-based genotype data. Am J Hum Genet. 2012;91(5):839-48.
18.   Manichaikul A, Mychaleckyj JC, Rich SS, Daly K, Sale M, Chen WM. Robust relationship inference in genome-wide association studies. Bioinformatics. 2010;26(22):2867-73.
19.   Povysil G, Chazara O, Carss KJ, Deevi SVV, Wang Q, Armisen J, et al. Assessing the Role of Rare Genetic Variation in Patients With Heart Failure. JAMA Cardiol. 2021;6(4):379-86.
20.   Petrovski S, Todd JL, Durheim MT, Wang Q, Chien JW, Kelly FL, et al. An Exome Sequencing Study to Assess the Role of Rare Genetic Variation in Pulmonary Fibrosis. Am J Respir Crit Care Med. 2017;196(1):82-93.

21.   Gregory JM, Fagegaltier D, Phatnani H, Harms MB. Genetics of Amyotrophic Lateral
Sclerosis. Current Genetic Medicine Reports. 2020;8(4):121-31.
22.   Kuang W, Jin H, Yang F, Chen X, Liu J, Li T, et al. ALKBH3-dependent m(1)A
demethylation of Aurora A mRNA inhibits ciliogenesis. Cell Discov. 2022;8(1):25.
23.   Konopka A, Atkin JD. DNA Damage, Defective DNA Repair, and Neurodegeneration in
Amyotrophic Lateral Sclerosis. Front Aging Neurosci. 2022;14:786420.
24.   van Rheenen W, van der Spek RAA, Bakker MK, van Vugt J, Hop PJ, Zwamborn RAJ,
et al. Common and rare variant association analyses in amyotrophic lateral sclerosis identify 15
risk loci with distinct genetic architectures and neuron-specific biology. Nat Genet.
2021;53(12):1636-48.
25.   Elsheikh SSM, Chimusa ER, Mulder NJ, Crimi A. Genome-Wide Association Study of
Brain Connectivity Changes for Alzheimer's Disease. Sci Rep. 2020;10(1):1433.
26.   Duncan T, Trewick SC, Koivisto P, Bates PA, Lindahl T, Sedgwick B. Reversal of DNA
alkylation damage by two human dioxygenases. Proc Natl Acad Sci U S A. 2002;99(26):16660-
5.
27.   Bian K, Lenz SAP, Tang Q, Chen F, Qi R, Jost M, et al. DNA repair enzymes ALKBH2,
ALKBH3, and AlkB oxidize 5-methylcytosine to 5-hydroxymethylcytosine, 5-formylcytosine and
5-carboxylcytosine in vitro. Nucleic Acids Res. 2019;47(11):5522-9.
28.   Jiang F, Wang L, Dong Y, Nie W, Zhou H, Gao J, et al. DPPA5A suppresses the
mutagenic TLS and MMEJ pathways by modulating the cryptic splicing of Rev1 and Polq in
mouse embryonic stem cells. Proc Natl Acad Sci U S A. 2023;120(30):e2305187120.
29.   Liefke R, Windhof-Jaidhauser IM, Gaedcke J, Salinas-Riester G, Wu F, Ghadimi M, et
al. The oxidative demethylase ALKBH3 marks hyperactive gene promoters in human cancer
cells. Genome Med. 2015;7(1):66.
30.   Sreedharan J, Blair IP, Tripathi VB, Hu X, Vance C, Rogelj B, et al. TDP-43 mutations in
familial and sporadic amyotrophic lateral sclerosis. Science. 2008;319(5870):1668-72.
31.   Kwiatkowski TJ, Jr., Bosco DA, Leclerc AL, Tamrazian E, Vanderburg CR, Russ C, et al.
Mutations in the FUS/TLS gene on chromosome 16 cause familial amyotrophic lateral sclerosis.
Science. 2009;323(5918):1205-8.
32.   Vance C, Rogelj B, Hortobágyi T, De Vos KJ, Nishimura AL, Sreedharan J, et al.
Mutations in FUS, an RNA processing protein, cause familial amyotrophic lateral sclerosis type
6. Science. 2009;323(5918):1208-11.
33.   Kenna KP, van Doormaal PT, Dekker AM, Ticozzi N, Kenna BJ, Diekstra FP, et al.
NEK1 variants confer susceptibility to amyotrophic lateral sclerosis. Nat Genet.
2016;48(9):1037-42.
34.   Denadai R, Raposo-Amaral CE, Bertola D, Kim C, Alonso N, Hart T, et al. Identification
of 2 novel ANTXR2 mutations in patients with hyaline fibromatosis syndrome and proposal of a
modified grading system. Am J Med Genet A. 2012;158A(4):732-42.
35.   Hanks S, Adams S, Douglas J, Arbour L, Atherton DJ, Balci S, et al. Mutations in the
gene encoding capillary morphogenesis protein 2 cause juvenile hyaline fibromatosis and
infantile systemic hyalinosis. Am J Hum Genet. 2003;73(4):791-800.
36.   Reeves CV, Dufraine J, Young JA, Kitajewski J. Anthrax toxin receptor 2 is expressed in
murine and tumor vasculature and functions in endothelial proliferation and morphogenesis.
Oncogene. 2010;29(6):789-801.
37.   Lambrechts D, Lafuste P, Carmeliet P, Conway EM. Another angiogenic gene linked to
amyotrophic lateral sclerosis. Trends Mol Med. 2006;12(8):345-7.

Supplemental Figures and Tables

Supplemental Figures

Supplemental Figure 1. **UMAP plot of ALS participants** (A) Cluster assignment (B) Predicted
ancestry

757   **Supplemental Figure 2.** Q-Q plots of gene level collapsing in a Synonymous model. Yellow
758   and green lines indicate the 2.5th and 97.5th percentile of expected p-values, respectively. The
759   genomic inflation factor, lambda (λ), is 0.97 indicating no inflation. We generated p values from
760   the exact two-sided Cochran-Mantel-Haenszel (CMH) test by gene by cluster to indicate a
761   different carrier status of affected individuals in comparison to control individuals.

Supplemental Figure 3. **UMAP plot of PLS participants** (A) Cluster assignment (B) Predicted
ancestry

792   **Supplemental Tables**

794   **Supplemental Table 1. Sequencing kits used for cohort**

Sample Type	Capture Kit	Cases	Controls
Exome	Agilent All Exon kits	0	1,272
	Nimblegen SeqCap EZ Exome Enrichment kits	2,185	11,201
	IDT Exome Enrichment panel	51	8,498
Genome		4,734	1,553
Total		6,970	22,524

813 **Supplemental Table 2. Description of collapsing models**

	Synonymous	PTV	Missense&PTV	Missense&PTV subRVIS
Inheritance	Dominant	Dominant	Dominant	Dominant
Functions	Synonymous	PTV	Missense + PTV	Missense + PTV
LOO MAF	0.001	0.001	0.001	0.001
External MAF	0.001	0.001	0.001	0.001
Additional Filters				subRVIS Domain Percentage <25

Supplemental Table 3. Cluster sizes for ALS cohort

Cluster	Ancestry	Cases	Controls	Ratio
0	European1	2,620	4,792	0.55
1	European2	2,502	3,989	0.63
2	European3	922	3,868	0.24
3	Latino1	104	2,788	0.037
4	African1	204	2,102	0.097
5	MiddleEastern1	294	1,924	0.15
6	Latino2	110	1,068	0.1
7	MiddleEastern2	86	669	0.13
8	EastAsian1	35	496	0.071
9	SouthAsian1	40	457	0.088
10	European4	36	265	0.14
11	Admixed1	17	106	0.15
Total		6,970	22,524	0.31

842 **Supplemental Table 4. Phenotypes of control participants**

Control Phenotype	Number of Participants
Kidney and urological disease	8,416 (37.36%)
Healthy family member of a different study	6,297 (27.96%)
Control of different study	3,156 (14.01%)
Epilepsy	2,892 (12.84%)
Obsessive compulsive disorder	721 (3.2%)
Pulmonary disease	403 (1.79%)
Hematological disease	256 (1.13%)
Schizophrenia	161 (0.71%)
Infectious disease	107 (0.48%)
Liver disease	82 (0.36%)
Primary immune deficiency	33 (0.15%)
Total	22,524

Supplemental Table 5. Cluster sizes for PLS cohort

Cluster	Ancestry	Cases	Controls	Ratio
0	European1	90	4,792	0.017
1	European2	19	3,989	0.0047
2	European3	41	3,868	0.012
3	Latino1	1	2,788	0.00039
4	African1	3	2,102	0.0013
5	MiddleEastern1	10	1,924	0.0052
6	Latino2	3	1,068	0.0027
7	MiddleEastern2	1	669	0.0016
8	SouthAsian1	2	496	0.004
9	EastAsian1	1	457	0.002
10	European4	1	265	0.0028
11	Admixed1	0	106	0
Total		172	22,530	0.0076

Supplemental Table 6. Description of PLS participants with *ANTXR2* PTV Qualifying Variants

ID	Sex	Age Range at Onset	Phenotype	Location Hg19	Effect	Depth	MAF	Alleles in Gnomad 2.1
PLS1	M	66-70	Leg onset	4-80990653-A-G	p.Met1?	39x	0.44	0
PLS2	F	41-45	Leg onset	4-80905972-C-T	c.1086+1G>A	29x	0.48	0
PLS3	M	61-65	Leg onset	4-80905984-CA-C	p.Ala359fs	20x	0.35	3
Ctrl1	F	Unk	OCD	4-80957125-C-T	c.697+1G>A	18x	0.67	1
Ctrl2	M	Unk	Kidney disease	4-80977127-TA-T	p.Ser113fs	20x	0.6	0