Endurance and power genes modify the protective effect of fast walking pace on lung cancer risk: a UK Biobank prospective cohort study

Mengying Wang[1,2], Xiangqi Meng[3], Weiliang Tian[4], Ruinan Sun[5], Siyue Wang[2,6], Janice M. Ranson[7], Hexiang Peng[2,6], Valerio Napolioni[8], Patrick W. C. Lau[9], Tao Wu[2,6], Jie Huang[3]

Department of Nutrition and Food Hygiene, School of Public Health, Peking University, Beijing, China

Key Laboratory of Epidemiology of Major Diseases (Peking University), Ministry of Education, China

School of Public Health and Emergency Management, Southern University of Science and Technology, Shenzhen, China

Department of Global Statistics, Eli Lilly and Company, Branchburg, New Jersey, USA

Department of Public and Ecosystem Health, Cornell University College of Veterinary Medicine, Ithaca, NY USA

Department of Epidemiology and Biostatistics, School of Public Health, Peking University, Beijing, China

College of Medicine and Health, University of Exeter, Exeter, UK

School of Biosciences and Veterinary Medicine, University of Camerino, Camerino, Italy

Department of Sport, Physical Education & Health, Hong Kong Baptist University, Kowloon Tong, Hong Kong SAR, China

Declaration of interests: All authors declare no competing interests.

Corresponding to:

Jie Huang, School of Public Health and Emergency Management, Southern University of Science and Technology, Shenzhen, China

E-mail: huangj@sustech.edu.cn

Tao Wu, Department of Epidemiology and Biostatistics, School of Public Health, Peking University, 38 Xueyuan Road, Haidian District, Beijing, China

E-mail: twu@bjmu.edu.cn

Abstract

Background: To investigate the association between walking pace and lung cancer risk, and explore whether the association is modified by endurance and power-related genes.

Methods: We analyzed data from 449,890 participants of the UK Biobank study, without cancer diagnosis at baseline. Data on self-reported walking pace were collected by touchscreen questionnaire at baseline. Blood samples were obtained for genotyping. Hazard ratios (HR) and 95% confidence intervals (CI) were calculated for lung cancer incidence and mortality, with slow walking pace as the reference.

Results: 4,087 lung cancer incident cases and 2,245 lung cancer deaths were identified during a median follow-up period of 12.1 and 12.7 years, respectively. For incident lung cancer, HR (95% CI) were 0.71 (0.65-0.78) and 0.55 (0.49-0.61) among participants with steady and brisk walking pace, respectively. For lung cancer mortality, steady and brisk walking pace were associated with 32% and 48% lower risks, respectively. Associations of walking pace with risks of lung cancer occurrence and mortality were modified by rs1815739 in *ACTN3* and rs7191721 in *RBFOX1*.The protective effect of faster walking pace was more evident among participants carrying a higher number of T allele for rs1815739 in *ACTN3* ($P_{interaction}$=0.04 for both lung cancer incidence and mortality) and A allele for rs7191721 in *RBFOX1* ($P_{interaction}$=0.01 for lung cancer incidence, $P_{interaction}$=0.004 for lung cancer mortality).

Conclusions: Faster walking pace is associated with lower risks of both lung cancer occurrence and mortality, and this protective effect is modulated by polymorphisms in endurance gene *RBFOX1* and in power gene (*ACTN3*).

Keywords: walking pace; endurance gene; power gene; lung cancer

Introduction

Lung cancer is the main contributor to cancer incidence and mortality in the world.[1] Previous studies have shown that healthy lifestyle behaviors are important for the primary prevention and mortality reduction of lung cancer.[2-4] In particular, physical activity has been recognized as a modifiable lifestyle factor for both incidence and prognosis of the disease.[5,6] The frequency and intensity of physical activity could be attributed to physical fitness, which is composed of cardiorespiratory fitness and muscle strength.[7] Therefore, physical fitness is reported to be an important risk factor for both of lung cancer occurrence and mortality.[8,9]

Walking pace is a key indicator of physical fitness. Epidemiological and genetic studies have suggested that walking pace is associated with cardiovascular disease,[7,10] type 2 diabetes,[11] and all-cause mortality.[12] However, the association between habitual walking pace and the risk of lung cancer incidence and mortality is less and inconsistently reported.[8,13] In addition, previous studies have demonstrated that the association between walking pace and cancer risk is likely to be modified by sociodemographic factors and lifestyles.[7,12,14,15] For example, Stamatakis et al. found that the association between walking pace and all-cause mortality was modified by age and physical activity.[12]

Genetic susceptibility also appears to interact with behavioral factors for lung cancer risk.[14,15] Notably, multiple endurance- and power-related genes have been identified in genome-wide association studies (GWAS).[16] For instance, *RBFOX1* is reported to be associated with muscular dystrophies.[17] Furthermore, the R577X variant (rs1815739) in α-actinin-3 (*ACTN3*)

gene is a well-recognized genetic biomarker in exercise studies.[18,19] The C to T base

substitution in *ACTN3* leads to the transformation of an arginine base (R) to a premature stop

codon (X) at amino acid position 577 (R577X).[20] Individuals with the XX genotype (TT)

completely lack the expression of *ACTN3,* which might influence fast fibers in the body.[20,21]

The polymorphism of this gene has been related to speed and cardiometabolic fitness.[19,22] To

the best of our knowledge, no study has evaluated the interaction between these genetic

polymorphisms and walking speed on lung cancer risk.

In the current study, we aimed to investigate the associations between walking pace and risks

of lung cancer occurrence and mortality in a large prospective cohort study. We further

assessed the potential modifying effects of demographics, lifestyles, and endurance- and

power-related genes on this association.

Methods

Study design and participants

The current analysis was based on data from the UK Biobank study. The detailed study

design and procedures were described previously.[23] In brief, over 0.5 million participants

aged 37 to73 years were assessed at one of the 22 centers across England, Wales, and

Scotland from 2006 to 2010. All participants completed a touchscreen questionnaire and

physical measurements as well as provided biological samples at the baseline survey. All

participants provided written informed consent during the baseline visit. Ethical approval of

the UK Biobank study was obtained from the National Information Governance Board for

Health and Social Care and the National Health Service North West Multi-Centre Research

Ethics Committee (Ref 11/NW/0382).

Among 502,618 participants with available data, 449,890 individuals were included in the

final analysis, after excluding those with any type of cancers (N=48,535) or missing data on

self-reported walking pace (N=4,193). For the genetic analysis, only participants with

complete data on genetic polymorphisms were included (N=411,647).

Assessment of outcomes

Lung cancer

Incident lung cancer was derived from hospital admission records via linkage to Health

Episode Statistics in England and Wales and the Scottish Morbidity Records in Scotland.

Incident lung cancer was defined as a hospital admission with International Classification of

Diseases, Tenth Revision (ICD-10) code of C34.

Lung Cancer Deaths

Causes and date of death were identified by using the death certificates held by the National

Health Service Information Center (England and Wales) and the National Health Service

Central Register (Scotland). Lung cancer deaths were ascertained with ICD-10 code of C34.

Assessment of walking pace

The information on self-reported walking pace was collected using the baseline touchscreen

questionnaire. All participants were asked about the question "How would you describe your usual walking pace?", with 'slow', 'steady', and 'brisk' pace as the response options. Slow pace was defined as <3 miles per hour; steady pace was defined as 3 to 4 miles per hour; and brisk pace was defined as >4 miles per hour.

Ascertainment of covariates

All models were adjusted for sociodemographic factors (age, sex, assessment center, ethnicity, UK Biobank assessment center, and Townsend Deprivation Index), lifestyles (smoking, alcohol consumption, diet, and total physical activity), health status (body mass index [BMI], grip strength, hypertension, diabetes, and cardiovascular disease), and family history of lung cancer.

The Townsend Deprivation Index is a comprehensive deprivation indicator comprising unemployment, overcrowded household, non-car ownership, and non-home ownership.[24] Smoking and alcohol consumption status were self-reported as never, former, or current. Dietary factors were assessed using a previously published healthy diet score. One point was given for each favorable dietary factor classified by the corresponding median intake of vegetables, fruits, fish, unprocessed red meat, and processed meat, with the total score ranging from 0 to 5.[25] Physical activity was classified as low, moderate, or high according to the frequency and intensity of walking, moderate activity, vigorous activity, and Metabolic Equivalent Task (MET)-minutes/week based on items from the short International Physical Activity Questionnaire form. The detailed information about physical activity measurements

is available online (https://biobank.ndph.ox.ac.uk/ukb/ukb/docs/ipaq_analysis.pdf). Height

and body weight were measured by trained investigators during the baseline visit. BMI was

calculated by dividing the weight (kg) by the square of the height (m). Grip strength was

measured using a Jamar J00105 hydraulic hand dynamometer. The mean of the three

measurements for each hand was calculated, and the maximum value of the two hands was

included in the analysis. History of hypertension, diabetes, and cardiovascular disease were

defined based on self-reported information and linked medical records at the time of the

baseline visit. Information on family history was also collected from the touchscreen

questionnaire, with family history of lung cancer being defined by an affirmative response to

any of the following three questions: "Has/did your father ever suffer from lung cancer?",

"Has/did your mother ever suffer from lung cancer?", and "Have any of your brothers or

sisters suffered from lung cancer?".

Genotyping

The genotyping process of the UK Biobank study has been described elsewhere.[26] In the

present study, only 411,647 individuals of European descent had available genotyping data

were included in the genetic analysis.

Statistical analyses

In the analysis for the association between walking pace and lung cancer occurrence, survival

time for each participant was calculated as the duration from the response date of baseline

survey through the time of incident lung cancer, death, or the censoring date (7 April 2021),

whichever occurred first. For lung cancer mortality, survival time was calculated as the duration from the baseline survey date through death date or the censoring date, whichever came first. The Cox proportional hazards regression model was adopted to calculate hazard ratios (HR) and 95% confidence intervals (CI) for each outcome. We divided participants into three categories based on their self-reported walking pace: slow, steady, and brisk, with the slow walking pace as the reference group. Several multivariable models were adopted to adjust for the potential confounding factors. In the first model, all HRs were adjusted for sociodemographic factors, namely age (continuous), sex (male, female), race (white European, mixed, South Asian, black, others), UK Biobank assessment center, and Townsend Deprivation Index (continuous). In the second model, we further adjusted for alcohol consumption (current, former, never), smoking status (current, former, never), physical activity (low, moderate, high), and healthy diet score (0, 1, 2, 3, 4, 5). Next, we also adjusted for health indicators including BMI (continuous), grip strength (continuous), hypertension (yes/no), diabetes (yes/no), cardiovascular disease (yes/no), and family history of lung cancer (yes/no). In the genetic analysis, we also adjusted for genotyping batch and the first ten genetic principal components of ancestry. Walking pace was included as a continuous variable when testing for the linear trend. To treat missing covariates, we used a missing indicator for categorical variables and imputed mean values for continuous variables, respectively.

Secondary analyses

Stratified analyses were performed a priori by treating walking pace as a continuous variable

according to age (<65 and ≥65 years), sex (male or female), Townsend Deprivation Index (<-2.1 or ≥-2.1), smoking status (non-current or current), alcohol consumption (non-current or current), physical activity (low, moderate, high), and BMI (18.5-24.9, 25.0-29.9, or ≥30 kg/m^2). In particular, we conducted stratified analyses by the genotype of endurance- and power-related genes. Tests for interaction were carried out by setting variable cross-product terms of walking pace with each category in the third multivariable model, and the likelihood ratio test was adopted to compare models with and without the cross-product terms.

We also conducted several sensitivity analyses to assess the robustness of the results: (1) additionally adjusted for particulate matter with diameters ≤2.5 µm ($PM_{2.5}$) exposure at baseline; (2) further adjusted for body fat-free mass, which was assessed by the sum of the muscle mass from four limbs divided by the squared height (kg/m^2); (3) excluded participants who had hypertension, diabetes, or cardiovascular disease at baseline; (4) excluded events occurring in the first two years of follow-up to minimize potential reverse causality.

All analyses were conducted using SAS software (version 9.4; SAS Institute Inc., Cary, NC, USA). All *P*-values for the tests were two-sided and *P*-values<0.05 were considered as statistically significant.

Patient and Public Involvement

Patients or the public were not involved in the design, or conduct, or reporting, or dissemination plans of our research.

Results

Table 1 shows the baseline characteristics of the study participants. In total, 449,890

participants were included in the primary analysis, among whom 7.9%, 52.7%, and 39.4%

individuals had slow, steady, and brisk walking pace, respectively. The mean age was 56.2

(SD=8.1 years) and 240,135 (53.4%) were females. Participants with faster walking pace had

lower Townsend Deprivation Index and were less likely to be current smokers, but more

likely to drink alcohol, eat healthy foods, or take exercise. Additionally, participants with

faster walking pace had lower BMI and stronger hand grip strength. Hypertension, diabetes,

cardiovascular disease, and family history of lung cancer was less prevalent among

participants with faster walking pace.

During a median of 12.1 years of follow-up (5,349,719 person-years), 4087 lung cancer

incident cases were identified, and 2245 lung cancer deaths were identified during a median

follow-up period of 12.7 years (5,586,963 person-years).

The association between walking pace and lung cancer occurrence is shown in **Table 2.** In

total, 765, 2331, and 991 lung cancer incident cases were identified in slow, steady, and brisk

walking pace groups, respectively. We found that faster walking pace was associated with a

lower risk of incident lung cancer in all multivariable adjusted models. In model 1 adjusted

for age, sex, race, UK Biobank assessment centre, and Townsend Deprivation Index, the HR

(95% CI) was 0.56 (0.52-0.61) of those with steady walking pace and 0.39 (0.35-0.42) of

those with brisk walking pace compared to those with slow walking pace (P trend <0.001).

After further adjustment for alcohol consumption, smoking status, physical activity, and the

healthy diet score, compared with slow walking pace, participants reported steady and brisk

walking pace had HRs of 0.70 (0.64-0.76) and 0.55 (0.50-0.61), respectively (P trend <0.001).

In model 3 with additional adjustment for BMI, hand grip strength, hypertension, diabetes,

cardiovascular diseases, and family history of lung cancer, the corresponding HRs were 0.71

(0.65-0.78) and 0.55 (0.49-0.61) among participants with steady and brisk walking pace,

respectively (P trend <0.001).

A similar association pattern between walking pace and lung cancer mortality was found

(**Table 3**). In total, 498, 1390, and 567 lung cancer deaths were identified in slow, steady, and

brisk walking pace groups, respectively. In model 1, we observed 48% and 66% lower risks

of lung cancer mortality in groups of steady and brisk walking pace when compared with

slow walking pace, respectively (P trend <0.001). In model 3 adjusted for age, sex, race, UK

Biobank assessment centre, Townsend Deprivation Index, alcohol consumption, smoking

status, physical activity, healthy diet score, BMI, hand grip strength, hypertension, diabetes,

cardiovascular diseases, and family history of lung cancer, lung cancer mortality rate was

reduced by 32% and 48% for those with steady and brisk walking pace, respectively when

compared with slow walking pace (P trend <0.001)

We examined the interactions between walking pace and various sociodemographic and

lifestyle factors. The associations of walking pace with lung cancer occurrence and mortality

were largely unchanged across subgroups of participants stratified by sex, age, Townsend Deprivation Index, smoking, alcohol consumption, physical activity, and BMI (**Supplementary Figure 1 and Figure 2**).

We further explored the interaction between walking pace and endurance- and power-related genes on the risk of lung cancer. We found that associations of walking pace with risks of lung cancer occurrence and mortality were modified by rs1815739 in *ACTN3* and rs7191721 in *RBFOX1*. For rs1815739 polymorphisms, the protective HR of faster walking pace was more evident among participants carrying a higher number of T allele ($P_{\text{interaction}}$=0.04 for both lung cancer incidence and mortality). The stratified analysis according to rs1815739 genotype showed that the HRs (95% CIs) of incident lung cancer associated with steady and brisk walking pace were 0.89 (0.75-1.06) and 0.65 (0.53-0.80) among participants with CC genotype; 0.67 (0.59-0.77) and 0.53 (0.46-0.63) among participants with CT genotype; and 0.60 (0.49-0.73) and 0.47 (0.36-0.60) among participants with TT genotype, respectively (**Figure 1**). Similar interaction patterns were observed in the analysis on lung cancer mortality. Within the brisk walking pace group, lung cancer mortality rate was reduced by 41%, 48%, and 57% for carriers of CC, CT, and TT genotype (Supplementary Figure 3). For rs7191721 polymorphisms, the protective HR associated with faster walking pace was stronger among participants carrying a higher number of A allele ($P_{\text{interaction}}$=0.01 for lung cancer incidence, $P_{\text{interaction}}$=0.004 for lung cancer mortality). The subgroup analysis by rs7191721 genotype showed that the HRs (95% CIs) of incident lung cancer associated with steady and brisk walking pace were 0.79 (0.66-0.95) and 0.63 (0.51-0.78) among participants

with GG genotype; 0.74 (0.64-0.84) and 0.57 (0.48-0.66) among participants with GA

genotype; and 0.58 (0.48-0.70) and 0.43 (0.34-0.54) among participants with AA genotype,

respectively (**Figure 2**). Similar interaction patterns were also observed for lung cancer

mortality. In the brisk walking pace group, lung cancer mortality rate was reduced by 40%,

43%, and 65% for GG, GA, and AA genotype carriers, respectively (**Supplementary Figure

4**).

The sensitivity analyses showed that additional adjustment for $PM_{2.5}$ might not appreciably

change the results (**Supplementary Table 1**). Additionally, the associations remained

significant with further adjustment for fat-free mass (**Supplementary Table 2**). Furthermore,

after excluding participants who had hypertension, diabetes, or cardiovascular disease at

baseline, the results were stable (**Supplementary Table 3**). In addition, the results were

largely unchanged after only including participants with a follow-up time of more than two

years (**Supplementary Table 4**).

Discussion

In this prospective cohort study, we observed that faster walking pace was associated with

lower risks of lung cancer occurrence and mortality, independent of lifestyles, BMI, and hand

grip strength. Of note, we found interactions of walking pace with rs1815739 in *ACTN3* and

rs7191721 in *RBFOX1* on incident lung cancer and the disease mortality, where the lower

risk of lung cancer risk associated with faster walking pace was more prominent in T allele

carriers in rs1815739 and A allele carriers in rs7191721, respectively. However, the

associations were not modified by sociodemographic or lifestyle factors.

The associations of walking pace with lung cancer occurrence and mortality have been documented in several epidemiological studies as well as meta-analyses of prospective cohorts, yet conflicting results were reported.[8,13,27,28] Similar to our findings, Smith et al found that slower walking pace was associated with a higher risk of lung cancer mortality compared with faster walking pace.[27] In addition, the results from the 40 years follow-up of the Whitehall study also indicated a significant linear trend, as faster walking pace was associated with a lower risk for lung cancer mortality compared with slower walking pace.[28] However, a recent meta-analysis summarizing the evidence from available cohort studies for walking pace and lung cancer mortality found that slower walking speed were not associated with lung cancer mortality.[8] Notably, another previous analysis in the UK Biobank did not find associations between walking pace and lung cancer occurrence and mortality. Considering the limited number of cases and a shorter follow-up time in the previous analysis,[7] the current analysis might provide more stable results for the association between walking pace and lung cancer risk.

The reasons underlying the observed inverse association between faster walking pace and reduced risks of lung cancer incidence and mortality remain unknown. As suggested by a meta-analysis of randomized controlled trials, brisk walking pace might enhance aerobic fitness,[29] thus to improve respiratory health. A previous study also demonstrates a strong dose-response relationship between walking pace and maximal oxygen uptake.[7] These data

indicate that walking pace is assumed to be an important marker of cardiorespiratory fitness, which has been shown to be associated with reduced lung cancer incidence and mortality.[8] Additionally, exercise including walking might reduce lung cancer risk through preventing inflammatory process.[30]

Walking is the most popular physical activity accepted and accessible by almost the entire population.[31] In addition, among individuals with adequate physical capacity, the change in walking pace is more actionable than an increase in walking time. Therefore, the study has important public health implications for physical activity recommendations in lung cancer prevention and control, suggesting that guidelines should encourage people a faster walking pace to get more benefits in preventing lung cancer and reducing mortality risk.[32]

We did not find evidence for interaction by sex, age, Townsend Deprivation Index, smoking status, alcohol consumption, physical activity, or BMI in the stratified analyses. Interestingly, we observed that there were significant interactions of walking pace with rs1815739 in *ACTN3* and rs7191721 in *RBFOX1* on lung cancer occurrence and mortality, as the inverse association between walking pace and lung cancer risk was stronger in participants carrying T allele in rs1815739 and A allele in rs7191721. The *ACTN3* gene rs1815739 polymorphism occurs in about 16% of people worldwide.[20,33] The muscle proteins of α-actinins 1-4 are important factors regulating the integrity of sarcomeres by binding to actin filaments in skeletal muscle. Furthermore, α-actinin proteins mediate the binding of several glycolytic enzymes to actin filaments.[34,35] α-actinin-3, encoded by *ACTN3* gene is only expressed in fast

fibers.[20,21] Several epidemiological and experimental studies have suggested that rs1815739 polymorphism in *ACTN3* is related to sports performance, especially speed.[18,19,22] *ACTN3* has been recognized as a "speed gene", which is related to both endurance and power.[16,33,36,37] In addition, rs7191721 in *RBFOX1* gene is a marker for aerobic performance and endurance athlete status.[16,38] RNA-binding protein, fox-1 homolog (C. elegans) 1 (RBFOX1) encoded by *RBFOX1* plays a key role in the development of muscle tissues.[39] RNA-binding proteins are important regulators of gene expression, and Rbfox1 knockdown might result in the inhibition of muscle differentiation.[39] These data indicate that the interactions between walking pace and these genes are biologically plausible. Our results may provide insights into the inconsistent observations on the relations between walking pace and lung cancer risk in previous observational studies, in which the genetic modifications were not considered.

Strengths and limitations

The main strengths of our study include the large population-based sample and a wealth of information on sociodemographic factors, lifestyles, and other covariates, which enabled us to conduct comprehensive stratified analyses and sensitivity analyses. Additionally, we for the first time considered the modification effects of the genetic variations related to exercise in the association between walking pace and lung cancer risk. Several potential limitations, however, warrant consideration. Although we controlled for demographic and lifestyle factors in the analyses, unknown residual confounding could still be existing. Walking pace was based on self-reported information, although self-reported walking pace is reported to be a good proxy of gait speed and a well-established predictor of health outcomes.[40] Because the

participants in the genetic analysis were limited to those of European descent, the generalizability of the observed interactions to other populations is unknown. Finally, due to the observational nature of this study, we cannot determine the causality of associations. A clinical trial for a walking pace intervention would be a valuable next step to ascertain causation and clinical relevance.

Conclusions

In this population-based study of adults in the UK, faster walking pace is associated with lower risks of lung cancer occurrence and mortality. The study provides novel evidence that the association was modified by rs1815739 in *ACTN3* and rs7191721 in *RBFOX1*, underscoring the importance of considering genetic modifications in epidemiological studies.

Funding

This work was supported by grants from the China Postdoctoral Science Foundation (Grant No. BX2021021, 2022M710249), Fujian Provincial Health Technology Project (Grant No. 2020CXB009), and the Natural Science Foundation of Fujian Province, China (Grant No. 2021J01352).

Authorships

MYW: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Software; Visualization; Roles/Writing - original draft; Writing - review & editing.

JH and TW: Conceptualization; Funding acquisition; Project administration; Software;

Supervision; Writing - review & editing.

XM, WT, HXP, SYW, JMR, and VN: Investigation; Methodology; Software; Supervision; Validation; Writing - review & editing.

All authors approved the final version of the manuscript.

Acknowledgements

This research was conducted using the UK Biobank resources under application 66137. We thank the participants for sharing their heath related information. Dr. Janice M. Ranson is supported by Alzheimer's Research UK and the Alan Turing Institute/Engineering and Physical Sciences Research Council.

Data Availability

Data for this study are available by request on the UK Biobank website (https://biobank.ctsu.ox.ac.uk/).

References

1 Bray, F. *et al.* Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* **68**, 394-424, doi:10.3322/caac.21492 (2018).

2 Arem, H. *et al.* Physical activity and cancer-specific mortality in the NIH-AARP Diet and Health Study cohort. *International journal of cancer* **135**, 423-431, doi:10.1002/ijc.28659 (2014).

3 Wei, X. *et al.* Diet and Risk of Incident Lung Cancer: A Large Prospective Cohort Study in UK

Biobank. *The American journal of clinical nutrition* **114**, 2043-2051, doi:10.1093/ajcn/nqab298 (2021).

Yang, J. J. *et al.* Association of Dietary Fiber and Yogurt Consumption With Lung Cancer Risk: A Pooled Analysis. *JAMA Oncol* **6**, e194107, doi:10.1001/jamaoncol.2019.4107 (2020).

Liu, Y., Li, Y., Bai, Y. P. & Fan, X. X. Association Between Physical Activity and Lower Risk of Lung Cancer: A Meta-Analysis of Cohort Studies. *Frontiers in oncology* **9**, 5, doi:10.3389/fonc.2019.00005 (2019).

Messina, G. *et al.* The Beneficial Effects of Physical Activity in Lung Cancer Prevention and/or Treatment. *Life (Basel)* **12**, doi:10.3390/life12060782 (2022).

Yates, T. *et al.* Association of walking pace and handgrip strength with all-cause, cardiovascular, and cancer mortality: a UK Biobank observational study. *European heart journal* **38**, 3232-3240, doi:10.1093/eurheartj/ehx449 (2017).

Lee, J. Cardiorespiratory Fitness, Physical Activity, Walking Speed, Lack of Participation in Leisure Activities, and Lung Cancer Mortality: A Systematic Review and Meta-Analysis of Prospective Cohort Studies. *Cancer Nurs* **44**, 453-464, doi:10.1097/NCC.0000000000000847 (2021).

Vainshelboim, B., Lima, R. M., Kokkinos, P. & Myers, J. Cardiorespiratory Fitness, Lung Cancer Incidence, and Cancer Mortality in Male Smokers. *American journal of preventive medicine* **57**, 659-666, doi:10.1016/j.amepre.2019.04.020 (2019).

Timmins, I. R. *et al.* Genome-wide association study of self-reported walking pace suggests beneficial effects of brisk walking on health and survival. *Commun Biol* **3**, 634, doi:10.1038/s42003-020-01357-7 (2020).

Boonpor, J., Ho, F. K., Gray, S. R. & Celis-Morales, C. A. Association of Self-reported Walking Pace With Type 2 Diabetes Incidence in the UK Biobank Prospective Cohort Study. *Mayo Clin Proc* **97**, 1631-1640, doi:10.1016/j.mayocp.2022.02.028 (2022).

Stamatakis, E. *et al.* Self-rated walking pace and all-cause, cardiovascular disease and cancer mortality: individual participant pooled analysis of 50 225 walkers from 11 population British cohorts. *Br J Sports Med* **52**, 761-768, doi:10.1136/bjsports-2017-098677 (2018).

Celis-Morales, C. A. *et al.* Walking Pace Is Associated with Lower Risk of All-Cause and Cause-Specific Mortality. *Med Sci Sports Exerc* **51**, 472-480, doi:10.1249/MSS.0000000000001795 (2019).

You, D. *et al.* Associations of genetic risk, BMI trajectories, and the risk of non-small cell lung cancer: a population-based cohort study. *BMC medicine* **20**, 203, doi:10.1186/s12916-022-02400-6 (2022).

Huang, Y. *et al.* Air Pollution, Genetic Factors, and the Risk of Lung Cancer: A Prospective Study in the UK Biobank. *Am J Respir Crit Care Med* **204**, 817-825, doi:10.1164/rccm.202011-4063OC (2021).

Varillas-Delgado, D. *et al.* Genetics and sports performance: the present and future in the identification of talent for sports based on DNA testing. *Eur J Appl Physiol* **122**, 1811-1830, doi:10.1007/s00421-022-04945-z (2022).

Ahmetov, II & Fedotovskaya, O. N. Current Progress in Sports Genomics. *Adv Clin Chem* **70**, 247-314, doi:10.1016/bs.acc.2015.03.003 (2015).

Wang, Z. *et al.* Genome-wide association analyses of physical activity and sedentary behavior provide insights into underlying mechanisms and roles in disease prevention. *Nat Genet* **54**,

1332-1344, doi:10.1038/s41588-022-01165-1 (2022).

Pickering, C. & Kiely, J. ACTN3: More than Just a Gene for Speed. *Front Physiol* **8**, 1080, doi:10.3389/fphys.2017.01080 (2017).

North, K. N. *et al.* A common nonsense mutation results in alpha-actinin-3 deficiency in the general population. *Nature genetics* **21**, 353-354, doi:10.1038/7675 (1999).

MacArthur, D. G. *et al.* Loss of ACTN3 gene function alters mouse muscle metabolism and shows evidence of positive selection in humans. *Nature genetics* **39**, 1261-1265, doi:10.1038/ng2122 (2007).

Deschamps, C. L. *et al.* The ACTN3 R577X Polymorphism Is Associated with Cardiometabolic Fitness in Healthy Young Adults. *PloS one* **10**, e0130644, doi:10.1371/journal.pone.0130644 (2015).

Sudlow, C. *et al.* UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS Med* **12**, e1001779, doi:10.1371/journal.pmed.1001779 (2015).

Adams, J., Ryan, V. & White, M. How accurate are Townsend Deprivation Scores as predictors of self-reported health? A comparison with individual level data. *J Public Health (Oxf)* **27**, 101-106, doi:10.1093/pubmed/fdh193 (2005).

Pazoki, R. *et al.* Genetic Predisposition to High Blood Pressure and Lifestyle Factors: Associations With Midlife Blood Pressure Levels and Cardiovascular Events. *Circulation* **137**, 653-661, doi:10.1161/CIRCULATIONAHA.117.030898 (2018).

Bycroft, C. *et al.* Genome-wide genetic data on ~500,000 UK Biobank participants [published online July 20, 2017]. *bioRxiv*, doi:10.1101/166298 (2017).

Davey Smith, G., Shipley, M. J., Batty, G. D., Morris, J. N. & Marmot, M. Physical activity and cause-specific mortality in the Whitehall study. *Public Health* **114**, 308-315, doi:10.1038/sj.ph.1900675 (2000).

Batty, G. D., Shipley, M. J., Kivimaki, M., Marmot, M. & Davey Smith, G. Walking pace, leisure time physical activity, and resting heart rate in relation to disease-specific mortality in London: 40 years follow-up of the original Whitehall study. An update of our work with professor Jerry N. Morris (1910-2009). *Ann Epidemiol* **20**, 661-669, doi:10.1016/j.annepidem.2010.03.014 (2010).

Murphy, M. H., Nevill, A. M., Murtagh, E. M. & Holder, R. L. The effect of walking on fitness, fatness and resting blood pressure: a meta-analysis of randomised, controlled trials. *Preventive medicine* **44**, 377-385, doi:10.1016/j.ypmed.2006.12.008 (2007).

Lin, Y. Y., Rau, K. M. & Lin, C. C. Longitudinal study on the impact of physical activity on the symptoms of lung cancer survivors. *Support Care Cancer* **23**, 3545-3553, doi:10.1007/s00520-015-2724-7 (2015).

Dai, S. *et al.* Participation in Types of Physical Activities Among US Adults--National Health and Nutrition Examination Survey 1999-2006. *J Phys Act Health* **12 Suppl 1**, S128-140, doi:10.1123/jpah.2015-0038 (2015).

Stamatakis, E. *et al.* Infographic. Self-rated walking pace and all-cause, cardiovascular disease and cancer mortality: individual participant pooled analysis of 50 225 walkers from 11 population British cohorts. *Br J Sports Med* **53**, 1381-1382, doi:10.1136/bjsports-2018-100468 (2019).

Berman, Y. & North, K. N. A gene for speed: the emerging role of alpha-actinin-3 in muscle

metabolism. *Physiology (Bethesda)* **25**, 250-259, doi:10.1152/physiol.00008.2010 (2010).

Sjoblom, B., Salmazo, A. & Djinovic-Carugo, K. Alpha-actinin structure and regulation. *Cell Mol Life Sci* **65**, 2688-2701, doi:10.1007/s00018-008-8080-8 (2008).

Mills, M. *et al.* Differential expression of the actin-binding proteins, alpha-actinin-2 and -3, in different species: implications for the evolution of functional redundancy. *Hum Mol Genet* **10**, 1335-1346, doi:10.1093/hmg/10.13.1335 (2001).

MacArthur, D. G. & North, K. N. A gene for speed? The evolution and function of alpha-actinin-3. *Bioessays* **26**, 786-795, doi:10.1002/bies.20061 (2004).

Chan, S. *et al.* A gene for speed: contractile properties of isolated whole EDL muscle from an alpha-actinin-3 knockout mouse. *Am J Physiol Cell Physiol* **295**, C897-904, doi:10.1152/ajpcell.00179.2008 (2008).

Ahmetov, I. *et al.* Genome-wide association study identifies three novel genetic markers associated with elite endurance performance. *Biol Sport* **32**, 3-9, doi:10.5604/20831862.1124568 (2015).

Fogel, B. L. *et al.* RBFOX1 regulates both splicing and transcriptional networks in human neuronal development. *Hum Mol Genet* **21**, 4171-4186, doi:10.1093/hmg/dds240 (2012).

Syddall, H. E., Westbury, L. D., Cooper, C. & Sayer, A. A. Self-reported walking speed: a useful marker of physical performance among community-dwelling older people? *Journal of the American Medical Directors Association* **16**, 323-328, doi:10.1016/j.jamda.2014.11.004 (2015).

Table 1. Baseline characteristics of participants according to walking pace in the UK

Biobank study

Characteristics	Walking pace		
	Slow	Steady	Brisk
N	35 450	237 271	177 169
Age (years)	58.5	56.8	54.9
Sex, male (%)	45.6	46.4	47.1
Race, white (%)	88.3	93.4	96.2
Townsend Deprivation Index	0.1	-1.3	-1.6
Current smokers (%)	17.7	10.9	8.8
Current drinkers (%)	81.3	91.9	94.3
Physical activity level (%)			
Low	31.2	15.5	11.1
Moderate	25.7	33.1	34.0
High	14.0	30.2	39.7
Healthy diet score			
0-1	16.1	13.2	9.9
2-3	52.7	50.7	46.0
4-5	31.3	36.2	44.1
Body mass index (kg/m^2)	31.2	28.0	25.9
Hand grip strength (kg)	28.3	32.5	34.4
Hypertension history (%)	68.9	57.0	46.8
Diabetes history (%)	15.3	5.6	2.6
Cardiovascular diseases history (%)	20.7	7.0	3.8
Family history of lung cancer (%)	14.8	12.6	11.2

Table 2. Adjusted HRs and 95% CI for walking pace with the risk of incident lung cancer in the UK Biobank study

	Walking pace			
	Slow	Steady	Brisk	*P* for trend
Cases/N	765/35 450	2331/237 271	991/177 169	
Model 1 [a]	1.00	0.56 (0.52-0.61)	0.39 (0.35-0.42)	<0.001
Model 2 [b]	1.00	0.70 (0.64-0.76)	0.55 (0.50-0.61)	<0.001
Model 3 [c]	1.00	0.71 (0.65-0.78)	0.55 (0.49-0.61)	<0.001

[a] Adjusted for age (continuous, years), sex (male, female), race (white European, mixed, South Asian, black, others), UK Biobank assessment centre, and Townsend Deprivation Index (continuous).

[b] Adjusted for age (continuous, years), sex (male, female), race (white European, mixed, South Asian, black, others), UK Biobank assessment centre, Townsend Deprivation Index (continuous), alcohol consumption (current, former, never), smoking status (current, former, never), physical activity (low, moderate, high), and healthy diet score (0, 1, 2, 3, 4, 5).

[c] Adjusted for age (continuous, years), sex (male, female), race (white European, mixed, South Asian, black, others), UK Biobank assessment centre, Townsend Deprivation Index (continuous), alcohol consumption (current, former, never), smoking status (current, former, never), physical activity (low, moderate, high), healthy diet score (0, 1, 2, 3, 4, 5), body mass index (continuous, kg/m^2), Hand grip strength (continuous, kg), baseline hypertension (yes/no), baseline diabetes (yes/no), baseline cardiovascular diseases (yes/no), and family history of lung cancer (yes/no).

Table 3. Adjusted HRs and 95% CI for walking pace with the lung cancer mortality in the UK Biobank study

	Walking pace			
	Slow	Steady	Brisk	*P* for trend
Cases/N	498/35 450	1390/237 271	567/177 169	
Model 1 [a]	1.00	0.52 (0.47-0.58)	0.34 (0.30-0.39)	<0.001
Model 2 [b]	1.00	0.68 (0.61-0.76)	0.52 (0.45-0.59)	<0.001
Model 3 [c]	1.00	0.68 (0.61-0.76)	0.52 (0.45-0.59)	<0.001

[a] Adjusted for age (continuous, years), sex (male, female), race (white European, mixed, South Asian, black, others), UK Biobank assessment centre, and Townsend Deprivation Index (continuous).

[b] Adjusted for age (continuous, years), sex (male, female), race (white European, mixed, South Asian, black, others), UK Biobank assessment centre, Townsend Deprivation Index (continuous), alcohol consumption (current, former, never), smoking status (current, former, never), physical activity (low, moderate, high), and healthy diet score (0, 1, 2, 3, 4, 5).

[c] Adjusted for age (continuous, years), sex (male, female), race (white European, mixed, South Asian, black, others), UK Biobank assessment centre, Townsend Deprivation Index (continuous), alcohol consumption (current, former, never), smoking status (current, former, never), physical activity (low, moderate, high), healthy diet score (0, 1, 2, 3, 4, 5), body mass index (continuous, kg/m^2), Hand grip strength (continuous, kg), baseline hypertension (yes/no), baseline diabetes (yes/no), baseline cardiovascular diseases (yes/no), and family history of lung cancer (yes/no).

Figure 1. The association between walking pace and the risk of incident lung cancer stratified by rs1815739 polymorphism in *ACTN3* gene. Adjusted for age (continuous, years), sex (male, female), race (white European, mixed, South Asian, black, others), UK Biobank assessment centre, Townsend Deprivation Index (continuous), alcohol consumption (current, former, never), smoking status (current, former, never), physical activity (low, moderate, high), healthy diet score (0, 1, 2, 3, 4, 5), body mass index (continuous, kg/m^2), Hand grip strength (continuous, kg), baseline hypertension (yes/no), baseline diabetes (yes/no), baseline cardiovascular diseases (yes/no), and family history of lung cancer (yes/no).

Figure 2 The association between walking pace and the risk of incident lung cancer stratified by rs7191721 polymorphism in *RBFOX1* gene. Adjusted for age (continuous, years), sex (male, female), race (white European, mixed, South Asian, black, others), UK Biobank assessment centre, Townsend Deprivation Index (continuous), alcohol consumption (current, former, never), smoking status (current, former, never), physical activity (low, moderate, high), healthy diet score (0, 1, 2, 3, 4, 5), body mass index (continuous, kg/m^2), Hand grip strength (continuous, kg), baseline hypertension (yes/no), baseline diabetes (yes/no), baseline cardiovascular diseases (yes/no), and family history of lung cancer (yes/no).

		N	Cases		HR (95% CI)	*P* for interaction
						0.01
GG genotype	Slow	9041	186		1.00	
	Steady	63289	659		0.79 (0.66-0.95)	
	Brisk	48347	286		0.63 (0.51-0.78)	
GA genotype	Slow	14967	348		1.00	
	Steady	107254	1073		0.74 (0.64-0.84)	
	Brisk	82989	458		0.57 (0.48-0.66)	
AA genotype	Slow	6151	177		1.00	
	Steady	44698	448		0.58 (0.48-0.70)	
	Brisk	34911	183		0.43 (0.34-0.54)	

Hazard ratio